

**PROCEEDINGS OF THE ZOOLOGICAL INSTITUTE
OF THE RUSSIAN ACADEMY OF SCIENCES**


**ANNOTATED CATALOGUE
OF THE HYMENOPTERA OF RUSSIA**

VOLUME I

SYMPHYTA AND APOCRITA: ACULEATA



SUPPLEMENT № 6, 2017



PROCEEDINGS OF THE ZOOLOGICAL INSTITUTE
OF THE RUSSIAN ACADEMY OF SCIENCES

SUPPLEMENT № 6

**ANNOTATED CATALOGUE
OF THE HYMENOPTERA OF RUSSIA**

VOLUME I

SYMPHYTA AND APOCRITA: ACULEATA



SAINT PETERSBURG
2017

PROCEEDINGS OF THE ZOOLOGICAL INSTITUTE
OF THE RUSSIAN ACADEMY OF SCIENCES

SUPPLEMENT № 6

ANNOTATED CATALOGUE
OF THE HYMENOPTERA OF RUSSIA

VOLUME I

SYMPHYTA AND APOCRITA: ACULEATA

Edited by S.A. Belokobylskij and A.S. Lelej

Editor-in-Chief – *Oleg N. Pugachev*

Editorial board:

*Alexander O. Averianov, Nikolay V. Aladin, Alexander F. Alimov, Tamara A. Asanovich, Arkadiy V. Balushkin,
Kirill V. Galaktionov, Andrey V. Gorokhov, Sergey D. Grebelnyi, Elena L. Markhaseva, Vladimir A. Paevsky,
Andrey A. Przhiboro, Sergey Yu. Sinev, Alexey V. Smirnov, Vladislav V. Khlebovich, Semen Ya. Tsalolikhin*

Managing editor:
Leonid N. Anisyutkin

Reviewers:

Professor *Alexander P. Rasnitsyn* (Moscow, Russia) and Dr *Serguei V. Triapitsyn* (Riverside, USA)

ТРУДЫ ЗООЛОГИЧЕСКОГО ИНСТИТУТА
РОССИЙСКОЙ АКАДЕМИИ НАУК

ПРИЛОЖЕНИЕ № 6

**АННОТИРОВАННЫЙ КАТАЛОГ
ПЕРЕПОНЧАТОКРЫЛЫХ НАСЕКОМЫХ РОССИИ**

ТОМ I

**СИДЯЧЕБРЮХИЕ (SYMPHYTA)
И ЖАЛОНОСНЫЕ (APOCRITA: ACULEATA)**



САНКТ-ПЕТЕРБУРГ
2017

ТРУДЫ ЗООЛОГИЧЕСКОГО ИНСТИТУТА
РОССИЙСКОЙ АКАДЕМИИ НАУК

ПРИЛОЖЕНИЕ № 6

АННОТИРОВАННЫЙ КАТАЛОГ
ПЕРЕПОНЧАТОКРЫЛЫХ НАСЕКОМЫХ РОССИИ

ТОМ I

СИДЯЧЕБРЮХИЕ (SYMPHYTA)
И ЖАЛОНОСНЫЕ (APOCRITA: ACULEATA)

Под общей редакцией С.А. Белокобыльского и А.С. Лелея

Главный редактор – академик РАН *О.Н. Пугачев*

Редколлегия:

*А.О. Аверьянов, Н.В. Аладин, А.Ф. Алимов, Т.А. Асанович, А.В. Балушкин, К.В. Галактионов,
А.В. Горохов, С.Д. Гребельный, Е.Л. Мархасева, В.А. Паевский, А.А. Пржеборо, С.Ю. Синёв,
А.В. Смирнов, В.В. Хлебович, С.Я. Цалолихин*

Ответственный редактор:
Л.Н. Анисюткин

Рецензенты:
профессор, д.б.н. *А.П. Расницын* (Москва, Россия) и д-р *С.В. Тряпицын* (Риверсайд, США)



УДК 595.79 (470+571)

ANNOTATED CATALOGUE OF THE HYMENOPTERA OF RUSSIA. VOLUME I. SYMPHYTA AND APOCRITA: ACULEATA

A.V. Antropov, Yu.V. Astafurova, S.A. Belokobylskij, A.M. Byvaltsev, Yu.N. Danilov, D.A. Dubovikoff, K.I. Fadeev, A.V. Fateryga, N.V. Kurzenko, A.S. Lelej, T.V. Levchenko, V.M. Loktionov, M.V. Mokrousov, P.G. Nemkov, M.Yu. Proshchalykin, P. Rosa, D.A. Sidorov, Yu.N. Sundukov, Z.M. Yusupov, L.A. Zaytseva

Editorial board of the first volume:

A.S. Lelej (executive editor), M.Yu. Proshchalykin and V.M. Loktionov

ABSTRACT

Hymenoptera is one of the largest insect orders, with a world fauna of 155.5 thousand species in 8423 genera from 94 extant families, a Palaearctic fauna of 50–60 thousand estimated species, and a Russian fauna of more than 15 thousand estimated species in more than 1500 genera from 78 families. The modern classification of Hymenoptera is given. The first volume treats the sawflies (13 families, 170 genera, 1546 species), wasps (15 families, 253 genera, 1695 species), ants (1 family, 44 genera, 264 species) and bees (6 families, 66 genera, 1216 species) which together, in Russia, comprise 4721 species in 533 genera. The sections for each of the 35 families include characteristics, number of taxa, references, and an annotated catalogue of the genera and species recorded from the fauna of Russia. For each valid genus the type species, synonymy, short characterization, and number of species are given. For each valid species the synonymy, known host plants, hosts or prey, and distribution (in Russia and general) are provided. The book is illustrated using geoschemes for Russia, Europe, and China. The large bibliography helps to assess the level of study of each family. The index of scientific names of Hymenoptera (more than 16000 names) will be useful for taxon searching.

Key words: sawflies, wasps, ants, bees, Xyeloidea, Tenthredinoidea, Pamphilioidea, Cephoidea, Siricoidea, Orussoidea, Chryridoidea, Scolioidea, Pompiloidea, Vespoidea, Formicoidea, Apoidea

АННОТИРОВАННЫЙ КАТАЛОГ ПЕРЕПОНЧАТОКРЫЛЫХ НАСЕКОМЫХ РОССИИ. ТОМ I. СИДЯЧЕБРЮХИЕ (SYMPHYTA) И ЖАЛОНОСНЫЕ (APOCRITA: ACULEATA)

А.В. Антропов, Ю.В. Астафурова, С.А. Белокобыльский, А.М. Бывальцев, Ю.Н. Данилов, Д.А. Дубовиков, К.И. Фадеев, А.В. Фатерыга, Н.В. Курзенко, А.С. Лелей, Т.В. Левченко, В.М. Локтионов, М.В. Мокроусов, П.Г. Немков, М.Ю. Прощалыкин, П. Роза, Д.А. Сидоров, Ю.Н. Сундуков, З.М. Юсупов, Л.А. Зайцева

Редакционная коллегия первого тома:

А.С. Лелей (ответственный редактор), М.Ю. Прощалыкин, В.М. Локтионов

РЕЗЮМЕ

Перепончатокрылые – один из самых больших отрядов насекомых, который насчитывает в мире 155.5 тыс. видов, 8423 рода из 94 recentных семейств, в Палеарктике ожидается 50–60 тыс. видов, в России – более 15 тыс. видов из более чем 1500 родов и 78 семейств. Приведена современная общая классификация отряда

Нутепортера. Первый том охватывает пилильщиков и рогохвостов (13 семейств, 170 родов, 1546 видов), ос (15 семейств, 253 рода, 1695 видов), муравьев (1 семейство, 44 рода, 264 вида) и пчел (6 семейств, 66 родов, 1216 видов); в целом в России насчитываются 4721 вид из 533 родов. Каждый раздел (35 семейств) включает краткую характеристику, число таксонов, литературу и аннотированный каталог родов и видов, известных из России. Для каждого валидного рода приведены типовой вид, синонимия, краткая характеристика и число видов. Для каждого валидного вида приведены синонимия, известные кормовые растения, хозяева или добыча и распространение (в России и общее). Книга иллюстрирована схематическими картами России, Европы и Китая. Большая библиография дает представление о степени изученности каждого семейства. Указатель латинских названий перепончатокрылых (в целом свыше 16000 названий) облегчит поиск необходимых названий.

Key words: пилильщики, осы, муравьи, пчелы, *Xyeloidea*, *Tenthredinoidea*, *Pamphilioidea*, *Cephoidea*, *Siricoidea*, *Orussoidea*, *Chryridoidea*, *Scolioidea*, *Pomipoidea*, *Vespoidea*, *Formicoidea*, *Apoidea*

CONTENTS

Preface (<i>S.A. Belokobylskij and A.S. Lelej</i>)	9
Authors	13
Order Hymenoptera	15
Introduction (<i>A.S. Lelej and S.A. Belokobylskij</i>)	15
Suborder Symphyta – Sawflies and wood wasps (<i>Yu.N. Sundukov</i>)	20
Superfamily Xyeloidea	20
1. Family Xyelidae	20
Superfamily Tenthredinoidea	22
2. Family Argidae	22
3. Family Blasticotomidae	27
4. Family Cimbicidae	28
5. Family Diprionidae	33
6. Family Heptamelidae	36
7. Family Tenthredinidae	37
Superfamily Pamphilioidea	101
8. Family Pamphiliidae	101
9. Family Megalodontesidae	107
Superfamily Cephoidea	108
10. Family Cephidae	108
Superfamily Siricoidea	112
11. Family Siricidae	112
12. Family Xiphydriidae	115
Superfamily Orussoidea	117
13. Family Orussidae	117
Suborder Apocrita	118
Superfamily Chrysidoidea	118
57. Family Dryinidae (<i>A.S. Lelej and V.M. Loktionov</i>)	118
58. Family Embolemidae (<i>S.A. Belokobylskij</i>)	122
59. Family Bethylidae (<i>A.S. Lelej and K.I. Fadeev</i>)	123
60. Family Chrysididae – Cuckoo wasps (<i>P. Rosa, A.S. Lelej, S.A. Belokobylskij, V.M. Loktionov and L.A. Zaytseva</i>)	126
Superfamily Scolioidea	145
61. Family Sapygidae (<i>A.S. Lelej and N.V. Kurzenko</i>)	145
62. Family Scoliidae (<i>M.V. Mokrousov and A.S. Lelej</i>)	147
63. Family Tiphiidae (<i>M.V. Mokrousov and A.S. Lelej</i>)	149
64. Family Sierolomorphidae (<i>A.S. Lelej and M.V. Mokrousov</i>)	151
65. Family Mutilidae – Velvet ants (<i>A.S. Lelej</i>)	152
66. Family Bradynobaenidae (<i>A.S. Lelej and M.V. Mokrousov</i>)	159

Superfamily Pompiloidea	160
67. Family Pompilidae – Spider wasps (<i>V.M. Loktionov and A.S. Lelej</i>)	160
Superfamily Vespoidea	175
68. Family Vespidae (<i>A.V. Antropov</i>)	175
Subfamily Masarinae – Pollen wasps (<i>A.V. Fateryga</i>)	177
Subfamily Eumeninae – Potter wasps (<i>A.V. Fateryga</i>)	178
Subfamily Polistinae – Paper wasps (<i>A.V. Antropov</i>)	192
Subfamily Vespinae – Hornets and Jellow Jackets (<i>A.V. Antropov</i>)	193
Superfamily Formicoidea	197
69. Family Formicidae – Ants (<i>D.A. Dubovikoff and Z.M. Yusupov</i>)	197
Superfamily Apoidea	211
Section Spheciformes – Digger wasps	211
70. Family Ampulicidae – Ampulicid digger wasps (<i>Yu.N. Danilov</i>)	211
71. Family Sphecidae – Sphecid digger wasps (<i>Yu.N. Danilov</i>)	212
72. Family Crabronidae – Crabronid digger wasps	217
Subfamily Astatinae (<i>P.G. Nemkov</i>)	218
Subfamily Bembicinae (<i>P.G. Nemkov</i>)	219
Subfamily Crabroninae (<i>A.V. Antropov</i>)	225
Subfamily Dinetinae (<i>P.G. Nemkov</i>)	245
Subfamily Eremiaspheiinae (<i>M.V. Mokrousov</i>)	245
Subfamily Mellininae (<i>P.G. Nemkov</i>)	246
Subfamily Pemphredoninae (<i>M.V. Mokrousov</i>)	246
Subfamily Philanthinae (<i>Yu.N. Danilov and M.V. Mokrousov</i>)	252
Section Apiformes – Bees	257
73. Family Colletidae (<i>M.Yu. Proshchalykin</i>)	257
74. Family Andrenidae (<i>M.Yu. Proshchalykin, Yu.V. Astafurova</i> and <i>D.A. Sidorov</i>)	263
75. Family Halictidae (<i>Yu.V. Astafurova and M.Yu. Proshchalykin</i>)	277
76. Family Melittidae (<i>M.Yu. Proshchalykin and Yu.V. Astafurova</i>)	293
77. Family Megachilidae (<i>M.Yu. Proshchalykin and A.V. Fateryga</i>)	295
78. Family Apidae (<i>T.V. Levchenko, A.M. Byvaltsev</i> and <i>M.Yu. Proshchalykin</i>)	309
References	333
Index of Latin names of Hymenoptera	389

PREFACE

S.A. BELOKOBILSKIJ AND A.S. LELEJ

Hymenoptera is one of the largest insect orders, with a world fauna 155.5 thousand species in 8423 genera from 94 extant families (Aguilar et al., 2013). Probably Hymenoptera in total includes 250–300 thousand species (Rasnitsyn, 2002; Gauld et al., 1988). According to our estimation, the Palaearctic fauna has about 50–60 thousand species; in Russia there are more than 15 thousand species in more than 1500 genera from 78 families of Hymenoptera (Lelej et al., 2012).

The Annotated Catalogue of Hymenoptera of Russia is based on six Hymenoptera books of the series “Keys to the insects of the European part of USSR” (1978a, 1978b; 1981; 1986a, 1978b; 1988, which treat 10748 recorded and potentially occurring species in 1938 genera) and five Hymenoptera books of the series “Keys to the insects of the Russian Far East” (1995a, 1995b; 1998; 2000; 2007, which include 4948 recorded and potentially occurring species in 1389 genera) as well as original papers published in the last decades. The last catalogue of Hymenoptera of the Russian Far East (Lelej et al., 2012) includes 7503 recorded species in 1363 genera from 69 families. Among the 78 families of Hymenoptera so far recorded from Russia, Stephanidae, Evaniiidae, Agaonidae, Signiphoridae, Bradynobaenidae and Ampulicidae are known to occur only in the European part of the country, while distribution of Roproniidae, Proctorenyxidae and Vanhorniidae is limited to south of the Russian Far East. The rare family Sierolomorphidae is known from south of the Russian Far East (Primorskiy Terr.), Western Siberia (Novosibirsk Prov.), and Ural (Orenburg Prov.).

Significance of such types of catalogues is in uniting comprehensive information about contents, biological peculiarity, distribution, and host specificity of the insect fauna of a country of interest. They are also extremely important for development of the Russian national insect collections.

The first volume of this Catalogue treats the sawflies (13 families, 170 genera, 1546 species), wasps (15 families, 253 genera, 1695 species), ants (1 family, 44 genera, 264 species), and bees (6 families, 66 genera, 1216 species); their total number in Russia is 4721 species in 533 genera. Sections for each of 35 families include brief data, number of taxa, references, and annotated catalogues of genera and species recorded from Russia. For each valid genus, the type species, synonymy, a brief characteristic, and number of species are

given. For each valid species, synonymy, known host plants, hosts or preys, and distribution (from Russia and general) are provided. The book is illustrated by geoschemes for Russia, Europe, and China (Figs 1–3).

The boundary of Russia and the adjacent countries is given in geoscheme (Fig. 1) using abbreviations. For better understanding of the species distribution, also provided are geoschemes of Europe (Fig. 2) and China (Fig. 3). For most families and genera, the current (as of October 2017) number of valid species in the World is given, followed by that in the Palaearctic region and Russia. For the species distribution, the Russian regions are listed first, followed by Europe and other countries (regions) in the following order: Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KA, MG, CH). – Europe (WE, NE, SE, EE), N Africa, Georgia, Armenia, Azerbaijan (or Caucasus – for all three countries), Turkey, Syria, Jordan, Lebanon, Israel, Iraq, Iran, Afghanistan, Pakistan, Turkmenistan, Tajikistan, Uzbekistan, Kyrgyzstan (or Central Asia – for all four countries), Kazakhstan, Mongolia, China (NE, NC, NW, CC, SW, WP, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu), N America, SE Asia, India, Africa, S America, Australia. The names of countries and Russian regions are given according to the All-Russian classifier of the countries of the world (<http://klassifikators.ru/oksm>), and English names of the Russian regions follows Gorskaya (1994).

Geoscheme for Europe according to United Nations geoscheme for Europe (https://en.wikipedia.org/wiki/United_Nations_geoscheme_for_Europe), including of United Kingdom and Republic of Ireland in Western Europe and excluding of Turkey and Kazakhstan (Fig. 2). **Western Europe** (WE): Austria, Belgium, France, Germany, Republic of Ireland, Liechtenstein, Luxembourg, Monaco, Netherlands, Switzerland, United Kingdom; **Northern Europe** (NE): Denmark, Estonia, Finland, Iceland, Latvia, Lithuania, Norway, Sweden; **Southern Europe** (SE): Albania, Andorra, Bosnia and Herzegovina, Croatia, Greece, Italy, Republic of Macedonia, Malta, Montenegro, Portugal, San Marino, Serbia, Slovenia, Spain; **Eastern Europe** (EE): Belarus, Bulgaria, Czech Republic, Hungary, Republic of Moldova, Poland, Romania, Slovakia, Ukraine; European part of Russia (treated separated from other countries of EE in structure of whole Russian Federation).

DZ – Algeria; **MA** – Morocco; **TN** – Tunisia; **TR** – Turkey. Codes for the names of countries follow the standard of the International Organization for Standardization (ISO 3166-1 alpha-2).




Fig. 1. Geoscheme for Russia. **EUROPEAN PART (RUEP = EP):** North (N): Murmansk Prov., Arkhangelsk Prov., Karelian Rep., Vologda Prov., Komi Rep.; North-West (NW): Kaliningrad Prov., Leningrad Prov., Pskov Prov., Novgorod Prov., Tver Prov., Yaroslavl Prov., Kostroma Prov., Smolensk Prov., Moscow Prov., Orel Prov., Lipetsk Prov., Tambov Prov., Penza Prov., Kursk Prov., Nizhniy Novgorod Prov., Kaluga Prov., Ryazan Prov., Mordovian Rep., Mari El Rep., Ulyanovsk Prov., Samara Prov., Saratov Prov.; South (S): Rostov Prov., Volgograd Prov., Astrakhan Prov., North Caucasian (NC); Krasnodar Terr., Stavropol Terr., Adygei Rep., Karachayevо-Cherkess Rep., Ingush Rep., Kabardino-Balkarian Rep., North Ossetian Rep., Chechen Rep., Dagestan Rep., Crimea (CR); Crimea (CR); Crimean Rep., Sebastopol, URAILS (UR); Perm Terr., Sverdlovsk Prov., Bashkir Rep., Chelyabinsk Prov., Orenburg Prov., Kurgan Prov., Altai (AL) (including Altai Terr. and Altai Terr.), EASTERN SIBERIA (ES): Khakass Rep. (KS), Tuva Rep. (TU), Krasnoyarsk Terr. (KR), Irkutsk Prov. (IR), Buryat Rep. (BR), Yakutsk Rep. (YA), Zabaykalskii Terr. (ZB); FAR EAST (FE): Amur Prov. (AM), Khabarovsk Terr. (including Jewish Autonomous Region) (KH), Primorskii Terr. (PR), Sakhalin (SA), Kuril Islands (KU), Kamchatka Terr. (KA), Magadan Prov. (MG), Chukot Autonomous Area (CH). **Other countries:** AM – Armenia; AZ – Azerbaijan; BY – Belarus; CN – China; EE – Estonia; FI – Finland; GE – Georgia; IR – Iran; JP – Japan; KP – North Korea (Democratic People’s Republic of Korea); KZ – Kazakhstan; LT – Lithuania; LV – Latvia; MN – Mongolia; NO – Norway; PL – Poland; RU – Russia; SE – Sweden; SY – Syria; TR – Turkey; UA – Ukraine; UZ – Uzbekistan.




Fig. 2. Geoscheme for Europe. European countries: **AD** – Andorra; **AL** – Albania; **AT** – Austria; **BA** – Bosnia and Herzegovina; **BE** – Belgium; **BG** – Bulgaria; **BY** – Belarus; **CH** – Switzerland; **CZ** – Czech Republic; **DE** – Germany; **DK** – Denmark; **EE** – Estonia; **ES** – Spain; **FI** – Finland; **FR** – France; **GB** – United Kingdom; **GR** – Greece; **HR** – Croatia; **HU** – Hungary; **IE** – Republic of Ireland; **IS** – Iceland; **IT** – Italy; **KS** – Kosovo; **LI** – Liechtenstein; **LT** – Lithuania; **LU** – Luxembourg; **LV** – Latvia; **MC** – Monaco; **MD** – Republic of Moldova; **ME** – Montenegro; **MK** – Republic of Macedonia; **MT** – Malta; **NL** – Netherlands; **NO** – Norway; **PL** – Poland; **PT** – Portugal; **RO** – Romania; **RS** – Serbia; **RUEP** – Russian Federation (European part); **SE** – Sweden; **SI** – Slovenia; **SM** – San-Marino; **SK** – Slovakia; **UA** – Ukraine. Other countries: **CY** – Cyprus; **DZ** – Algeria; **MA** – Morocco; **TN** – Tunisia; **TR** – Turkey. Codes for the names of countries follow the standard of the International Organization for Standardization (ISO 3166-1 alpha-2).




Fig. 3. Geoscheme for China. *Regions of China*: Northeastern Territory (NE), Northern Territory (NC), Northwestern Territory (NW), the Western Plateau (WP), Central Territory (CC), Southwestern Territory (SW), Southeastern Territory (SE).

Acknowledgments

We thank Academician Oleg N. Pugachev (Director of Zoological Institute, St Petersburg, Russia) for his comprehensive support of our work on the Catalogue, Alexander P. Rasnitsyn (Paleontological Institute, Moscow, Russia) and Serguei V. Triapitsyn (University of California, Riverside, USA) for critically reviewing the manuscript and making important suggestions which have improved it. We are grateful to Andreas Taeger (Senckenberg Deutsches Entomologisches Institut, Müncheberg, Germany) for the review of the Symphyta sections and James M. Carpenter

(American Museum of Natural History, New York, USA) for the help with preparing of Vespidae section.

The work was supported by the Russian Fund of Basic Research (№№ 15–29–02466, 16–54–00041, 16–04–00197, 17–04–00259), the Additional State Task of the Russian FASO “Zoological Institute collection inventory and development, 2017 [Inventarizatsiya i razvitie fondovykh kollektsiy Zoologicheskogo Instituta, 2017]” and the Program by Presidium of the Russian Academy of Sciences № 1.21P “Biodiversity of Natural Systems. Biological Resources of Russia: Assessment of Condition and Fundamental Basis of Monitoring”.

AUTHORS

Alexander V. Antropov

Zoological Museum of Moscow Lomonosov State University
Bolshaya Nikitskaya Str., 2, Moscow 125009, Russia
E-mail: antropov@zmmu.msu.ru

Yulia V. Astafurova

Zoological Institute, RAS
Universitetskaya naberezhnaya, 1,
St Petersburg 199034, Russia
E-mail: yulia.astafurova@zin.ru

Sergey A. Belokobylskij

Zoological Institute, RAS
Universitetskaya naberezhnaya, 1,
St Petersburg 199034, Russia;
Museum and Institute of Zoology, PAS
Wilcza, 64, Warsaw 00–679, Poland
E-mail: doryctes@gmail.com

Alexander M. Byvaltsev

Novosibirsk State University
Pirogova str., 2,
Novosibirsk 630090, Russia
E-mail: ByvAM@yandex.ru

Yuriy N. Danilov

Institute of Systematics and Ecology of Animals,
SB RAS
Frunze Street 11,
Novosibirsk 630091, Russia.
E-mail: prionyx@mail.ru

Dmitriy A. Dubovikoff

Saint Petersburg State University
16th line of Vasilevskiy Island, 29,
St Petersburg 199178, Russia
E-mail: dubovikoff@gmail.com

Kirill I. Fadeev

Zoological Institute, RAS
Universitetskaya naberezhnaya, 1,
St Petersburg 199034, Russia
E-mail: kfadeev66eumen@list.ru

Alexander V. Fateryga

T.I. Vyazemsky Karadag Scientific Station – Nature Reserve, RAS
Nauki Str., 24, Kurortnoe, Feodosiya 298188, Russia
E-mail: fater_84@list.ru

Nikolay V. Kurzenko

Falls Church, Virginia, 22042, USA
E-mail: kurzenko.nv@gmail.com

Arkadiy S. Lelej

Federal Scientific Center of the East Asia Terrestrial Biodiversity, FEB RAS
100-letiya Vladivostoka Ave., 159, Vladivostok 690022, Russia
E-mail: lelej@biosoil.ru

Timofey V. Levchenko

Department of Collections,
Moscow State Darwin Museum
Vavilova st., 57/1, Moscow 117292, Russia
E-mail: antimofa1@yandex.ru

Valeriy M. Loktionov

Federal Scientific Center of the East Asia Terrestrial Biodiversity, FEB RAS
100-letiya Vladivostoka Ave., 159, Vladivostok 690022, Russia
E-mail: pompilidaefer@mail.ru

Mikhail V. Mokrousov

Lobachevsky State University of Nizhni Novgorod
Gagarina Str., 23, Nizhni Novgorod 603950, Russia
E-mail: sphecid@inbox.ru

Pavel G. Nemkov

Federal Scientific Center of the East Asia Terrestrial Biodiversity, FEB RAS
100-letiya Vladivostoka Ave., 159, Vladivostok 690022, Russia
E-mail: p_nemkov@mail.ru

Maxim Yu. Proshchalykin

Federal Scientific Center of the East Asia Terrestrial Biodiversity, FEB RAS
100-letiya Vladivostoka Ave., 159, Vladivostok 690022, Russia
E-mail: proshchalikin@biosoil.ru

Paolo Rosa

Via Belvedere, 8/d, I-20881, Bernareggio (MB), Italy
E-mail: rosa@chrysis.net

Dmitriy A. Sidorov

Kemerovo State University
Krasnaya str. 6, Kemerovo 650000, Russia
E-mail: *raddimus@yandex.ru*

Yuriy N. Sundukov

State Nature Reserve Kurilskiy
Zarechnaya Str., 5, Yuzhno-Kurilsk,
Kunashir Island 694500, Russia
E-mail: *yun-sundukov@mail.ru*

Zalim M. Yusupov

Institute of Ecology of Mountain Territories, Kabardino-Balkarian Scientific Centre RAS
I. Armand Str., 37a, Nalchik, Kabardino-Balkar Republic 360051, Russia
E-mail: *yzalim@mail.ru*

Liuba A. Zaytseva

Via Belvedere, 8/d, I-20881, Bernareggio (MB), Italy
E-mail: *rosa@chrysis.net*

ORDER HYMENOPTERA

INTRODUCTION

A.S. LELEJ AND S.A. BELOKOBILSKIJ

The first volume of the Annotated Catalogue of the Hymenoptera of Russia (Table) treats the sawflies (13 families, 170 genera, 1546 species), wasps (15 families, 253 genera, 1695 species), ants (1 family, 44 genera, 264 species), and bees (6 families, 66 genera, 1216 species); their total number in Russia is 4721 species in 533 genera. The large bibliography helps to document the level of knowledge of each family. Index of Latin names of Hymenoptera (more than 16000 names total) can be useful for searching particular taxa.

Classification

In this book we followed the classification of Hymenoptera as proposed by Sharkey (2007) and Sharkey et al. (2011) for higher-level groupings and the survey of taxonomic richness by Aguiar et al. (2013), complemented by the recent new descriptions (Yu, 2012; Li et al., 2013; Engel et al., 2013, 2016a, b; Engel, 2015; Rodriguez et al., 2016; Rasnitsyn et al., 2017; Wang et al., 2017; Zhang et al., 2017). Also used were classifications by Rasnitsyn (1988) and Rasnitsyn & Zhang (2010) regarding the infraorders but with some changes. Infraorder Orussomorpha is a transition between the suborders Symphyta and Apocrita, but is traditionally placed in Symphyta.

Recently, Pilgrim et al. (2008) undertook a molecular analysis based on four nuclear genes of 64 taxa across the entire Vespoidea, in which they concluded that this superfamily and some included families were paraphyletic, and proposed recognition of six “vespid” superfamilies (Formicoidea, Pompiloidea, Scolioidea, Tiphioidae, Thynnoidea and Vespoidea) and transfer of Myrmosinae from Mutillidae to its own family (Myrmosidae), both within Pompiloidea (with Pompilidae and Sapygidae), but they lacked specimens of most of the mutillid subfamilies and had only one of Myrmosinae, so the basis for their results was limited.

The families recorded from Russia are numbered (under these numbers they are placed in the taxonomic part of this catalogue; these are marked by a point (•) if they are not registered in Russia, and by a special symbol (†) if they are fossil).

Suborder SYMPHYTA

Xyelomorpha

Xyeloidea Newman, 1834

1. Xyelidae Newman, 1834

† Syspastoxylidae Engel et Huang, 2016

Tenthredinomorpha

Tenthredinoidea Latreille, 1803

† Electrotomidae Rasnitsyn, 1977

† Xyelotomidae Rasnitsyn, 1968

2. Argidae Konow, 1890

• Pergidae Rohwer, 1911

3. Blasticotomidae Thomson, 1871

4. Cimbicidae W. Kirby, 1837

5. Diprionidae Rohwer, 1910

6. Heptamelidae Benson, 1938

7. Tenthredinidae Latreille, 1803

Siricomorpha

Pamphilioidea Cameron, 1890

† Xyelydidae Rasnitsyn, 1986

† Mirolydidae Wang, Rasnitsyn et Ren, 2017

8. Pamphiliidae Cameron, 1890

9. Megalodontesidae Konow, 1897

Cephidoidea Newman, 1834

† Sepulcidae Rasnitsyn, 1968

10. Cephidae Newman, 1834

Siricoidea Billberg, 1820

† Daohugoidae Rasnitsyn et Zhang, 2004

† Protosiricidae Rasnitsyn et Zhang, 2004

† Pseudosiricidae Handlirsch, 1908

11. Siricidae Billberg, 1820

12. Xiphydriidae Leach, 1819

Anaxyloidea Martynov, 1925

• Anaxyelidae Martynov, 1925

Orussomorpha

Orusoidea Newman, 1834

† Karatavitidae Rasnitsyn, 1963

† Paroryssidae Martynov, 1925

13. Orussidae Newman, 1834

Suborder APOCRITA

Parasitica¹

Stephanomorpha

Stephanoidea Leach, 1815

† Eostephanitidae Hong, 2002²

† Ephialtitidae Handlirsch, 1906

† Myanmarinidae Zhang et Rasnitsyn, 2017

¹ Group of non-fixed rank

² *Eostephanites* Hong, 2002 [type genus of Eostephanitidae] has been placed to Stephanoidea: Eostephanitidae (Yu, 2012) but A.P. Rasnitsyn considers (pers. comm.) that it belongs to the Braconidae

Table. Hymenoptera of Russia: number of taxa of Symphyta and Apocrita: Aculeata

Family	World		Palaearctic		Russia	
	Genera	Species	Genera	Species	Genera	Species
Symphyta						
1. Xyelidae	8	85	4	40	3	13
2. Argidae	60	920	17	310	7	88
3. Blasticomidae	2	12	2	8	2	3
4. Cimbicidae	15	190	10	150	8	72
5. Diprionidae	11	140	9	70	8	28
6. Heptamelidae	6	50	2	13	2	5
7. Tenthredinidae	400	5600	220	3500	109	1124
8. Pamphiliidae	10	300	10	200	9	99
9. Megalodontesidae	1	37	1	37	1	12
10. Cephidae	24	165	16	120	10	57
11. Siricidae	11	130	7	75	5	20
12. Xiphydriidae	29	150	14	50	4	21
13. Orussidae	17	90	5	20	2	4
Total	594	7879	317	4603	170	1546
Apocrita: Aculeata						
57. Dryinidae	50	1827	24	257	8	63
58. Embolenidae	3	39	3	10	2	5
59. Bethylidae	84	2340	20	244	20	34
60. Chrysididae	95	2815	36	1557	23	330
61. Sapygidae	12	66	6	31	5	9
62. Scoliidae	43	560	11	100	5	20
63. Tiphidae	120	2000	17	100	6	28
64. Sierolomorphidae	2	12	2	5	1	2
65. Mutillidae	216	4300	61	525	20	74
66. Bradynobaenidae	10	188	4	43	1	2
67. Pompilidae	254	4855	65	1000	37	246
68. Vespidae	256	5274	90	1181	38	197
69. Formicidae	485	13978	94	1350	44	264
70. Ampulicidae	6	202	3	38	2	4
71. Sphecidae	19	779	13	260	11	68
72. Crabronidae	250	8900	84	2000	74	613
73. Colletidae	86	2630	2	432	2	100
74. Andrenidae	77	2917	13	1100	5	244
75. Halictidae	79	4420	23	950	13	263
76. Melittidae	15	198	5	85	3	25
77. Megachilidae	76	4096	36	1100	18	198
78. Apidae	176	5614	33	1358	25	386
Total	2414	68010	645	13726	363	3175

Note. Data for the World are given from Aguiar et al. (2013), and for the Palaearctic region and Russia from this book.

14. Stephanidae Leach, 1815
- E v a n i o m o r p h a**
- Evanoidea** Latreille, 1802
- † Andreneliidae Rasnitsyn et Martínez-Declòs, 2000
 - † Anomopterellidae Rasnitsyn, 1975
 - † Othniodellithidae Engel et Huang, 2016
 - † Praeaulacidae Rasnitsyn, 1972
 - 15. Aulacidae Shuckard, 1841
 - 16. Gasteruptiidae Ashmead, 1900
 - 17. Esviidae Latreille, 1802
- C e r a p h r o n o m o r p h a**
- Ceraphronoidea** Haliday, 1833
- † Radiophronidae Ortega-Blanco, Rasnitsyn et Declòs, 2010
 - † Stigmaphronidae Kozlov, 1975
 - † Aptenoperissidae Rasnitsyn & Poinar, 2017
 - 18. Megaspilidae Ashmead, 1893
 - 19. Ceraphronidae Haliday, 1833
- Megalyroidea** Schletterer, 1889
- † Maimetshidae Rasnitsyn, 1975
 - Megalyridae Schletterer, 1889
- Trigonalyoidea** Cresson, 1887
- 20. Trigonalyidae Cresson, 1887
- P r o c t o t r u p o m o r p h a**
- Proctotrupoidea** Latreille, 1802
- † Jurapriidae Rasnitsyn, 1983
 - † Mesoserphidae Kozlov, 1970
 - Austroniidae Kozlov, 1975
 - Pelecinidae Haliday, 1839
 - Peradeniidae Naumann et Masner, 1985
 - 21. Heloridae Förster, 1856
 - 22. Proctotrupidae Latreille, 1802
 - 23. Roproniidae Bradley, 1905
 - 24. Proctorenyxidae Lelej et Kozlov, 1999
 - 25. Vanhorniidae Crawford, 1909
- Diapiroidea** Haliday, 1833
- 26. Diapriidae Haliday, 1833
 - 27. Ismaridae Thomson, 1858
 - Maamingidae Early, Masner, Naumann et Austin, 2001
 - Monomachidae Ashmead, 1902
 - † Spathopterygidae Engel et Ortega-Blanco, 2013
- Platygastroidea** Haliday, 1833
- 28. Platygastridae Haliday, 1833
 - 29. Scelionidae Haliday, 1839
- † **Serphitoidea** Brues, 1937
- † Serphitidae Brues, 1937
 - † Archaeoserphitidae Engel, 2015

- Cynipoidea** Latreille, 1802
- † Gerocynipidae Liu et Engel, 2007
 - † Protimaspidae Liu et Engel, 2007
 - † Stolamissidae Liu et Engel, 2007
 - Austrocynipidae Riek, 1971
 - 30. Ibaliiidae Thomson, 1862
 - 31. Liopteridae Ashmead, 1895
 - 32. Figitidae Hartig, 1840
 - 33. Cynipidae Latreille, 1802
- Chalcidoidea** Latreille, 1817
- † Khutelchalcididae Rasnitsyn, Basibuyuk et Quicke, 2004
 - 34. Chalcididae Latreille, 1817
 - 35. Leucospidae Walker, 1834
 - 36. Perilampidae Förster, 1856
 - 37. Eucharitidae Walker, 1846
 - 38. Pteromalidae Dalman, 1820
 - 39. Eupelmidae Walker, 1833
 - 40. Encyrtidae Walker, 1837
 - 41. Eurytomidae Walker, 1832
 - 42. Torymidae Walker, 1833
 - 43. Ormyridae Förster, 1856
 - 44. Agaonidae Walker, 1846
 - 45. Tetracampidae Förster, 1856
 - 46. Eulophidae Westwood, 1829
 - 47. Aphelinidae Thomson, 1876
 - 48. Azotidae Nikol'skaya et Yasnosh, 1966
 - 49. Cynipencyrtidae Trjapitzin, 1973
 - Eriaporidae Ghesquère, 1955
 - 50. Trichogrammatidae Haliday, 1851
 - 51. Signiphoridae Howard, 1894
 - 52. Mymaridae Haliday, 1833
 - Rotoitidae Bouček et Noyes, 1987
 - Tanaostigmatidae Ashmead, 1904
- Mymaromatoidea** Debauche, 1948
- † Alavarommatidae Ortega-Blanco, Peñalver, Delclòs et Engel, 2011
 - † Gallorommatidae Gibson, Read et Huber, 2007
 - 53. Mymarommatidae Debauche, 1948
- I c h n e u m o n o m o r p h a**
- Ichneumonoidea** Latreille, 1802
- † Praeichneumonidae Rasnitsyn, 1983
 - 54. Ichneumonidae Latreille, 1802
 - 55. Braconidae Nees, 1811
 - 56. Aphidiidae Haliday, 1833
- Aculeata**³
- V e s p o m o r p h a**
- † **Bethylonymoidea** Rasnitsyn, 1975
- † Bethylonymidae Rasnitsyn, 1975

³ Group of non-fixed rank

Chrysidoidea Latreille, 1802

- † Falsiformicidae Rasnitsyn, 1975
- † Plumalexiidae Brothers, 2011
- Plumariidae Bischoff, 1914
- Sclerogibbidae Ashmead, 1902
- Scolebythidae Evans, 1963
- 57. Dryinidae Haliday, 1833
- 58. Embolemididae Förster, 1856
- 59. Bethylidae Haliday, 1839
- 60. Chrysidiidae Latreille, 1802

Scolioidea Latreille, 1802

- † Bryopompilidae Engel et Grimaldi, 2006
- 61. Sapygidae Latreille, 1810
- 62. Scoliidae Latreille, 1802
- 63. Tiphidae Leach, 1815
- 64. Sierolomorphidae Krombein, 1951
- 65. Mutillidae Latreille, 1802
- 66. Bradyobaenidae de Saussure, 1892
- Rhopalosomatidae Ashmead, 1896

Pompiloidea Latreille, 1804

- 67. Pompilidae Latreille, 1804

Vespoidea Latreille, 1802

- 68. Vespidae Latreille, 1802

Formicoidea Latreille, 1802

- 69. Formicidae Latreille, 1802

Apoidea Latreille, 1802Section Spheciformes⁴

- † Angarosphecidae Rasnitsyn, 1975
- Heterogynaidae Nagy, 1969
- 70. Ampulicidae Shuckard, 1840
- 71. Sphecidae Latreille, 1802
- 72. Crabronidae Latreille, 1802

Section Apiformes⁵

- † Mellitosphecidae Poinar et Danforth, 2006
- † Paleomelittidae Engel, 2001
- 73. Colletidae Lepeletier de Saint Fargeau, 1841
- 74. Andrenidae Latreille, 1802
- 75. Halictidae Thomson, 1869
- 76. Melittidae Schenck, 1860
- 77. Megachilidae Latreille, 1802
- 78. Apidae Latreille, 1802
- Stenotritidae Cockerell, 1934

Apocrita incertae sedis

- † Archaeocynipidae Rasnitsyn et Kovalev, 1988
- † Kuafuidae Rasnitsyn et Zhang, 2010

References

- Aguiar A.P., Deans A.R., Engel M.S., Forshage M., Huber J.T., Jennings J.T., Johnson N.F., Lelej A.S., Longino J.T., Lohrmann V., Mikó I., Ohl M., Rasmussen C., Taeger A. and Yu D.S.K. 2013. Order Hymenoptera Linnaeus, 1758. In: Zhang Z.-Q. (Ed.). *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. Zootaxa, **3703**(1): 51–62.
- Engel M.S. 2015. A new family of primitive serphitoid wasps in Labenese amber (Hymenoptera Serphitoidea). *Novitates Paleontologicae*, **13**: 1–22.
- Engel M.S., Grimaldi D.A. and Ortega-Blanco J. 2013. *Zoropelecinus zigrasi*, a pelecinid wasp in mid-Cretaceous amber from Myanmar (Hymenoptera: Pelecinidae). *Novitates Paleontologicae*, **4**: 1–10.
- Engel M.S., Huang D., Alqarni A.S. and Cai C.-y. 2016a. A remarkable evanioid wasp in mid-Cretaceous amber from northern Myanmar (Hymenoptera: Evanioidea). *Cretaceous Research*, **60**: 121–127.
- Engel M.S., Huang D., Alqarni A.S. and Cai C.-y. 2016b. An unusual new lineage of sawflies (Hymenoptera) in Upper Cretaceous amber from northern Myanmar. *Cretaceous Research*, **60**: 281–286.
- Gauld I.D. and Bolton B. (Eds.) 1988. *The Hymenoptera*. New York, 1988. XI + 332 p.
- Gorskaya M.V. 1994. *English-Russian and Russian-English geographical dictionary*. 2nd Edition. Russky Yazyk Publishers, Moscow, 272 p. [In Russian].
- Hong Y. 2002. *Amber insects of China*. Beijing: Beijing Science and Technology Press, 653 p. [In Chinese].
- Keys to the insects of the USSR European part. (Ed. Medvedev G.S.) 1978–1988. Leningrad: Nauka. **1978a**, 3(1), 584 p.; **1978b**, 3(2), 758 p.; **1981**, 3(3), 668 p.; **1986a**, 3(4), 502 p.; **1986b**, 3(5), 310 p.; **1988**, 3(6), 268 p. [In Russian].
- Keys to the insects of the Russian Far East (Ed. Lehr P.A.) 1995–2007. **1995a**, 4(1), Leningrad: Nauka. 606 p.; **1995b**, 4(2), Vladivostok: Dalnauka. 600 p.; **1998**, 4(3), 708 p.; **2000**, 4(4), 651 p.; **2007**, 4(5), 1052 p. [In Russian].
- Lelej A.S., Kupianskaya A.N., Proshchalykin M.Yu. and Loktionov V.M. (Eds.) 2012. *Annotated catalogue of the insects of Russian Far East. Vol. I. Hymenoptera*. Dalnauka, Vladivostok, 635 p. [In Russian].
- Li L., Rasnitsyn A.P., Shih C. and Ren D. 2013. Anomopterellidae restored, with two new genera and its phylogeny in Evanioidea (Hymenoptera). *PLoS ONE*, **8**(12): e82587.
- Pilgrim E.M., von Dohlen C.D. and Pitts J.P. 2008. Molecular phylogenetics of Vespoidea indicate paraphyly of the superfamily and novel relationships of its component families and subfamilies. *Zoologica Scripta*, **37**: 539–560.
- Rasnitsyn A.P. 1988. An outline of evolution of the hymenopterous insects (order Vespoidea). *Oriental Insects*, **22**: 115–145.

^{4, 5} Group of non-fixed rank

- Rasnitsyn A.P.** 2002. Order Hymenoptera Linné, 1758. In: Rasnitsyn A.P., Quicke D.L.J. (Eds.). *History of Insects*. Dordrecht, Boston, London: 242–254.
- Rasnitsyn A.P., Poinar Jr.G. and Brown A.E.** 2017. Bizarre wingless parasitic wasp from mid-Cretaceous Burmese amber (Hymenoptera, Ceraphronoidea, Aptenoperissidae fam. nov.). *Cretaceous Research*, **69**: 113–118.
- Rasnitsyn A.P. and Zhang H.-C.** 2010. Early evolution of Apocrita (Insecta, Hymenoptera) as indicated by new findings in the Middle Jurassic of Daohugou, Northeast China. *Acta Geologica Sinica*, **84**: 834–873.
- Rodriguez J., Waichert C., von Dohlen C.D., Poinar G. Jr. and Pitts J.P.** 2016. Eocene and not Cretaceous origin of spider wasps: Fossil evidence from amber. *Acta Palaeontologica Polonica*, **61** (1): 89–96.
- Sharkey M.J.** 2007. Phylogeny and Classification of Hymenoptera. *Zootaxa*, **1668**: 521–548.
- Sharkey M.J., Carpenter J.M., Vilhelmsen L., Heraty J., Liljeblad J., Dowling A.P.G., Schulmeister S., Murray D., Deans A.R., Ronquist F., Krogmann L. and Wheeler W.C.** 2011. Phylogenetic relationships among superfamilies of Hymenoptera. *Cladistics*, **27**: 1–33.
- Wang M., Rasnitsyn A.P., Yang Zh., Shih Ch., Wang H. and Ren D.** 2017. Mirolydidae, a new family of Jurassic pamphilioid sawfly (Hymenoptera) highlighting mosaic evolution of lower Hymenoptera. *Scientific Reports*, **7:43944**: P. 1–9.
- Yu D.S.K.** 2012. Home of Icnheumonoidea. Available at <http://www.taxapad.com/local.php?taxonidLC=90240608>
- Zhang Q., Rasnitsyn A.P., Wang B. and Zhang H.** 2017. Myanmarinidae, a new family of basal Apocrita (Hymenoptera: Stephanoidea) from mid-Cretaceous Burmese amber. *Cretaceous Research*, **81**: 86–92.

SUBORDER SYMPHYTA – SAWFLIES AND WOOD WASPS

Yu.N. SUNDUKOV

Sawflies are distinguished from other Hymenoptera by the anterior margin of the tergum 1, which appears as part of abdomen rather than thorax, though it is fused to the postnotum of the metathorax. The suborder is distributed worldwide. Number of recent taxa: World – 15 families, 45 subfamilies, about 650 genera, about 8250 species; Palaearctic – 13/24/>310/about 4400; Russia – 13/22/170/1546.

The classification used here reflects classification, derived from the “World catalog of Symphyta” (Taeger et al., 2010), taking into account the latest taxonomic changes (see the list of references).

INFRAORDER XYELOMORPHA

SUPERFAMILY XYELOIDEA

One family: Xyelidae.

1. FAMILY XYELIDAE

Xyelidae is the earliest known family of Hymenoptera according to the fossil record, with the oldest members dating from the Early Triassic some 220 million years ago (Blank, 2002b). Xyelidae is a small family of herbivorous wasps. These are small (Xyelinae) or medium sized (Macroxyelinae) insects (3–17 mm). Characterized by very long 3th antennal segment (actually formed by fusion of at least 8 segments) surmounted by a slender filament of 9 or more tiny segments. The larvae of subfamily Xyelinae live in the male cones of pines (*Pinus* spp.), where they feed on the sporophylls, or bores in the shoots of conifers (*Abies*, *Picea*). Larvae of Macroxyelinae – in the inflorescences of deciduous trees.

The family is distributed in the Holarctic, Oriental and Neotropical regions, mainly in their temperate zone. Number of recent taxa: World – 2 subfamilies, 8 genera, about 85 species; Palaearctic – 2/4/> 40; Russia – 2/3/13.

R e f e r e n c e s . Gussakovskij, 1935; Rasnitsyn, 1965; Shinohara, 1992b, 1995; Blank et al., 2005, 2013, 2017; Taeger et al., 2006, 2010; Sundukov, Lelej, 2012; Vasilenko, Korshunov, 2012; Kostyunin, 2015; Sundukov, 2015.

Subfamily MACROXYELINAE

MEGAXYELA Ashmead, 1898 (*Odontophyes* Konow, 1899; *Megaloxyla* Schulz, 1906; *Paraxyela* MacGillivray, 1912). The larvae feed on the Juglandaceae. Type species *Xyela major* Cresson, 1880. The genus is distributed in East Palaearctic and Nearctic region. Number of species: World – 11; Palaearctic – 6; Russia – 2.

Megaxyela gigantea Mocsàry, 1909 (*M. mikado* Sato, 1930). Host plant: *Juglans mandshurica*, *J. ailanthifolia*, *J. regia* (Juglandaceae). Russia: **FE** (KH, PR). – Korean Peninsula.

Megaxyela pulchra Blank, Shinohara et Sundukov, 2017. Host plant: *Juglans mandshurica*, ?*J. ailanthifolia*, ?*J. regia*, ?*Pterocarya* spp. (Juglandaceae). Russia: **FE** (PR). – China (NE, WP, SE), Korean Peninsula.

Subfamily XYELINAE

PLERONEURA Konow, 1897 (*Manoxyela* Ashmead, 1898). Type species *Xyela dahli* Hartig, 1837. The larvae of species bores in the shoot of *Abies* and *Picea* (Pinaceae). The genus is distributed in the Holarctic and Neotropical regions. Number of species: World – 12; Palaearctic – 7; Russia – 3.

Pleroneura dahlii (Hartig, 1837) [*Xyela*]. Host plant: *Abies* spp. (Pinaceae). Russia: **EP** (NC). – Europe (WE, EE).

Pleroneura piceae Shinohara et Hara, 1995. Host plant: *Picea glehnii* (Pinaceae). Russia: **FE** (KU). – Japan (Hok).

Pleroneura subulata Shinohara, 1995. Russia: **FE** (KH, PR). – Korean Peninsula, Japan (Hon, Shi).

XYELA Dalman, 1819 (*Pinicola* Brébisson, 1818; *Mastigocerus* Latreille, 1818; *Tritikreion* Schilling, 1826; *Mastigocera* Berthold, 1827; *Pinicolites* Meunier, 1920; *Neoxyela* Curran, 1923; *Xyelatana* Benson, 1938; *Mesoxyela* Rasnitsyn, 1965; *Magnixyela* Rasnitsyn, 1971; *Alpigenixyela* Rasnitsyn, 1971; *Desertixyela* Rasnitsyn, 1971; *Concavixyela* Rasnitsyn, 1971; *Linsleyixyela* Rasnitsyn, 1971; *Minorixyela* Rasnitsyn, 1971). Type species *Xyela pusilla* Dalman, 1819. The larvae of most species live in the male cones of pines (Pinaceae) where they feed on the sporophylls. The genus is distributed in the Holarctic, Oriental and

Neotropical regions. Number of species: World – near 50; Palaearctic – 29; Russia – 8.

Xyela julii (Brébisson, 1818) [Pinicola] (*X. pusilla* Dalman, 1819; *X. henschii* Mocsáry, 1912). Host plant: *Pinus nigra*, *P. rotundata*, *P. sylvestris* (Pinaceae). Russia: **EP** (N, NW, C), **WS** (KM, AL), **ES** (IR). – Europe (WE, NE, SE, EE), Georgia, Turkey, Mongolia.

Xyela kamtshatica Gussakovskij, 1935. Host plant: *Pinus pumila* (Pinaceae). Russia: **FE** (KA). – Japan (Hok, Hon).

Xyela koraiensis Blank et Shinohara, 2013. Host plant: *Pinus koraiensis* (Pinaceae). Russia: **FE** (PR). – Korean Peninsula.

Xyela longula Dalman, 1819 (*X. piliserra* C.G. Thomson, 1871). Host plant: *Pinus sylvestris* (Pinaceae). Russia: **EP** (NW, C). – Europe (WE, NE, EE).

Xyela pumilae Blank et Shinohara, 2013. Host plant: *Pinus pumila* (Pinaceae). Russia: **FE** (KU). – Japan (Hok).

Xyela rasnitsyni Blank et Shinohara, 2013. Host plant: *Pinus koraiensis* (Pinaceae). Russia: **FE** (PR). – China (NE), Korean Peninsula.

Xyela sibirica Blank, 2013. Host plant: *Pinus sibirica* (Pinaceae). Russia: **WS** (KM), **ES** (KR, IR). – Mongolia.

Xyela ussuriensis Rasnitsyn, 1965 (*X. suwonae* Ryu et Lee, 1992). Host plant: *Pinus koraiensis* (Pinaceae). Russia: **FE** (PR). – Korean Peninsula.

INFRAORDER TENTHREDINOMORPHA

SUPERFAMILY TENTHREDINOIDEA

Head usually open without hypostomal bridge. Pronotum very short medially and strongly emarginate behind; mesopleura with epimeron divided into upper convex and lower concave portion; mesosternum trapezoidal without presternal bridge and sutures often obsolete. Male genitalia strophandrious; female saw-sheath usually abbreviated, but sometimes exserted. The larvae are polypodous or oligopodous.

The superfamily is distributed worldwide. Number of taxa: World – 7 families, 33 subfamilies, > 550 genera, > 7300 species; Palaearctic – 6/15/> 250/about 4000; Russia – 6/13/134/1318 (+ 34 spp.).

2. FAMILY ARGIDAE

Slow, heavily built insects. Distinguished from all other families by the flagellum of its antenna being fused into a single long segment; this segment is sub-clavate in the female, but in the male is either uniform in thickness with a brush of short setae below (Arginae), or bifid like a tuning-fork (Sterictophorinae). Fore wing with vein $2r$ absent. Cenchri very large. Tibiae with or without preapical spines; front tibia with two unmodified apical spurs. The larvae are free feeding, polypodous, with 6–8 pairs of abdominal legs; they feed predominantly on the leaves of various trees and shrubs.

The family is distributed worldwide. Number of taxa: World – 6 subfamilies, 60 genera, about 920 species; Palaearctic – 3/17/about 310; Russia – 2/7/88 (+ 2 spp.).

References. Zaddach, 1859; André, 1881; Dalla Torre, 1894; Jakovlev, 1900; Konow, 1902b; Dovnar-Zapskij, 1926, 1930; Kuznetsov-Ugamskij, 1927; Takeuchi, 1927, 1932; Gussakovskij, 1935; Zhelochovtsev, 1939, 1976, 1988; Stroganova, 1966, 1970, 1978a; Muche, 1972, 1977; Verzhutskii, 1973, 1974, 1981; Ermolenko, 1975, 1976, 1984; Zhelochovtsev, Prochorova, 1976; Koch, 1982, 1988a; Liston, 1992, 1995; Zhelochovtsev, Zinovjev, 1992, 1995; Pesarini et al., 1995; Blank, 1996; Blank, Taeger, 1998a,b; Wen et al., 1998; Wei, Wen, 2002; Vasilenko, 2004, 2009a, 2010b,c; Vikberg, 2004; Taeger, 2005; Vasilenko, Dolgov, 2005; Hara, Shinohara, 2006, 2008, 2012, 2014; Kajmuk, Popov, 2006; Liston et al., 2006, 2014a; Taeger et al., 2006, 2010; Wei et al., 2006, 2017; Hara et al., 2007, 2012; Popov, Kajmuk, 2007, 2009a; Shinohara, Hara, 2007; Sundukov, 2007,

2009, 2011b, 2013, 2015, 2017; Koch, Goergen, 2008; Roller, Haris, 2008; Humala, Polevoi, 2009; Schedl, 2009a; Shinohara et al., 2009, 2011; Sundukov, Lelej, 2009, 2012; Blank et al., 2011, 2014; Taeger, Blank, 2011; Artokhin et al., 2012; Macek, 2012; Vasilenko, Korshunov, 2012; Beneš, 2014b; Beneš, Holuša, 2015; Choi et al., 2015, 2016; Kostyunin, 2015; Noblecourt, 2016.

Subfamily ARGINAE

ARGE Schrank, 1802 (*Cryptus* Jurine, 1801; *Hylotoma* Latreille, 1803; *Corynia* Labram et Imhoff, 1836; *Acanthoptenos* Ashmead, 1898; *Bathyblepta* Konow, 1906; *Didocha* Konow, 1907; *Miocephala* Konow, 1907; *Calarge* Enslin, 1911; *Alloscenia* Enderlein, 1919; *Rhopalospiria* Enderlein, 1919; *Fivaldzskyiformis* Ermolenko, 1975; *Ochropusiformis* Ermolenko, 1975; *Paganiformis* Ermolenko, 1975; *Melanochroiformis* Ermolenko, 1975; *Pleuriticiformis* Ermolenko, 1975; *Ustilatiformis* Ermolenko, 1975; *Rusticiformis* Ermolenko, 1975; *Berberidiformis* Ermolenko, 1975). Type species *Tenthredo enodis* Linnaeus, 1767. The larvae feed on the Rosaceae, Salicaceae, Betulaceae. The genus is distributed worldwide. Number of species: World – > 350; Palaearctic – about 170; Russia – 55.

Arge altaica Gussakovskij, 1935. Russia: **WS** (AL). – Korean Peninsula.

Arge aurata (Zaddach, 1864) [Hylotoma]. Russia: **EP** (NC). – Caucasus, Turkey, Syria.

Arge auripennis Konow, 1891 (*Hylotoma graeffei* Kriechbaumer, 1892). Russia: **EP** (NC). – Europe (EE), Caucasus, Turkey, Syria.

Arge azhendarovensis Vasilenko, 2010. Russia: **WS** (KM).

Arge beckeri Tournier, 1889 (*Hylotoma rufescens* Zaddach, 1864; *H. pallipes* Kriechbaumer, 1892; *H. pallidipes* Dalla Torre, 1894). Russia: **EP** (S, NC, CR). – Europe (EE, SE), Turkey.

Arge berberidis Schrank, 1802 (*Tenthredo xanthopyga* Vallot, 1836). Host plant: Berberis spp. (Berberidaceae). Russia: **EP** (C, S, NC, CR). – Europe (WE, EE, SE), Caucasus, Turkey, Japan (Hok).

Arge captiva (F. Smith, 1874) [Hylotoma] (*Hylotoma flavidollis* Cameron, 1876; *Arge sanguinolenta* Mocsáry, 1909; *A. kolthoffi* Forsius, 1927; *A. captiva rufoscutellata* Takeuchi, 1927; *A. captiva watanabei* Takeuchi, 1932). Host plant: Ulmus spp. (Ulmaceae); flies into the light. Russia: **WS** (NS, KM, AL), **ES** (ZB), **FE** (KH, PR, KU). – Kazakhstan (N),

- Mongolia, China (NE, CC, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Tsushima), India (N), SE Asia.
- Arge carinifrons** Enslin, 1917. Russia: **EP** (NC). – Georgia, Armenia.
- Arge caucasica** Tournier, 1889 (*A. halinae* Kuznetzov-Ugamskij, 1927). Russia: **EP** (E, S, NC), **UR**, **WS** (OM, NS, AL). – Europe (EE, SE), Armenia, Kyrgyzstan, Kazakhstan.
- Arge ciliaris** (Linnaeus, 1767) [Tenthredo] (*Hylotoma coerulea* Klug, 1814; *H. corusca* Zaddach, 1859; *Arge fuscipes bleusei* Pic, 1917). Host plant: *Filipendula* spp., *Spiraea* spp. (Rosaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TK, NS, KM, AL), **ES** (KR, IR, BR, YA), **FE** (AM, KH, PR, KA). – Europe (WE, EE, NE, SE), Turkey, Mongolia, China (NE, NC).
- Arge cinnabarina** Gussakovskij, 1935. Russia: **FE** (PR). – China (SW).
- Arge corallina** Gussakovskij, 1935. Russia: **FE** (AM, PR). – China (SW).
- Arge cyanocrocea** (Förster, 1771) [Tenthredo] (*Tenthredo coerulescens* Fabricius, 1775; *T. bicolor* Schrank, 1776; *Hylotoma syriaca* Mocsáry, 1880; *H. cyanocrocea messanensis* De-Stefani, 1885; *H. syriaca damascena* Magretti, 1890). Host plant: *Rubus* spp., *Sanguisorba* spp. (Rosaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (NS, KM). – Europe (WE, EE, NE, SE), Caucasus, Asia Minor, Iran, Turkmenistan.
- Arge dimidiata** (Fallén, 1808) [Hylotoma] (*Hylotoma rufescens* Drapiez, 1819; *H. fasciata* Lepeletier, 1823; *H. fasciata* Serville, 1823; *H. mediata* Fallén, 1829; *H. aenescens* Förster, 1854; *H. confusa* Dietrich, 1868; *Arge kongauensis* Rohwer, 1925; *A. dimidiata aeneoviridescens* Bischoff, 1925; *A. pravei* Dovnar-Zapolskij, 1926). Host plant: *Alnus* spp., *Betula* spp. (Betulaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TK, NS, KM, AL), **ES** (KS, KR, IR, BR, YA), **FE** (AM, PR, SA). – Europe (WE, EE, NE, SE), Georgia, Kazakhstan, Mongolia, China (NE, CC, SE), Korean Peninsula.
- Arge disparilis** (W.F. Kirby, 1882) [Hylotoma] (*Hylotoma humeralis* F. Smith, 1874). Russia: **FE** (PR). – Japan (Hon).
- Arge enodis** (Linnaeus, 1767) [Tenthredo] (*Tenthredo violacea* Fabricius, 1779; *T. coeruleipennis* Retzius, 1783; *Hylotoma atrata* Klug, 1814; *H. amethystina* Klug, 1814; *H. vulgaris* Klug, 1834). Host plant: the smooth-leaved species of *Salix* (Salicaceae). Russia: **EP** (NW, C, E, S, NC), **UR**, **WS** (TK, NS, KM, AL), **ES** (KR, BR, YA), **FE** (PR). – Europe (WE, EE, NE, SE), Georgia, Armenia, Turkey, Iran, Kyrgyzstan, Kazakhstan, Korean Peninsula, Japan (Hon).
- Arge expansa** (Klug, 1834) [Hylotoma]. Host plant: *Salix* spp., *Populus* spp. (Salicaceae). Russia: **WS** (TM, KM, AL), **FE** (AM, KU, KA). – Mongolia, China (NC).
- Arge flavomixta** (André, 1881) [Hylotoma]. Russia: “Sibiria or.” (André, 1881), **FE** (PR). – Korean Peninsula.
- Arge frivaldszkyi** (Tischbein, 1852) [Hylotoma] (*Hylotoma schmiedeknechti* Costa, 1890). Russia: **EP** (NC, CR). – Europe (EE, SE), Caucasus, Turkey.
- Arge fuscipennis** (Herrich-Schäffer, 1835) [Hylotoma]. Host plant: *Betula* spp. (Betulaceae), ?*Lythrum* spp. (Lythraceae), ?*Solidago* spp. (Asteraceae). Russia: **EP** (C, E), **UR**. – Europe (WE, EE, SE).
- Arge fuscipes fuscipes** (Fallén, 1808) [Hylotoma] (*Hylotoma violacea* Klug, 1814; *H. atrocoerulea* Serville, 1823; *H. atrocoerulea* Lepeletier, 1823; *Arge fuscinervis* Lindqvist, 1974). Host plant: *Betula* spp. (Betulaceae), *Salix* spp., *Populus* spp. (Salicaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TK, NS, KM, AL), **ES** (KR, ZB), **FE** (KH, PR, KU, KA, MG). – Europe (WE, EE, NE, SE), Kazakhstan, Mongolia, China (NE), Korean Peninsula.
- Arge gracilicornis** (Klug, 1814) [Hylotoma] (*Tenthredo coerulescens* Geoffroy, 1785; *T. incolorata* Christ, 1791; *Hylotoma pilicornis* Leach, 1817; *H. cyamella* Klug, 1834; *H. crassa* Konow, 1884). Host plant: *Rosa* spp., *Rubus* spp. (Rosaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, TK, NS, KM, AL), **ES** (TU, KR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KA, MG, CH). – Europe (WE, EE, NE, SE), Caucasus, Turkey, Mongolia, China (Wei et al., 2006), Korean Peninsula, Japan (Hok, Hon, Shi).
- Arge hasegawae** Takeuchi, 1927. Russia: **FE** (PR). – Korean Peninsula.
- Arge jonasi** (W.F. Kirby, 1882) [Hylotoma] (*Hylotoma nigritarsis* F. Smith, 1874; *H. japonica* Marlatt, 1898). Host plant: *Pourthiae* spp., *Aria* spp., *Crataegus* spp., *Sorbus* spp., *Cotoneaster* spp. (Rosaceae). Russia: **FE** (KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Arge kobayashii** Takeuchi, 1931. Host plant: *Betula* spp. (Betulaceae). Russia: **FE** (SA, KU). – Japan (Hok).
- Arge longicornis** Kuznetzov-Ugamskij, 1927 (*A. ciliaris pacifica* Malaise, 1931). Russia: **FE** (PR). – Korean Peninsula.

- Arge macrops** Shinohara, Hara et Kim, 2009. Host plant: *Ulmus* spp. (Ulmaceae). Russia: **WS** (AL), **FE** (PR). – China (NE, NC, CC), Korean Peninsula.
- Arge mali** (Uchiyama, 1906) [Hylotoma] (*A. xanthocera* Mocsáry, 1909; *Hylotoma mali* Matsumura, 1912). Host plant: *Malus* spp. (Rosaceae). Russia: **FE** (AM, KH, PR). – China (NE), Korean Peninsula, Japan (Hok, Hon).
- Arge melanochra** (Gmelin, 1790) [Tenthredo] (*Hylotoma femoralis* Klug, 1814; *H. dimidiata* Lepoletier, 1823; *H. dimidiata* Serville, 1823; *H. nigritarsis* Klug, 1834; *H. similis* Rudow, 1871; *Arge fuliginata* Konow, 1907). Host plant: *Crataegus* spp. (Rosaceae). Russia: **EP** (S, NC, CR), **UR** (S), **WS** (KM). – Europe (WE, EE, SE), Armenia, Asia Minor, Iran, Turkmenistan.
- Arge nigripes nigripes** (Retzius, 1783) [Tenthredo] (*Hylotoma anglica* Leach, 1817). Host plant: *Rosa* spp., *Fragaria* spp., *Rubus* spp. (Rosaceae); live in colonies. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, TK, NS, KM, AL), **ES** (TU, KR, IR, BR, YA), **FE** (AM, PR, SA, KA, MG). – Europe (WE, EE, NE, SE), Georgia, Turkey, Kyrgyzstan, Kazakhstan, Mongolia.
- Arge nigronodosa** (Motschulsky, 1860) [Hylotoma] (*Hylotoma flava* Snellen van Vollenhoven, 1860; *H. trinotata* F. Smith, 1874; *H. similis* F. Smith, 1874; *H. dubia* W.F. Kirby, 1882; *H. quadripunctata* W.F. Kirby, 1882). Host plant: *Rosa* spp. (Rosaceae). Russia: **FE** (AM, KH, PR, SA). – China (NE), Japan (Hok, Hon, Shi, Kyu), Nepal.
- Arge nigrovaginata** Malaise, 1931. Russia: **FE** (AM, PR). – China (NE, CC, SE), Korean Peninsula, Japan (Hon).
- Arge nipponensis** Rohwer, 1910. Host plant: *Rosa* spp. (Rosaceae); 3-4 generations per year. Russia: **FE** (SA, KU). – China (SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Sado).
- Arge obesa** Hara et Shinohara, 2012. Host plant: *Carpinus* spp., *Ostrya japonica* (Betulaceae). Russia: **FE** (KU). – Japan (Hok, Hon, Shi, Kyu).
- Arge ochropus** (Gmelin, 1790) [Tenthredo] (*A. rosincola* Schrank, 1802; *Hylotoma rosarum* Klug, 1814; *H. pyrenaica* André, 1879; *Arge soror* Konow, 1890; *A. pyrenaica nigripes* Konow, 1895; *A. modesta* Konow, 1905; *A. pyrenaica mauritanica* Schulz, 1906; *Hylotoma rosae diversicolor* Pic, 1917; *Arge pyrenaica luteola* Muche, 1977). Host plant: *Rosa* spp. (Rosaceae). Russia: **EP** (N, NE, C, E, S, NC, CR), **UR**, **WS** (OM, TK, NS, KM, AL), **ES** (TU, KR, IR, YA). – Europe (WE, EE, NE, SE), N Africa, Georgia, Armenia, Asia Minor, Iran, Turkmenistan, Kazakhstan; introduced into N America.
- Arge pagana pagana** (Panzer, 1797) [Tenthredo] (*Tenthredo tricolor* Gmelin, 1790; *T. nigripennis* Panzer, 1804; *Hylotoma flaviventris* Fallén, 1807; *H. assimilis* Radoszkowsky, 1889). Host plant: *Rosa* spp. (Rosaceae); live in colonies. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TK, NS, KM, AL), **ES** (TU, KR, IR, BR, YA), **FE** (AM, PR, KA). – Europe (WE, EE, NE, SE), N Africa, Mongolia, China (NE, NC, CC, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu), India (NE).
- Arge paganiformis** Rohwer, 1925. Host plant: *Rosa* spp. (Rosaceae). Russia: **FE** (KH, PR).
- Arge pallidinervis** Gussakovskij, 1935. Russia: **EP** (CR). – Europe (EE), Armenia.
- Arge pleuritica** (Klug, 1834) [Hylotoma] (*Hylotoma sanguinicollis* André, 1880). Russia: **EP** (E, S, NC, CR). – Europe (EE, SE), Caucasus, Asia Minor, Uzbekistan.
- Arge potanini** Jakovlev, 1891 (*Hylotoma zonalis* Matsumura, 1911). Host plant: *Acer mono*, *A. mayrii* (Sapindaceae). Russia: **ES** (IR, BR, ZB), **FE** (AM, KH, PR, SA). – Mongolia, China (NE, NC), Korean Peninsula.
- Arge pseudorejecta** Wei et Lee, 2016. Russia: **FE** (PR). – China (NE), Korean Peninsula.
- Arge pullata** (Zaddach, 1859) [Hylotoma] (*Hylotoma ventriosa* Zaddach, 1864; *Arge nyemitawa* Rohwer, 1925). Host plant: *Betula* spp. (Betulaceae). Russia: **EP** (NW, C, E, S, NC), **UR**, **WS** (TM, TK, NS, AL), **ES** (KS, KR, ZB), **FE** (PR, SA, KU). – Europe (WE, EE, NE, SE), Armenia, Turkey, China (NE, NC), Korean Peninsula, Japan (Hok, Hon).
- Arge rufocincta** Gussakovskij, 1935. Russia: **FE** (PR). – China (NE), Japan (Hok).
- Arge rustica** (Linnaeus, 1758) [Tenthredo] (*Tenthredo atrata* Forster, 1771; *Cryptus segmentarius* Panzer, 1803; *Hylotoma klugii* Leach, 1817; *H. albicruris* Brullé, 1832; *H. leachii* Stephens, 1835; *H. marginata* Boheman, 1852; *H. discus* Costa, 1858; *H. saliceti* Rudow, 1871; *H. thomsoni* Konow, 1884; *Arge segmentaria rufiventris* Konow, 1899). Host plant: *Quercus* spp. (Fagaceae). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, EE, NE, SE), Armenia, Turkey, Syria.
- Arge scita** (Mocsáry, 1880) [Hylotoma] (*Hylotoma proxima* André, 1881; *Arge debilis* Konow, 1887; *A. zarudnyi* Gussakovskij, 1935). Russia: **EP**

- (Taeger, Blank, 2011). – Europe (SE, EE), Armenia, Asia Minor, Iran, Turkmenistan.
- Arge semicoerulea** (Klug, 1814) [Tenthredo] (*Hylotoma dimidiata* Klug, 1814). Russia: **EP** (E), **ES** (KS, KR, ZB), **FE** (AM, PR). – Mongolia, China (NE).
- Arge similis** (Snellen van Vollenhoven, 1860) [*Hylotoma*] (*Hylotoma imperator* F. Smith, 1874). Host plant: Rhododendron spp. (Ericaceae); up to 3 generations per year. Russia: **FE** (PR, KU). – China (NE, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu), Nepal, India.
- Arge simillima** (F. Smith, 1874) [*Hylotoma*] (*A. forcifula* Jakovlev, 1891; *A. coriacea* Jakovlev, 1891; *A. simillima asahi* Tskeuchi, 1932). Russia: **FE** (KH, PR, KU). – China (NE, SE, CC, SW), Korean Peninsula, Japan (Hon).
- Arge simulatrix** Konow, 1887. Russia: **EP** (NC). – Georgia, Asia Minor, Iran.
- Arge solowiyofka** (Matsumura, 1911) [*Hylotoma*]. Host plant: Betula spp. (Betulaceae). Russia: **FE** (PR, SA). – Mongolia, China (NE, CC, SE), Japan (Hok, Hon).
- Arge stroganova** Sundukov, 2007 (*A. altaica* Stroganova, 1977). Russia: **WS** (AL).
- Arge suspicax** Konow, 1908 (*A. punctifrons* Kuznetzov-Ugamskij, 1927; *A. masudai* Takeuchi, 1932). Russia: **ES** (IR, BR, ZB), **FE** (AM, PR). – Mongolia, China (NE), Korean Peninsula, Japan (Hon, Shi).
- Arge ustulata** (Linnaeus, 1758) [Tenthredo] (*Tenthredo flavipes* Retzius, 1783; *T. ochroptera* Geoffroy, 1785; *T. crocipennis* Christ, 1791; *T. pilicornis* Preyssler, 1793; *Hylotoma claripennis* Rudow, 1871; *H. distinguenda* Konow, 1884; *Arge monostega* Konow, 1886; *Hylotoma anceps* Radoszkowsky, 1889; *Arge ustulata stigmatica* Pic, 1927). Host plant: Salix spp. (Salicaceae), Betula spp. (Betulaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TK, NS, KM, AL), **ES** (KS, KR, IR, BR, YA, ZB), **FE** (AM, PR, SA, KU, KA). – Europe (WE, EE, NE, SE), Georgia, Turkey, Kyrgyzstan, Mongolia, Korean Peninsula.
- Arge verticosa** Konow, 1907. Russia: **WS** (AL). – Kazakhstan, China (NW).
- Arge zhelochovzevi** Stroganova, 1966. Russia: **ES** (KR, IR).
- KOKUJEWIA** Konow, 1902. Type species *Kokujewia ectrapela* Konow, 1902. The larvae feed on the Polygonaceae. The genus is distributed in West Palaearctic. Number of species: World and Palaearctic – 3; Russia – 1.
- Kokujewia ectrapela** Konow, 1902 (*K. ectrapela clarescens* Zirngiebl, 1949). Host plant: Rumex spp. (Polygonaceae). Russia: **EP** (NC). – Caucasus, Turkey, Iran.
- PSEUDARGE** Gussakovskij, 1935. Type species *Pseudarge rubicunda* Gussakovskij, 1935. The genus is distributed in the Palaearctic and Oriental regions. Number of species: World – 6; Palaearctic – 4; Russia – 1.
- Pseudarge eversmanni** Gussakovskij, 1935. Russia: **EP** (E, S). – Europe (EE).
- SPINARGE** Wei, 1998. Type species *Spinarge sichuanensis* Wei, 1998. The larvae feed on the Betulaceae. The genus is distributed in East Asia and Oriental region. Number of species: World – 11; Palaearctic – 9; Russia – 2.
- Spinarge fulvicornis** (Mocsáry, 1909) [Arge]. Host plant: Betula spp. (Betulaceae). Russia: **ES** (YA), **FE** (AM, KH, PR, SA, KU). – China (NE, CC), Korean Peninsula, Japan (Hok, Hon, Shi).
- Spinarge metallica** (Klug, 1834) [*Hylotoma*]. Host plant: Betula spp. (Betulaceae). Russia: **EP** (NW, C, E), **UR**, **WS** (NS, KM, AL), **ES** (IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KA). – Europe (WE, EE, NE, SE), Mongolia, Korean Peninsula.

Subfamily STERICTIPHORINAE

- APROCEROS** Malaise, 1931. Type species *Aproceros umbricola* Malaise, 1931. The larvae feed on the Ulmus spp. (Ulmaceae). The genus is distributed in East Asia; one species in the Oriental region, and one species was introduced into West Europe. Number of species: World – 10; Palaearctic – 9; Russia – 3.
- Aproceros hakusanus** Togashi, 1962. Host plant: Ulmus spp. (Ulmaceae). Russia: **FE** (PR, SA, KU). – Japan (Hon).
- Aproceros leucopoda** Takeuchi, 1939. Host plant: Ulmus spp. (Ulmaceae). Russia: **EP** (C, S, NC), **WS** (KM), **FE** (PR, SA). – Europe (WE, EE, SE), Kazakhstan, China (CC), Japan (Hok).
- Aproceros pallidicornis** (Mocsáry, 1909) [Aprosthemma] (*A. umbricola* Malaise, 1931). Host plant: Ulmus spp. (Ulmaceae). Russia: **FE** (AM, KH, PR). – Korean Peninsula, Japan (Hon).
- APROSTHEMA** Konow, 1899 (*Copidoceros* Forsius, 1921; *Lyrola* Ross, 1937). Type species

- Hylotoma brevicornis* Fallén, 1808. The larvae feed on the Fabaceae. The genus is distributed in the Holarctic and Oriental regions, mainly in the Palaearctic region. Number of species: World – about 55; Palaearctic – > 50; Russia – 22 (+ 2 ssp.).
- Aprosthemma austriacum** (Konow, 1892) [Schizocera]. Russia: **EP** (CR). – Europe (WE, EE, NE, SE), Kazakhstan.
- Aprosthemma axillare** (Zaddach, 1863) [Schizocera] (*Schizocera zaddachi* André, 1880). Russia: **EP** (NW, NC, CR). – Europe (WE, EE, SE).
- Aprosthemma bifidum** (Klug, 1834) [Hylotoma] (*Schizocera flavipes* Zaddach, 1864; *S. geniculata* Thomson, 1871). Russia: **EP** (NW, NC). – Europe (WE, EE, NE, SE).
- Aprosthemma bifurca** (Klug, 1834) [Hylotoma] (*Schizocera fusca* Zaddach, 1863). Russia: **EP** (NW, C). – Europe (WE, EE, NE, SE), Turkey.
- Aprosthemma brevicorne** (Fallén, 1808) [Hylotoma] (*Schizocera pallipes* Bremi-Wolf, 1849). Russia: **EP** (C, E, S). – Europe (WE, EE, NE).
- Aprosthemma enslini** Dovnar-Zapolskij, 1930. Russia: **EP** (NC).
- Aprosthemma fulvum** Vasilenko, 2009. Russia: **WS** (NS).
- Aprosthemma fuscicorne** (Thomson, 1871) [Schizocera]. **EP** (NW). – Europe (WE, EE, NE, SE).
- Aprosthemma humeratum** (Konow, 1892) [Schizocera]. Russia: **EP** (Zhelochovtsev, Zinovjev, 1995). – Europe (EE, SE).
- Aprosthemma instratum instratum** (Zaddach, 1859) [Schizocera]. Russia: **EP** (NW, C). – Europe (WE, EE, SE).
- Aprosthemma intermedium** (Zaddach, 1864) [Schizocera] (*A. hyalinoptera* Conde, 1934; *A. volgensis* Gussakovskij, 1935). Host plant: *Lathyrus vernus* (Fabaceae). Russia: **EP** (NW, C, S). – Europe (WE, EE, NE).
- Aprosthemma maculatum maculatum** (Jurine, 1807) [Cryptus] (*Hylotoma scutellaris* Herrich-Schäffer, 1835; *Schizocera scutellaris dorsalis* Konow, 1892; *Aprosthemma maculata moravica* Gregor, 1940). Russia: **EP** (CR). – Europe (WE, EE, SE).
- Aprosthemma maculatum nigripenne** Muche, 1972. Russia: **EP** (NC). – Georgia.
- Aprosthemma melanurum** (Klug, 1814) [Hylotoma] (*Schizocera cylindricornis* Thomson, 1871; *S. alfskeni* Konow, 1895; *S. friesei* Konow, 1895). Host plant: *Lathyrus* spp., *Vicia* spp. (Fabaceae); in East Siberia 2 generations. Russia: **EP** (NW, C, E, S, NC), **UR**, **WS** (TK, NS, KM), **ES** (KR, IR). – Europe (WE, EE, NE, SE), Central Asia, China (NC).
- Aprosthemma peletieri peletieri** (Villaret, 1832) [Schizocerus] (*Schizocera bleusei* Pic, 1918). Russia: **EP** (C, S, NC). – Europe (WE, EE, SE).
- Aprosthemma peletieri melanoptera** Muche, 1972. Russia: **EP** (NC). – Georgia.
- Aprosthemma sibiricum** Gussakovskij, 1935. Russia: **ES** (ZB).
- Aprosthemma staudingeri georgiense** Muche, 1972. Russia: **EP** (NC). – Georgia.
- Aprosthemma stroganovae** Vasilenko, 2009. Russia: **WS** (KM).
- Aprosthemma syrmiente** (Mocsáry, 1897) [Schizocera]. Russia: **EP** (NC). – Europe (SE), Turkey.
- Aprosthemma tardum** (Klug, 1814) [Hylotoma] (*A. carpentieri* Konow, 1902). Host plant: *Lathyrus* spp., *Vicia* spp. (Fabaceae). Russia: **EP** (C, E, S, NC), **UR**, **WS** (KM, AL), **ES** (KS). – Europe (WE, EE, SE), Turkey, Kyrgyzstan, Kazakhstan.
- Aprosthemma tauricum** Gussakovskij, 1935. **EP** (S, CR). – Europe (EE).
- Aprosthemma vittatum** (Mocsáry, 1879) [Schizocera]. Russia: **EP** (CR). – Europe (EE, SE).
- Aprosthemma xanthurum** Gussakovskij, 1935. Russia: **WS** (TK), **FE** (AM).
- SERICTIPHORA** Billberg, 1820 (*Schizocerus* Berthold, 1827; *Schizocerus* Lepeletier et Serville, 1828; *Schyzocera* Latreille, 1829; *Schizocera* Guérin, 1834; *Cyphona* Dahlbom, 1835; *Schizoceros* Konow, 1899). Type species *Tenthredo furcata* Villers, 1789. The larvae feed on the Rosaceae. The genus is distributed in the Holarctic; one species in Afrotropical, and two species in Oriental regions. Number of species: World – > 40; Palaearctic – about 35; Russia – 4.
- Sterictiphora caspica** Koch, 1988. Russia: **EP** (NC). – Iran.
- Sterictiphora furcata** (Villers, 1789) [*Tenthredo*] (*Tenthredo rubi idaei* Rossi, 1790; *Schizocera inaequalis* Bremi-Wolf, 1849; *Aprosthemma terebralis flavipes* Enslin, 1917). Host plant: *Rubus* spp., *Prunus* spp. (Rosaceae). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, EE, SE), N Africa, Azerbaijan, Armenia, Turkey, Syria, Iran, Kazakhstan.
- Sterictiphora geminata** (Gmelin, 1790) [*Tenthredo*]. Host plant: *Rosa* spp., *?Sorbus* spp. (Rosaceae). Russia: **EP** (C, NC), **UR**, **WS** (NS, KM), **ES** (BR, ZB), **FE** (PR, KA). – Europe (WE, EE, NE, SE), Armenia, Turkey, Turkmenistan, Kyrgyzstan.
- Sterictiphora orientalis** (Zhelochovtsev, 1939) [*Schizocera*]. Russia: **FE** (PR).

3. FAMILY BLASTICOTOMIDAE

Adults are characterized by their short 4-segmented antennae, of which an elongate 3rd segment forms almost the entire flagellum except for the minute (and sometimes obsolete) 4th segment. Scutellum without a post-tergite; the wing venation is unique in having the pear-shaped cell *IM* of the fore wing, rounded apically; the stigma is large and almost semi-circular in shape; vein *Sc* without a transverse branch joining *C*; and the anal cell has an oblique cross-vein; the wing membranes are corrugated at the apices. The legs are without preapical tibial spines and the front tibia has 2 apical spurs, the inner one of which is bifid apically. The abdomen is carinate laterally, and the ovipositor is of the exserted type, about half as long as the abdomen, though the apical projecting part, the sawsheath, is much shorter than the basal plate. The larvae are stem-borers in ferns and of a oligopodous type, with no abdominal legs.

The family is distributed in the Palaearctic and Oriental regions. Number of taxa: World – 2 genera, 12 species; Palaearctic – 2/8; Russia – 2/3 (+ 1 ssp.).

R e f e r e n c e s. Gussakovskij, 1935; Zhelochovtsev, 1935a, 1988; Ermolenko, 1972; Togashi, 1989; Zhelochovtsev, Zinovjev, 1995; Kim et al., 1997; Wei, Nie, 1999; Liston et al., 2006, 2014a; Shcherbakov, 2006; Taeger et al., 2006, 2010; Wei et al., 2006; Biryukova, Novgorodova, 2008; Schedl, 2009; Novgorodova, Biryukova, 2011; Taeger, Blank, 2011; Vasilenko, 2011b; Sundukov, Lelej, 2012; Kostyunin, 2015; Sundukov, 2015; Noblecourt, 2016.

BLASTICOTOMA Klug, 1834. Type species

Blasticotoma filiceti Klug, 1834. The larvae in the petioles of ferns (Polypodiophyta). The genus is distributed in the Palaearctic region; 1 species in the Oriental region. Number of species: World – 5; Palaearctic – 4; Russia – 2 (+ 1 ssp.).

Blasticotoma atra Zhelochovtsev, 1935. Host plant: ferns of families of Dryopteridaceae, Onocleaceae, Osmundaceae, Dennstaedtiaceae. Russia: **FE** (KH, PR, KU). – Japan (Hok, Hon).

Blasticotoma filiceti filiceti Klug, 1834. Host plant: *Athyrium* spp., *Dryopteris* spp. (Dryopteridaceae), *Matteuccia* spp. (Onocleaceae), *Osmunda* spp. (Osmundaceae), *Pteridium* spp. (Dennstaedtiaceae). Russia: **EP** (N, C, E), **UR**, **WS** (NS, KM, AL), **ES** (KR, IR, BR). – Europe (WE, EE, NE).

Blasticotoma filiceti pacifica Malaise, 1931. Host plant: ferns of families of Dryopteridaceae, Onocleaceae, Osmundaceae, Dennstaedtiaceae. Russia: **FE** (KH, PR, SA, KU). – China (NE, CC), Korean Peninsula, Japan (Hok, Hon).

RUNARIA Malaise, 1931 (*Bohea* Maa, 1944). Type species *Runaria reducta* Malaise, 1931. The genus is distributed in East Asia and Oriental region. Number of species: World – 7; Palaearctic – 4; Russia – 1.

Runaria reducta Malaise, 1931. Host plant: ferns of families of Dryopteridaceae, Onocleaceae, Osmundaceae, Dennstaedtiaceae. Russia: **FE** (AM, PR, KU). – Japan (Hok, Hon).

4. FAMILY CIMBICIDAE

Stout fast-flying insects with strongly clubbed antennae and broad, laterally carinate abdomens, arched above and flat beneath, so that they can be curled right down under the thorax at rest. Fore wing vein $2r$ presents. Tibiae without preapical spines and front tibia with pair of unmodified apical spurs. The larvae are free-feeding, with 8 pairs of abdominal legs. The larvae feed on Salicaceae, Betulaceae, Caprifoliaceae and other plant families. Pupation usually occurs in a cocoon with a dense wall in the soil or on branches of trees and shrubs.

The family is distributed in the Northern Hemisphere and Neotropical region. Number of taxa: World – 4 subfamilies, 15 genera, about 190 species; Palaearctic – 3/10/about 150; Russia – 3/8/72 (+ 2 spp.).

R e f e r e n c e s. Konow, 1906; Mocsáry, 1909; Kuznetsov-Ugamskij, 1927; Zhelochovtsev, 1935a, 1968, 1976, 1988; Malaise, 1939; Gussakovskij, 1947; Saarenen, 1950b; Ermolenko, 1972, 1973, 1976, 1984, 1992; Verzhutskii, 1974, 1981; Zhelochovtsev, Prochorova, 1976; Stroganova, 1978a; Viitasaari, 1989; Shinohara, 1990; Vikberg, Viitasaari, 1991; Liston, 1995; Zhelochovtsev, Zinovjev, 1995; Kim, Shinohara, 1997; Togashi, 1998; Lee, Jung, 1999; Hara, Shinohara, 2000, 2017; Vasilenko, 2004, 2009b, 2010b,d, 2011c; Taeger, 2005; Vasilenko, Dolgov, 2005; Kajmuk, Popov, 2006; Liston et al., 2006, 2014a,b; Taeger et al., 2006, 2010; Wei et al., 2006; Roller, Haris, 2008; Humala, Polevoi, 2009; Schedl, 2009; Sundukov, Lelej, 2009, 2012; Popov, Kajmuk, 2010a; Blank, 2011; Sundukov, 2011b, 2013, 2015; Taeger, Blank, 2011; Vasilenko, Korshunov, 2012; Beneš, 2014b; Noblecourt, 2016.

Subfamily ABIINAE

ABIA Leach, 1817 (*Zaraea* Leach, 1817; *Parabia* Semenov, 1891; *Orientabia* Malaise, 1934; *Hemibia* Semenov, 1937; *Auroabia* Kangas, 1946; *Aenoabia* Kangas, 1946; *Procimbex* Hong, 1984). Ty p e s p e c i e s *Tenthredo sericea* Linnaeus, 1767. The larvae on the Caprifoliaceae. The genus is distributed in the Holarctic and Oriental regions. Number of species: World – 52; Palaearctic – 43; Russia – 24.

Abia aenea (Klug, 1820) [Cimbex] (*A. bifida* Thomson, 1871; *A. bigens* Kangas, 1946). Host plant: Lonicera spp., Symphoricarpos spp. (Caprifoliaceae), Samucus spp. (Adoxaceae). Russia: **EP** (NW, C, E, S,

NC), **WS** (TK, KM). – Europe (WE, EE, NE, SE), Georgia, Turkey.

Abia aurata (Takeuchi, 1931) [*Zaraea*]. Russia: **FE** (KU). – Korean Peninsula, Japan (Hok).

Abia berezowskii Semenov, 1896. Russia: **FE** (AM, PR). – Mongolia, China (SW), Korean Peninsula, Japan (Hon).

Abia candens Konow, 1887 (*A. symbalophthalma* Semenov, 1892; *A. candens subopaca* Kangas, 1946). Host plant: Knautia arvensis (Caprifoliaceae). Russia: **EP** (NW, C, E, NC). – Europe (WE, EE, NE, SE).

Abia coreana Takeuchi, 1927 (*Orientabia pacifica* Semenov et Gussakovskij, 1937). Russia: **FE** (PR). – Korean Peninsula.

Abia dubiosa (Gussakovskij, 1947) [*Orientabia*]. Russia: **FE** (SA). – Japan (Hon).

Abia egregia Kuznetsov-Ugamskij, 1927. Russia: **FE** (PR).

Abia fasciata (Linnaeus, 1758) [*Tenthredo*]. Host plant: Lonicera spp., Symphoricarpos spp., Linnaea spp., Leycesteria spp. (Caprifoliaceae). Russia: **EP** (N, NW, C, E), **UR**, **WS** (TM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, SA, KA, MG). – Europe (WE, EE, NE, SE), Kazakhstan, China (NE, NW), Korean Peninsula, Japan (Hok, Hon).

Abia fulgens Zaddach, 1863 (*A. fulgens unicoloricornis* Pic, 1933; *A. plana* Benson, 1954). Host plant: Lonicera spp. (Caprifoliaceae). Russia: **EP** (NC). – Europe (WE, EE, SE).

Abia gussakovskii Semenov, 1935. Russia: **EP** (NC).

Abia japonica Cameron, 1887 (*A. pilosa* Konow, 1896; *A. bantaizana* Matsumura, 1912; *Orientabia finitima* Semenov et Gussakovskij, 1937). Host plant: Weigela spp. (Caprifoliaceae). Russia: **FE** (PR, KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Sado).

Abia lewisi Cameron, 1887 (*Zaraea lewisi matsu-*
murai Takeuchi, 1931). Host plant: Lonicera spp. (Caprifoliaceae). Russia: **FE** (KU). – Japan (Hon).

Abia marginata Moscáry, 1909. Host plant: Lonicera spp. (Caprifoliaceae). Russia: **FE** (KH, PR). – Korean Peninsula.

Abia metallica Moscáry, 1909. Host plant: Lonicera spp. (Caprifoliaceae). Russia: **FE** (KH, PR). – China (NE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Abia mutica Thomson, 1871. Host plant: Lonicera spp., Knautia arvensis (Caprifoliaceae). Russia: **EP** (N, NW, C), **UR**, **WS** (NS, AL), **ES** (TU, KR, IR, BR, YA, ZB), **FE** (PR). – Europe (WE, EE, NE, SE), Kazakhstan, Mongolia.

- Abia ochotica** (Semenov et Gussakovskij, 1937) [Orientalia]. Russia: **FE** (KH).
- Abia sachalinensis** Takeuchi, 1931. Russia: **FE** (PR, SA). – Japan (Hok).
- Abia semenoviiana** Gussakovskij, 1947. Host plant: Lonicera spp. (Caprifoliaceae). Russia: **ES** (BR, ZB), **FE** (PR). – Korean Peninsula.
- Abia sericata** Zhelochovtsev, 1935. Russia: **FE** (AM, KH, PR).
- Abia sericea** (Linnaeus, 1767) [Tenthredo] (*A. dorsalis* Costa, 1859). Host plant: Succica spp., Knautia spp., Dipsacus spp. (Caprifoliaceae), ?Fragaria spp. (Rosaceae). Russia: **EP** (NW, C, E, S, NC, CR). – Europe (WE, EE, NE, SE), Caucasus, Turkey, China (NW).
- Abia sibirica** Mocsáry, 1883. Host plant: Lonicera spp. (Caprifoliaceae). Russia: **WS** (OM, NS, AL), **ES** (KS, BR). – Kazakhstan.
- Abia skorikovi** Semenov, 1935. Russia: **EP** (NC).
- Abia tshershkii** Semenov, 1935. Russia: **FE** (KH).
- Abia zhelochovtzevi** (Gussakovskij, 1947) [Zaraea]. Host plant: Lonicera spp. (Caprifoliaceae). Russia: **FE** (KH, PR).

Subfamily CIMBICINAE

AGENOCIMBEX Rohwer, 1910. Type species *Cimbex maculata* Marlatt, 1898. The genus is distributed in East Asia; one species in the Oriental region. Number of species: World and Palaearctic – 6; Russia – 2.

Agenocimbex eous Semenov, 1935. Russia: **FE** (PR). – Korean Peninsula, Japan (Kyu).

Agenocimbex malaisei Gussakovskij, 1947. Russia: **FE** (PR).

CIMBEX Olivier, 1791 (*Crabro* Geoffroy, 1762; *Clavellarius* Olivier, 1789; *Clavellaria* Lamarck, 1801; *Palaeocimbex* Semenov, 1935; *Deuterocimbex* Semenov, 1935; *Neocimbex* Malaise, 1937; *Allocimbex* Zirngiebl, 1953). Type species *Tenthredo lutea* Linnaeus, 1758. The larvae mainly on the Salicaceae and Betulaceae. The genus is distributed in the Holarctic region; one species in the Oriental region. Number of species: World – 14; Palaearctic – 10; Russia – 6 (+ 2 spp.).

Cimbex amurensis Forsius, 1930. Host plant: Salix spp. (Salicaceae), Betula spp. (Betulaceae). Russia: **FE** (AM, PR).

Cimbex carinulatus Konow, 1897 (*C. nomurae* Marlatt, 1898; *C. suzukii* Matsumura, 1912). Host

plant: Pyrus spp., Amelanchier asiatica, Prunus spp., Chaenomeles spp. (Rosaceae). Russia: **FE** (PR, KU). – Korean Peninsula, Japan (Hok, Hon).

Cimbex connatus connatus (Schrank, 1776) [Tenthredo] (*Crabro maculatus* Geoffroy, 1785; *Tenthredo montana* Panzer, 1801; *T. alnivaga* Bechstein et Scharfenberg, 1805; *Cimbex ornata* Serville, 1823; *C. luteola* Serville, 1823; *C. ornata* Lepeletier, 1823; *C. luteola* Lepeletier, 1823; *C. humboldtii* Ratzeburg, 1844; *C. variabilis nigroviolacea* Eversmann, 1847; *C. violascens* Thomson, 1871). Host plant: Salix spp., Populus spp. (Salicaceae), Alnus spp. (Betulaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (NS, KM), **ES** (IR). – Europe (WE, EE, NE, SE), Caucasus, Kazakhstan.

Cimbex connatus japonicus W.F. Kirby, 1882 (*C. saporensis* Matsumura, 1912). Host plant: Salix spp. (Salicaceae). Russia: **FE** (KH, PR, SA, KU). – China (NE, CC), Japan (Hok, Hon).

Cimbex connatus taukushi Marlatt, 1898 (*C. taukushiensis* Konow, 1905; *C. castanea* Matsumura et Ushida, 1927; *Neocimbex taukushi carbonaria* Malaise, 1939). Host plant: Alnus spp. (Betulaceae), Salix spp. (Salicaceae). Russia: **ES** (IR, BR, YA, ZB), **FE** (AM, KH, PR, KU). – Mongolia, China (NE), Korean Peninsula, Japan (Hon, Shi, Kyu).

Cimbex femoratus (Linnaeus, 1758) [Tenthredo] (*Tenthredo tristis* Fabricius, 1779; *Crabro lunulatus* Geoffroy, 1785; *C. annulatus* Geoffroy, 1785; *Tenthredo sylvarum* Fabricius, 1793; *Cimbex europaea* Leach, 1817; *C. varians* Leach, 1817; *C. variabilis* Klug, 1820; *Tenthredo russa* Klug, 1820; *Cimbex schaefferi* Lepeletier, 1823; *C. pallens* Lepeletier, 1823; *C. schaefferi* Serville, 1823; *C. pallens* Serville, 1823; *C. pallidus* Stephens, 1829; *C. venusta* Perty, 1833; *C. biguetina* Lepeletier, 1834; *C. pallida* Stephens, 1835; *C. betulae* Zaddach, 1863; *C. betulae pulla* Zaddach, 1863; *C. betulae lutescens* Zaddach, 1863; *C. betulae nigra* Zaddach, 1863; *C. tonnaitchana* Matsumura, 1911; *C. femorata unicolor* Enslin, 1917; *C. femorata abdominalis* Enslin, 1917; *C. femorata ornata* Uchida, 1927; *C. quadrimaculata sachalinensis* Uchida, 1927; *Crabro uchidai* Takeuchi, 1931). Host plant: Alnus spp. (Betulaceae), Salix spp. (Salicaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KA, MG). – Europe (WE, EE, NE, SE), Turkey, Kyrgyzstan, Uzbekistan, Kazakhstan, Mongolia, China (NE, NC), Korean Peninsula, Japan (Hok, Hon).

Cimbex luteus (Linnaeus, 1758) [Tenthredo] (*C. grifinii* Leach, 1817; *C. saliceti* Zaddach, 1863; *C. variabilis salicis* Zaddach, 1863; *C. brevispina* Thomson, 1871; *C. lutea testacea* Konow, 1890; *C. saliceti testacea* Dalla Torre, 1894; *C. capreae* Konow, 1897; *C. femorata coreana* Uchida, 1927). Host plant: Salix spp., Populus spp. (Salicaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, KR, IR, BR, YA), **FE** (KH, PR, SA, MG). – Europe (WE, EE, NE, SE), Kyrgyzstan, Uzbekistan, Kazakhstan, China (NC, NW), Korean Peninsula, Japan (Hok, Hon).

Cimbex sibiricus W.F. Kirby, 1882. Russia: **ES** (IR, BR).

LEPTOCIMBEX Semenov, 1896 (*Euclavellaria* Enslin, 1911; *Cimbicisoma* Rohwer, 1915; *Okamotonius* Sato, 1928). Type species *Leptocimbex potanini* Semenov, 1896. The genus is distributed in East Asia and Oriental region. Number of species: World – 26; Palaearctic – 21; Russia – 4.

Leptocimbex allantiformis (Mocsáry, 1909) [Clavelaria] (*L. petrimagni* Malaise, 1931). Russia: **FE** (KH, PR). – China (NE), Korean Peninsula.

Leptocimbex gracilentus (Mocsáry, 1904) [Clavelaria] (*Okamotonius kurisuei* Sato, 1928). Russia: **FE** (KH, PR). – China (SE), Korean Peninsula.

Leptocimbex potanini Semenov, 1896 (*Clavellaria marginata* R.E. Turner, 1920; *Leptocimbex potanini sinobirmanica* Malaise, 1939). Russia: **FE** (PR). – China (SW), India (NE), SE Asia.

Leptocimbex terrificus Malaise, 1931. Russia: **FE** (PR).

PRAIA Wankowicz, 1880. Type species *Praia taczanowskii* Wankowicz, 1880. The larvae feed on the Betulaceae and Rosaceae. The genus is distributed in the Palaearctic region. Number of species: World, Palaearctic and Russia – 2.

Praia taczanowskii Wankowicz, 1880. Host plant: Betula spp. (Betulaceae). Russia: **EP** (NW, C, E), **UR**, **WS** (TK, NS, KM), **ES** (IR, BR, YA), **FE** (AM, PR, KA). – Europe (WE, EE, NE), Mongolia.

Praia ussuriensis Malaise, 1939. Host plant: Betula spp. (Betulaceae), Prunus spp. (Rosaceae). Russia: **ES** (Zhelochovtsev, Zinovjev, 1995), **FE** (PR, SA, KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

PSEUDOCLAVELLARIA Schulz, 1906. Type species *Tenthredo amerinae* Linnaeus, 1758. The larvae feed on the Salicaceae. The genus is distributed in Palaearctic. Number of species: World and Palaearctic – 3; Russia – 2.

Pseudoclavellaria amerinae (Linnaeus, 1758) [Tenthredo] (*Tenthredo marginata* Linnaeus, 1767; *T. 4-fasciata* Degeer, 1773; *T. rufa* Retzius, 1783; *Cimbex saliceti* Sasaki, 1902; *Clavellaria nitobei* Matsumura, 1912; *Trichosoma albopilosum* Matsumura, 1912; *Clavellaria amerinae delagrangei* Pic, 1920; *C. amerinae anatolica* Pic, 1920). Host plant: Salix spp., Populus spp. (Salicaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, MG). – Europe (WE, EE, NE, SE), Caucasus, Turkey, Mongolia, China (NE), Korean Peninsula.

Pseudoclavellaria semenovi (Gussakovskij, 1947) [Clavellaria]. Russia: **ES** (ZB), **FE** (PR).

TRICHIOSOMA Leach, 1817 (*Asitrichiosoma* Malaise, 1939; *Lucorumiformis* Ermolenko, 1972; *Vitelliniformis* Ermolenko, 1972; *Sericiformis* Ermolenko, 1972; *Sorbiformis* Ermolenko, 1972). Type species *Tenthredo lucorum* Linnaeus, 1758. The larvae feed on the deciduous of trees and shrubs. The genus is distributed in Holarctic and Oriental region. Number of species: World – 38; Palaearctic – 33; Russia – 24.

Trichiosoma aenescens Gussakovskij, 1947. Host plant: Prunus spp., Padus spp. (Rosaceae). Russia: **EP** (N, NW, C, E), **UR**, **WS** (NS, KM, AL), **ES** (KS, IR, YA), **FE** (AM, PR). – Europe (EE, NE), Japan (Hok).

Trichiosoma anthracinum Forsius, 1930. Russia: **WS** (AL). – China (SW).

Trichiosoma asachalinense Verzhutskii, 1973. Host plant: Spiraea spp., Dasiphora fruticosa (Rosaceae). Russia: **ES** (IR), **FE** (KA).

Trichiosoma bombiforme Takeuchi, 1939. Russia: **FE** (PR). – China (SE), Japan (Hon).

Trichiosoma crassum W.F. Kirby, 1882 (*T. arcticum* W.F. Kirby, 1882; *T. arctica* Konow, 1903; *T. jakovlevi* Konow, 1905; *T. jakovlevi tinctipennis* Enslin, 1919; *T. jakovlevi* Gussakovskij, 1947). Host plant: Betula spp. (Betulaceae). Russia: **EP** (N), **WS** (TM, AL), **ES** (KR, YA), **FE** (KH, PR, SA, KA, MG, CH). – Europe (NE), Mongolia, Japan (Hok, Hon), N America.

Trichiosoma hirtellum Gussakovskij, 1947. Host plant: Salix spp. (Salicaceae), Betula spp. (Betulaceae). Russia: **ES** (KR, IR, YA, ZB), **FE** (AM, KH, MG).

Trichiosoma latreillii Leach, 1817. Host plant: Salix spp. (Salicaceae). Russia: **EP** (N, NW, C, E), **UR**, **WS** (TK, NS, KM, AL), **ES** (KR, IR, BR, YA, ZB),

- FE** (AM, KH, PR, KA, MG). – Europe (WE, EE), Turkey, Mongolia, China (NE, NC), Japan (Hon).
- Trichiosoma lucorum** (Linnaeus, 1758) [Tenthredo] (*T. unidentatum* Leach, 1817; *Cimbex betuleti* Klug, 1834; *Trichiosoma salicis* Hartig, 1840). Host plant: Betula spp. (Betulaceae), Salix spp. (Salicaceae). Russia: **EP** (N, NW, C, E), **UR**, **WS** (TM, TK, NS, KM, AL), **ES** (KR, IR, YA), **FE** (SA). – Europe (WE, EE, NE, SE), Kazakhstan, Mongolia, China (NC), Japan (Hok, Hon).
- Trichiosoma malaisei** Saarinen, 1950. Host plant: Betula spp. (Betulaceae). Russia: **EP** (N), **FE** (MG). – Europe (NE).
- Trichiosoma melanopygum** Zhelochovtsev, 1968. Host plant: Betula spp. (Betulaceae). Russia: **ES** (IR, BR).
- Trichiosoma nanae** Vikberg et Viitasaari, 1991. Host plant: Betula nana (Betulaceae). Russia: **ES** (KR). – Europe (NE).
- Trichiosoma nigricoma** Konow, 1906. Host plant: Salix spp. (Salicaceae), Betula spp. (Betulaceae). Russia: **EP** (N, NW), **WS** (AL), **ES** (TU, KR). – Europe (NE).
- Trichiosoma opacum** Konow, 1906 (*T. jozankeanum* Matsumura, 1912). Host plant: Salix spp. (Salicaceae), Betula spp. (Betulaceae). Russia: **ES** (KR, IR, YA, ZB), **FE** (KH, PR, SA, MG). – Korean Peninsula, Japan (Hok, Hon).
- Trichiosoma pusillum** Stephens, 1835 (*T. biverrucatum* Stephens, 1835; *T. nigripes* Gussakovskij, 1947; *T. groenblomi* Saarinen, 1947). Host plant: Populus spp., Salix spp. (Salicaceae). Russia: **EP** (N), **WS** (NS, AL), **ES** (KR, IR, BR, ZB), **FE** (AM). – Europe (WE, NE).
- Trichiosoma sachalinense** Matsumura, 1911 (*T. bilineata* Malaise, 1931). Host plant: Salix spp. (Salicaceae). Russia: **EP** (N, NW), **WS** (NS, KM, AL), **ES** (IR, BR, YA, ZB), **FE** (KH, SA, KA, MG, CH). – Mongolia, Korean Peninsula, Japan (Hok).
- Trichiosoma scalesii** Leach, 1817 (*T. fennicum* Saarinen, 1947). Russia: **ES** (YA). – Europe (WE, NE).
- Trichiosoma sericeum** Konow, 1903. Host plant: Betula spp. (Betulaceae), Populus spp., Salix spp. (Salicaceae). Russia: **EP** (N, NW, E), **UR**, **WS** (TK, NS, KM, AL), **ES** (KS, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR). – Europe (NE), Mongolia, China (NE), Korean Peninsula.
- Trichiosoma sibiricum** Gussakovskij, 1947. Russia: **WS** (AL), **ES** (KR, IR, YA), **FE** (AM, KH, KA). – Europe (NE), China (NE).
- Trichiosoma sorbi** Hartig, 1840. Host plant: Sorbus spp. (Rosaceae). Russia: **EP** (N, NW, C, E), **UR**, **WS** (KM, AL), **ES** (YA). – Europe (WE, EE, NE, SE).
- Trichiosoma sylvaticum** Leach, 1817. Host plant: Populus spp., Salix spp. (Salicaceae). Russia: **EP** (N, NW, C, E), **WS** (TK, NS, KM, AL), **ES** (KS, BR, YA). – Europe (EE, SE).
- Trichiosoma tibiale** Stephens, 1835 (*Cimbex crataegi* Zaddach, 1863; *Tenthredo crataegi* Newman, 1870; *Trichiosoma oxyacanthae* Morley, 1907; *T. tibiale decipiens* Enslin, 1917). Host plant: Salix spp. (Salicaceae), Sorbus spp., Crataegus spp. (Rosaceae). Russia: **EP** (N, NW, C), **WS** (AL), **ES** (IR, YA), **FE** (KU, KA, MG, CH). – Europe (WE, EE, NE, SE), Japan (Hok, Hon).
- Trichiosoma ushinskii** Ermolenko, 1973. Russia: **FE** (KU).
- Trichiosoma villosum** (Motschulsky, 1860) [*Cimbex*] (*T. pubescens* Jakovlev, 1891; *T. relictum* Saarinen, 1950). Host plant: Populus spp., Salix spp. (Salicaceae). Russia: **EP** (NW, C), **WS** (NS, KM, AL), **ES** (KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, KA). – Europe (NE), Kazakhstan, China (NE).
- Trichiosoma vitellina** (Linnaeus, 1760) [Tenthredo] (*Cynips betulae* Strøm, 1762; *Trichiosoma marginale* Leach, 1817; *Cimbex trichiosoma* Lepeletier, 1823; *Trichiosoma boreale* Gussakovskij, 1947). Host plant: Alnus spp., Betula spp. (Betulaceae), Salix spp., Populus spp. (Salicaceae). Russia: **EP** (N, NW, C, E, S), **UR**, **WS** (TK, NS, KM, AL), **ES** (KS, KR, IR, BR, YA, ZB), **FE** (KH, PR). – Europe (WE, EE, NE, SE), Korean Peninsula.

Subfamily CORYNIDINAE

CORYNIS Thunberg, 1789 (*Amasis* Leach, 1817; *Amoeniformis* Ermolenko, 1972; *Lateralisiformis* Ermolenko, 1972; *Obscuriformis* Ermolenko, 1972). Type species *Tenthredo obscura* Fabricius, 1775. The larvae feed on the herbaceous plants. The genus is distributed in the Palaearctic region. Number of species: World and Palaearctic – 28; Russia – 8.

Corynis amoena (Klug, 1834) [*Cimbex*. Adults on the Ranunculus (Ranunculaceae). Russia: **EP** (N, NW, C, E, S, NC). – Europe (WE, EE, NE), Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia.

Corynis caucasica (Mocsáry, 1886) [*Amasis caspica* Konow, 1886]. Russia: **EP** (NC). – Europe (EE), Armenia, Azerbaijan, Turkey, Iran.

Corynis crassicornis (Rossi, 1790) [Tenthredo] (*Cimbex sylvatica* Olivier, 1791; *Tenthredo laeta* Fabricius, 1798; *T. ranunculi* Panzer, 1804; *Cimbex jurinae* Serville, 1823; *C. jurinae* Lepeletier, 1823; *C. olivieri* Lepeletier, 1823; *Amasis crassicornis nigriventris* Soldanski, 1916; *A. crassicornis algerica* Pic, 1916; *A. crassicornis gallica* Pic, 1918). Host plant: Sedum spp. (Crassulaceae), Ranunculus spp. (Ranunculaceae). Russia: **EP** (CR). – Europe (WE, EE, SE), N Africa, Georgia, Azerbaijan.

Corynis frontina (Konow, 1905) [Amasis]. Russia: **EP** (NC). – Europe (SE), Azerbaijan, Turkey.

Corynis lateralis (Brullé, 1832) [Amasis] (*Amasis lateralis sibirica* Kuznetzov-Ugamskij, 1925; *Corynis dominiquei* Chevin, 1988). Russia: **EP** (C, E, S,

NC, CR), **UR**. – Europe (EE, SE), Caucasus, Turkey, Kazakhstan.

Corynis mutabilis (Tischbein, 1852) [Abia] (*Amasis concinna* J.P.E.F. Stein, 1876; *A. bleyli* Muche, 1964). Adults on the Ranunculus (Ranunculaceae). Russia: **EP** (CR). – Europe (EE, SE), Armenia, Turkey.

Corynis obscura (Fabricius, 1775) [Tenthredo] (*Amasis helvetica* Konow, 1886; *A. obscura adusta* Zirngiebl, 1953; *A. valkanovi* Vassilev, 1969). Host plant: Geranium spp. (Geraniaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TK, NS, KM), **ES** (IR). – Europe (WE, EE, NE, SE), Caucasus, Turkey, Iran.

Corynis ovicula Zhelochovtsev, 1968. Russia: **ES** (BR, ZB).

5. FAMILY DIPRIONIDAE

Stout, slow-flying insects associated with conifers. They are characterized by plumose male and serrate female antennae of more than 9 segments and by absence of vein $2r$ on the fore wing; scutellum without a transverse furrow cutting off a post-tergite; hind wing with both middle closed cells RS and M present; tibiae without preapical spines and front tibiae with simple unmodified spurs.

The larvae are free-living, often gregariously on needles of conifers. They have 8 pairs of abdominal legs (segments 2-8 and 10). The cocoons are double, and upon emerging the adult severs a cap at one end of the cocoon and this cap is left hanging by a few threads.

The family is distributed in the Holarctic, Oriental and Neotropical regions. Number of taxa: World – 2 subfamilies, 11 genera, about 140 species; Palaearctic – 2/9/about 70; Russia – 2/7 (+ 1 unplaced)/26 (+ 2 unplaced).

References. Fischer von Waldheim, 1806; Gussakovskij, 1947; Verzhutskii, 1974, 1981; Ermolenko, 1975; Zhelochovtsev, Prochorova, 1976; Ermolenko, Zemkova, 1979; Viitasaari, Varama, 1987; Zhelochovtsev, 1988; Liston, 1995; Zhelochovtsev, Zinovjev, 1995; Liston et al., 2006, 2014a; Taeger et al., 2006, 2010; Wei et al., 2006; Roller, Haris, 2008; Blank et al., 2009; Gnienko, Seryi, 2009; Humala, Polevoi, 2009; Schedl, 2009; Taeger, Blank, 2011; Vasilenko, 2011b; Hara, Smith, 2012; Macek, 2012; Sundukov, Lelej, 2012; Vasilenko, Korshunov, 2012; Kostyunin, 2013, 2015; Popov, 2013; Beneš, 2014b; Borowski, Marczak, 2015; Hara, Shinohara, 2015; Sundukov, 2015, 2017; Noblecourt, 2016.

Subfamily DIPRIONINAE

DIPRION Schrank, 1802 (*Pteronus* Jurine, 1801; *Lophyrus* Latreille, 1803; *Pteronus* Panzer, 1806; *Anchoreta* Gistel, 1848; *Cristiger* Gistel, 1848). Type species *Tenthredo pini* Linnaeus, 1758. The larvae feed on the conifers. The genus is distributed in the Palaearctic and Oriental regions; one species was introduced into the Nearctic region. Number of species: World – 13; Palaearctic – 9; Russia – 3.

Diprion koreanus Takagi, 1931. Host plant: *Larix* spp. (Pinaceae); live in groups. Russia: **UR**, **ES** (IR, YA, ZB), **FE** (AM, KH). – Korean Peninsula.

Diprion pini (Linnaeus, 1758) [Tenthredo] (*Tenthredo dorsata* Fabricius, 1781; *T. eques* Schrank, 1782;

T. pineti Bechstein et Scharfenberg, 1805; *Lophyrus pini nigripectus* Matsumura, 1912; *Diprion pini nigroscutellatum* Enslin, 1916; *D. pini nigristernis* Enslin, 1916; *D. pini klugi* Enslin, 1916; *D. butovitschi* Hedqvist, 1967). Host plant: *Pinus* spp. (Pinaceae); live in groups, 2 generation per year. Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TK, NS, KM), **ES** (IR, YA). – Europe (WE, EE, NE, SE), N Africa, Georgia, Cyprus, Turkey.

Diprion similis (Hartig, 1836) [*Lophyrus*] (*Lophyrus eremita* Thomson, 1871; *Diprion simile claristernis* Enslin, 1916; *D. simile atroscutellatum* Enslin, 1916). Host plant: *Pinus* spp. (Pinaceae); live alone, 2 generation per year. Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TM, AL), **ES** (IR, YA). – Europe (WE, EE, NE, SE); introduced into N America.

GILPINIA Benson, 1939. Type species *Lophyrus polytomus* Hartig, 1834. Larvae feed on conifers. The genus is distributed in the Palaearctic and Oriental regions; two species were introduced to the Nearctic region. Number of species: World – about 40; Palaearctic – > 30; Russia – 15.

Gilpinia abieticola (Dalla Torre, 1894) [*Lophyrus*] (*Lophyrus abietis* Stein, 1866). Host plant: *Picea* spp. (Pinaceae). Russia: **EP** (C), **WS** (TM), **ES** (IR, YA). – Europe (WE, EE, NE, SE).

Gilpinia catocala (Snellen van Vollenhoven, 1858) [*Lophyrus*] (*Gilpinia verticalis* Gussakovskij, 1947). Host plant: *Pinus* spp. (Pinaceae). Russia: **EP** (NW, C, E), **WS** (TK, NS, KM). – Europe (EE, NE).

Gilpinia daisetusana Takeuchi, 1940. Host plant: *Picea* spp. (Pinaceae). Russia: **FE** (KU). – Japan (Hok).

Gilpinia fennica (Forsius, 1911) [Diprion]. Host plant: *Picea* spp. (Pinaceae). Russia: **ES** (TU, YA). – Europe (EE, NE), China (NC).

Gilpinia frutetorum (Fabricius, 1793) [Tenthredo] (*Lophyrus anachoreta* Costa, 1894; *Diprion frutetorum lutescens* Enslin, 1916; *D. frutetorum luteolum* Enslin, 1916). Host plant: *Pinus* spp. (Pinaceae); live alone, 2 generation per year. Russia: **EP** (NW, C, E, S, NC), **UR**, **WS** (TK, NS, AL), **ES** (IR, YA). – Europe (WE, EE, NE, SE), Turkey; introduced into N America.

Gilpinia hercyniae (Hartig, 1837) [*Lophyrus*]. Host plant: *Picea* spp. (Pinaceae). Russia: **EP** (NW, C, NC), **UR**, **WS** (TM), **ES** (KR, IR, BR, YA), **FE** (PR). – Europe (WE, EE, NE, SE), Pakistan, Mongolia, Korean Peninsula, Japan (Hok, Hon); introduced into N America.

Gilpinia hokkaidoensis Hara et Shinozaki, 2015. Host plant: *Picea* spp., *Abies* spp. (Pinaceae). Russia: **FE** (KU). – Japan (Hok).

Gilpinia laricis (Jurine, 1807) [Pteronus] (*Diprion laricis fenestratum* Enslin, 1916; *D. laricis flavomarginatum* Enslin, 1916; *D. laricis lugens* Enslin, 1916). Host plant: *Pinus* spp. (Pinaceae); live alone, 1-2 generation per year. Russia: **EP** (NW, C, E), **WS** (TM, TK, KM, AL), **ES** (IR, YA). – Europe (WE, EE, NE, SE).

Gilpinia pallida (Klug, 1812) [Lophyrus] (*Lophyrus minor* Serville, 1823; *L. minor* Lepeletier, 1823). Host plant: *Pinus* spp. (Pinaceae); live in groups, 2 generation per year. Russia: **EP** (NW, C, S), **WS** (TK), **ES** (IR, YA). – Europe (WE, EE, NE, SE).

Gilpinia polytoma (Hartig, 1834) [Lophyrus] (*Diprion polytomum pseudopallidum* Enslin, 1916). Host plant: *Picea* spp., *Abies* spp., *Larix* spp. (Pinaceae); live alone, 1-2 generation per year. Russia: **EP** (NW, C), **UR**, **WS** (TM, TK, NS), **ES** (IR, BR, YA), **FE** (KH, PR, KU). – Europe (WE, EE, NE, SE), Korean Peninsula, Japan (Hok, Hon), Pakistan, India.

Gilpinia sachalinensis Takeuchi, 1940. Russia: **FE** (SA).

Gilpinia socia (Klug, 1812) [Lophyrus] (*Lophyrus pineti* Hartig, 1837). Host plant: *Pinus* spp. (Pinaceae); live in groups, 2 generation per year. Russia: **EP** (NW), **ES** (KR). – Europe (WE, EE, NE, SE).

Gilpinia tohi Takeuchi, 1940. Host plant: *Picea* spp., *Abies* spp. (Pinaceae). Russia: **FE** (PR, KU). – China (NE), Japan (Hok).

Gilpinia variegata (Hartig, 1834) [Lophyrus] (*Lophyrus thomsoni* Konow, 1884). Host plant: *Pinus* spp. (Pinaceae); live alone, 2 generation per year. Russia: **EP** (NW, C), **WS** (TK, NS, AL), **ES** (KR, IR). – Europe (WE, EE, NE, SE), Kazakhstan.

Gilpinia virens (Klug, 1812) [Lophyrus] (*Diprion virens atramentarium* Enslin, 1916). Host plant: *Pinus* spp. (Pinaceae); live alone, 1-2 generation per year. Russia: **EP** (NW, C), **WS** (TM, NS, AL), **ES** (IR, YA). – Europe (WE, EE, NE, SE), Mongolia.

MACRODIPRION Enslin, 1914. Type species *Lophyrus nemoralis* Enslin, 1917. The larvae feed on the *Pinus* spp. (Pinaceae). The genus is distributed in Palaearctic and Oriental region. Number of species: World – 2; Palaearctic – 1; Russia – 1.

Macrodrion nemoralis (Enslin, 1917) [Lophyrus] (*Tenthredo nemorum* Fabricius, 1793). Host plant: *Pinus* spp. (Pinaceae); live alone, 1-2 generation per

year. Russia: **EP** (N, NW, C), **WS** (NS), **ES** (KR, IR, BR). – Europe (WE, EE, NE, SE).

MICRODIPRION Enslin, 1914. Type species *Hylotoma pallipes* Fallén, 1808. The larvae on the *Pinus* spp. (Pinaceae). The genus is distributed in Palaearctic and Oriental region. Number of species: World – 5; Palaearctic – 3; Russia – 2.

Microdiprion fuscipennis (Forsius, 1911) [Diprion]. Host plant: *Picea* spp. (Pinaceae); live in groups in the younger ages, singly in the senior, 2 generation per year. Russia: **EP** (C, E), **ES** (KR). – Europe (EE, NE).

Microdiprion pallipes (Fallén, 1808) [Hylotoma] (*Lophyrus elongatulus* Klug, 1812; *L. politus* Klug, 1812; *L. pulchricornis* Bremer-Wolf, 1849; *L. pallidipes* Dalla Torre, 1894; *Diprion pallipes aterrimum* Enslin, 1916). Host plant: *Pinus* spp. (Pinaceae); live in groups, 1-2 generation per year. Russia: **EP** (N, NW, C, NC), **WS** (TK), **ES** (KR, IR, BR, YA), **FE** (AM, PR, KU). – Europe (WE, EE, NE, SE), Kazakhstan.

NEODIPRION Rohwer, 1918. Type species *Lophyrus lecontei* Fitch, 1858. The larvae feed on the *Pinus* spp. (Pinaceae). The genus is distributed in Holarctic, Oriental and Neotropical regions. Number of species: World – > 45; Palaearctic – 2; Russia – 1.

Neodiprion sertifer (Geoffroy, 1785) [Tenthredo] (*Tenthredo pectinata rufa* Retzius, 1783; *T. pinirufa* Villers, 1789; *Hylotoma rufa* Fallén, 1808; *Lophyrus piceae* Lepeletier, 1823; *L. piceae* Serville, 1823; *L. basalis* Matsumura, 1915). Host plant: *Pinus* spp. (Pinaceae); live in groups, often severely pest. Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TK, NS, KM, AL), **ES** (IR), **FE** (AM). – Europe (WE, EE, NE, SE), Caucasus, Turkey, Kazakhstan, Korean Peninsula, Japan (Hok, Hon, Kyu), N America.

NESODIPRION Rohwer, 1910. Type species *Lophyrus japonicus* Marlatt, 1898. The larvae feed on the *Pinus* (Pinaceae). The genus is distributed in East Palaearctic and Oriental region. Number of species: World – 13; Palaearctic – 11; Russia – 1.

Nesodiprion japonicus (Marlatt, 1898) [Lophyrus]. Host plant: *Pinus* spp., *Larix* spp., *Cedrus* spp. (Pinaceae). Russia: **FE** (PR). – China (SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu), N America (California).

Subfamily MONOCTENINAE

MONOCTENUS Dahlbom, 1835. Type species *Tenthredo juniperi* Linnaeus, 1758. Larvae feed on the Juniperus (Cupressaceae). The genus occurs in the Holarctic and Neotropical regions. Number of species: World – 12; Palaearctic – 8; Russia – 3.

Monocetus juniperi (Linnaeus, 1758) [Tenthredo (*Tenthredo pterophorus* Sulzer, 1761; *Monocetus andrei* Mocsáry, 1880)]. Host plant: Juniperus spp. (Cupressaceae). Russia: **EP** (NW, C, E), **UR**. – Europe (WE, EE, NE, SE), Turkey, Syria.

Monocetus nipponicus Takeuchi, 1940. Host plant: Juniperus spp. (Cupressaceae). Russia: **FE** (PR). – Japan (Hon).

Monocetus obscuratus (Hartig, 1837) [Lophyrus (*M. obscuratus intermedius* Zombori, 1975)]. Host

plant: Juniperus spp. (Cupressaceae). Russia: **WS** (AL), **ES** (YA). – Europe (WE, EE, NE, SE).

Unplaced Diprionidae

NYCTERIDIUM Fischer von Waldheim, 1806. Type species *Nycteridium fischeri* Fischer von Waldheim, 1806. The genus with incomprehensible systematic status (see Blank et al., 2009). The genus distribution in Palaearctic. Number of species: World and Palaearctic – 2; Russia – 2.

Nycteridium fischeri Fischer von Waldheim, 1806. Russia: without region (Fischer von Waldheim, 1806).

Nycteridium mosquense Fischer von Waldheim, 1806. Russia: **EP** (C).

6. FAMILY HEPTAMELIDAE

Have previously been classified into the subfamily Selandriinae within Tenthredinidae. Now treated as a separate family, within the “higher” Tenthredinoidea (Tenthredinidae, Cimbicidae, and Diprionidae), thus not forcing the inclusion of the well defined families Diprionidae and Cimbicidae into Tenthredinidae (Liston et al., 2014a; Malm, Nyman, 2015; Schmidt et al., 2016; Schneider, 2016). All known larvae feed on ferns. No reports exist of economic damage caused by Heptamelidae.

The family is distributed in the Holarctic and Oriental regions. Number of taxa: World – 6 genera, about 50 species; Palaearctic – 2/13; Russia – 2/5.

R e f e r e n c e s. Verzhutskii, 1974; Zhelochovtsev, 1988; Zhelochovtsev, Zinovjev, 1992, 1996; Vikberg, Liston, 2009; Sundukov, Lelej, 2012; Liston et al., 2014a; Malm, Nyman, 2015; Schmidt et al., 2016; Schneider, 2016; Vikberg, 2017.

HEPTAMELUS Haliday, 1855 (*Melicerta* Stephens, 1835; *Caenoneura* Thomson, 1870). Type species *Melicerta ochroleuca* Stephens, 1835. The larvae feed on the ferns. The genus is distributed in Palaearctic and Oriental region; one species was

introduced into N America. Number of species: World – 37; Palaearctic – 11; Russia – 4.

Heptamelus dahlbomi (Thomson, 1870) [Caenoneura]. Host plant: *Athyrium filix-femina* (Dryopteridaceae). Russia: **EP** (N). – Europe (WE, NE); introduced into N America.

Heptamelus magnocularis Malaise, 1931. Russia: **FE** (SA, KA).

Heptamelus ochroleucus (Stephens, 1835) [Meli certa]. Host plant: Dryopteridaceae, Blechnaceae, Onocleaceae, Polypodiaceae. Russia: **EP** (NW, C). – Europe (WE, EE, NE), Japan (Hon).

Heptamelus ussuriensis Malaise, 1931. Russia: **FE** (AM, PR).

PSEUDOHEPTAMELUS Conde, 1932. Type species *Pseudoheptamelus runari* Conde, 1932. The larvae feed on the ferns. The genus is distributed in Palaearctic. Number of species: World and Palaearctic – 2; Russia – 1.

Pseudoheptamelus runari Conde, 1932. Host plant: *Athyrium* spp. (Dryopteridaceae); larva in the ferns stem. Russia: **EP** (N, NW, C), **ES** (IR), **FE** (PR). – Europe (NE).

7. FAMILY TENTHREDINIDAE

Adults of Tenthredinidae are distinguished from those of other families by the presence of 7-10 flagellomeres, by the metapleuron clearly separated from the tergum 1, and by the long and clearly outlined (at least laterally) mesoscutellar appendage. Adults are herbivorous or do not feed, few predators. The larvae of most species are external feeders on foliage, but those of some species are leaf miners, gall formers, or shoot borers; usually oligophages, less often monophages or polyphages. Host plants are horsetails, ferns, conifers or angiosperms (especially many species on willows, poplars, birches and rosaceous). It usually hibernates a prepupa, sometimes a pupa, less often eggs. Part of the species are pests of fruit and berry and forest plants.

The family is distributed worldwide. Number of taxa: World – 6 subfamilies, about 400 genera, > 5600 species; Palaearctic – 6/about 220/about 3500; Russia – 6/109/1124 (+ 29 ssp.).

R e f e r e n c e s. Hartig, 1837; Eversmann, 1847; Erichson, 1851; Zaddach, 1859; Kawall, 1864; Kirby, 1882; Brischke, 1883, 1884; Holmgren, 1883; Konow, 1886, 1891, 1895, 1897, 1900, 1902a,c, 1904, 1907, 1908; Jakovlev, 1888, 1891, 1900; Dalla Torre, 1894; Mocsáry, 1909; Rohwer, 1910, 1925; Matsumura, 1911; Enslin, 1915, 1919, 1920, 1927; Dovnar-Zapolskij, 1925, 1929, 1930, 1931b; Kokujev, 1927; Kuznetsov-Ugamskij, 1927, 1929; Forsius, 1928, 1929; Zhelochovtsev, 1928, 1935b, 1939, 1941, 1951, 1952, 1968, 1976, 1988; Malaise, 1931a,b, 1945, 1947, 1963, 1964; Takeuchi, 1931, 1933, 1936, 1937, 1938, 1939, 1940, 1941, 1955, 1956; Conde, 1935; Hellén, 1948, 1956; Saarinen, 1950a; Ermolenko, 1955, 1959, 1968, 1971a, 1975, 1981, 1984, 1992; Stroganova, 1960, 1973, 1978b, 1980, 1988; Benson, 1962, 1965b; Muche, 1965a,b, 1968a,b, 1969, 1971a,b, 1973a,b,c,d, 1975a; Kuwayama, 1967; Lindqvist, 1968, 1970, 1971, 1972, 1975, 1977; Inomata, 1970; Naito, 1971a,b, 1980, 1990, 2004; Okutani, 1972; Verzhutskii, 1973, 1974, 1981; Zombori, 1973, 1974, 1978; Schtundjuk, Zelochovtsev, 1974; Tomilova, 1974; Smith, 1975, 1981; Zhelochovtsev, Prochorova, 1976; Bogacheva, 1977; Zinovjev, 1978, 1981, 1985, 1986, 1993a,b,c,d, 1994, 2000, 2010; Abe, 1981; Seiyama, 1981; Zhelochovtsev, Zinovjev, 1981, 1992, 1994, 1995, 1996; Koch, 1982, 1988b,c, 1989; Taeger, 1985, 1986, 1988, 1989, 1991, 1992, 2005, 2013; Goulet, 1986; Kajmuk, 1986, 1988; Altenhofer, Zombori, 1987; Zerova et al., 1988; Viitasaari, Zinovjev, 1991; Blank, 1993, 1996, 2002a; Liston, 1994, 1995, 2012; Shinohara, 1994a,b, 1997c, 2002a, 2005b, 2015; Zinovjev,

Schmidt, 1994; Pesarini et al., 1995; Haris, Blank, 1996; Lee, Ryu, 1996; Schmidt, 1997; Wei, 1997a,b, 1998a,b, 2006; Wei, Nie, 1997b, 2008; Andreeva, 1998; Blank, Ritzau, 1998; Blank, Taeger, 1998a, 2006; Lacourt, 1999; Zinovjev, Vikberg, 1999; Haris, 2000, 2001, 2002, 2003, 2006a,b, 2009; Lee et al., 2000; Shinohara et al., 2000, 2000, 2013, 2015; Togashi, 2000; Nie, Wei, 2001, 2004a,b; Altenhofer, Pschorn-Walcher, 2003; Kajmuk, Popov, 2003; Magis, 2003; Vikberg, 2003, 2009, 2010, 2013; Heidemaa et al., 2004, 2012; Heidemaa, Viitasaari, 2004; Heidemaa, Zinovjev, 2004; Vasilenko, 2004, 2006, 2008, 2009b, 2010a,b, 2011a; Astapenko et al., 2006; Averenskiy et al., 2006; Heidemaa, Prous, 2006; Liston et al., 2006, 2014a, 2017; Saini et al., 2006; Shilenkov, Tolstonogova, 2006; Taeger et al., 2006, 2010, 2016, 2017; Vikberg, Zinovjev, 2006, 2014; Wei et al., 2006; Haris, Zsolnai, 2007; Kopelke, 2007a,b,c; Lelej, Taeger, 2007b; Stepanov et al., 2007; Zabroda, Sirenko, 2007, 2008; Popov, Kajmuk, 2008, 2009a,b, 2010b; Roller, Haris, 2008; Shcherbakov, 2008; Arutyunyan et al., 2009; Blank et al., 2009; Schedl, 2009; Sundukov, Lelej, 2009, 2012; Macek, 2010, 2012; Sundukov, 2010, 2011b, 2013, 2017; Wei, Niu, 2010; Yegorenkova et al., 2010; Astapenko, Pal'nikova, 2011; Belov, 2011; Li et al., 2011, 2014, 2016; Popov, 2011a,b, 2016; Prous et al., 2011, 2011, 2014, 2016, 2017; Smith, Altenhofer, 2011a,b; Taeger, Blank, 2011; Nie et al., 2012; Niu et al., 2012; Ahn et al., 2013; Niu, Wei, 2013; Petrov, Sautkin, 2013; Beneš, 2014a,b,c, 2015; Heibo et al., 2014; Setrakova, 2014; Beneš, Holuša, 2015; Liu et al., 2015; Qi et al., 2015; Shinohara, Hara, 2015; Shinohara, Yoshida, 2015; Smith, Wei, 2015; Taeger, Viitasaari, 2015; Noblecourt, 2016; Smith, Gourlay, 2016; Wu et al., 2016; Hara, Shinohara, 2017; Taeger, Kramp, 2017.

Subfamily ALLANTINAE

ALLANTUS Panzer, 1801 (*Emphytus* Klug, 1815; *Anemphytus* Dovnar-Zapolskij, 1931; *Synemphytus* Malaise, 1945). **T y p e s p e c i e s** *Tenthredo togata* Panzer, 1801. The larvae feed on the deciduous trees and shrubs; when pupation is carried into dry branches and bark. The genus is distributed in Holarctic and Oriental region. Number of species: World – > 45; Palaearctic – about 35; Russia – 17 from 2 subgenera.

Allantus (Allantus) calliblepharus (Konow, 1900) [*Emphytus*] (*Emphytus karaefutonis* Matsumura, 1911). Host plant: *Salix* spp. (Salicaceae). Russia: WS

- (KM, AL), **ES** (IR, YA), **FE** (KH, PR, SA, KU). – Europe (NE), Korean Peninsula, Japan (Hok, Hon).
- Allantus (Allantus) luctifer luctifer** (F. Smith, 1874) [Macrophyta]. Host plant: *Salix* spp. (Salicaceae), *Rumex* spp., *Polygonum* spp. (Polygonaceae). Russia: **FE** (PR, SA). – China (CC, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Allantus (Allantus) togatus** (Panzer, 1801) [Tenthredo] (*Tenthredo ferruginosa* Gmelin, 1790; *T. succincta* Klug, 1818; *Emphytus succinctus steini* Schmiedeknecht, 1881; *E. leucostomus* Costa, 1890; *E. succinctus ruficornis* Costa, 1890). Host plant: *Salix* spp. (Salicaceae), *Betula* spp. (Betulaceae), *Quercus* spp. (Fagaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (NS), **ES** (YA), **FE** (SA). – Europe (WE, EE, NE, SE), Turkey.
- Allantus (Allantus) viennensis** (Schrank, 1781) [Tenthredo] (*Emphytus viennensis nigricoxis* De-Stefani, 1883; *E. viennensis medinae* Konow, 1894; *Allantus viennensis uberior* Enslin, 1914; *Emphytus viennensis atricornis* Pic, 1948). Host plant: *Rosa* spp. (Rosaceae). Russia: **EP** (S, NC). – Europe (WE, EE, SE), Georgia, Armenia, Turkey, Central Asia, N America.
- Allantus (Emphytus) albocinctus** (Matsumura, 1912) [Dolerus]. Host plant: *Filipendula* spp., *Rubus* spp., *Rosa* spp. (Rosaceae). Russia: **FE** (SA, KU). – China (NC), Japan (Hok).
- Allantus (Emphytus) basalis basalis** (Klug, 1818) [Tenthredo] (*Emphytus basalis masculus* Zirngiebl, 1937). Host plant: *Rosa* spp., *Dasiphora* spp., *Potentilla* spp. (Rosaceae), *Betula* spp. (Betulaceae). Russia: **EP** (N, NW, C, E, S), **UR**, **WS** (NS), **ES** (IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KA, MG). – Europe (WE, EE, NE, SE), Turkey, Mongolia, China (NE, NC), Korean Peninsula, Japan (Hok, Hon), N America.
- Allantus (Emphytus) calceatus** (Klug, 1818) [Tenthredo] (*Dolerus vicinus* Lepeletier, 1823; *D. vicinus* Serville, 1823; *Tenthredo vitis* Vallot, 1845; *Emphytus dissimilis* Dietrich, 1868). Host plant: *Filipendula* spp., *Spiraea* spp., *Rubus* spp. and others Rosaceae. Russia: **EP** (N, NW, C, S, E, NC), **UR**, **WS** (Zhelochovtsev, Zinovjev, 1996), **FE** (AM, PR, YA). – Europe (WE, EE, NE, SE), Georgia, Kazakhstan, Mongolia, China (Wei et al., 2006).
- Allantus (Emphytus) cinctus** (Linnaeus, 1758) [Tenthredo] (*Tenthredo cordigera* Geoffroy, 1785; *Dolerus cinctus* Jurine, 1807; *D. varipes* Lepeletier, 1823; *D. cingulatus* Serville, 1823; *D. varipes* Serville, 1823; *Emphytus neglectus* Zaddach, 1859; *E. cinctipes* Norton, 1867; *E. infasciatus* Pic, 1948). Host plant: *Fragaria* spp., *Rosa* spp., *Rubus* spp. (Rosaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **WS** (NS), **ES** (IR, BR, YA, ZB), **FE** (AM, KH, PR, KA, MG). – Europe (WE, EE, NE, SE), Armenia, Turkey, Iran, Kazakhstan, Mongolia, China (Wei et al., 2006), N America.
- Allantus (Emphytus) cingillipes** (Kontuniemi, 1947) [Emphytus]. Russia: **FE** (SA). – Europe (NE).
- Allantus (Emphytus) cingillum** (Klug, 1818) [Tenthredo]. Host plant: *Betula* spp. (Betulaceae), ?*Rosa* spp. (Rosaceae). Russia: **EP** (NW, C, E), **ES** (KR, IR, BR, YA). – Europe (WE, EE, NE).
- Allantus (Emphytus) cingulatus** (Scopoli, 1763) [Tenthredo] (*Dolerus cingulatus* Lepeletier, 1823; *Tenthredo tenius* Lepeletier, 1823; *Emphytus elegans* Costa, 1859; *Allantus cingulatus muliebris* Enslin, 1914). Host plant: *Fragaria* spp., *Rosa* spp. (Rosaceae). Russia: **EP** (NW, NC), **ES** (TU), **FE** (KA). – Europe (WE, EE, NE, SE), Georgia, Armenia, Turkey.
- Allantus (Emphytus) didymus didymus** (Klug, 1818) [Tenthredo]. Host plant: *Sanguisorba* spp., *Rosa* spp. (Rosaceae). Russia: **EP** (C, E, S, NC), **WS** (NS). – Europe (WE, EE, SE), N Africa, Caucasus, SW Asia, Iran, Kazakhstan.
- Allantus (Emphytus) melanarius** (Klug, 1818) [Tenthredo] (*Emphytus tricoloripes* Costa, 1859). Host plant: *Cornus sanguinea* (Cornaceae). Russia: **EP** (NC). – Europe (WE, EE, SE), Iran.
- Allantus (Emphytus) ruficeps** (Konow, 1899) [Emphytus]. **EP** (NC). – Georgia.
- Allantus (Emphytus) rufocinctus** (Retzius, 1783) [Tenthredo] (*Tenthredo zonata* Christ, 1791; *T. rufocincta* Klug, 1818; *T. cingulum* Klug, 1818; *T. agilis* Drapiez, 1820; *Dolerus fasciatus* Lepeletier, 1823; *D. fasciatus* Serville, 1823; *Tenthredo ruralis* Dahlbom, 1835; *Emphytus rufocinctus ruralis* Thomson, 1871). Host plant: *Rosa* spp., *Rubus* spp. (Rosaceae). Russia: **EP** (N, NW, C). – Europe (WE, EE, SE), Mongolia.
- Allantus (Emphytus) truncatoides** Koch, 1988. Russia: **ES** (IR).
- Allantus (Emphytus) truncatus** (Klug, 1818) [Tenthredo] (*Tenthredo fulvipes* Fallén, 1829; *T. schoenherri* Dahlbom, 1835; *T. angustata* Zetterstedt, 1838). Host plant: *Filipendula* spp., *Fragaria* spp., *Potentilla* spp. and others (Rosaceae). Russia: **EP** (NW, C, E, NC), **UR**, **WS** (NS), **ES** (YA), **FE** (KA). – Europe (WE, EE, NE), Kyrgyzstan, Kazakhstan, Mongolia, China (NC, NW).

AMETASTEGIA Costa, 1882 (*Aomodyctium* Ashmead, 1898; *Protemphytus* Rohwer, 1909; *Emphytina* Rohwer, 1911; *Simplemphytus* MacGillivray, 1914; *Unitaxonus* MacGillivray, 1921; *Ocla* Malaise, 1957). Type species *Ametastegia fulvipes* Costa, 1882. The larvae before pupation borer branches, stems, bark or fruit. The genus distribution in Holarctic, Oriental and Neotropical regions. Number of species: World – about 60; Palaearctic - > 30; Russia – 13 from 2 subgenera.

Ametastegia (Ametastegia) albipes (Thomson, 1871) [Taxonus] (*Taxonus fletcheri* Cameron, 1878; *Eriocampa alabastripes* Costa, 1890). Russia: **EP** (N, NW, C, E, NC), **FE** (KA). – Europe (WE, EE, NE, SE).

Ametastegia (Ametastegia) equiseti (Fallén, 1808) [Tenthredo] (*Tenthredo bicolor* Klug, 1817; *T. gracilis* Serville, 1823; *T. gracilis* Lepeletier, 1823; *T. fuscitarsis* Hummel, 1827; *T. coxalis* Hartig, 1837; *T. bizonata* Zetterstedt, 1838; *Taxonus minutus* Costa, 1859; *Tenthredo angustula* Eversmann, 1864; *Taxonus innominatus* MacGillivray, 1901; *Unitaxonus repentinus* MacGillivray, 1921; *U. rumicis* MacGillivray, 1921; *Strongylogastroidea depressata* MacGillivray, 1921; *S. rufocinctella* MacGillivray, 1923; *Hemitaxonus dediticius* MacGillivray, 1923; *Ametastegia equiseti stitia* Zirngiebel, 1954). Host plant: *Polygonum* spp., *Rumex* spp. (Polygonaceae), *Chenopodium* (Amaranthaceae), etc. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **ES** (IR, BR, YA). – Europe (WE, EE, NE, SE), Georgia, Armenia, Turkey, Mongolia, China (Wei et al., 2006), N America.

Ametastegia (Ametastegia) glabrata (Fallén, 1808) [Tenthredo] (*Tenthredo violacea* Christ, 1791; *T. agilis* Klug, 1817; *T. rufipes* Lepeletier, 1823; *T. rufipes* Serville, 1823; *T. nigrisoma* Norton, 1862; *Ametastegia fulvipes* Costa, 1882; *Strongylogaster abnormis* Provancher, 1885; *Strongylogastroidea potulenta* MacGillivray, 1923). Host plant: *Polygonum* spp., *Rumex* spp. (Polygonaceae), *Chenopodium* (Amaranthaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **WS** (KM). – Europe (WE, EE, NE, SE), N Africa, SW Asia, Kazakhstan, Central Asia, Mongolia, China (NW), N America, S America, Australia.

Ametastegia (Ametastegia) kirishimensis Takeuchi, 1929. Russia: **FE** (PR, SA). – Japan (Kyu).

Ametastegia (Ametastegia) wuorentausi Forsius, 1928 (*A. montana* Malaise, 1931). Russia: **FE** (KA). – China (NE).

Ametastegia (Protemphytus) albovaria (Takeuchi, 1936) [Emphytina]. Russia: **FE** (SA, KU). – Japan (Hok, Hon).

Ametastegia (Protemphytus) carpini (Hartig, 1837) [Emphytus] (*Taxonus glottianus* Cameron, 1874; *Emphytus arcticus* Nerén, 1892; *Allantus carpini decipiens* Enslin, 1914; *Emphytus perla obscura* Zirngiebel, 1954). Host plant: *Geranium* spp. (Geraniaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (TU, BR), **FE** (AM, PR, KA). – Europe (WE, EE, NE, SE), Armenia, Turkey, Central Asia, Kazakhstan.

Ametastegia (Protemphytus) pallipes (Spinola, 1808) [Tenthredo] (*Tenthredo grossulariae* Klug, 1818; *Dolerus leucopodus* Lepeletier, 1823; *D. leucopodus* Serville, 1823; *Tenthredo lapponica* Zetterstedt, 1838; *Emphytus pallipes* Provancher, 1878; *E. canadensis* W.F. Kirby, 1882; *E. pallidipes* Dalla Torre, 1894; *Empria cavata* MacGillivray, 1911; *E. cetaria* MacGillivray, 1921; *Emphytus hyacinthus* MacGillivray, 1923; *E. hospitus* MacGillivray, 1923; *E. halesus* MacGillivray, 1923; *E. heroicus* MacGillivray, 1923; *E. hiatus* MacGillivray, 1923). Host plant: *Viola* spp. (Violaceae), *Vicia* spp. (Fabaceae). Russia: **EP** (NW, C, E, NC, CR), **UR**, **ES** (TU), **FE** (KU, KA). – Europe (WE, EE, NE, SE), Caucasus, Turkey, Iran, N America.

Ametastegia (Protemphytus) perla (Klug, 1818) [Tenthredo] (*Tenthredo bohemanni* Dahlbom, 1835). Host plant: *Salix* spp., *Populus* spp. (Salicaceae), *Quercus* spp. (Fagaceae), *Polygonum* spp. (Polygonaceae). Russia: **EP** (NW, C), **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (IR), **FE** (KA). – Europe (WE, EE, NE).

Ametastegia (Protemphytus) tenera (Fallén, 1808) [Tenthredo] (*Tenthredo patellata* Klug, 1818; *Doleres luctuosus* Lepeletier, 1823; *D. nigritus* Lepeletier, 1823; *D. luctuosus* Serville, 1823; *D. nigritus* Serville, 1823; *Tenthredo trunculi* Vallot, 1845; *Simplemphytus pacificus* MacGillivray, 1914; *Emphytina vanduzeei* Rohwer, 1915; *Emphytus haustus* MacGillivray, 1923; *E. haliartus* MacGillivray, 1923; *Empria columna* MacGillivray, 1923). Host plant: *Acetosa* spp., *Rumex* spp. (Polygonaceae), *Filipendula* spp. (Rosaceae), *Cirsium* spp. (Asteraceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (IR, BR), **FE** (AM, KH, MG). – Europe (WE, EE, NE, SE), Armenia, Turkey, Iran, Kyrgyzstan, Kazakhstan, Mongolia, China (NW), N America.

Ametastegia geranii (Takeuchi, 1929) [Emphytina].
Host plant: *Geranium* spp. (Geraniaceae). Russia: **FE** (KU). – Japan (Hon, Shi).

Ametastegia longicornis (Takeuchi, 1929) [Emphytina]. Russia: **FE** (KU). – Japan (Hon, Shi).

Ametastegia polygoni Takeuchi, 1929. Host plant: *Polygonum* spp. (Polygonaceae). Russia: **FE** (SA, KU). – China (NE), Korean Peninsula, Japan (Hon, Kyu).

APETHYmus Benson, 1939 (*Kjellia* Malaise, 1947). Type species *Dolerus abdominalis* Lepeletier, 1823. The larvae feed on deciduous trees and shrubs. The genus is distributed in Palaearctic and Oriental region. Number of species: World – about 25; Palaearctic – about 24; Russia – 7.

Apethymus apicalis (Klug, 1818) [Tenthredo] (*Emphytus klugii* Thomson, 1871). Host plant: *Rosa* spp. (Rosaceae). Russia: **EP** (Liston, 1995). – Europe (WE, EE, NE), Turkey.

Apethymus filiformis (Klug, 1818) [Tenthredo] (*Tenthredo serotina* Klug, 1818; *T. tarsata* Zetterstedt, 1819; *Dolerus abdominalis* Lepeletier, 1823; *D. abdominalis* Serville, 1823; *Emphytus temesiensis* Mocsáry, 1879; *E. serotinus melanopus* Ulbricht, 1913; *E. autumnalis* Forsius, 1933). Host plant: *Quercus* spp. (Fagaceae). Russia: **EP** (NW). – Europe (WE, EE, NE, SE).

Apethymus kunugi Togashi, 2005 (*A. proceratis*: Sundukov, Lelej, 2012). Host plant: *Quercus acutissima*, *Q. mongolicus* (Fagaceae). Russia: **FE** (PR). – Japan (Hon).

Apethymus kuri Takeuchi, 1952. Host plant: *Castanea crenata*, *Quercus crispula* (Fagaceae). Russia: **FE** (KU). – Japan (Hok, Hon, Shi, Kyu).

Apethymus parallelus (Eversmann, 1847) [Emphytus]. Russia: **UR, FE** (MG).

Apethymus serotinus (O.F. Müller, 1776) [Tenthredo] (*Tenthredo braccata* Gmelin, 1790; *T. varicornis* Gmelin, 1790; *T. tibialis* Panzer, 1799; *Emphytus caligatus* Eversmann, 1847; *E. panzeri* W.F. Kirby, 1882). Host plant: *Quercus* spp. (Fagaceae), *Rosa* spp. (Rosaceae). Russia: **EP** (NW, C, E, NC), **UR, WS** (NS). – Europe (WE, EE, NE, SE), Georgia.

Apethymus sidorenkoi Sundukov, 2010. Host plant: *Quercus mongolicus* (Fagaceae). Russia: **FE** (PR).

ASIEMPHYTUS Malaise, 1947. Type species *Macremphytus deutziae* Takeuchi, 1929. The larvae feed on the deciduous trees and shrubs; when pupation is carried into dry branches and bark.

The genus is distributed in East Asia and Oriental region. Number of species: World – 10; Palaearctic – 8; Russia – 2.

Asiemphytus dellei (Conde, 1935) [Macremphytus]. Host plant: *Deutzia crenata* (Hydrangeaceae). Russia: **FE** (AM, PR, KU). – China (NE, SE).

Asiemphytus fasciatus (Takeuchi, 1929) [Macremphytus]. Host plant: *Cornus controversa* (Cornaceae). Russia: **FE** (SA). – Japan (Hok, Hon, Shi, Kyu).

BELESES Cameron, 1877 (*Anisoneura* Cameron, 1876; *Belesidea* Rohwer, 1916). Type species *Anisoneura stigmatica* Cameron, 1876. The genus is distributed in East Asia and Oriental region. Number of species: World – 22; Palaearctic – 14; Russia – 1.

Beleses satonis (Takeuchi, 1929) [Abeleses]. Host plant: *Actinidia arguta* (Actinidiaceae). Russia: **FE** (KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

DINAX Konow, 1897 (*Adamas* Malaise, 1945). Type species *Dinax jakowleffi* Konow, 1897. The larvae feed on the Rosaceae. The genus is distributed mainly in East Asia; one species in Oriental region, and one species up to the Urals. Number of species: World – 6; Palaearctic – 5; Russia – 2.

Dinax ermak (Zhelochovtsev, 1968) [Adamas]. Host plant: *Spiraea* spp. (Rosaceae). Russia: **EP** (E), **UR, ES** (ZB), **FE** (MG). – Europe (EE).

Dinax jakowleffi Konow, 1897. Host plant: *Spiraea* spp. (Rosaceae). Russia: **ES** (IR, YA), **FE** (AM, KH, PR). – China (NE).

EMPRIA Lepeletier et Serville, 1828 (*Poecilostoma* Dahlbom, 1835; *Proscris* Gistel, 1848; *Poecilosoma* Thomson, 1870; *Poecilostomidea* Ashmead, 1898; *Tetratneura* Ashmead, 1898; *Parataxonous* MacGillivray, 1908; *Leucempria* Enslin, 1913; *Triempria* Enslin, 1914). Type species *Dolerus pallimacula* Lepeletier, 1823. The larvae mainly feed on Rosaceae, some on *Betula* (Betulaceae) and *Salix* (Salicaceae). The genus is distributed in Holarctic, Oriental and Neotropical regions. Number of species: World – about 50; Palaearctic – about 40; Russia – 28.

Empria alector Benson, 1938. Host plant: *Filipendula* spp. (Rosaceae). Russia: **EP** (N, NW, C, E), **WS** (Zhelochovtsev, 1988). – Europe (WE, EE, NE).

Empria alpina Benson, 1938. Host plant: *Dryas* spp. (Rosaceae). Russia: **EP** (NW). – Europe (WE, NE).

- Empria archangelskii** Dovnar-Zapolskij, 1929. Russia: **EP** (NC). – Caucasus, SW Asia.
- Empria basalis** Lindqvist, 1968. Host plant: Geum spp. (Rosaceae). Russia: **EP** (N). – Europe (WE, EE, NE).
- Empria camtschatica** Forsius, 1928. Host plant: Salix spp. (Salicaceae). Russia: **FE** (KA). – Europe (NE).
- Empria candidata** (Fallén, 1808) [Tenthredo] (*Tenthredo repanda* Klug, 1816). Host plant: Betula spp. (Betulaceae). Russia: **EP** (NW, C, E), **WS** (NS), **ES** (IR, BR, YA), **FE** (AM, KH, PR, KA). – Europe (WE, EE, NE, SE), Kazakhstan, Mongolia, China (NE), Korean Peninsula, Japan (Hok), N America.
- Empria excisa** (Thomson, 1871) [Poecilosoma] (*E. africana* Forsius, 1918). Host plant: Filipendula spp. (Rosaceae). Russia: **EP** (N). – Europe (WE, EE, NE, SE), N Africa, Turkey.
- Empria fletcheri** (Cameron, 1878) [Poecilosoma]. Host plant: Betula spp. (Betulaceae). Russia: **EP** (N), **WS** (TM), **ES** (TU, BR). – Europe (WE, NE), Mongolia.
- Empria gelida** (Erichson, 1851) [Tenthredo]. Russia: **FE** (KH).
- Empria hungarica** (Konow, 1895) [Poecilosoma] (*E. zacharovi* Dovnar-Zapolskij, 1929). Host plant: Filipendula spp. (Rosaceae). Russia: **EP** (C, S, NC). – Europe (WE, EE, NE, SE), Kazakhstan.
- Empria immersa** (Klug, 1818) [Tenthredo] (*Selandria labialis* Brullé, 1832; *Tenthredo opacomaculata* Eversmann, 1847; *Emphytus majalis* Snellen van Vollenhoven, 1869; *Empria tirolensis* Enslin, 1914). Host plant: Salix spp. (Salicaceae). Russia: **EP** (NW, E), **UR**, **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (IR, BR, YA), **FE** (AM, KH, MG). – Europe (WE, EE, NE, SE), Mongolia.
- Empria liturata** (Gmelin, 1790) [Tenthredo] (*Tenthredo guttata* Fallén, 1808; *T. impressa* Klug, 1816; *Emphytus pallimaculatus* Stephens, 1835; *Tenthredo spuria* Zetterstedt, 1838; *Emphytus fenestratus* Eversmann, 1847; *E. vernalis* Dietrich, 1868; *Poecilosoma submutica* Thomson, 1871; *P. undulata* Konow, 1885; *Harpiphorus calderoni* Medina, 1888). Host plant: Filipendula spp., Fragaria spp., Geum spp., Rubus spp. (Rosaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (NS, KM), **ES** (KR, IR, YA), **FE** (MG). – Europe (WE, EE, NE, SE), Turkey, Mongolia, Japan (Hok).
- Empria loktini** Ermolenko, 1971. Host plant: Geum japonicum, G. calthifolium (Rosaceae). Russia: **FE** (SA). – Japan (Hok).
- Empria longicornis** (Thomson, 1871) [Poecilosoma] (*E. rubi* Kontuniemi, 1951). Host plant: Rubus spp. (Rosaceae). Russia: **EP** (NW, C, E), **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (Zhelochovtsev, Zinovjev, 1996). – Europe (WE, EE, NE, SE), Mongolia.
- Empria mongolica** (Konow, 1895) [Poecilosoma]. Russia: **ES** (TU). – Mongolia, China (NC).
- Empria pallimacula** (Serville, 1823) [Dolerus] (*Dolerus pallimacula* Lepeletier, 1823; *Empria baltica* Conde, 1937). Host plant: Filipendula spp. (Rosaceae). Russia: **EP** (NW, C, E, S, NC), **UR**, **WS** (NS, KM), **ES** (YA). – Europe (WE, EE, NE, SE), Armenia.
- Empria parvula** (Konow, 1892) [Poecilosoma] (*E. pseudodoklugi* Dovnar-Zapolskij, 1929; *E. kuznetzovi* Dovnar-Zapolskij, 1929). Host plant: Geum spp. (Rosaceae). Russia: **EP** (N, NW, C, E, S), **FE** (AM, PR). – Europe (WE, EE, NE, SE), Mongolia.
- Empria plana** (Jakovlev, 1891) [Poecilosoma] (*Tenthredo hybrida* Erichson, 1851; *Empria itelmena* Malaise, 1931; *E. erichsoni* Liston, 1995). Host plant: Salix spp. (Salicaceae). Russia: **WS** (Lelej, Taeger, 2007), **ES** (IR), **FE** (AM, KH, PR, KA, MG). – Mongolia, Japan (Hok, Hon).
- Empria pravei** Dovnar-Zapolskij, 1925. Russia: **EP** (NC). – Armenia, Iran.
- Empria pumila** (Konow, 1896) [Poecilosoma]. Host plant: Filipendula spp. (Rosaceae). Russia: **EP** (NW, E), **WS** (KM). – Europe (WE, EE, NE, SE), Kazakhstan.
- Empria pumiloides** Lindqvist, 1968 (*E. tricornis* Lindqvist, 1968). Host plant: Filipendula spp. (Rosaceae). Russia: **EP** (Heidemaa, Prous, 2006). – Europe (WE, NE).
- Empria quadrimaculata** Takeuchi, 1952. Host plant: Rosa multiflora, Potentilla indica, ?Rubus parvifolius (Rosaceae). Russia: **FE** (KU). – Japan (Hon, Shi, Kyu).
- Empria rubicola** Ermolenko, 1971. Russia: **FE** (SA). – Japan (Hok).
- Empria sexpunctata** (Serville, 1823) [Tenthredo] (*Tenthredo sexpunctata* Lepeletier, 1823; *Selandria klugii* Stephens, 1835; *Poecilosoma guttata* Thomson, 1871; *P. carbonarium* Konow, 1884; *Empria waldstaettrense* Liston, 1980). Host plant: Geum spp. (Rosaceae). Russia: **EP** (NW, C, E, S, NC), **UR**. – Europe (WE, EE, NE, SE), Georgia, Armenia.
- Empria testaceipes** (Konow, 1896) [Poecilosoma]. Host plant: Sanguisorba officinalis (Rosaceae).

- Russia: **EP (E), ES (IR), FE (AM)**. – Europe (WE, EE, NE), SW Asia, Mongolia.
- Empria tridens** (Konow, 1896) [Poecilosoma] (*E. caucasicum* Dovnar-Zapolskij, 1929; *E. gussakovskii* Dovnar-Zapolskij, 1929; *E. konowi* Dovnar-Zapolskij, 1929). Host plant: Geum japonicum, Filipendula camtschatica, Rubus spp. (Rosaceae). Russia: **EP (NW, C, E, S, NC), UR, ES (TU, IR), FE (AM, PR, SA, KA)**. – Europe (WE, EE, NE, SE), Turkey, Mongolia, Japan (Hok).
- Empria tridentis** Lee et Ryu, 1996. Host plant: Geum japonicum, Filipendula camtschatica (Rosaceae). Russia: **FE (KH, PR, SA)**. – Korean Peninsula, Japan (Hok).
- Empria ushinskii** Dovnar-Zapolskij, 1929. **EP (NC, CR)**. – Europe (EE).
- EOPSIS** Benson, 1959. Type species *Eopsis beaumonti* Benson, 1959. The genus is distributed in West Palaearctic. Number of species: one West Palaearctic species.
- Eopsis beaumonti** Benson, 1959 (*Monosoma mosquense* Zhelochovtsev, 1981). Host plant: Rumex spp., Polygonum spp. (Polygonaceae). Russia: **EP (C)**. – Europe (WE, EE).
- ERIOCAMP** Hartig, 1837 (*Brachycampa* Zirngiebl, 1956). Type species *Tenthredo ovata* Linnaeus, 1760. The genus is distributed in Holarctic and Oriental region; one species was introduced into North America. Number of species: World – about 20; Palaearctic – 15; Russia – 6.
- Eriocampa albipes** Matsumura, 1912. Host plant: Alnus spp. (Betulaceae). Russia: **FE (AM, KH, SA)**. – Korean Peninsula, Japan (Hok, Hon, Shi).
- Eriocampa dorpatica** Konow, 1887. Host plant: black currant – Ribes nigrum and others (Grossulariaceae). Russia: **EP (NW, C, E), UR, ES (KR, IR)**. – Europe (NE).
- Eriocampa mitsukurii** Rohwer, 1910. Host plant: Alnus spp. (Betulaceae). Russia: **FE (AM, KH, PR, SA, KU)**. – China (SE), Japan (Hok, Hon, Shi).
- Eriocampa ovata ovata** (Linnaeus, 1760) [*Tenthredo*] (*Tenthredo gossypina* Retzius, 1783; *T. vernalis* Geoffroy, 1785; *T. leucogona* Schrank, 1802; *Eriocampa ovata nigritibialis* Enslin, 1914). Host plant: Alnus spp. (Betulaceae), Rhamnus spp., Frangula spp. (Rhamnaceae), Salix spp. (Salicaceae). Russia: **EP (N, NW, C, E, S, NC), UR**. – Europe (WE, EE, NE, SE), Georgia, Turkey; introduced into N America.
- Eriocampa peineae** Zirngiebl, 1956. Russia: **EP (Lacourt, 1999)**. – Iran.
- Eriocampa umbratica** (Klug, 1816) [*Tenthredo*]. Host plant: Alnus spp. (Betulaceae). Russia: **EP (N, NW, C, E, S, NC)**. – Europe (WE, EE, NE, SE), Georgia, Turkey.
- ERIOCAMPOPSIS** Takeuchi, 1952. Type species *Eriocampopsis subtruncata* Takeuchi, 1952. The genus is distributed in East Asia. Number of species: World and Palaearctic – 2; Russia – 1.
- Eriocampopsis subtruncata** Takeuchi, 1952. Host plant: Actaea spp. (Ranunculaceae). Russia: **FE (PR)**. – Japan (Hon).
- FILIXUNGULIA** Wei, 1997. Type species *Filixungulia alboclypea* Wei, 1997. East Palaearctic genus. Palaearctic – 3 species, Russia – 1.
- Filixungulia alboclypea** Wei, 1997. Russia: **FE (PR)**. – China (NE).
- HARPIPHORUS** Hartig, 1837 (*Asticta* Newman, 1838; *Harpophorus* Agassiz, 1848). Type species *Tenthredo lepida* Klug, 1818. The larvae feed on the Quercus (Fagaceae). The genus is distributed in West Palaearctic. Number of species: one species.
- Harpiphorus lepidus** (Klug, 1818) [*Tenthredo*] (*Fenusianthe* Newman, 1837). Host plant: Quercus robur (Fagaceae). Russia: **EP (NW, C)**. – Europe (WE, EE, NE, SE).
- HEMIBELESES** Takeuchi, 1929. Type species *Hemibeleses nigriceps* Takeuchi, 1929. The genus is distributed in East Asia and Oriental region. Number of species: World – about 20; Palaearctic – 14; Russia – 1.
- Hemibeleses ventralis** Takeuchi, 1952. Russia: **FE (PR)**. – Korean Peninsula, Japan (Hon, Shi, Kyu).
- HEPTAPOTAMIUS** Malaise, 1935. Type species *Heptapotamius simini* Malaise, 1935. The genus is distributed in Central Eurasia. Number of species: one species.
- Heptapotamius simini** Malaise, 1935. Russia: **EP (S)**. – Uzbekistan, Kyrgyzstan, Kazakhstan.
- MONOSTEGIA** Costa, 1859 (*Nematoceros* Konow, 1896). Type species *Tenthredo luteola* Klug, 1816. The larvae pupate in the soil. The genus is distributed in Palaearctic; one species was introduced

into N America. Number of species: World and Palaearctic – 5; Russia – 2.

Monostegia abdominalis (Fabricius, 1798) [Tenthredo] (*Tenthredo luteola* Klug, 1816; *Monostegia martini* MacGillivray, 1908; *M. nearctica* Rohwer, 1912; *Empria abdominalis rufinotis* Enslin, 1914). Host plant: *Lysimachia* spp., *Anagallis* spp. (Primulaceae), *Lactuca* spp. (Asteraceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (IR, YA), **FE** (AM, PR). – Europe (WE, EE, NE, SE), Azerbaijan, Armenia, SW Asia, Kyrgyzstan, Kazakhstan, China (Wei et al., 2006); introduced into N America.

Monostegia cingulata (Konow, 1891) [Poecilosoma]. Russia: **EP** (NC). – Georgia.

MONSOMA MacGillivray, 1908 (*Monosoma* Viereck, 1910). Type species *Poecilostoma inferentia* Norton, 1868. The larvae feed on Betulaceae. The genus is distributed in Holarctic. Number of species: World – 4; Palaearctic – 3; Russia – 2.

Monsoma pallipes (Matsumura, 1912) [Poecilosoma]. Host plant: ?*Alnus* spp. (Betulaceae). Russia: **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (Zhelochovtsev, Zinovjev, 1996), **FE** (PR). – Japan (Hok).

Monsoma pulveratum (Retzius, 1783) [Tenthredo] (*Tenthredo limbata* Gmelin, 1790; *T. pulverulenta* Christ, 1791; *T. obesa* Klug, 1817; *T. obtusa* Klug, 1817; *Selandria pulchella* Stephens, 1835; *Tenthredo leuczonias* Hartig, 1837; *T. segmentata* Zetterstedt, 1838; *Harpiphorus taeniatus* Costa, 1869; *Empria pulverata obtusalis* Enslin, 1914). Host plant: *Alnus* spp. (Betulaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **WS** (KM). – Europe (WE, EE, NE, SE), N Africa, Caucasus, SW Asia; introduced into N America.

STENEMPHYTUS Wei et Nie, 1999. Type species *Stenemphytus superbus* Wei et Nie, 1999. The genus is distributed in East Asia and Oriental region. Number of species: World – 4; Palaearctic – 2; Russia – 1.

Stenemphytus nakabusensis (Takeuchi, 1929) [Allantus]. Host plant: *Prunus* spp. (Rosaceae). Russia: **FE** (KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

TAXOBLENUS Wei et Nie, 1999. Type species *Taxoblenus longicornis* Wei et Nie, 1999. The genus is distributed in Palaearctic and Oriental region. Number of species: World – 11; Palaearctic – 8; Russia – 1.

Taxoblenus latus (Jakovlev, 1888) [Taxonus] (*Taxonius ballioni* Konow, 1891). Russia: **EP** (S), **ES** (IR, ZB).

TAXONUS Hartig, 1837 (*Ermilia* Costa, 1859; *Parasibla* Ashmead, 1898; *Strongylogastroidea* Ashmead, 1898; *Hypotaxonus* Ashmead, 1898). Type species *Tenthredo nitida* Klug, 1817. The larvae pupate in the soil. The genus is distributed in Holarctic and Oriental region. Number of species: World – 60; Palaearctic – about 45; Russia – 7.

Taxonus abdominalis Lee et Ryu, 1996. Russia: **FE** (PR). – Korean Peninsula.

Taxonus agrorum (Fallén, 1808) [Tenthredo] (*Tenthredo nitida* Klug, 1817; *Perineura dualis* Förster, 1844; *Tenthredo anomala* Eversmann, 1847; *T. subiecta* Eversmann, 1847; *Ermilia pulchella* Costa, 1859; *Strongylogaster rubrofasciatus* Palma, 1861; *Taxonus creperus* Konow, 1900). Host plant: *Rubus* spp. (Rosaceae). Russia: **EP** (NW, C, E), **UR**, **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (IR, BR, YA, ZB), **FE** (KH). – Europe (WE, EE, NE, SE), Turkey, Japan (Hok, Hon, Shi, Kyu).

Taxonus carbonarius Takeuchi, 1929. Russia: **FE** (PR). – China (SE).

Taxonus delumbis Konow, 1900 (*T. kamtchaticus* Malaise, 1931). Host plant: *Populus* spp. (Salicaceae). Russia: **ES** (KR, IR, BR, YA), **FE** (PR, SA, KA, MG). – China (Wei et al., 2006), Korean Peninsula, Japan (Togashi, 2000).

Taxonus fluvicornis Matsumura, 1912 (*Conaspidia bergrothi* Forsius, 1925). Host plant: *Polygonum cuspidatum* (Polygonaceae). Russia: **FE** (SA, KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Taxonus sticticus (Klug, 1817) [Tenthredo]. Host plant: *Rosa* spp. (Rosaceae). Russia: **EP** (S, NC), **ES** (IR). – Europe (WE, EE, SE), Turkey.

Taxonus zhelochovtsevi Viitasaari et Zinovjev, 1991. Russia: **FE** (KH, PR). – China (NE).

Unplaced Allantinae

Emphytus albisternus André, 1882. Russia: “Amur” (?*Ametastegia* Costa, 1882; see Blank et al., 2009).

Subfamily ATHALIINAE

ATHALIA Leach, 1817 (*Dentathalia* Benson, 1931). Type species *Tenthredo spinarum* Fabricius, 1793. The larvae feed on the Brassicaceae and Lamiaceae. The genus is distributed in Palaearctic,

- Oriental and Afrotropic regions. Number of species: World – about 95; Palaearctic - > 45; Russia – 18 (+ 3 ssp.).
- Athalia ahngeri** Kokujev, 1910. Russia: **EP** (Lacourt, 1999). – Azerbaijan, Armenia, Uzbekistan, Kazakhstan.
- Athalia ancilla ancilla** Serville, 1823 (*A. ancilla* Lepetier, 1823; *A. glabricollis* Thomson, 1870; *A. glabricollis bulgarica* Gregor, 1932). Host plant: *Daucus* spp. (Apiaceae). Russia: **EP** (C, NC, CR), **UR**, **WS** (NS). – Europe (WE, EE, NE, SE), N Africa, Georgia, Turkey.
- Athalia bicolor** Serville, 1823 (*Tenthredo annulata* Fabricius, 1787; *Athalia richardi* Serville, 1823; *A. richardi* Lepeletier, 1823; *A. bicolor* Lepeletier, 1823; *A. bicolor nigritegulis* Enslin, 1918). Host plant: ?*Ranunculus* spp. (Ranunculaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **WS** (NS). – Europe (WE, EE, NE, SE), N Africa, Caucasus, Turkey, Turkmenistan.
- Athalia circularis circularis** (Klug, 1815) [*Tenthredo* (*Tenthredo annulata* Gmelin, 1790; *Athalia lineolata* Lepeletier, 1823; *A. lineolata* Serville, 1823; *A. rosae sternalis* Costa, 1894; *A. bolivari* Dusmet, 1896; ?*A. lineolata niezabitowskii* Szulczewski, 1938; ?*A. lineolata gorkaensis* Szulczewski, 1938; *A. cordatoides* Kontuniemi, 1951; *A. alpina* Chevin, 1969). Host plant: Plantaginaceae, Asteraceae, Lamiaceae, etc. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (NS, KM, AL), **ES** (KS, KR). – Europe (WE, EE, NE, SE), N Africa, Caucasus, Turkey, Pakistan, Uzbekistan, Tajikistan, Kazakhstan, India (NW).
- Athalia circularis melanoptera** Benson, 1962. Host plant: Plantaginaceae, Asteraceae, Lamiaceae, etc. Russia: **ES** (IR, ZB), **FE** (AM, SA, KU, KA). – Mongolia, China (NE), Korean Peninsula.
- Athalia cordata** Serville, 1823 (*A. suessionensis* Serville, 1823; *A. suessionensis* Lepeletier, 1823; *A. cordata* Lepeletier, 1823; *A. blanchardi* Brullé, 1846; *A. rosae obscura* Konow, 1884; *A. lineolata analis* Enslin, 1913). Host plant: Plantaginaceae and Lamiaceae. Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, EE, NE, SE), N Africa, Caucasus, SW Asia.
- Athalia cornubiae** Benson, 1931. Host plant: *Sedum* spp. (Crassulaceae). Russia: **EP** (CR). – Europe (WE, EE, SE), Georgia, Armenia, Turkey, Uzbekistan; introduced into N America.
- Athalia decorata** Konow, 1900. Russia: **ES** (KS, KR, IR, BR, ZB), **FE** (AM, PR). – China (SW).
- Athalia dimidiata** Konow, 1891. Russia: **EP** (Lacourt, 1999). – Georgia, Armenia, Turkey.
- Athalia infumata** (Marlatt, 1898) [*Phyllotoma* (*A. nigrinotum* Matsumura, 1912)]. Host plant: *Cardamine* spp., *Nasturtium* spp., *Rorippa* spp., *Brassica* spp. (Brassicaceae). Russia: **FE** (PR, SA, KU). – China (SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu), India.
- Athalia japonica** (Klug, 1815) [*Tenthredo*] (*A. novitella* Kokujev, 1927). Host plant: Brassicaceae. Russia: **WS** (AL), **ES** (KS, KR, IR), **FE** (AM, KH, PR, SA, KU). – Kazakhstan, China (NE, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu), India.
- Athalia liberta** (Klug, 1815) [*Tenthredo*] (*A. rosae immaculata* Konow, 1884). Host plant: Brassicaceae. Russia: **EP** (NW, C, E, S, NC), **UR**, **ES** (KR). – Europe (WE, EE, NE, SE), Caucasus, Turkey, Iran, Tajikistan, Uzbekistan, Kyrgyzstan, China (NW, SW), India.
- Athalia longifoliae** Kontuniemi, 1951. Host plant: *Veronica longifolia* (Plantaginaceae). Russia: **EP** (NW, C, S, NC). – Europe (WE, NE), Central Asia, Kazakhstan.
- Athalia lugens** (Klug, 1815) [*Tenthredo*] (*A. lugens flavoscutellata* Enslin, 1918). Host plant: *Ajuga reptans* (Lamiaceae), ?Brassicaceae. Russia: **EP** (NW, C, NC, CR), **WS** (NS), **ES** (BR), **FE** (PR, KA). – Europe (WE, EE, NE, SE), Georgia, China (NE), Korean Peninsula, Japan (Hon), India, SE Asia.
- Athalia proxima** (Klug, 1815) [*Tenthredo*] (*A. tibialis* Cameron, 1876; *A. spinarum orientalis* Cameron, 1877; *A. proxima funebris* Forsius, 1925; *A. colibri kuroiwae* Matsumura et Uchida, 1926; *A. lugens camtschatica* Forsius, 1928; *A. lugens tristis* Forsius, 1934). Host plant: Brassicaceae. Russia: **FE** (KH, PR, SA, KU, KA). – China (SE), Korean Peninsula, Japan (Ryu), India, SE Asia.
- Athalia rosae rosae** (Linnaeus, 1758) [*Tenthredo* (*Tenthredo colibri* Christ, 1791; *T. spinarum* Fabricius, 1793; *T. centifoliae* Panzer, 1797)]. Host plant: Brassicaceae. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (NS, KM, AL), **ES** (KS, KR). – Europe (WE, EE, NE, SE), N Africa, Caucasus, SW Asia, Kazakhstan.
- Athalia rosae ruficornis** Jakovlev, 1888 (*A. leucostoma* Cameron, 1904; *A. spinarum japanensis* Rohwer, 1910). Host plant: Brassicaceae. Russia: **ES** (IR, BR), **FE** (AM, KH, PR, SA, KU). – Mongolia, China (NE, NC, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu), India.

Athalia rufoscutellata Mocsáry, 1879 (*A. maculata* Mocsáry, 1879; *A. maritima* W.F. Kirby, 1884; *A. rufoscutellata mocsaryi* Konow, 1892; *A. rufoscutellata nigroscutellata* Konow, 1892; *A. rufoscutellata pleuralis* Konow, 1898; *A. rufoscutellata obscurata* Konow, 1904). Host plant: *Teucrum* spp. (Lamiaceae). Russia: **EP** (S, NC, CR). – Europe (WE, EE, SE), Georgia, Armenia, Turkey, China (Wei et al., 2006).

Athalia scutellariae scutellariae Cameron, 1880 (*Dentathalia galericulatae* Kontuniemi, 1951). Host plant: *Scutellaria* spp., *Lycopus* spp. (Lamiaceae). Russia: **EP** (NW, C, E). – Europe (WE, EE, NE, SE), Turkey, Kyrgyzstan.

Athalia scutellariae flammma Zhelochovtsev, 1927. Host plant: *Scutellaria* spp. (Lamiaceae). Russia: **ES** (IR, ZB), **FE** (AM, KH, PR). – Mongolia, China (NC).

Athalia yanoi Takeuchi, 1952. Russia: **FE** (KH, PR, SA, KU). – Japan (Hok, Hon).

Subfamily BLENNOCAMPINAE (HETERARTHINAE)

ANAFENUSA Benson, 1959. Type species *Entodecta impropria* Malaise, 1931. The genus is distributed in East Asia and Oriental region. Number of species: World – 4; Palaearctic – 3; Russia – 1.

Anafenusa impropria (Malaise, 1931) [Entodecta]. Russia: **FE** (PR).

APAREOPHORA Sato, 1928. Type species *Apareophora forsythiae* Sato, 1928. The genus is distributed in Holarctic and Oriental region. Number of species: World – 8; Palaearctic – 6; Russia – 1.

Apareophora coreana Sato, 1928. Russia: **FE** (PR). – China (NC), Korean Peninsula.

ARDIS Konow, 1886 (*Valco* Ross, 1937). Type species *Tenthredo bipunctata* Klug, 1817. The larvae without spines on the body; borer the branches of *Rosa* (Rosaceae); go down the branch. The genus is distributed in Holarctic. Number of species: World – 5; Palaearctic – 4; Russia – 1.

Ardis pallipes (Serville, 1823) [Dolerus] (*Tenthredo bipunctata* Klug, 1817; *Dolerus pallipes* Lepeletier, 1823; *Tenthredo brunniventris* Hartig, 1837; *Monophadnus dissimilis* Costa, 1859; *Selandria irrogata* Cresson, 1880; *Emphytus dubius* W.F. Kirby, 1882; *Aphanisus odoratus* MacGillivray, 1908;

A. parallelus MacGillivray, 1923). Host plant: *Rosa* spp. (Rosaceae). Russia: **EP** (N, NW, C, E, S, NC), **FE** (MG). – Europe (WE, EE, NE, SE), Central Asia, China (Wei et al., 2006), Japan (Hon); introduced into N America.

BLENNOCAMPA Hartig, 1837. Type species *Tenthredo pusilla* Klug, 1816. The larvae with branched spines on the body; feeding on Rosaceae. The genus is distributed in Palaearctic, Oriental and Neotropical regions. Number of species: World – 10; Palaearctic – 3; Russia – 2.

Blennocampa phyllocolpa Viitalaari et Vikberg, 1985 (*Tenthredo pusilla* Klug, 1816). Host plant: *Rosa* spp. (Rosaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **ES** (IR), **FE** (AM, SA, KA, MG). – Europe (WE, EE, NE, SE), Turkey, Central Asia, Kazakhstan, China (NW).

Blennocampa sibirica Jakovlev, 1891. Russia: **ES** (IR).

CALIROA Costa, 1859 (*Eriocampoides* Konow, 1890; *Periclistoptera* Ashmead, 1898). Type species *Caliroa sebetia* Costa, 1859. The larvae are covered with mucus; skeletonize of leaves of trees and shrubs, pupate in the soil. The genus is distributed in Holarctic and Oriental region; was introduced into the Neotropical, Ethiopian and Australasian regions. Number of species: World – about 60; Palaearctic – 30; Russia – 7.

Caliroa angustata Forsius, 1927. Russia: **FE** (AM). – China (CC, SE).

Caliroa annulipes (Klug, 1816) [Tenthredo] (*Selandria atra* Stephens, 1835; *Eriocampa myrtilli* Brischke, 1889). Host plant: Salicaceae, Betulaceae, Fagaceae, Cannabaceae, etc. Russia: **EP** (N, NW, C, E), **UR**, **ES** (IR, BR, YA), **FE** (KA, MG). – Europe (WE, EE, NE, SE), Turkey, China (Wei et al., 2006), Japan (Hon), N America.

Caliroa cerasi (Linnaeus, 1758) [Tenthredo] (*Tenthredo flavipes* Schrank, 1781; *T. limacina* Retzius, 1783; *T. adumbrata* Klug, 1816; *Monostegia antipoda* W.F. Kirby, 1881; *Caliroa laudata* MacGillivray, 1909; *C. lacinata* MacGillivray, 1909). Host plant: Rosaceae, Salicaceae, Fagaceae, etc. Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (NS), **FE** (AM, KH, PR). – Europe (WE, EE, NE, SE), N Africa, SW Asia, Uzbekistan, Kyrgyzstan, Kazakhstan, China (NE, CC, SW, SE); introduced into N America, S America, S Africa, Australia, New Zealand.

Caliroa cinctia (Klug, 1816) [Tenthredo]. Host plant: Quercus spp. (Fagaceae). Russia: **EP** (C, S, NC). – Europe (WE, EE, NE, SE).

Caliroa oishii (Takeuchi, 1933) [Eriocampoides]. Host plant: Quercus spp. (Fagaceae). Russia: **FE** (PR, SA). – China (NE), Korean Peninsula, Japan (Hon).

Caliroa tremulae Chevin, 1974. Host plant: Populus tremula (Salicaceae). Russia: **EP** (NW). – Europe (WE, EE, NE).

Caliroa varipes (Klug, 1816) [Tenthredo] (*Eriocampa crassicornis* Tischbein, 1846; *Eriocampoides variipes* Dalla Torre, 1894). Host plant: Salix spp. (Salicaceae), Betula spp. (Betulaceae), Quercus robur (Fagaceae). Russia: **EP** (NW, C, S, NC), **WS** (Zhelochovtsev, Zinovjev, 1996), **FE** (AM, PR). – Europe (WE, EE, NE), Turkey, Japan (Hok).

CLADARDIS Benson, 1952. Type species *Tenthredo elongatula* Klug, 1817. The larvae bore the branches of Rosa (Rosaceae); move from top to bottom along the branch. The genus is distributed in Palaearctic; one species in Oriental region. Number of species: World and Palaearctic – 4; Russia – 2.

Cladardis elongatula (Klug, 1817) [Tenthredo]. Host plant: Rosa spp. (Rosaceae). Russia: **EP** (NW, C, E), **FE** (PR). – Europe (WE, EE, NE, SE), Turkey, Mongolia.

Cladardis hartigi Liston, 1995 (*Tenthredo semicincta* Hartig, 1837). Host plant: Rosa spp. (Rosaceae). Russia: **EP** (NW). – Europe (WE, EE, NE).

CLAREMONTIA Rohwer, 1909 (*Pseudoblennocampa* Malaise, 1935). Type species *Claremontia typica* Rohwer, 1909. The larvae feed on herbaceous of Rosaceae. The genus is distributed in Holarctic; one species in Oriental region. Number of species: World – 16; Palaearctic – 9; Russia – 7.

Claremontia alternipes (Klug, 1816) [Tenthredo] (*Tenthredo maura* Schrank, 1802; *Blennocampa divisa* Konow, 1884; *B. intermedia* Kriechbaumer, 1885; *B. tergestina* Kriechbaumer, 1888; *B. finitima* Konow, 1900). Host plant: Rubus spp. (Rosaceae). Russia: **EP** (NW, C, S, NC, CR), **WS** (Lelej, Taeger, 2007), **ES** (IR), **FE** (MG). – Europe (WE, EE, NE, SE), Mongolia.

Claremontia brevicornis (Brischke, 1883) [Blennocampa] (*Blennocampa confusa* Konow, 1886; *B. mixta* Malaise, 1920). Host plant: Fragaria spp., Potentilla spp., Sanguisorba spp. (Rosaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (KM), **ES** (IR). – Europe (WE, EE, NE, SE), Mongolia.

Claremontia puncticeps (Konow, 1886) [Blennocampa].

Host plant: Poterium sanguisorba (Rosaceae). Russia: **EP** (C, E), **WS** (NS), **ES** (IR), **FE** (AM, PR). – Europe (WE, SE), Turkey, Kyrgyzstan, Kazakhstan.

Claremontia springatei Koch, 1988. Russia: **ES** (IR).

Claremontia tenuicornis (Klug, 1816) [Tenthredo] (*Phyllotoma conformis* Fallén, 1829; *Selandria tibialis* Stephens, 1835; *Blennocampa spiraeae* Brischke, 1883). The larvae feed on the Spiraea spp., Filipendula spp., Alchemilla spp. (Rosaceae). Russia: **EP** (N, NW, C, E, NC), **UR**, **ES** (IR), **FE** (PR, MG). – Europe (WE, EE, NE, SE), Armenia, Turkey, China (Wei et al., 2006).

Claremontia uncta (Klug, 1816) [Tenthredo] (*Selandria humeralis* Snellen van Vollenhoven, 1869; *Blennocampa lioparea* Konow, 1884). Russia: **EP** (Liston, 1995). – Europe (WE, EE, NE).

Claremontia waldheimii (Gimmerthal, 1847) [Tenthredo] (*Selandria subcana* Zaddach, 1859; *Blennocampa subserrata* Thomson, 1870; *B. subcana angularis* Konow, 1895). Host plant: Geum spp. (Rosaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (NS), **ES** (Zhelochovtsev, Zinovjev, 1996), **FE** (AM, PR). – Europe (WE, EE, NE, SE), Armenia, Turkey, Mongolia.

ENDELOMYIA Ashmead, 1898. Type species *Selandria rosae* T.W. Harris, 1841. The larvae skeletonize of leaves of trees and shrubs, pupate in the soil. The genus is distributed in Palaearctic and Oriental region; one wasp species introduced into N America. Number of species: World and Palaearctic – 3; Russia – 1.

Endelomyia aethiops (Gmelin, 1790) [Tenthredo] (*Tenthredo morio* Fabricius, 1781; *T. atratula* Dahlbom, 1835; *Selandria rosae* T.W. Harris, 1841; *Eriocampa livonensis* Gimmerthal, 1844; *E. nitida* Tischbein, 1846; *Tenthredo dolosa* Eversmann, 1847; *Emphytus radialis* Eversmann, 1847; *Selandria soror* Snellen van Vollenhoven, 1869; *Eriocampa atratula* Thomson, 1870; *E. testaceipes* Cameron, 1874; *E. caninae* Cameron, 1878; *Poecilosoma nigricolle* Cameron, 1882). Host plant: Rosa spp. (Rosaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (AL), **FE** (PR). – Europe (WE, EE, NE, SE), Georgia, Armenia, Kyrgyzstan; introduced into N America.

ENDEMOLYIA Wei, 1998. Type species *Endemolygia genata* Wei, 1998. The larvae skeletonize of leaves of trees and shrubs, pupate in the soil. The genus is distributed in East Asia and Oriental

region. Number of species: World – 6; Palaearctic – 1; Russia – 1.

Endemyolia negativa (Malaise, 1931) [Caliroa]. Russia: **FE** (PR). – China (Wei et al., 2006).

EUPAREOPHORA Enslin, 1914. Type species *Blennocampa exarmata* Thomson, 1871. The genus is distributed in West Palaearctic and Nearctic. Number of species: World – 2; Palaearctic – 1; Russia - 1.

Eupareophora exarmata (Thomson, 1871) [*Blennocampa*]. Host plant: *Fraxinus* spp. (Oleaceae). Russia: **EP** (NC). – Europe (WE, EE, NE), Georgia.

EURHADINOCERAEA Enslin, 1920 (*Pseudoblennocampa* Malaise, 1935). Type species *Rhadinoceraea roseni* Enslin, 1920. The larvae feed on the Ranunculaceae. The genus is distributed in Palaearctic and Oriental region. Number of species: World – 16; Palaearctic – 15; Russia – 4.

Eurhadinoceraea athaliooides (Jakovlev, 1891) [Monophaenus] (*Rhadinoceraea singularis* Konow, 1897). Host plant: *Clematis* spp., *Pulsatella* spp. (Ranunculaceae). Russia: **EP** (NC), **FE** (PR). – Europe (WE), Armenia, China (NC).

Eurhadinoceraea roseni (Enslin, 1920) [*Rhadinoceraea*]. Host plant: *Clematis* spp. (Ranunculaceae). Russia: **FE** (KH). – China.

Eurhadinoceraea sanguinicollis (Mocsáry, 1880) [*Blennocampa*] (*Rhadinoceraea insignis* Konow, 1896). Host plant: *Clematis* spp. (Ranunculaceae). Russia: **ES** (IR). – Europe (EE).

Eurhadinoceraea ventralis (Panzer, 1799) [Tenthredo] (*Monophaenus rufoniger* Tischbein, 1852; *Blennocampa coronata* André, 1881). Host plant: *Clematis recta*, *C. vitalba* (Ranunculaceae). Russia: **EP** (C), **WS** (NS), **ES** (TU). – Europe (WE, SE), Kyrgyzstan, Kazakhstan.

EUTOMOSTETHUS Enslin, 1914 (*Atomostethus* Enslin, 1914; *Tomostethopsis* Sato, 1928; *Forsia* Malaise, 1931). Type species *Tenthredo luteiventris* Klug, 1816. The larvae feed on the Juncaceae, Poaceae, Cyperaceae. The genus is distributed in Palaearctic and Oriental region; two species were introduced into N America. Number of species: World – > 100; Palaearctic – > 60; Russia – 10 (+ 1 ssp.).

Eutomostethus apicalis (Matsumura, 1912) [Monophaenus] (*Tomostethus juncivorus* Rohwer, 1924). Host plant: *Juncus* spp. (Juncaceae). Russia: **FE** (SA, KU). – Japan (Hok, Hon, Shi).

Eutomostethus ephippium ephippium (Panzer, 1798) [Tenthredo] (*Tenthredo dubia* Gmelin, 1790; *Sealandria inhabilis* Norton, 1861). Host plant: *Poa* spp. (Poaceae). Russia: **EP** (N, NW, C, E, S), **UR**, **FE** (PR). – Europe (WE, EE, NE, SE), N Africa, Caucasus, Turkey, India (NW); introduced into N America.

Eutomostethus ephippium vopiscus (Konow, 1899) [Tomostethus]. Host plant: *Poa* spp. (Poaceae). Russia: **EP** (NC, CR). – Europe (EE), Georgia, Armenia.

Eutomostethus flavipes (Matsumura, 1911) [Tomostethus]. Russia: **FE** (SA).

Eutomostethus gagathinus (Klug, 1816) [Tenthredo] (*E. gagathinus meridionalis* Benson, 1954). Host plant: *Juncus* spp. (Juncaceae), *Carex* spp. (Cyperaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (OM, NS, AL), **ES** (IR). – Europe (WE, EE, NE, SE), N Africa, SW Asia, Kazakhstan, Mongolia.

Eutomostethus luteiventris (Klug, 1816) [Tenthredo] (*Tenthredo fuscipennis* Serville, 1823; *T. fuscipennis* Lepeletier, 1823; *Phyllotoma fuscipennis* Fallén, 1829). Host plant: *Juncus* spp. (Juncaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**. – Europe (WE, EE, NE, SE), Caucasus; introduced into N America.

Eutomostethus metallicus (Sato, 1928) [Tomostethopsis]. Russia: **FE** (PR). – Korean Peninsula, China (NE, SE).

Eutomostethus punctatus (Konow, 1887) [Tomostethus] (*Tomostethus brachycera* Cameron, 1893). Host plant: *Carex* spp. (Cyperaceae). Russia: **EP** (E), **UR**, **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (TU, IR), **FE** (PR). – Europe (WE, EE, NE, SE), Mongolia, China (NC).

Eutomostethus teberdicus Muche, 1969. Russia: **EP** (NC).

Eutomostethus tomostethi (Malaise, 1931) [Forsia]. Russia: **FE** (PR, SA, KA).

Eutomostethus tricolor (Malaise, 1934) [Tomostethus] (*E. hyalinus* Takeuchi, 1936). Russia: **FE** (PR, SA). – China (SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

FENELLA Westwood, 1839 (*Melinia* Costa, 1859; *Paraphyllotoma* Forsius, 1930). Type species *Fenella nigrita* Westwood, 1839. The larvae in mines on leaves of herbaceous plants. The genus is distributed in Palaearctic; one species was introduced into N America. Number of species: World and Palaearctic – 11; Russia – 3.

Fenella minuta (Dahlbom, 1835) [Phyllotoma] (*F. voigti* Hering, 1932). Host plant: *Geranium*

spp. (Geraniaceae). Russia: **EP** (NC), **ES** (IR), **FE** (AM). – Europe (WE, EE, NE, SE), Kyrgyzstan, Kazakhstan, Mongolia.

Fenella monilicornis (Dahlbom, 1835) [Phyllotoma] (*F. famosa* Benson, 1950). Host plant: *Geranium* spp. (Geraniaceae). Russia: **WS** (AL). – Europe (WE, EE, NE), Mongolia, China (NW).

Fenella nigrita Westwood, 1839 (*Melinia minutissima* Costa, 1859; *Phyllotoma tormentillae* Healy, 1868; *Fenella agrimoniae* Brischke, 1888). Host plant: *Agrimonia* spp., *Potentilla* spp. (Rosaceae). Russia: **EP** (NW), **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (YA). – Europe (WE, EE, NE, SE), Kyrgyzstan, Kazakhstan, Mongolia, Korean Peninsula, Japan (Hon); introduced into N America.

FENUSA Leach, 1817 (*Kaliosysphinga* Tischbein, 1846; *Aphadnurus* Costa, 1859; *Phaenus* Cameron, 1875; *Phoenusa* André, 1880; *Caliosysphinga* Konow, 1905; *Lithoryssus* Brues, 1906; *Kaliofenus* Viereck, 1910). Type species *Tenthredo pumila* Klug, 1818. The larvae in mines on leaves of trees. The genus is distributed in Holarctic and Oriental region; one species was introduced into the Afrotropical region. Number of species: World – 10; Palaearctic – 8; Russia – 6 from 2 subgenera.

Fenusa (Fenusa) dohrnii (Tischbein, 1846) [*Kaliosysphinga*] (*F. curtus* Norton, 1862; *Phaenus* *melanopoda* Cameron, 1876; *Fenella westwoodi* Cameron, 1882). Host plant: *Alnus* spp., *Betula* spp. (Betulaceae). Russia: **EP** (N, NW, C, E, S), **UR**, **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (TU, IR, BR), **FE** (KU). – Europe (WE, EE, NE, SE), Turkey, Kazakhstan, Japan (Hok, Hon, Shi); introduced into N America and Tropical Africa.

Fenusa (Fenusa) pumila Leach, 1817 (*Tenthredo pumila* Klug, 1818; *Aphadnurus tantillus* Costa, 1859; *Fenusa fuliginosa* Healy, 1869; *F. minima* Brischke, 1883). Host plant: *Betula* spp., *Alnus* spp. (Betulaceae), *Ulmus* spp. (Ulmaceae). Russia: **EP** (N, NW, C, E, NC), **WS** (Lelej, Taeger, 2007), **ES** (IR, YA), **FE** (KH, SA, KA). – Europe (WE, EE, NE, SE), Turkey, Kyrgyzstan, Kazakhstan, Mongolia, China (SW), Japan (Hok, Hon); introduced into N America.

Fenusa (Kaliofenus) ulmi Sundevall, 1847 (*F. intermedia* Thomson, 1871; *Messa alsia* MacGillivray, 1923). Host plant: *Ulmus* spp. (Ulmaceae). Russia: **EP** (C, E, S, NC), **ES** (IR, BR), **FE** (KH, PR). – Europe (WE, EE, NE, SE), Georgia, Armenia,

Turkey, Japan (Smith, 1981); introduced into N America.

Fenusa (Kaliofenus) zinojevi (Liston, 1994) [Kaliofenus]. Russia: **FE** (PR).

Fenusa absens D.R. Smith et Altenhofer, 2011. Host plant: *Ulmus pumila* (Ulmaceae). Russia: **WS** (NS).

Fenusa ewaldi D.R. Smith, 2011. Host plant: *Rosa* spp. (Rosaceae). Russia: **WS** (NS).

FENUSELLA Enslin, 1912 (*Melanobates* MacGillivray, 1916). Type species *Fenusa wuestneii* Konow, 1894. The larvae in mines on leaves of Betulaceae, Salicaceae, etc. The genus is distributed in Holarctic. Number of species: World – 11; Palaearctic – 8; Russia – 4.

Fesunella glaucopis (Konow, 1907) [*Phoenusa*] (*Phoenusa tictinensis* Magretti, 1886). Host plant: *Populus* spp. (Salicaceae). Russia: **EP** (C, E, S), **UR**, **ES** (IR). – Europe (WE, EE, NE, SE), Kazakhstan.

Fesunella hortulana (Klug, 1818) [*Tenthredo*] (*Phoenusa doderleini* De-Stefani, 1883; *Fenusella soenderupi* Hering, 1935). Host plant: *Populus* spp. (Salicaceae). Russia: **EP** (NC). – Europe (WE, EE, SE), N Africa, Azerbaijan, Turkey, Central Asia; introduced into N America.

Fenusella nana (Klug, 1816) [*Tenthredo*] (*Phyllotoma mellita* Newman, 1870; *Fenusa quercus* Cameron, 1885; *Scolioneura laeta* Enslin, 1918). Host plant: *Betula* spp. (Betulaceae). Russia: **EP** (NW, C, E), **UR**, **WS** (NS), **ES** (IR, BR), **FE** (AM, PR). – Europe (WE, EE, NE), Kyrgyzstan, Kazakhstan, Mongolia, Japan (Hok); introduced into N America.

Fenusella wuestneii (Konow, 1894) [*Fenusa*]. Host plant: *Salix* spp. (Salicaceae). Russia: **ES** (IR, BR), **FE** (KH). – Europe (WE, EE, NE), Kyrgyzstan, Kazakhstan, Japan (Hon); introduced into N America.

HALIDAMIA Benson, 1939. Type species *Hylotoma affinis* Fallén, 1807. The genus is distributed in West Palaearctic, introduced into the Nearctic. Number of species: Russia – 1.

Halidamia affinis (Fallén, 1807) [*Hylotoma*] (*Tenthredo hyalina* Klug, 1816; *Blennocampa assimilis* Thomson, 1870; *B. formosella* Costa, 1882; *B. affinis pleuritica* Enslin, 1914). Host plant: *Galium* spp. (Rubiaceae). Russia: **EP** (NW, C, E, S, NC, CR), **UR**. – Europe (WE, EE, NE, SE), Georgia, Armenia, Turkey; introduced into N America.

HETERARTHUS Stephens, 1835 (*Phyllotoma* Falén, 1829; *Decatria* Stephens, 1835; *Druida* Newman, 1838; *Phlebatrophia* MacGillivray, 1909). Type species *Tenthredo ochropoda* Klug, 1818. The larvae in mines on leaves of Acer (Sapindaceae), Betulaceae, Salicaceae. The genus is distributed in Palaearctic; one species in Oriental region; one species introduced to North America. Number of species: World – 22; Palaearctic – 21; Russia – 11.

Heterarthrus aceris (Kaltenbach, 1856) [*Phyllotoma*] (*Phyllotoma fumipennis* Cameron, 1888). Host plant: Acer platanoides (Sapindaceae). Russia: **EP** (C, S). – Europe (WE, EE, NE, SE), Armenia.

Heterarthrus aceroides (Malaise, 1931) [*Phyllotoma*]. Russia: **FE** (KA).

Heterarthrus aihinoensis Haris, 2006. Russia: **FE** (KU).

Heterarthrus alnivorus Togashi, 1992. Host plant: Alnus spp. (Betulaceae). Russia: **FE** (PR). – Japan (Hon).

Heterarthrus fasciatus (Malaise, 1931) [*Phyllotoma*]. Host plant: Populus spp. (Salicaceae). Russia: **ES** (BR), **FE** (KA).

Heterarthrus flavigollis (Gussakovskij, 1947) [*Phyllotoma*]. Host plant: Acer platanoides (Sapindaceae). Russia: **EP** (NC). – Europe (WE, EE, NE), Georgia.

Heterarthrus microcephalus (Klug, 1818) [*Tenthredo*]. Host plant: Salix spp. (Salicaceae). Russia: **EP** (NW, C, NC), **WS** (NS), **ES** (IR), **FE** (KH). – Europe (WE, EE, NE, SE), Armenia, Kyrgyzstan, Kazakhstan.

Heterarthrus nemoratus (Fallén, 1808) [*Hylotoma*] (*Fenusia parviceps* Newman, 1837; *Phyllotoma tenella* Zaddach, 1859; *Phlebatrophia mathesonii* MacGillivray, 1909). Host plant: Betula spp., Alnus spp. (Betulaceae). Russia: **EP** (N, NW, C), **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (IR). – Europe (WE, EE, NE), Armenia; introduced into N America.

Heterarthrus ochropoda (Klug, 1818) [*Tenthredo*] (*Decatria fuscipennis* Stephens, 1835; *Phyllotoma pinguis* Snellen van Vollenhoven, 1869; *Ph. nigrescens* Grädl, 1881; *Ph. costae* Costa, 1894; *Ph. maxima* Strobl, 1896). Host plant: Populus spp. (Salicaceae), Betula spp. (Betulaceae). Russia: **EP** (NW, C, S, NC, CR), **UR**, **WS** (NS), **ES** (IR, BR), **FE** (AM, KH). – Europe (WE, EE, NE), Turkey, Kyrgyzstan.

Heterarthrus vagans (Fallén, 1808) [*Hylotoma*] (*Tenthredo melanopyga* Klug, 1818; *T. amaura* Klug,

1818; *Phyllotoma leucopoda* Dahlbom, 1835; *Ph. kamtchatica* Malaise, 1931; *Heterarthrus vagans anatolicus* Muche, 1983). Host plant: Alnus spp., Betula spp. (Betulaceae). Russia: **EP** (N, NW, C, E, NC, CR), **UR**, **ES** (IR), **FE** (PR, KU, KA). – Europe (WE, EE, NE, SE), Azerbaijan, SW Asia, China (NE); introduced into N America.

Heterarthrus wuestneii (Konow, 1905) [*Phyllotoma*] (*Phyllotoma aceris* McLachlan, 1867; *Heterarthrus tauricus* Ermolenko, 1984; *H. healyi* Altenhofer et Zombori, 1987). Host plant: Acer spp. (Sapindaceae), Alnus spp. (Betulaceae). Russia: **EP** (CR). – Europe (WE, EE), N America.

HINATARA Benson, 1936. Type species *Fenusia excisa* Konow, 1885. The larvae in mines on leaves of trees. The genus is distributed in Palaearctic. Number of species: World and Palaearctic – 4; Russia – 3.

Hinatara hakusana Togashi, 2002. Russia: **FE** (PR). – Japan (Hon).

Hinatara nigripes (Konow, 1907) [*Fenusia*]. Host plant: Acer campestre (Sapindaceae). Russia: **EP** (S). – Europe (WE, EE, SE).

Hinatara recta (Thomson, 1871) [*Blennocampa*]. Host plant: Acer platanoides (Sapindaceae). Russia: **EP** (NW, C, S). – Europe (WE, EE, NE), Armenia.

HOPLOCAMPOIDES Enslin, 1914. Type species *Selandria xylostei* Giraud, 1863. The genus is distributed in Palaearctic. Number of species: World and Palaearctic – 2; Russia – 1.

Hoplocampoides xylostei (Vallot, 1836) [*Tenthredo*] (*Selandria xylostei* Giraud, 1863). Host plant: Lonicera spp. (Caprifoliaceae). Russia: **EP** (N, NW, C), **ES** (KR). – Europe (WE, EE, NE, SE).

LAGONIS Ross, 1937. Type species *Selandria nevadensis* Cresson, 1880. The genus is distributed in East Palaearctic and Nearctic. Number of species: World – 3; Palaearctic – 2; Russia – 1.

Lagonis opacicollis (Malaise, 1931) [*Rhadinoceraea*]. Host plant: Sambucus sieboldiana (Adoxaceae). Russia: **FE** (KU). – Japan (Hok, Hon, Shi).

MASAAKIA Takeuchi, 1950. Type species *Masaakia longivaginata* Takeuchi, 1950. The genus is distributed in East Asia. Number of species: World and Palaearctic – 5; Russia – 2.

Masaakia kichizoi Haris, 2006. Russia: **FE** (KU).

Masaakia longivaginata Takeuchi, 1950. Russia: **FE** (KU). – Japan (Hon).

METALLUS Forbes, 1885 (*Entodecta* Konow, 1886; *Polybates* MacGillivray, 1909). Type species *Metallus rubi* Forbes, 1885. The larvae in mines on leaves of Rosaceae. The genus is distributed in Holarctic and Oriental region. Number of species: World – 21; Palaearctic – 7; Russia – 5.

Metallus adamantis Lee et Ryu, 1996. Russia: **FE** (PR). – Korean Peninsula.

Metallus albipes (Cameron, 1875) [*Phaenusa*] (*Entodecta tenuicornis* Hellén, 1935). Host plant: Rubus spp. (Rosaceae). Russia: **EP** (C), **FE** (SA, KA, MG). – Europe (WE, EE, NE), Japan (Hon).

Metallus beckeri (Konow, 1904) [*Entodecta*]. Russia: **EP** (NC). – Georgia, Turkey.

Metallus lanceolatus (Thomson, 1870) [*Blennocampa*] (*Fenusae gei* Brischke, 1883; *Entodecta decolor* Konow, 1886; *Metallus bensoni* D.R. Smith, 1971). Host plant: Geum spp. (Rosaceae). Russia: **EP** (NW, C, NC), **UR**, **FE** (AM). – Europe (WE, EE, NE), N America.

Metallus pumilus (Klug, 1816) [*Tenthredo*] (*Emphytus pumilio* Hartig, 1837; *Fenusae rubi* Boie, 1848). Host plant: Rubus spp. (Rosaceae). Russia: **EP** (NW, C, NC), **ES** (IR, BR), **FE** (KH, PR). – Europe (WE, EE, NE, SE), Japan (Hon).

MONARDIS Enslin, 1914. Type species *Tenthredo plana* Klug, 1817. The genus is distributed in Holarctic and Oriental region. Number of species: World – 8; Palaearctic – 4; Russia – 1.

Monardis plana (Klug, 1817) [*Tenthredo*] (*Tenthredo sericans* Hartig, 1837; *Blennocampa rosarum* Brischke, 1883). Host plant: Rosa spp. (Rosaceae); larvae in the buds and shoots. Russia: **EP** (NW, C, E, S, NC), **UR**, **WS** (KM). – Europe (WE, EE, NE, SE), Caucasus.

MONOPHADNOIDES Ashmead, 1898 (*Pseudomonophadnus* Malaise, 1935). Type species *Selandria rubi* T.W. Harris, 1845. The genus is distributed in Holarctic and Oriental region. Number of species: World and Palaearctic – 9; Russia – 4.

Monophadnoides klausnitzeri (Muche, 1973) [Monophadnus]. Russia: **EP** (NC).

Monophadnoides rubi (T.W. Harris, 1845) [*Selandria*] (*Tenthredo geniculata* Hartig, 1837; *T. longicornis* Hartig, 1837; *Selandria nigella* Cresson, 1880;

Monophadnus hudsonicus W.F. Kirby, 1882; *M. atricornus* MacGillivray, 1893; *M. atricornis* Konow, 1905; *Blennocampa gilletti* Weldon, 1907; *Monophadnoides consobrinus* MacGillivray, 1908; *M. costalis* MacGillivray, 1908; *M. coracinus* MacGillivray, 1908; *M. collaris* MacGillivray, 1908; *M. conspicuus* MacGillivray, 1908; *M. conspersus* MacGillivray, 1908; *M. crassus* MacGillivray, 1908; *M. concessus* MacGillivray, 1908; *Aphanisus nigritus* MacGillivray, 1908; *Aphanisus lenis* Rohwer, 1909; *Monophadnoides costatus* MacGillivray, 1916; *M. kincaidi* MacGillivray, 1923; *M. shawi* MacGillivray, 1923; *M. consonus* MacGillivray, 1923; *M. constitutus* MacGillivray, 1923; *M. corythus* MacGillivray, 1923; *M. curiosus* MacGillivray, 1923; *Paracharactus obversus* MacGillivray, 1923). Host plant: Geum spp., Filipendula spp., Rubus spp. (Rosaceae). Russia: **EP** (NW, C, E), **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (KR, BR). – Europe (WE, EE, NE, SE), Mongolia, N America; introduced into S America.

Monophadnoides ruficruris (Brullé, 1832) [*Selandria*] (*Blennocampa flavigollis* Jacobs, 1884; *Monophadnus dispar* Konow, 1886; *M. ruficruris plagiatus* Enslin, 1914). Host plant: Rubus idaeus (Rosaceae). Russia: **EP** (NC). – Europe (WE, EE, SE).

Monophadnoides tuberculatus Smith et Wei, 2015. Host plant: Rosaceae. Russia: **FE** (PR). – China (NE), Korean Peninsula.

MONOPHADNUS Hartig, 1837 (*Corporaalinus* Forsius, 1925; *Doderia* Malaise, 1935). Type species *Tenthredo albipes* Gmelin, 1790. The larvae without branched spines; feed on Ranunculaceae. The genus is distributed in Holarctic, Oriental and Neotropical regions. Number of species: World – about 25; Palaearctic – about 10; Russia – 3.

Monophadnus monticola (Hartig, 1837) [*Tenthredo*] (*Selandria feriata* Zaddach, 1859; *Monophadnus fuliginipennis* Costa, 1859; *M. fulgidipennis* Dalla Torre, 1894). Russia: **EP** (NW, C, CR). – Europe (WE, EE, NE, SE).

Monophadnus pallescens (Gmelin, 1790) [*Tenthredo*] (*Tenthredo albipes* Gmelin, 1790; *Selandria tiliae* Norton, 1861; *Blennocampa emarginata* Thomson, 1871; *Hoplocampa calceolata* Costa, 1881; *Monophadnus bipunctatus* MacGillivray, 1908; *M. furvus* Benson, 1930). Host plant: Ranunculus spp., Anemone spp. (Ranunculaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (KM, AL), **ES** (IR), **FE** (PR). – Europe (WE, EE, NE, SE), Caucasus,

Turkey, Kyrgyzstan, Kazakhstan; introduced into N America.

Monophadnus spinolae (Klug, 1816) [Tenthredo] (*Hylotoma ventralis* Spinola, 1806; *Tenthredo croceiventris* Klug, 1816; *T. bipartita* Lepeletier, 1823; *T. bipartita* Serville, 1823; *Monophadnus pleuriticus* Costa, 1859; *M. gastricus* Costa, 1859; *Blennocampa scutellaris* André, 1881; *Monophadnus spinolae pectoralis* Konow, 1886; *Periclista nigricarpa* Costa, 1894; *Monophadnus spinolae ruberrima* Enslin, 1914; *M. spinolae rufipictus* Enslin, 1914). Host plant: Clematis spp. (Ranunculaceae). Russia: **EP** (CR). – Europe (WE, EE, SE), Georgia; introduced into New Zealand.

OBTUSIA Malaise, 1964. Type species *Obtusia ussuriensis* Malaise, 1964. The genus is distributed in East Palaearctic. Number of species: one East Asian species.

Obtusia ussuriensis Malaise, 1964. Russia: **FE** (PR).

OKUTANIUS D.R. Smith, 1981. Type species *Okutanius lobatus* D.R. Smith, 1981. The genus is distributed in East Palaearctic. Number of species: World – 2; Palaearctic – 2; Russia – 1.

Okutanius lobatus D.R. Smith, 1981. Russia: **FE** (PR). – Korean Peninsula.

PARACHARACTUS MacGillivray, 1908 (*Dicrostema* Benson, 1952). Type species *Paracharactus obscuratus* MacGillivray, 1908. The genus is distributed in Holarctic. Number of species: World – 7; Palaearctic – 4; Russia – 2.

Paracharactus gracilicornis (Zaddach, 1859) [Selandria]. Host plant: Adoxa spp. (Adoxaceae). Russia: **EP** (NW, C, E), **UR**, **WS** (AL). – Europe (WE, EE, NE, SE), Kyrgyzstan, Kazakhstan.

Paracharactus japonicus (Malaise, 1931) [Neotomostethus]. Russia: **FE** (SA, KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

PARAMASAAKIA Ermolenko, 1971. Type species *Paramasaakia ajnu* Ermolenko, 1971. The genus is distributed in East Asia. Number of species: one species.

Paramasaakia ajnu Ermolenko, 1971. Russia: **FE** (SA). – China (SE).

PAREOPHORA Konow, 1886. Type species *Tenthredo luridiventris* Klug, 1816. The genus is

distributed in Palaearctic. Number of species: World and Palaearctic – 4; Russia – 1.

Pareophora pruni (Linnaeus, 1758) [Tenthredo] (*Tenthredo luridiventris* Klug, 1816; *T. nigripes* Klug, 1816; *Blennocampa bicolor* Tischbein, 1852; *Monophadnus nigripes maculiventris* Costa, 1894). Host plant: *Prunus spinosa*, *P. tenella* (Rosaceae). Russia: **EP** (S, NC, CR). – Europe (WE, EE, NE, SE).

PARNA Benson, 1936. Type species *Tenthredo tenella* Klug, 1816. The larvae in mines on leaves of trees. The genus is distributed in Palaearctic. Number of species: World and Palaearctic – 7; Russia – 2.

Parna apicalis (Brischke, 1888) [Blennocampa] (*P. re-seri* Liston, 1993). Host plant: *Tilia* spp. (Tiliaceae). Russia: **EP** (NW). – Europe (WE, EE, NE).

Parna tenella (Klug, 1816) [Tenthredo] (*Blennocampa tiliae* Kaltenbach, 1874). Host plant: *Tilia* spp. (Tiliaceae). Russia: **EP** (C, E, NC), **FE** (PR). – Europe (WE, EE, NE), Armenia, Japan (Hon).

PERICLISTA Konow, 1886 (*Mogerus* MacGillivray, 1895; *Isodyctium* Ashmead, 1898; *Aphanisus* MacGillivray, 1908; *Neocharactus* MacGillivray, 1908; *Apericlista* Enslin, 1914; *Neoclista* Malaise, 1964). Type species *Tenthredo lineolata* Klug, 1816. The genus is distributed in Holarctic, Oriental and Neotropical regions. Number of species: World – about 55; Palaearctic – about 30; Russia – 4 from 2 subgenera.

Periclista (Periclista) albida (Klug, 1816) [Tenthredo] (*Tenthredo melanocephala* Fabricius, 1798; *T. testacea* Serville, 1823; *T. testacea* Lepeletier, 1823; *Selandria fulvicollis* Stephens, 1835; *S. versicolor* Newman, 1837; *Monophadnus inquilinus* Förster, 1844; *Periclista albida orbitalis* Enslin, 1914). Host plant: *Quercus robur*, *Q. mongolica* (Fagaceae). Russia: **EP** (S, NC, CR), **FE** (KH). – Europe (WE, EE, NE, SE).

Periclista (Periclista) lineolata (Klug, 1816) [Tenthredo] (*Monophadnus tenuicingulatus* Costa, 1859). Host plant: *Quercus robur* (Fagaceae). Russia: **EP** (NW, NC). – Europe (WE, EE, NE, SE).

Periclista (Periclista) pubescens (Zaddach, 1859) [Selandria] (*Periclista pubescens analis* Konow, 1886). Host plant: *Quercus* spp. (Fagaceae). Russia: **EP** (NW, S, CR). – Europe (WE, EE, SE).

Periclista (Neocharactus) albipennis (Zaddach, 1859) [Selandria]. Russia: **EP** (NW). – Europe (WE, EE).

PHYMATOCERA Dahlbom, 1835 (*Pectinia* Brullé, 1846; *Phymatoceros* Konow, 1905; *Hypargyricus* MacGillivray, 1908; *Melanoselandria* MacGillivray, 1908). Type species *Tenthredo aterrima* Klug, 1816. The larvae feed on the monocotyledonous plants (Liliales). The genus is distributed in Holarctic and Oriental region. Number of species: World – 13; Palaearctic – 6; Russia – 2.

Phymatocera aterrima (Klug, 1816) [Tenthredo] (*Tenthredo trichocera* Serville, 1823; *T. trichocera* Lepeletier, 1823; *Selandria robinsoni* Curtis, 1850). Host plant: *Polygonatum* spp. (Asparagaceae), *Convallaria* spp., *Smilacina* spp. (Convallariaceae). Russia: **EP** (NW, C, E, S, NC, CR), **WS** (NS). – Europe (WE, EE, NE, SE), Armenia, Kazakhstan, China (NW), Korean Peninsula.

Phymatocera nipponica Togashi, 1958. Host plant: Liliaceae. Russia: **FE** (SA, KU). – Korean Peninsula, Japan (Hon, Shi, Kyu).

PHYMATOCERIOLA Sato, 1928. Type species *Phymatoceriola suigenensis* Sato, 1928. The genus is distributed in East Asia. Number of species: World and Palaearctic – 4; Russia – 2.

Phymatoceriola suigenensis Sato, 1928. Russia: **ES** (IR), **FE** (KH). – Korean Peninsula.

Phymatoceriola ussuriensis Malaise, 1964. Russia: **FE** (PR).

PHYMATOCEROPTYSIS Rohwer, 1916. Type species *Phymatoceropsis fulvocincta* Rohwer, 1916. The genus is distributed in East Asia and Oriental region. Number of species: World – 9; Palaearctic – 5; Russia – 2.

Phymatoceropsis melanogaster He, Wei et Zhang, 2005. Russia: **FE** (PR). – China (CC, SE).

Phymatoceropsis peregrinator (Malaise, 1931) [Neotomostethus]. Russia: **FE** (KA). – Japan (Hok, Hon).

PROFENUSA MacGillivray, 1914 (*Gunnea* Malaise, 1964). Type species *Profenusa collaris* MacGillivray, 1914. The larvae in mines on leaves of trees. The genus is distributed in Holarctic and Oriental region. Number of species: World – 14; Palaearctic – 8; Russia – 2.

Profenusa pygmaea (Klug, 1816) [Tenthredo]. Host plant: *Quercus* spp., *Castanea sativa* (Fagaceae). Russia: **EP** (NW, C, S, NC). – Europe (WE, EE, NE, SE), Turkey.

Profenusa thomsoni (Konow, 1886) [Fenusia]. Host plant: *Betula* spp. (Betulaceae). Russia: **EP** (NW, C, NC), **ES** (BR), **FE** (AM, KH). – Europe (WE, EE, NE), Armenia, Turkey, Kyrgyzstan, Kazakhstan, Japan (Hok, Hon); introduced into N America.

RHADINOCERAEA Konow, 1886 (*Veratra* D.R. Smith, 1969). Type species *Tenthredo micans* Klug, 1816. The larvae feed on the Liliaceae, Iridaceae, Adoxaceae. The genus is distributed in Holarctic and Oriental region. Number of species: World – about 20; Palaearctic – about 10; Russia – 8 from 2 subgenera.

Rhadinoceraea (Rhadinoceraea) bensoni Beneš, 1961. Host plant: *Lilium* spp. (Liliaceae). Russia: **EP** (Zhelochovtsev, Zinovjev, 1996), **ES** (IR). – Europe (WE, EE, SE).

Rhadinoceraea (Rhadinoceraea) caucasica Dovnar-Zapolskij, 1930. Russia: **EP** (NC).

Rhadinoceraea (Rhadinoceraea) micans (Klug, 1816) [Tenthredo] (*Monophadnus iridis* Kaltenbach, 1862). Host plant: *Iris* spp. (Iridaceae). Russia: **EP** (NW, C, E), **UR**, **WS** (AL). – Europe (WE, EE, NE, SE).

Rhadinoceraea (Rhadinoceraea) reitteri Konow, 1890. Host plant: *Iris* spp. (Iridaceae). Russia: **EP** (C). – Europe (WE, EE, SE).

Rhadinoceraea (Rhadinoceraea) sachalinensis Ermolenko, 1971. Russia: **FE** (SA).

Rhadinoceraea (Veratra) nodicornis Konow, 1886 (*Rh. heydeni* Konow, 1887). Host plant: *Veratrum* spp. (Liliaceae). Russia: **EP** (C, S), **UR**, **WS** (KM, AL), **ES** (KS, KR, IR, BR). – Europe (WE, EE, SE), China (NW).

Rhadinoceraea (Veratra) subarctica Hellén, 1948. Russia: **EP** (N). – Europe (NE).

Rhadinoceraea sibiricola Zhelochovtsev, 1939. Russia: **ES** (KR).

SCOLIONEURA Konow, 1890. Type species *Tenthredo betuleti* Klug, 1816. The genus is distributed in Holarctic, mainly in West Palaearctic; one species was introduced into N America. Number of species: World and Palaearctic – 4; Russia – 1.

Scolioneura betuleti (Klug, 1816) [Tenthredo] (*Tenthredo nigricans* Klug, 1818; *Fenusia betulae* Zaddach, 1859). Host plant: *Betula* spp. (Betulaceae). Russia: **EP** (NW, C, E), **UR**, **WS** (AL), **ES** (TU, IR, BR). – Europe (WE, EE, NE, SE), Armenia,

Kyrgyzstan, Kazakhstan, Mongolia; introduced into N America.

SENOCLIDEA Rohwer, 1912. Type species *Senoclidea amala* Rohwer, 1912. The genus is distributed in East Palaearctic, Oriental and Australian regions. Number of species: World – 17; Palaearctic – 6; Russia – 1.

Senoclidea koreana (Konow, 1898) [Monophadnus]. Russia: **FE** (PR). – Korean Peninsula, China (Wei et al., 2006).

STETHOMOSTUS Benson, 1939. Type species *Tenthredo fuliginosa* Schrank, 1781. The larvae feed on Ranunculus (Ranunculaceae). The genus is distributed in Palaearctic and Oriental region; one species was introduced into North America. Number of species: World – 6; Palaearctic – 4; Russia – 3.

Stethomostus flavigollaris (Sato, 1928) [Atomostethus]. Host plant: Ranunculus spp. (Ranunculaceae). Russia: **ES** (BR), **FE** (AM, PR). – China (NE), Korean Peninsula.

Stethomostus fuliginosus (Schrank, 1781) [Tenthredo] (*Tenthredo fusca* Serville, 1823; *T. fusca* Lepeletier, 1823; *Selandria geniculata* Stephens, 1835; *Blennocampa croceipes* Costa, 1864; *Monophadnus fukaii* Rohwer, 1910). Host plant: Ranunculus spp. (Ranunculaceae), Kerria spp. (Rosaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (KM). – Europe (WE, EE, NE, SE), Turkey, Kazakhstan, Japan (Hon, Shi); introduced into N America.

Stethomostus funereus (Klug, 1816) [Tenthredo]. Host plant: Ranunculus spp. (Ranunculaceae). Russia: **EP** (N, NW, C, E), **UR**, **ES** (BR). – Europe (WE, EE, NE), Turkey, Kazakhstan.

TOMOSTETHUS Konow, 1886. Type species *Tenthredo nigrita* Fabricius, 1804. The larvae feed on the woody plants. The genus is distributed in Holarctic, Oriental and Neotropical regions. Number of species: World – 8; Palaearctic – 6; Russia – 3.

Tomostethus juglans Takeuchi, 1929. Host plant: Juglans spp. (Juglandaceae). Russia: **FE** (PR). – Korean Peninsula.

Tomostethus nigritus (Fabricius, 1804) [Tenthredo] (*Tenthredo nigerrima* Klug, 1816; *T. brevicornis* Klug, 1816; *T. fraxini* Serville, 1823; *T. fraxini* Lepeletier, 1823; *Monophadnus latus* Costa, 1894). Host

plant: Fraxinus spp. (Oleaceae). Russia: **EP** (NW, C, S, NC), **ES** (BR), **FE** (PR, SA). – Europe (WE, EE, NE, SE), N Africa, Caucasus, Turkey, China (NE, CC), Korean Peninsula, Japan (Hon).

Tomostethus veles Konow, 1900. Russia: **ES** (IR).

ZAPHYMATOCERA Sato, 1928. Type species *Zaphymatocera typica* Sato, 1928. The genus is distributed in East Asia. Number of species: World and Palaearctic – 2; Russia – 1.

Zaphymatocera typica Sato, 1928. Russia: **FE** (KU, KA). – Korean Peninsula.

Subfamily NEMATINAE

ANOPLONYX Marlatt, 1896 (*Marlattia* Ashmead, 1898). Type species *Nematus lariciphagus* Zaddach, 1883. The larvae feed on Larix (Pinaceae). The genus is distributed in Holarctic. Number of species: World – 12; Palaearctic – 8; Russia – 7.

Anoplonyx apicalis (Brischke, 1883) [Leptopus] (*Leptopus curinthisicus* Zaddach, 1883; *Camponiscus braunsi* Konow, 1884; *C. tscheki* Rogenhofer et Kohl, 1885; *C. carinthiacus* Dalla Torre, 1894). Host plant: Larix spp. (Pinaceae). Russia: **EP** (NW), **UR**, **ES** (KR, IR, BR, YA, ZB), **FE** (AM). – Europe (WE, EE, NE), Mongolia.

Anoplonyx bilineatus Verzhutskii, 1981. Host plant: Larix spp. (Pinaceae). Russia: **ES** (TU, IR, YA, ZB), **FE** (MG). – Mongolia.

Anoplonyx destructor Benson, 1952. Host plant: Larix spp. (Pinaceae). Russia: **EP** (NW). – Europe (WE, EE, NE), Korean Peninsula, Japan (Hon).

Anoplonyx lariciphagus (Zaddach, 1883) [Nematus] (*Camponiscus luridus* Konow, 1884; *Platycampus pectoralis obscurus* Enslin, 1915). Host plant: Larix spp. (Pinaceae). Russia: **ES** (KR, IR, BR, ZB). – Europe (WE, EE), Korean Peninsula.

Anoplonyx orientis D.R. Smith, 1988. Host plant: Larix leptolepis, L. kurilensis (Pinaceae). Russia: **FE** (KU). – Japan (Hok, Hon).

Anoplonyx ovatus (Zaddach, 1883) [Leptopus]. Host plant: Larix spp. (Pinaceae). Russia: **EP** (NW), **ES** (YA, ZB). – Europe (WE, EE, NE).

Anoplonyx versicolor Vikberg, 1975 (*A. ornatus* Verzhutskii, 1981). Host plant: Larix spp. (Pinaceae). Russia: **EP** (NW), **ES** (IR, BR, YA), **FE** (AM). – Europe (NE), Mongolia, Korean Peninsula.

CLADIUS Illiger, 1807 (*Priophorus* Dahlbom, 1835; *Trichiocampus* Hartig, 1837; *Stevenia* Brullé, 1846;

Eudryas Gistel, 1848; *Prionophorus* Agassiz, 1848). Type species *Tenthredo difformis* Panzer, 1799. The larvae feed on the woody plants. The genus is distributed in Holarctic and Oriental region; one species was introduced into Neotropical and Australasian regions. Number of species: World – about 45; Palaearctic - > 35; Russia – 15 from 3 subgenera.

Cladius (Cladius) pectinicornis (Geoffroy, 1785) [Tenthredo] (*Tenthredo alces* Thunberg, 1789; *T. difformis* Panzer, 1799; *Cladius geoffroyi* Serville, 1823; *C. morio* Serville, 1823; *C. geoffroyi* Lepeletier, 1823; *C. morio* Lepeletier, 1823; *Nematus crassicornis* Stephens, 1835; *Cladius isomera* Norton, 1861; *C. ramicornis* André, 1880; *C. gracilicornis* Konow, 1884; *C. crassicornis* Konow, 1884; *C. comari* Stein, 1886; *C. hyalinopterus* Konow, 1886; *C. palmicornis* Konow, 1892; *C. ordubadensis* Konow, 1892; *C. major* Cobelli, 1892; *C. orientalis* Cameron, 1902; *C. tibialatus* Konow, 1906). Host plant: Rosa spp., Fragaria spp., Potentilla spp., Filipendula spp. (Rosaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, KM, AL), **ES** (IR, BR, YA, ZB), **FE** (AM, PR, SA, KA, MG). – Europe (WE, EE, NE, SE), Georgia, Azerbaijan, SW Asia, Afghanistan, Central Asia, Kazakhstan, China (NE, NC), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu), India (NW), N America.

Cladius (Priophorus) brullei (Dahlbom, 1835) [Priophorus] (*C. immunis* Stephens, 1835; *Nematus melanostigma* Stephens, 1835; *Cladius tener* Zaddach, 1859; *C. tristis* Zaddach, 1859; *C. parvus* Zaddach, 1883; *Priophorus rubivorus* Rohwer, 1922; *P. montanus* Rohwer, 1922; *P. rubi* Rohwer, 1922; *P. foveivaginatus* Malaise, 1931). Host plant: Rubus spp. (Rosaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (AL), **ES** (KR, IR, ZB), **FE** (PR, SA, KA). – Europe (WE, EE, NE, SE), Central Asia, China (NC), Korean Peninsula, Japan (Hok, Hon), India, SE Asia, N America; introduced into S America, Australia, New Zealand.

Cladius (Priophorus) compressicornis (Fabricius, 1804) [Tenthredo] (*Tenthredo albipes* Fallén, 1808; *Cladius pallipes* Lepeletier, 1823; *C. pallipes* Serville, 1823; *C. albipes* Hartig, 1837; *C. aequalis* Norton, 1872; *Priophorus discors* Konow, 1894; *Cladius solitarius* Dyar, 1895; *Priophorus padi collaris* Enslin, 1915; *P. modestius* MacGillivray, 1921; *P. munditus* MacGillivray, 1921; *P. crataegi* Rohwer, 1922; *P. pruni* Rohwer, 1922; *P. plesius* Rohwer, 1922;

P. virginianus Rohwer, 1922). Host plant: Salix spp., Populus spp. (Salicaceae), Rubus spp., Prunus spp. (Rosaceae), and others. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **ES** (KR, IR, YA, ZB), **FE** (KH, PR, SA, KA, MG). – Europe (WE, EE, NE, SE), Caucasus, Turkey, Iran, Kyrgyzstan, Kazakhstan, China (CC), Korean Peninsula, Japan (Hok, Hon), N America.

Cladius (Priophorus) corniger Zhelochovtsev, 1952. Russia: **ES** (ZB). – China (Wei et al., 2006).

Cladius (Priophorus) hyalopterus (Jakovlev, 1891) [Priophorus] (*Priophorus triangulifer* Konow, 1902). Russia: **EP** (E), **UR**, **WS** (KM), **ES** (IR, ZB), **FE** (AM). – Kyrgyzstan, Kazakhstan, Mongolia, China (NW, NC).

Cladius (Priophorus) nubilus (Konow, 1897) [Trichiocampus] (*Priophorus dellei* Conde, 1935). Russia: **EP** (E), **UR**, **WS** (AL), **ES** (IR), **FE** (AM, PR, KU). – China (NE).

Cladius (Priophorus) rufipes Serville, 1823 (*C. rufipes* Lepeletier, 1823; *C. uncinnata* Hartig, 1837; *C. discrepans* Costa, 1859; *Trichiocampus garbiglietii* Costa, 1864; *Priophorus phaeopterus* Costa, 1894). Host plant: Ulmus spp. (Ulmaceae). Russia: **EP** (C, S, NC). – Europe (WE, EE, NE), Turkey, Iran.

Cladius (Trichiocampus) aeneus Zaddach, 1859. Host plant: Salix spp., Populus spp. (Salicaceae). Russia: **EP** (N, NW), **ES** (KR, IR, YA). – Europe (EE, NE).

Cladius (Trichiocampus) cannabis (Xiao et Huang, 1986) [Trichiocampus]. Host plant: Cannabis sativa (Cannabaceae). Russia: **EP** (E), **ES** (IR), **FE** (AM). – China (NC, CC).

Cladius (Trichiocampus) grandis (Serville, 1823) [Nematus] (*Tenthredo viminalis* Fallén, 1808; *Nematus grandis* Lepeletier, 1823; *Cladius eucera* Bouché, 1834; *C. luteicornis* Stephens, 1835; *C. luteiventris* Dahlbom, 1836; *Aulacomerus lutescens* Lintner, 1888; *Platycampus victoria* MacGillivray, 1920). Host plant: Populus spp., Salix spp. (Salicaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (Lelej, Taeger, 2007), **ES** (IR, BR, YA), **FE** (MG). – Europe (WE, EE, NE, SE), Georgia, Armenia, Turkey, Kazakhstan; introduced into N America.

Cladius (Trichiocampus) pilicornis Curtis, 1833. Host plant: Crataegus spp. (Rosaceae). Russia: **ES** (YA). – Europe (WE, EE, NE, SE).

Cladius (Trichiocampus) populi (Okamoto, 1912) [Trichiocampus]. Host plant: Populus spp. (Salicaceae). Russia: **FE** (SA). – Japan (Hok, Hon).

Cladius (Trichiocampus) rufus Verzhutskii, 1966 [*Trichiocampus pseudoviminalis* Huang et Wang,

- 1993). Host plant: *Populus* spp. (Salicaceae). Russia: **ES** (IR), **FE** (AM). – China (NE).
- Cladius (Trichiocampus) takeuchii** Liston, Taeger et Blank, 2009 (*Trichiocampus pruni* Takeuchi, 1956). Host plant: *Prunus* spp. (Rosaceae). Russia: **FE** (KU). – China (SE), Japan (Hok, Hon), SE Asia.
- Cladius (Trichiocampus) ulmi** (Linnaeus, 1758) [Tenthredo] (*C. eradiatus* Hartig, 1837; *C. drewseni* Thomson, 1871; *Priophorus distinguendus* Enslin, 1915; *P. laevifrons* Benson, 1936). Host plant: *Ulmus* spp. (Ulmaceae). Russia: **EP** (C, NC, CR), **WS** (AL). – Europe (WE, EE, NE, SE), Armenia.
- DINEURA** Dahlbom, 1835 (*Leachia* Brullé, 1846; *Dinevra* Agassiz, 1848; *Varna* Ross, 1937). Type species *Tenthredo degeeri* Klug, 1817. The larvae feed on the woody plants. The genus is distributed in Holarctic. Number of species: World and Palaearctic – 9; Russia – 5.
- Dineura parcivalvis** (Konow, 1901) [Pontania]. Russia: **ES** (IR). – Europe (NE).
- Dineura pullior** Schmidt et Walter, 1995. Russia: **EP** (N). – Europe (NE).
- Dineura stilata** (Klug, 1816) [Tenthredo] (*Tenthredo mediocris* Lepeletier et Serville, 1828; *Nematus apicalis* Stephens, 1835; *Selandria scapularis* Stephens, 1835; *Nematus xanthocerus* Hartig, 1840; *Dineura flaveola* Eversmann, 1847; *Nematus posticus* Förster, 1854; *N. xanthopus* Zaddach, 1876; *N. v-flavum* Cameron, 1882; *Dineura melanoxantha* Zaddach, 1883; *D. sulcifrons* Konow, 1885; *D. nigroflava* Magretti, 1886; *Pristiphora oblita* Costa, 1894; *Pontania ancilla* Konow, 1901; *Dineura stilata virilis* Enslin, 1918). Host plant: *Sorbus* spp., *Crataegus* spp., *Cotoneaster* spp. (Rosaceae). Russia: **EP** (E). – Europe (WE, EE, NE, SE).
- Dineura testaceipes** (Klug, 1816) [Tenthredo] (*D. ventralis* Zaddach, 1859; *D. nigra* Zaddach, 1883; *D. testaceipes nigriventris* Enslin, 1915). Host plant: *Sorbus* spp., *Crataegus* spp., *Cotoneaster* spp. (Rosaceae). Russia: **EP** (NW, C, E), **ES** (IR, BR, YA), **FE** (KH, PR, KA). – Europe (WE, EE, NE).
- Dineura virididorsata** (Retzius, 1783) [Tenthredo] (*Tenthredo nigricans* Christ, 1791; *T. degeeri* Klug, 1817; *Nematus varius* Lepeletier, 1823; *N. varius* Serville, 1823; *Dineura hartigii* Gimmerthal, 1844; *D. virididorsata dorsalis* Enslin, 1915). Host plant: *Betula* spp. (Betulaceae). Russia: **EP** (N, NW, C, E), **UR**, **ES** (IR, BR, YA), **FE** (AM, PR, SA, KA). – Europe (WE, EE, NE), China (Wei et al., 2006), Japan (Hok).
- ENDOPHYTUS** Hering, 1934 (*Neopelmatopus* Conde, 1934; *Verna* Kaisila, 1960; *Verna* Kontuniemi, 1960). Type species *Pelmatopus anemones* Hering, 1924. The larvae feed on *Anemone nemorosa* (Ranunculaceae). The genus is distributed in West Palaearctic. Number of species: one species.
- Endophytus anemones** (Hering, 1924) [*Pelmatopus*] (*Pelmatopus tenuiserra* Lindqvist, 1949). Host plant: *Anemone nemorosa* (Ranunculaceae). Russia: **EP** (NW). – Europe (WE, EE, NE).
- EUURA** Newman, 1837 (*Cryptocampus* Hartig, 1837; *Evura* Agassiz, 1848; *Pontania* Costa, 1852; *Epitactus* Förster, 1854; *Amauronematus* Konow, 1890; *Holcocneme* Konow, 1890; *Pachynematus* Konow, 1890; *Brachycolus* Konow, 1895; *Holcocnema* Schulz, 1906; *Holcocnemis* Konow, 1907; *Pteronidea* Rohwer, 1911; *Pontopristia* Malaise, 1921; *Brachycolumna* Strand, 1929; *Decanematus* Malaise, 1931; *Pikonema* Ross, 1937; *Phyllocolpa* Benson, 1960; *Eitelius* Kontuniemi, 1966; *Gemmura* E.L. Smith, 1968; *Ribinematus* Kontuniemi, 1975; *Eupontania* Zinovjev, 1985; *Larinematus* Zhelochovtsev, 1988; *Polynematus* Zhelochovtsev, 1988; *Bacconematus* Zhelochovtsev, 1988; *Alpinematus* Lacourt, 1996; *Epacenematus* Lacourt, 1998; *Kontuniemiana* Lacourt, 1998; *Lindqvistia* Lacourt, 1998; *Luea* Wei et Nie, 1998; *Tubpontania* Vikberg, 2010). Type species *Euura gallae* Newman, 1837. The genus is distributed in Holarctic and Oriental region; introduced into the Neotropical, Ethiopian and Australasian regions. Number of species: World – about 450; Palaearctic – about 300; Russia – 258 (+ 1 ssp.).
- Euura abnormis** (Holmgren, 1883) [*Nematus*] (*Amauronematus tolli* Konow, 1907; *A. aulatus* MacGillivray, 1919). Russia: **EP** (N), **WS** (TM), **ES** (KR, YA). – Europe (WE, NE), N America.
- Euura aborigensis** (Vikberg, 2003) [*Eupontania*]. Host plant: *Salix dshugdshurica* (Salicaceae). Russia: **FE** (MG).
- Euura acuta** (Lindqvist, 1970) [*Amauronematus*]. Host plant: *Dasiphora* spp. (Rosaceae). Russia: **ES** (IR, ZB), **FE** (MG).
- Euura acutifoliae** (Zinovjev, 1985) [*Pontania*] (*Pontania acutifoliae daphnoides* Zinovjev, 1993; *Eupontania acutifoliae baltica* Vikberg et Zinovjev, 2006). Host plant: *Salix acutifolia*, *S. daphnoides* (Salicaceae). Russia: **EP** (N, NW, C, E, S). – Europe (WE, EE, NE).

- Euura acutiserra** (Lindqvist, 1949) [Pontania]. Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (N). – Europe (WE, NE).
- Euura aeger** (Konow, 1895) [Amauronematus] (*Amauronematus dalecarlicus* Malaise, 1921). Russia: **ES** (Zhelochovtsev, Zinovjev, 1995). – Europe (WE, EE, NE).
- Euura albipennis** (Hartig, 1837) [Nematus] (*Nematus vernalis* Hartig, 1840; *N. obscuratus* Zaddach, 1883; *Polynematus legirupus* Konow, 1903; *P. sannio* Konow, 1903). Host plant: *Polygonum* spp. (Polygonaceae). Russia: **EP** (NW, C, E, S), **WS** (TK), **ES** (IR), **FE** (AM). – Europe (WE, EE, NE), Kazakhstan, Korean Peninsula.
- Euura albopicta** (Malaise, 1931) [Pontania]. Host plant: *Salix* spp. (Salicaceae). Russia: **FE** (KA).
- Euura alpicola** (Konow, 1895) [Amauronematus]. Host plant: *Salix phylicifolia* (Salicaceae). Russia: **EP** (NW). – Europe (WE).
- Euura amentorum** (Förster, 1854) [Nematus] (*Nematus suavis* Ruthe, 1859; *N. laevigatus* Zaddach, 1883; *Pontania atra* Marlatt, 1896; *P. maura* Rohwer, 1908; *P. amentivora* Rohwer, 1915; *Pristiphora amentorum nigripleuris* Enslin, 1916; *Pontopristia kamtchatica* Malaise, 1931). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (Taeger et al., 2006), **FE** (KA, CH). – Europe (WE, EE, NE), Greenland, N America.
- Euura amerinae** (Linnaeus, 1758) [Cynips] (*Tenthredo salicispentandrae* Retzius, 1783; *Tenthredo gallarum* Latreille, 1804; *T. saliceti* Fallén, 1808; *Nematus pentandrae* Dahlbom, 1835; *Nematus populi* Hartig, 1837; *N. medullarius* Hartig, 1837; *N. impunctatus* Herrich-Schäffer, 1840; *N. buccatus* Thomson, 1863; *Pontania joergensei* Strand, 1908). The larvae form a nutlike gall on the branches of *Salix pentandra* (Salicaceae). Russia: **EP** (N, NW, C, E), **UR**, **WS** (Zhelochovtsev, Zinovjev, 1995), **ES** (YA), **FE** (KA, MG). – Europe (WE, EE, NE, SE).
- Euura ampla** (Konow, 1895) [Amauronematus]. Host plant: *Betula* spp. (Betulaceae). Russia: **EP** (N, NW, C, E), **UR**, **WS** (Zhelochovtsev, Zinovjev, 1995), **ES** (IR, ZB). – Europe (WE, EE, NE).
- Euura amurensis** (Zinovjev, 1985) [Pontania]. Host plant: *Salix miyabeana* (Salicaceae). Russia: **ES** (ZB), **FE** (AM, KH, PR).
- Euura annulata** (Gimmerthal, 1834) [Nematus] (*Nematus xanthopterus* Dahlbom, 1835; *N. flavipennis* Cameron, 1876; *Pachynematus freyi* Lindqvist, 1949; *P. lacteipennis* Benson, 1963). Host plant: Rumex spp. (Polygonaceae). Russia: **EP** (E, NC), **ES** (IR, BR), **FE** (KH, SA). – Europe (WE, EE, NE), Central Asia.
- Euura anomaloptera** (Förster, 1854) [Nematus] (*Pontania joergensei* Enslin, 1916; *Amauronematus maidli* Zirngiebl, 1937; *Pontania cyrnea* Liston, 2005). Host plant: *Salix carpea* (Salicaceae). Russia: **EP** (N, NW). – Europe (WE, EE, NE, SE).
- Euura arbusculae** (Benson, 1941) [Pontania]. Host plant: *Salix arbuscula* (Salicaceae). Russia: **EP** (N). – Europe (WE).
- Euura arctica** MacGillivray, 1919 (*Pontania deminuta* MacGillivray, 1919; *P. delicatula* MacGillivray, 1919; *Euura abortiva* MacGillivray, 1919). Host plant: *Salix reticulata* (Salicaceae). Russia: **ES** (YA), **FE** (CH). – N America.
- Euura arcticornis** (Konow, 1904) [Pontania] (*Pontania phylicifoliae* Forsius, 1919; *P. viminalis hepaticulae* Malaise, 1920). Host plant: *Salix phyticifolia*, *S. lapponicum* (Salicaceae). Russia: **EP** (N, NW, E), **UR**, **WS** (TM, AL), **ES** (KR, IR, ZB, BR, YA), **FE** (AM, PR, MG). – Europe (WE, EE, NE), Korean Peninsula.
- Euura atra** (Jurine, 1807) [Pteronus] (*Nematus angustus* Hartig, 1837; *E. salicicola* E.A. Smith, 1879; *Cryptocampus helveticus* Zaddach, 1883; *C. robustus* Zaddach, 1883; *Euura nigra* Provancher, 1888). Host plant: *Salix* spp. (Salicaceae); larvae in galls on shoots. Russia: **EP** (NC), **ES** (IR, BR, YA, ZB), **FE** (MG). – Europe (WE, EE, NE, SE), Armenia, Cyprus, Kyrgyzstan, Kazakhstan; introduced into N America.
- Euura auritae** Kopalke, 2000 (*E. cinereae* Kopalke, 1996; *E. lapponica* Kopalke, 1996; *E. salicislapponicae* Kopalke, 2014). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (N). – Europe (WE, EE, NE).
- Euura bergmani** (Malaise, 1931) [Amauronematus] (*Pteronidea kamtchatica* Lindqvist, 1971). Russia: **FE** (KA).
- Euura bergmanni** (Dahlbom, 1835) [Nematus] (*Nematus virescens* Hartig, 1837; *N. pallicarpus* Hartig, 1837; *N. curtispina* Thomson, 1871; *N. varipictus* Holmgren, 1883; *N. anthophilus* Zaddach, 1884; *N. pallidicarpus* Dalla Torre, 1894; *Amauronematus longicornis* Konow, 1897; *Pteronus curtispinus* Konow, 1903; *Amauronematus trigonicus* Konow, 1905; *Lygaeonematus pallens* Enslin, 1916; *Pteronidea curtispina luctuosa* Enslin, 1916; *P. vernalis* Lindqvist, 1937). Host plant: *Salix* spp. (Salicaceae); up to 5 generations per year. Russia: **EP** (N, NW,

- C, E, S, NC), **UR, ES** (IR, YA). – Europe (WE, EE, NE, SE), Kyrgyzstan.
- Euura bertilpoppii** Heibo et Liston, 2014 (*Amauronematus poppii* Konow, 1904). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (N, NW), **ES** (KR), **FE** (KA). – Europe (NE).
- Euura betulae** (Schmidt, 1997) [*Amauronematus*]. Host plant: *Betula* spp. (Betulaceae). Russia: **FE** (MG). – Europe (NE).
- Euura bicolor** (Jakovlev, 1891) [*Amauronematus*]. Russia: **ES** (IR), **FE** (AM).
- Euura bigallae** (Vikberg et Zinovjev, 2014) [*Pontania*]. Host plant: *Salix* spp. (Salicaceae); larvae in galls on shoots. Russia: **EP** (N). – Europe (NE).
- Euura bipartita** (Serville, 1823) [*Nematus*] (*Nematus bipartitus* Lepeletier, 1823; *N. aurantiacus* Hartig, 1837; *N. confusus* Förster, 1854; *N. xanthogaster* Förster, 1854; *N. ochropus* Thomson, 1863; *Pachynematus albipennis salicicola* Enslin, 1916). Host plant: *Salix* spp., *Populus* spp. (Salicaceae). Russia: **EP** (N, NW, C, E, S), **UR, WS** (TM, OM). – Europe (WE, EE, NE, SE), Kazakhstan.
- Euura bohemani** (Thomson, 1871) [*Nematus*] (*Pteronidea pseudonotabilis* Enslin, 1916; *P. pseudonotabilis simplex* Forsius, 1921; *P. pseudonotabilis t-magnum* Malaise, 1931). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (N, C), **FE** (KA). – Europe (EE, NE).
- Euura boreophila** (Lindqvist, 1971) [*Pteronidea*]. Host plant: *Salix* spp. (Salicaceae). Russia: **ES** (IR, BR). – Europe (NE).
- Euura borisi** (Vikberg, 2003) [*Eupontania*]. Host plant: *Salix rhamnifolia* (Salicaceae). Russia: **ES** (BR).
- Euura brevicera** Taeger et Blank, 2014 (*Pteronidea brachycera* Lindqvist, 1975). Host plant: *Salix* spp. (Salicaceae). Russia: **ES** (IR).
- Euura breviseta** (Lindqvist, 1949) [*Pteronidea*]. Host plant: *Betula* spp. (Betulaceae). Russia: **EP** (N). – Europe (NE).
- Euura brevivalva** (Thomson, 1871) [*Nematus*] (*Pteronidea kriegeri* Konow, 1903; *Amauronematus spurcus* Konow, 1904; *Pteronidea assimilis* Lindqvist, 1949). Host plant: *Betula* spp. (Betulaceae). Russia: **EP** (N, NW), **ES** (IR, ZB), **FE** (AM, MG). – Europe (WE, NE).
- Euura bridgmanii** (Cameron, 1883) [*Nematus*] (*Pontania kirchneri* Zirngiebl, 1959). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (Zhelochovtsev, Zinovjev, 1995), “Siberia” (Zhelochovtsev, Zinovjev, 1995). – Europe (WE, NE).
- Euura brunneipicta** (Lindqvist, 1975) [*Amauronematus*]. Russia: **ES** (IR).
- Euura cadderensis** (Cameron, 1875) [*Nematus*] (*Pteronidea macroserrata* Lindqvist, 1943). Host plant: *Betula* spp. (Betulaceae), *Salix* spp. (Salicaceae). Russia: **EP** (N, E), **UR**. – Europe (WE, EE, NE).
- Euura caeruleocarpa** Hartig, 1837 (*Nematus propinquus* Dahlbom, 1835; *N. sulcipes* Hartig, 1837; *N. brevispinis* Förster, 1854; *N. brachyacanthus* Thomson, 1863; *N. insubricus* Cobelli, 1892). Host plant: *Salix* spp., *Populus* spp. (Salicaceae), *Paeonia*ceae, *Ranunculaceae*. Russia: **EP** (NW, C, NC), **ES** (IR, BR, YA, ZB), **FE** (KH, PR, MG). – Europe (WE, EE, NE), China (Wei et al., 2006), Japan (Hok, Hon).
- Euura camtschaticalis** (Enslin, 1927) [*Amauronematus*]. Russia: **FE** (KA).
- Euura camtschatica** (Forsius, 1928) [*Pontania*]. Host plant: *Salix* spp. (Salicaceae). Russia: **FE** (KA).
- Euura capito** (Konow, 1903) [*Pteronus*]. Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (N, C), **UR, FE** (KA). – Europe (WE, EE, NE, SE), Kazakhstan, China (NW).
- Euura carelica** (Hellén, 1948) [*Nematus*] (*Pteronidea winteri* Lindqvist, 1958). Russia: **EP** (N). – Europe (NE).
- Euura caucasica** (Liston, Taeger et Blank, 2009) [*Pachynematus*] (*Pachynematus alticola* Muche, 1973). Russia: **EP** (NC). – Georgia, Mongolia.
- Euura caudalis** (Eversmann, 1847) [*Nematus*]. Russia: **EP** (E).
- Euura clibrichella** (Cameron, 1878) [*Nematus*] (*Nematus thomsoni* Cameron, 1882; *Pachynematus pubescens* Marlatt, 1896; *P. gotarus* Kincaid, 1900; *P. venustus* MacGillivray, 1921; *P. vernus* MacGillivray, 1921). Host plant: *Carex* spp. (Cyperaceae). Russia: **EP** (N). – Europe (WE, NE), N America.
- Euura clitellata** (Serville, 1823) [*Nematus*] (*Nematus clitellatus* Lepeletier, 1823; *N. kirbyi* Dahlbom, 1835; *N. leucocarpus* Hartig, 1840; *N. einersbergensis* Hartig, 1840; *N. flaviventris* Hartig, 1840; *N. palliceps* Hartig, 1840; *N. diaphanus* Eversmann, 1847; *N. umbripennis* Eversmann, 1847; *N. griseus* Eversmann, 1847; *Epitactus praecox* Förster, 1854; *Nematus trisignatus* Förster, 1854; *N. microps* Förster, 1854; *N. pectoralis* Snellen van Vollenhoven, 1867; *N. turgidus* Zaddach, 1876; *N. leucocarpus* André, 1880; *N. valesiacus* W.F. Kirby, 1882; *N. tener* Zaddach, 1884; *N. pallidiceps* Dalla Torre, 1894; *Pachynematus infumatus* Marlatt, 1896; *P. zadachi* Konow, 1901; *P. latus* Konow, 1903; *P. glesipennis* Konow, 1903; *P. foveolatus* Konow, 1903; *P. clitellatus transigens* Enslin, 1916; *P. clitellatus*

flavissimus Enslin, 1916; *P. refractarius* MacGillivray, 1921; *P. truncatus* Benson, 1948). Host plant: Poaceae. Russia: **EP** (NW, C, E, S, NC), **UR**, **WS** (Zhelochovtsev, 1988), **ES** (IR, BR), **FE** (SA, MG). – Europe (WE, EE, NE), Kazakhstan, China (Wei et al., 2006), Korean Peninsula, Japan (Hok).

Euura clypeata (Lindqvist, 1975) [Amauronematus]. Russia: **ES** (IR).

Euura collectanea (Förster, 1854) [Nematus] (*Nematus deficiens* Förster, 1854; *N. crassipes vacciniellus* Cameron, 1876; *N. xanthostylos* Zaddach, 1884; *Pontania viminalis lugubris* Enslin, 1918; *P. collectanea relicta* Weiffenbach, 1962; *Eupontania collectanea rosmarinifolia* Vikberg et Zinovjev, 2006). Host plant: *Salix repens* repens, *S. repens rosmarinifolia* (Salicaceae). Russia: **EP** (NW). – Europe (WE, EE, NE).

Euura compacta (Bogacheva, 1977) [Amauronematus]. Russia: **WS** (TM).

Euura cornuta (Lindqvist, 1962) [Amauronematus]. Russia: **FE** (MG). – Europe (NE).

Euura crassidens (Lindqvist, 1975) [Amauronematus]. Russia: **ES** (IR).

Euura crassipes (Thomson, 1871) [Nematus] (*Pontania lapponica* Malaise, 1920; *P. enslini* Zirngiebel, 1937). Host plant: *Salix herbacea*, *S. lapporum*, *S. polaris* (Salicaceae); larvae in galls on the central vein of the leaf. Russia: **EP** (N, C), **WS** (TM), **ES** (IR, YA), **FE** (KU, MG). – Europe (WE, NE).

Euura crassispina (Thomson, 1871) [Nematus]. Host plant: *Salix glauca*, *S. arctica*, *S. sphenophylla* (Salicaceae). Russia: **ES** (YA), **FE** (KA, MG). – Europe (NE).

Euura dentata (Lindqvist, 1937) [Pachynematus]. Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (NW). – Europe (NE).

Euura dispar (Zaddach, 1876) [Nematus]. Host plant: *Betula* spp. (Betulaceae). Russia: **EP** (N, NW, C, E), **ES** (IR). – Europe (WE, EE, NE).

Euura distinguenda (Enslin, 1915) [Amauronematus]. Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (NW). – Europe (WE, EE, NE).

Euura dolichura (Thomson, 1871) [Nematus] (*Nematus femoralis* Cameron, 1876; *Pontania Robbinsi* Benson, 1935; *P. lapponicola* Kopalke, 1994). Host plant: *Salix lapporum*, *S. phylicifolia*, ?*S. iliensis* (Salicaceae); larvae form galls. Russia: **EP** (N, NC), **WS** (AL), **ES** (KR, IR, BR, YA), **FE** (PR, KA, MG). – Europe (WE, EE, NE, SE), Kyrgyzstan, N America.

Euura duplex (Serville, 1823) [Pristiphora] (*Pristiphora duplex* Lepeletier, 1823). Host plant: Poaceae.

Russia: **EP** (NW), **UR**, **ES** (KR, IR, BR, YA, ZB), **FE** (AM). – Europe (WE, EE, NE), Mongolia.

Euura edentata (Lindqvist, 1975) [Pachynematus]. Russia: **ES** (IR).

Euura edenticulata (Lindqvist, 1970) [Amauronematus]. Russia: **ES** (IR).

Euura eiteli (Saarinen, 1948) [Amauronematus]. Russia: **EP** (N), **ES** (ZB). – Europe (NE).

Euura elaeagnocola (Kopalke, 1994) [Pontania] (*Nematus parallelus* Kriechbaumer, 1876). Host plant: *Salix elaeagnos*, *S. lapponicum*, *S. alaxensis* (Salicaceae). Russia: **FE** (MG). – Europe (WE).

Euura elbrus Sundukov, 2017 (*Nematus nivalis* Muche, 1973). Russia: **EP** (NC). – Georgia.

Euura pilosa (Lindqvist, 1970) [Pteronidea]. Host plant: *Spiraea* spp. (Rosaceae). Russia: **ES** (IR).

Euura erecta (Lindqvist, 1945) [Amauronematus]. Russia: **WS** (TM). – Europe (WE, NE).

Euura ermolenkoi (Haris, 2006) [Nematus]. Russia: **FE** (KU).

Euura erythropus (Lindqvist, 1975) [Amauronematus]. Russia: **ES** (IR).

Euura evenensis (Zhelochovtsev, 1981) [Nematus]. Host plant: ?*Chosenia arbutifolia* (Salicaceae). Russia: **FE** (MG).

Euura excavata (Marlatt, 1896) [Pontania] (*Pontania megacephala* Rohwer, 1908; *P. distracta* MacGillivray, 1923; *P. apicifrons* Malaise, 1931). Host plant: *Populus suaveolens*, *Salix pentandra*, *S. pseudopentandra* (Salicaceae). Russia: **EP** (Zhelochovtsev, Zinovjev, 1995), **ES** (Zhelochovtsev, Zinovjev, 1995), **FE** (KA). – Europe (WE, NE), N America.

Euura excisa (Thomson, 1863) [Nematus] (*Lygaeone-matus strandi* Konow, 1901). Host plant: *Geranium* spp. (Geraniaceae). Russia: **ES** (IR). – Europe (NE), Mongolia, N America.

Euura extensicornis (Norton, 1861) [Nematus] (*Nematus marylandicus* Norton, 1864; *N. aureopectus* Norton, 1867; *N. suadus* Cresson, 1880; *Pachynematus affinis* Marlatt, 1896; *P. tritici* Marlatt, 1896). Host plant: Poaceae. Russia: **ES** (IR), **FE** (MG). – N America.

Euura facialis (Lindqvist, 1969) [Pteronidea]. Host plant: *Alnus* spp. (Betulaceae). Russia: **ES** (IR). – Europe (NE).

Euura fahraei (Thomson, 1863) [Nematus] (*Nematus melanodus* Zaddach, 1884). Host plant: *Populus* spp. (Salicaceae). Russia: **ES** (IR, YA). – Europe (WE, EE, NE).

Euura fallax (Serville, 1823) [Nematus] (*Nematus fal-lax* Lepeletier, 1823; *N. melanocerus* Hartig, 1840;

N. xanthocarpus Hartig, 1840; *N. eupodius* Hartig, 1840; *N. haemorrhoidalis* Hartig, 1840; *N. brachycercus* Hartig, 1840; *N. schmidtii* Gimmerthal, 1844; *N. eversmanni* Förster, 1854; *N. emarginatus* Förster, 1854; *N. circumscriptus* Förster, 1854; *N. declinatus* Förster, 1854; *N. incertus* W.F. Kirby, 1882; *N. erythropareus* Zaddach, 1884; *Pachynematus sulcatus* Benson, 1948; *P. chambersi* Benson, 1948; *P. virginialis* Liston, 1980). Host plant: Salix spp. (Salicaceae). Russia: **EP** (NW, NC), **FE** (PR, KA, MG). – Europe (WE, EE, NE, SE), Turkey, Kyrgyzstan, Kazakhstan, Mongolia, China (NW), Korean Peninsula, Japan (Hon).

Euura fasciata (Konow, 1897) [Amauronematus] (*Amauronematus perkinsi* Benson, 1933). Host plant: Salix spp. (Salicaceae). Russia: **EP** (NW), **ES** (IR). – Europe (WE, EE, NE).

Euura ferruginea (Förster, 1854) [Nematus] (*Nematus glottianus* Cameron, 1882; *Amauronematus moricei* Konow, 1902; *Pteronidea ferruginea plagiata* Enslin, 1916). Host plant: Salix spp., Populus spp. (Salicaceae). Russia: **EP** (N), **ES** (IR). – Europe (WE, EE, NE).

Euura filiformis (Lindqvist, 1975) [Pachynematus]. Russia: **ES** (IR).

Euura flavescens (Stephens, 1835) [Nematus] (*Nematus dilutus* Brischke, 1883). Host plant: Salix spp. (Salicaceae). Russia: **EP** (NW), **ES** (YA), **FE** (KA). – Europe (WE, EE, NE), Central Asia.

Euura foetidae (Kopelke, 1989) [Pontania]. Host plant: Salix foetida, S. saposhnikovii (Salicaceae); larvae form galls. Russia: **WS** (AL). – Europe (WE).

Euura frenalis (Thomson, 1888) [Nematus] (*Pteronus fastosus* Konow, 1904). Host plant: Salix spp. (Salicaceae). Russia: **EP** (N), “Siberia” (Zhelochovtsev, Zinovjev, 1995). – Europe (WE, EE, NE), China (Wei et al., 2006).

Euura freyja Liston, Taeger et Blank, 2009 (*Pontopristia analis* Lindqvist, 1961). Host plant: Salix spp. (Salicaceae). Russia: **ES** (IR). – Europe (NE).

Euura fujiiense (Zinovjev, 1993) [Pikonema]. Host plant: ?Abies spp., ?Picea spp. (Pinaceae). Russia: **FE** (KU). – Japan (Hon).

Euura fulvescens (Lindqvist, 1949) [Pteronidea]. Russia: **EP** (N). – Europe (NE).

Euura fuscomaculata (Förster, 1854) [Nematus] (*Nematus strongylogaster* Cameron, 1878; *Pteronus dossuarius* Konow, 1904). Host plant: Populus spp. (Salicaceae). Russia: **EP** (NW, C), **ES** (YA). – Europe (WE, EE, NE).

Euura gehrsi (Konow, 1903) [Pachynematus]. Host plant: Polygonum spp. (Polygonaceae). Russia: **EP** (NC), **ES** (IR). – Europe (WE).

Euura glabrifrons (Benson, 1960) [Pontania]. Host plant: Salix lanata (Salicaceae); larvae form galls. Russia: **EP** (N), **WS** (TM), **ES** (YA), **FE** (MG). – Europe (NE).

Euura glacialis (Jakovlev, 1891) [Amauronematus]. Russia: **EP** (N).

Euura glaucae (Kopelke, 1994) [Pontania]. Host plant: Salix glauca, S. alpina, S. saxatilis (Salicaceae); larvae form galls. Russia: **ES** (YA), **FE** (MG). – Europe (WE, NE).

Euura gracilidentata (Viitasaari, 1980) [Pteronidea]. Host plant: Ribes alpinum (Grossulariaceae). Russia: **EP** (NW). – Europe (WE, NE, SE).

Euura groenlandica (Malaise, 1933) [Amauronematus]. Russia: **EP** (N). – Europe (NE), Greenland, N America.

Euura harpicola (Bogacheva, 1977) [Amauronematus]. Russia: **WS** (TM).

Euura hartigi (Saarinen, 1950) [Amauronematus]. Russia: **EP** (NW). – Europe (WE).

Euura hastatae Malaise, 1920. Host plant: Salix hastata (Salicaceae). Russia: **EP** (N). – Europe (WE, EE, NE).

Euura hastatavora Vikberg, 2014 (*Pontania hastatae* Vikberg, 1970). Host plant: Salix hastata (Salicaceae). Russia: **EP** (N), **WS** (TM), **ES** (YA), **FE** (MG, CH). – Europe (WE, NE).

Euura hebes (Konow, 1907) [Amauronematus] (*Amauronematus pristiphorinus* Malaise, 1921). Russia: **WS** (TM). – Europe (NE), N America.

Euura hedstroemi (Malaise, 1931) [Amauronematus] (*Amauronematus rex* Benson, 1948). Russia: **EP** (N), **FE** (KA, CH). – Europe (WE, NE), N America.

Euura helleni (Lindqvist, 1941) [Amauronematus]. Russia: **EP** (N), **FE** (MG). – Europe (NE), N America.

Euura helveticae (Kopelke, 1986) [Pontania]. Host plant: Salix helvetica, S. lanata, S. brachypoda (Salicaceae); larvae form galls. Russia: **EP** (NW), **WS** (AL), **ES** (YA). – Europe (WE).

Euura herbacea (Cameron, 1876) [Nematus] (*Pontania polaris* Malaise, 1920). Host plant: Salix herbacea, S. polaris, S. nummularia (Salicaceae). Russia: **EP** (N), **WS** (TM), **ES** (KR, BR, YA), **FE** (MG, CH). – Europe (WE, NE).

Euura hirowatarii (Haris et Zsolnai, 2007) [Pachynematus]. Russia: **FE** (SA). – Korean Peninsula.

- Euura histrio** (Serville, 1823) [Nematus] (*Nematus histrio* Lepeletier, 1823; *N. rufescens* Hartig, 1837; *N. fur* Walsh, 1866; *N. glenelgensis* Cameron, 1882). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (NW, C, E, NC), **UR**, **ES** (IR, ZB). – Europe (WE, EE, NE, SE), N America.
- Euura hulteni** (Malaise, 1931) [Amauronematus]. Russia: **FE** (KA). – N America.
- Euura humeralis** (Serville, 1823) [Nematus] (*Nematus humeralis* Lepeletier, 1823; *N. squalidus* Eversmann, 1847; *N. incanus* Förster, 1854; *Amauronematus terminalis* Malaise, 1931). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (NW, C, E), **UR**, **WS** (Zhelochovtsev, Zinovjev, 1995), **ES** (IR), **FE** (KA, MG). – Europe (WE, EE, NE).
- Euura hypoxantha** (Förster, 1854) [Nematus] (*Nematus palliatus* Thomson, 1863; *N. orbitalis* Cameron, 1884; *Pteronidea nigronota* Lindqvist, 1958). Host plant: *Salix* spp., *Populus* spp. (Salicaceae). Russia: **EP** (NW, C, S), **ES** (IR). – Europe (WE, EE, NE, SE), Kazakhstan, Japan (Hon).
- Euura imperfecta** (Zaddach, 1876) [Nematus] (*Pachynematus imperfectus claristernis* Enslin, 1916). Host plant: *Larix* spp. (Pinaceae). Russia: **EP** (NW), **ES** (IR, BR, YA, ZB). – Europe (WE, EE, NE).
- Euura incompleta** (Förster, 1854) [Nematus] (*Nematus smaragdinus* Stein, 1881; *N. pulchellus* Cameron, 1882; *N. chlorogaster* Zaddach, 1884; *Pteronidea segmentaria signata* Enslin, 1916; *P. segmentaria tessonensis* Enslin, 1916). Host plant: *Lathyrus vernus* (Fabaceae). Russia: **EP** (N, NW, C, E, S, NC), **ES** (ZB). – Europe (WE, EE, NE, SE), Georgia, Kazakhstan.
- Euura insigne** (Hartig, 1840) [Nematus] (*Nematus superbus* Gradl, 1881; *N. nobilis* W.F. Kirby, 1882). Host plant: *Abies* spp., *Picea* spp. (Pinaceae). Russia: **EP** (NW), **ES** (YA), **FE** (PR). – Europe (WE, EE, NE).
- Euura integra** (Zinovjev, 1985) [Pontania]. Host plant: *Salix integra* (Salicaceae). Russia: **FE** (KH, PR). – Japan (Zinovjev, 2010).
- Euura itelmena** (Malaise, 1931) [Pontopristia]. Host plant: ?*Salix alaxensis* (Salicaceae). Russia: **FE** (KA). – N America.
- Euura itoi** (Okutani, 1955) [Pachynematus]. Host plant: *Larix* spp. (Pinaceae). Russia: **UR**, **ES** (TU, IR, BR, YA, ZB), **FE** (AM, MG). – Europe (WE), China (NE), Korean Peninsula, Japan (Hon).
- Euura jugicola** (Thomson, 1871) [Nematus] (*Pteronidea karvoneni* Lindqvist, 1969). Host plant: *Salix* aurita (Salicaceae). Russia: **EP** (NW). – Europe (WE, NE).
- Euura krausi** (Taeger et Blank, 1998) [Amauronematus]. Host plant: *Populus tremula* (Salicaceae). Russia: **EP** (N, NW, C), **ES** (IR). – Europe (WE, EE, NE, SE).
- Euura kunasirensis** (Haris, 2006) [Nematus]. Russia: **FE** (KU).
- Euura laeta** (Brischke, 1883) [Cryptocampus] (*E. viminalis* Kopelke, 2001). Host plant: *Salix viminalis* (Salicaceae); larvae in the galls at the base of the leaf petiole. Russia: **EP** (N, NW, C, E), **FE** (KA). – Europe (WE, EE, NE).
- Euura lanatae** Malaise, 1920. Host plant: *Salix lanata* (Salicaceae); larvae in enlarged buds. Russia: **WS** (TM), **ES** (KR). – Europe (NE), N America.
- Euura lanceata** (Hellén, 1970) [Amauronematus]. Russia: **FE** (MG). – Europe (NE).
- Euura lappo** Malaise, 1920. Host plant: *Salix lapponum* (Salicaceae). Russia: **EP** (N). – Europe (EE, NE).
- Euura latiserra** (Malaise, 1921) [Pontopristia]. Host plant: *Salix* spp. (Salicaceae). Russia: **WS** (TM), **ES** (KR), **FE** (CH). – Europe (NE).
- Euura latisulcata** (Malaise, 1931) [Amauronematus]. Russia: **FE** (KA).
- Euura leionota** (Benson, 1933) [Pteronidea]. Host plant: *Betula* spp. (Betulaceae). Russia: **EP** (N). – Europe (WE, EE, NE).
- Euura leptcephala** (Thomson, 1863) [Nematus] (*Amauronematus morawitsi* Jakovlev, 1891). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (NW). – Europe (NE), N America.
- Euura lethe** Prous et Liston, 2014 (*Amauronematus atratus* Lindqvist, 1961). Russia: **ES** (IR). – Europe (NE).
- Euura leucapsis** (Tischbein, 1846) [Nematus] (*Nematus viminalis* Hartig, 1840; *N. alienatus* Förster, 1854; *N. coriaceus* Benson, 1953; *Phyllocolpa rolleri* Liston, 2005). Host plant: *Salix cinerea*, *S. aurita*, *S. silesiaca* (Salicaceae). Russia: **EP** (NW), **ES** (IR, BR, YA), **FE** (AM, KA, MG). – Europe (WE, EE, NE), Tajikistan, Kyrgyzstan, Canada.
- Euura leucolena** (Brischke, 1883) [Nematus] (*Pontania unga* Kincaid, 1900; *Pteronidea egeria* MacGillivray, 1923). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (N, NW), **UR**, **WS** (TM), **ES** (KR, IR, BR, ZB), **FE** (KA, MG). – Europe (WE, EE, NE), N America.
- Euura leucopyga** (Lindqvist, 1949) [Amauronematus] (*Pteronidea bipicta* Lindqvist, 1959). Host plant:

- Alnus* spp. (Betulaceae). Russia: **EP** (N), **ES** (IR). – Europe (NE).
- Euura leucosticta** (Hartig, 1837) [Nematus] (*Nematus klugii* Gimmerthal, 1844; *N. crassulus* Thomson, 1863; *N. sharpi* Cameron, 1876; *Pontania ischnoceros nigrifrons* Konow, 1897; *P. auberti* Zirngiebl, 1957). Host plant: *Salix carpea*, *S. aurita*, *S. atrocincta* (Salicaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (Zhelochovtsev, Zinovjev, 1995). – Europe (WE, EE, NE, SE), N Africa, Armenia, Turkey, Kazakhstan.
- Euura leucotrocha** (Hartig, 1837) [Nematus] (*Nematus cylindricus* Hartig, 1840; *N. maculiventris* Hartig, 1840; *N. approximatus* Förster, 1854; *N. consobrinus* Snellen van Vollenhoven, 1871; *N. umbrenus* Zaddach, 1876; *N. sauterianus* Zaddach, 1883; *N. multiplex* Zaddach, 1883; *Pteronidea cognata* Lindqvist, 1958). Host plant: *Ribes* spp. (Grossulariaceae). Russia: **EP** (NW, C, E), **UR**, **WS** (Zhelochovtsev, Zinovjev, 1995). – Europe (WE, EE, NE), China (Wei et al., 2006).
- Euura lichtwardti** (Konow, 1903) [Pachynematus] (*Nematus apicalis* Hartig, 1837; *Pachynematus kubesi* Konow, 1905). Host plant: Poaceae. Russia: **EP** (N), **ES** (IR). – Europe (WE, EE, NE), Korean Peninsula.
- Euura lienterica** (Holmgren, 1883) [Nematus]. Host plant: *Vaccinium* spp. (Ericaceae). Russia: **EP** (N), **WS** (TM), **ES** (KR), **FE** (MG, CH). – Europe (NE).
- Euura lindqvisti** (Hellén, 1951) [Amauronematus]. Russia: **ES** (Zhelochovtsev, Zinovjev, 1995). – Europe (NE).
- Euura longicauda** (Hellén, 1948) [Nematus]. Russia: **EP** (N), **FE** (MG). – Europe (NE), N America.
- Euura longiserra** (Thomson, 1863) [Nematus] (*Nematus leucodous* Zaddach, 1884). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (N, C), **WS** (AL), **ES** (IR, YA), **FE** (KA). – Europe (WE, NE).
- Euura magadanensis** (Vikberg et Zinovjev, 2014) [Pontania]. Host plant: *Salix* spp. (Salicaceae); larvae form galls. Russia: **ES** (YA), **FE** (KA, MG, CH).
- Euura malaisei** (Hellén, 1970) [Amauronematus] (*Decanematus longiserra* Malaise, 1931). Russia: **FE** (KA). – Europe (WE, EE, NE).
- Euura mandshurica** (Zinovjev, 1981) [Pontania]. Host plant: *Salix pierotii*, *S. babylonica* (Salicaceae); larvae form galls. Russia: **FE** (AM, PR). – Japan (Zinovjev, 2010).
- Euura mcluckiei** (Benson, 1935) [Amauronematus] (*Amauronematus pustulatus* Lindqvist, 1962). Host plant: *Salix reticulata* (Salicaceae). Russia: **EP** (N), “Siberia” (Zhelochovtsev, Zinovjev, 1995). – Europe (WE, NE).
- Euura melanocephala** (Hartig, 1837) [Nematus] (*Tenthredo salicis* Retzius, 1783; *Nematus perspicillaris* Hartig, 1840; *N. breviusculus* Eversmann, 1847; *N. pyrrhonotus* Zaddach, 1883; *Pteronus xanthostomus* Rohwer, 1910; *Pteronidea melanocephala mesothoracica* Enslin, 1916). Host plant: *Salix* spp., *Populus* spp. (Salicaceae). Russia: **EP** (NW, C). – Europe (WE, EE, NE, SE), Armenia.
- Euura microphyes** (Förster, 1854) [Nematus] (*Pontopristia romani* Malaise, 1921; *P. nana* Lindqvist, 1949; *P. fennicus* Lindqvist, 1949). Host plant: *Salix* spp. (Salicaceae); larvae in the inflorescence. Russia: **EP** (N), **FE** (PR). – Europe (WE, NE), N America.
- Euura miliaris** (Panzer, 1797) [Tenthredo] (*Nematus dorsalis* Serville, 1823; *N. dorsalis* Lepeletier, 1823; *N. testaceus* Stephens, 1835; *N. fulvus* Hartig, 1837; *N. purus* Förster, 1854; *N. validicornis* Förster, 1854; *N. fulvus basalis* Costa, 1859; *N. fulvus exoleta* Costa, 1859; *N. trimaculatus* Snellen van Vollenhoven, 1862; *N. solitarius* Zaddach, 1883; *Pteronidea miliaris nigronotata* Enslin, 1916). Host plant: *Salix* spp. (Salicaceae); larvae form a galls on the leaves. Russia: **EP** (N, NW, C, E, S, NC), **UR**, **ES** (KR, IR, BR, YA), **FE** (SA). – Europe (WE, EE, NE, SE), SW Asia, Central Asia, China (Wei et al., 2006), Korean Peninsula.
- Euura miltonota** (Zaddach, 1883) [Nematus] (*Amauronematus jaroslawensis* Jakovlev, 1891; *A. hercyniae* Konow, 1896). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (N, NW, C), **FE** (KA). – Europe (WE, EE).
- Euura mimator** Schmidt, 2014 (*Amauronematus mimus* Schmidt, 1997). Russia: **EP** (NW). – Europe (WE, NE).
- Euura mimus** (Konow, 1903) [Pteronus]. Russia: **EP** (NW). – Europe (WE, EE, SE).
- Euura mirabilis** (Zinovjev, 1985) [Pontania]. Host plant: *Salix cardiophyla* (Salicaceae). Russia: **FE** (PR, SA).
- Euura montana** (Zaddach, 1883) [Nematus] (*Nematus jemilleri* Stein, 1894). Host plant: *Picea* spp., *Abies* spp. (Pinaceae). Russia: **EP** (C), **ES** (YA). – Europe (WE, EE, NE).
- Euura monticola** (Thomson, 1871) [Nematus] (*Pachynematus perkioemaekii* Lindqvist, 1960; *Nematus pschorwalcheri* Muche, 1972). Russia: **EP** (N). – Europe (WE, EE, NE, SE).

Euura mucronata (Hartig, 1837) [Nematus] (*Cryptocampus pictus* Zaddach, 1883; *C. gemmarum* Brischke, 1883; *Euura insularis* Kincaid, 1900). Host plant: *Salix caprea*, *S. aurita* (Salicaceae); larvae in bugs. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **ES** (IR, YA), **FE** (AM, KA). – Europe (WE, EE, NE, SE), Armenia, Japan (Hon), N America.

Euura mundus (Konow, 1895) [Amauronematus]. Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (NW), **ES** (IR). – Europe (WE, NE), Mongolia, China (Wei et al., 2006).

Euura myosotidis (Fabricius, 1804) [Tenthredo] (*Nematus interruptus* Serville, 1823; *N. interruptus* Lepeletier, 1823; *N. segmentarius* Förster, 1854; *N. ambiguus* Förster, 1854; *N. decipiens* W.F. Kirby, 1882; *Pteronius myosotidis fallaciosus* Konow, 1903; *Pteronidea myosotidis pseudopavida* Enslin, 1916). Host plant: Fabaceae, Rosaceae. Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (AL), **ES** (KR, IR). – Europe (WE, EE, NE, SE), Armenia, Turkey, Kyrgyzstan, Kazakhstan.

Euura myrsiniticola (Kopelke, 1991) [Pontania]. Host plant: *Salix myrsinoides* (Salicaceae). Russia: **EP** (N). – Europe (NE).

Euura myrtilloidica (Kopelke, 1991) [Pontania]. Host plant: *Salix myrtilloides* (Salicaceae). Russia: **EP** (N, NW), **FE** (AM, MG). – Europe (WE, NE).

Euura neglecta (W.F. Kirby, 1882) [Nematus] (*Amauronematus spiniserra* Malaise, 1920; *A. intermedius* Malaise, 1931). Host plant: *Salix* spp. (Salicaceae). Russia: **UR**, **ES** (TU, YA), **FE** (KA). – Europe (NE), N America.

Euura nigricornis (Serville, 1823) [Nematus] (*Nematus nigricornis* Lepeletier, 1823; *N. zetterstedti* Dahlbom, 1835; *N. miniatus* Hartig, 1837; *N. sibiricus* Jakovlev, 1888; *N. biannulatus* Costa, 1890). Host plant: *Salix* spp., *Populus* spp. (Salicaceae). Russia: **EP** (NW, C, E), **ES** (IR). – Europe (WE, EE, NE, SE), Japan (Hon).

Euura nigriventris (Holmgren, 1883) [Nematus] (*Amauronematus varians* MacGillivray, 1919; *Pteronidea melanostoma* Rohwer, 1920). Russia: **EP** (N). – N America.

Euura nitidipleuris (Malaise, 1931) [Amauronematus]. Russia: **FE** (KA). – Europe (NE), N America.

Euura nivalis (Vikberg, 1970) [Pontania]. Host plant: *Salix glauca* (Salicaceae). Russia: **EP** (N), **WS** (TM, AL). – Europe (WE, NE), Canada.

Euura nudipectus (Vikberg, 1965) [Pontania]. Host plant: *Salix phylicifolia* (Salicaceae). Russia: **EP** (N, NW), **FE** (MG). – Europe (NE).

Euura nuortevai (Lindqvist, 1958) [Pteronidea]. Russia: **ES** (IR). – Europe (NE).

Euura obducta (Hartig, 1837) [Nematus] (*Nematus conductus* Ruthe, 1859; *N. graminis* Cameron, 1874; *N. palliventris* Cresson, 1880; *N. pallidiventralis* Dalla Torre, 1894; *Pachynematus boulderensis* Rohwer, 1920). Host plant: Poaceae. Russia: **EP** (N, NW, C, NC), **ES** (TU, IR), **FE** (SA, KU, KA, MG). – Europe (WE, EE, NE, SE), Turkey, Kyrgyzstan, N America.

Euura oblita (Serville, 1823) [Nematus] (*Nematus oblitus* Lepeletier, 1823; *N. pineti* Hartig, 1837; *N. pleurostictus* Förster, 1854; *N. lepidus* Förster, 1854; *N. puella* Thomson, 1871; *N. elegans* Zaddach, 1884). Host plant: *Salix triandra*, *S. alba*, *S. fragilis* (Salicaceae). Russia: **EP** (NW, C, S), **ES** (ZB). – Europe (WE, EE, NE), Armenia, Turkey.

Euura obscuripes (Holmgren, 1883) [Nematus]. Russia: **EP** (N).

Euura olfaciens olfaciens (Benson, 1953) [Nematus]. Host plant: *Ribes* spp. (Grossulariaceae). Russia: **EP** (NW, C). – Europe (WE, EE, NE).

Euura olfaciens kolymensis (Zhelochovtsev, 1981) [Nematus]. Host plant: *Ribes* spp. (Grossulariaceae). Russia: **ES** (YA), **FE** (MG).

Euura oligospila (Förster, 1854) [Nematus] (*Nematus mendicus* Walsh, 1866; *N. trivittatus* Norton, 1867; *N. microcercus* Thomson, 1871; *N. dorsivittatus* Cresson, 1880; *N. salicivorus* Cameron, 1882; *Pteronius koebelei* Marlatt, 1896; *Pteronidea vanduzeei* Rohwer, 1913; *P. elelea* MacGillivray, 1923; *Nematus desantisi* D.R. Smith, 1983). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **ES** (TU, IR). – Europe (WE, EE, NE, SE), Caucasus, Central Asia, Pakistan, N America; introduced into Tropical Africa, S America, Australia, New Zealand.

Euura pallens (Konow, 1903) [Pteronus] (*Pteronidea straminea* Lindqvist, 1958). Russia: **ES** (IR). – Europe (NE).

Euura pallescens (Hartig, 1837) [Nematus] (*Nematus olivaceus* Thomson, 1871; *N. ruficeps* Zaddach, 1883; *N. pachycerus* Borries, 1896). Host plant: *Picea* spp. (Pinaceae). Russia: **EP** (C). – Europe (WE, EE, NE).

Euura pallidior (Hellén, 1970) [Amauronematus]. Russia: **WS** (TM). – Europe (NE).

Euura papillosa (Retzius, 1783) [Tenthredo] (*Tenthredo fuscata* Christ, 1791; *Nematus melanaspis* Hartig, 1840; *N. lacteus* Thomson, 1871; *N. sulphureus* Zaddach, 1876; *N. citreus* André, 1880; *N. maculiger* Cameron, 1882; *N. brunnicornis* Zaddach, 1883; *Pteronidea sveae* Lindqvist, 1958). Host plant: Salix spp., Populus spp. (Salicaceae), Betula spp. (Betulaceae). Russia: **EP** (N, NW, C), **WS** (Zhelochovtsev, 1988), **ES** (IR, BR, YA). – Europe (WE, EE, NE, SE), Pakistan, Kyrgyzstan, Kazakhstan, Japan (Takeuchi, 1952).

Euura parvula (Holmgren, 1883) [Nematus] (*Pontania poppii* Konow, 1904; *Pteronidea gracilicornis* Lindqvist, 1962). Host plant: Vaccinium spp. (Ericaceae). Russia: **EP** (N), **ES** (IR). – Europe (NE).

Euura pavida (Serville, 1823) [Nematus] (*Nematus pavidus* Lepeletier, 1823; *N. ochraceus* Hartig, 1837; *N. semiorbitalis* Förster, 1854; *Cryptocampus quadrum* Costa, 1859; *Nematus wtewaalli* Snellen van Vollenhoven, 1862; *N. cameronii* Dalla Torre, 1894; *Pteronidea pavida notanda* Enslin, 1916). Host plant: Salix spp., Populus spp. (Salicaceae). Russia: **EP** (NW, C), **WS** (Zhelochovtsev, Zinovjev, 1995). – Europe (WE, EE, NE, SE).

Euura pedunculi (Hartig, 1837) [Nematus] (*Nematus bellus* Zaddach, 1876; *N. baccarum* Cameron, 1876; *N. curticornis* Cameron, 1885; *Pontania bella nigrescens* Enslin, 1915; *P. pusilla* Lindqvist, 1964). Host plant: Salix spp. (Salicaceae); larvae form galls. Russia: **EP** (NW, C), **WS** (Zhelochovtsev, Zinovjev, 1995), **ES** (YA), **FE** (SA, KA, MG). – Europe (WE, EE, NE, SE), Korean Peninsula.

Euura piliserra (Thomson, 1863) [Nematus] (*Pontania piliserra mascula* Enslin, 1915; *P. piliserra tristis* Enslin, 1915; *Pteronidea weiffenbachi* Lindqvist, 1958). Host plant: Salix viminalis (Salicaceae). Russia: **EP** (Zhelochovtsev, Zinovjev, 1995), **WS** (Zhelochovtsev, Zinovjev, 1995). – Europe (WE, EE, NE).

Euura platystigma (Lindqvist, 1949) [Pteronidea]. Russia: **ES** (IR). – Europe (NE).

Euura poecilonota (Zaddach, 1876) [Nematus] (*Nematus viridescens* Cameron, 1885; *Pteronus hyalinus* Marlatt, 1896; *Pteronidea subnitens* Lindqvist, 1958). Host plant: Betula spp. (Betulaceae). Russia: **EP** (NW, C, E, S), **WS** (AL), **ES** (TU, IR, BR), **FE** (MG). – Europe (WE, EE, NE, SE), Turkey, Kyrgyzstan, Korean Peninsula.

Euura polaris (Holmgren, 1883) [Nematus] (*Nematus morionellus* Holmgren, 1883; *Pteronus shumagensis*

Kincaid, 1900; *Pontania islandica* Kincaid, 1900; *P. birulae* Konow, 1907). Host plant: Salix spp. (Salicaceae). Russia: **EP** (N), **WS** (TM), **FE** (MG). – Europe (WE, NE), N America.

Euura polita (Zaddach, 1883) [Nematus] (*Nematus sieboldii* Zaddach, 1884; *Pontania leucapsis connata* Enslin, 1915). Russia: **EP** (NW). – Europe (WE, EE).

Euura pravus (Konow, 1895) [Amauronematus]. Host plant: Salix spp. (Salicaceae). Russia: **EP** (NW), **ES** (YA). – Europe (WE, NE), N America.

Euura pribaicalica (Lindqvist, 1970) [Amauronematus]. Host plant: Salix spp. (Salicaceae). Russia: **ES** (IR).

Euura pristiphoroides (Lindqvist, 1972) [Amauronematus]. Host plant: Salix spp. (Salicaceae). Russia: **ES** (IR), **FE** (KA).

Euura proxima (Serville, 1823) [Nematus] (*Nematus proximus* Lepeletier, 1823; *N. gallicola* Stephens, 1835; *N. vallisnieri* Hartig, 1837; *N. redii* Contarinia, 1852; *Pontania gallicola* Costa, 1852; *Nematus albicarpus* Costa, 1859; *Messa hyalina* Norton, 1864; *Nematus festivus* Zaddach, 1884; *Euura flavipes* Cameron, 1885; *Pontania daedala* MacGillivray, 1921). Host plant: Salix spp. (Salicaceae); larvae form galls on lateral veins. Russia: **EP** (N, NW, C, E, S), **WS** (AL), **ES** (IR, BR), **FE** (KU, KA, MG). – Europe (WE, EE, NE, SE), Cyprus, Kazakhstan; introduced into N America, Australia, New Zealand.

Euura pseudodispar (Lindqvist, 1969) [Pteronidea]. Host plant: Betula spp. (Betulaceae). Russia: **ES** (YA). – Europe (NE).

Euura pseudoleptocephala (Haris, 2006) [Amauronematus]. Russia: **FE** (SA).

Euura pseudotorneensis (Conde, 1935) [Amauronematus]. Russia: **FE** (SA).

Euura pumilio (Konow, 1903) [Pachynematus]. Host plant: Ribes spp. (Grossulariaceae); larvae in black currant berries. Russia: **EP** (NW, C, E), **WS** (TM, TK, AL), **ES** (KR). – Europe (WE, EE, NE).

Euura punctifrons (Malaise, 1931) [Pontania]. Russia: **FE** (KA).

Euura punicea (Christ, 1791) [Tenthredo] (*Tenthredo nigrata* Retzius, 1783; *T. crocea* Fabricius, 1804; *Nematus melanosternus* Serville, 1823; *N. melanosternus* Lepeletier, 1823; *N. prototypus* Förster, 1854; *Amauronematus excellens* Forsius, 1921). Host plant: Populus spp. (Salicaceae). Russia: **EP** (N, NW, C), **ES** (IR, YA). – Europe (WE, EE, NE), Korean Peninsula.

Euura purpureae (Cameron, 1884) [Nematus] (*E. acuminata* Enslin, 1915). Host plant: *Salix purpurea*, *S. acutifolia* (Salicaceae). Russia: **EP** (C). – Europe (WE, EE, NE, SE).

Euura pustulator (Forsius, 1923) [Pontania]. Host plant: *Salix phyllicifolia*, *S. pulchra* (Salicaceae); larvae form galls. Russia: **EP** (N). – Europe (WE, EE, NE).

Euura pygmaea Brischke, 1883 (Cryptocampus). Russia: **EP** (NW). – Europe (EE).

Euura quieta (Eversmann, 1847) [Nematus]. Russia: **EP** (E).

Euura quinquemontana (Jakovlev, 1891) [Pachyne-matus]. Russia: **EP** (NC).

Euura ranini (Lindqvist, 1960) [Amauronematus]. Host plant: *Populus* spp. (Salicaceae). Russia: **ES** (IR). – Europe (NE).

Euura relictana (Zinovjev, 1981) [Pontania]. Host plant: *Chosenia arbutifolia* (Salicaceae); larvae form galls. Russia: **FE** (KH, PR, KA, MG).

Euura respondens (Förster, 1854) [Nematus] (*Nematus nitens* Thomson, 1888; *Pteronus balassagloii* Jakovlev, 1891). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (E). – Europe (WE, EE, NE), Uzbekistan, Kazakhstan.

Euura reticulata (Holmgren, 1883) [Nematus] (*Nematus arcticus* Thomson, 1871; *N. mysticus* Holmgren, 1883; *N. anceps* Holmgren, 1883; *N. occipitalis* Holmgren, 1883; *N. picticollis* Holmgren, 1883; *N. arcticola* Dalla Torre, 1894; *Pontania popofiana* Kincaid, 1900; *P. glinka* Kancaid, 1900; *P. forsiusi* Enslin, 1915; *Amauronematus arcticola* Enslin, 1915; *A. alsius* Benson, 1935). Host plant: *Vaccinium* spp. (Ericaceae). Russia: **EP** (N), **WS** (TM), **ES** (KR, YA), **FE** (MG). – Europe (WE, NE), N America.

Euura ribesicola (Lindqvist, 1949) [Pteronidea] (*Nematus beybienkoi* Skorikova, 1952). Host plant: *Ribes* spp. (Grossulariaceae). Russia: **EP** (N), **WS** (Zhelochovtsev, Zinovjev, 1995), **ES** (IR, YA), **FE** (KH). – Europe (NE).

Euura ribesii (Scopoli, 1763) [Tenthredo] (*Nematus dimidiatus* Lepeletier, 1823; *N. trimaculatus* Lepeletier, 1823; *N. dimidiatus* Serville, 1823; *N. trimaculatus* Serville, 1823; *N. ribesii* Stephens, 1829; *Tenthredo grossulariae* folii Moore, 1831; *T. ventricosa* Bouché, 1834; *Nematus grossulariae* Dahlbom, 1835; *N. ventricosus* Hartig, 1837; *N. flavipes* Hartig, 1840; *N. xanthophorus* Hartig, 1840; *N. macrocerus* Hartig, 1840; *N. ribis* Leduc, 1842; *Tenthredo gros-sulariae* Fischer von Waldheim, 1843; *Pteronus*

longicornis Marlatt, 1896; *P. arapahonum* Cockerell, 1906; *Pteronidea aceris* Rohwer, 1915; *P. ribesii feminina* Enslin, 1916; *P. ribesii konowi* Enslin, 1916; *P. equatia* MacGillivray, 1923). Host plant: *Ribes* spp. (Grossulariaceae). Russia: **EP** (NW, C, E, S, NC), **WS** (Zhelochovtsev, Zinovjev, 1995), **ES** (IR, YA, ZB), **FE** (KH, PR). – Europe (WE, EE, NE, SE), Central Asia, Kazakhstan, Korean Peninsula, Japan (Hon); introduced into N America.

Euura rubiginosa (Hellén, 1951) [Amauronematus]. Russia: **ES** (KR).

Euura rufomacula (Malaise, 1931) [Amauronematus] (*Amauronematus obesus* Lindqvist, 1959). Russia: **FE** (KA). – Europe (NE).

Euura rufus (Konow, 1896) [Amauronematus]. Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (N), **ES** (IR, ZB), **FE** (AM, KA, MG). – Europe (NE).

Euura sagmarius (Konow, 1895) [Amauronematus]. Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (N, NW, C, S), **WS** (TM), **ES** (KR). – Europe (WE, NE), N America.

Euura saliceti sibirica Stroganova, 1985. Russia: **WS**.

Euura salicis (Linnaeus, 1758) [Tenthredo] (*Tenthredo capreae* Linnaeus, 1758; *T. notata* O.F. Müller, 1776; *Cynips salicis capreae* Christ, 1791; *Nematus klugi* Dahlbom, 1835; *N. immaculatus* Snellen van Vollen-hoven, 1869; *N. inflatus* Thomson, 1871; *Pteronidea salicis obscurior* Enslin, 1916). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (N, NW, C, S, NC). – Europe (WE, EE, NE, SE), Caucasus.

Euura saliciscinereae (Retzius, 1783) [Tenthredo] (*Tenthredo salicis* Christ, 1791; *Nematus gallarum* Hartig, 1837; *N. acerosus* Hartig, 1840; *N. saliceti* Förster, 1854; *N. aestivus* Thomson, 1863; *N. cine-reae* Thomson, 1871; *Pontania harrisoni* Benson, 1940; *P. varia* Kopelke, 1991; *P. norvegica* Kopelke, 1991). Host plant: *Salix myrsinifolia* (Salicaceae); larvae form galls. Russia: **EP** (C, E), **UR**. – Europe (WE, EE, NE).

Euura samolad (Malaise, 1920) [Pontania]. Host plant: *Salix lapponum* (Salicaceae); larvae form galls. Russia: **EP** (N, NW). – Europe (WE, NE).

Euura schlueteri (Enslin, 1915) [Amauronematus] (*Amauronematus lundbohmi* Malaise, 1920; *A. uliginosae* Malaise, 1920; *A. squamotus* Lindqvist, 1959). Host plant: *Vaccinium* spp. (Ericaceae). Russia: **EP** (N), **WS** (TM), **ES** (KR, IR, BR, YA, ZB), **FE** (AM, MG). – Europe (NE).

Euura scotapsis (Förster, 1854) [Nematus] (*Nematus westermanni* Thomson, 1863; *N. anglicus* Cameron, 1877; *N. nigrolineatus* Cameron, 1879). Host

- plant: *Salix viminalis*, *S. schwerinii* and others (Salicaceae). Russia: **EP** (N, NW, C), **ES** (KR, IR, YA), **FE** (KA). – Europe (WE, EE, NE), Armenia, Kyrgyzstan.
- Euura scutellata** (Hartig, 1837) [Nematus] (*Nematus immundus* Thomson, 1863). Host plant: *Abies* spp., *Picea* spp., *Larix* spp. (Pinaceae). Russia: **EP** (N, NW, C), **ES** (KR, IR, YA). – Europe (WE, EE, NE).
- Euura semilactea** (Zaddach, 1983) [Nematus]. Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (NW). – Europe (WE, EE, NE), N America.
- Euura semiopaca** (Lindqvist, 1958) [Pteronidea]. Russia: **EP** (N, NC), **WS** (TM). – Europe (NE).
- Euura sempersolis** (Kiær, 1898) [Amauronematus] (*Amauronematus tristis* Lindqvist, 1959). Russia: **EP** (N), **WS** (TM), **ES** (YA), **FE** (KA, MG). – Europe (NE).
- Euura septentrionalis** (Saarinen, 1950) [Amauronematus] (*Nematus taiganus* Zhelochovtsev, 1988). Russia: **FE** (MG). – Europe (NE).
- Euura seriepunctata** (Malaise, 1921) [Pteronidea]. Russia: **EP** (N), **FE** (MG). – Europe (NE).
- Euura sibirica** (Malaise, 1931) [Pontania]. Host plant: *Salix* spp. (Salicaceae). Russia: **FE** (KA).
- Euura sjoeblomi** (Malaise, 1931) [Amauronematus] (*Amauronematus tridentatus* Malaise, 1931). Russia: **FE** (KA).
- Euura smithae** (Ross, 1945) [Pachynematus] (*Pachynematus angustatus* Lindqvist, 1949). Russia: **EP** (Taeger et al., 2006). – Europe (WE, NE), N America.
- Euura speciosa** (Hellén, 1951) [Amauronematus]. Russia: **FE** (MG). – Europe (NE).
- Euura spiraeae** (Zaddach, 1883) [Nematus]. Host plant: *Spiraea* spp., *?Aruncus* spp. (Rosaceae). Russia: **EP** (NW), **ES** (IR, BR). – Europe (WE, NE).
- Euura stenogaster** (Förster, 1854) [Nematus] (*Nematus trifurcatus* W.F. Kirby, 1882; *Amauronematus analis* Konow, 1897; *A. vescus* MacGillivray, 1921; *A. veneficus* MacGillivray, 1923; *A. festivus* Saarinen, 1950). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (Zhelochovtsev, Zinovjev, 1995), **ES** (IR, YA). – Europe (WE, EE, NE, SE), N America.
- Euura stichi** (Enslin, 1913) [Pteronidea] (*Nematus testaceus* Thomson, 1871; *Pteronidea fuscårina* Benson, 1933). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (N). – Europe (WE, EE, NE, SE).
- Euura styx** (Benson, 1958) [Pachynematus]. Host plant: *Picea* spp. (Pinaceae). Russia: **ES** (KR). – Europe (WE, EE, NE).
- Euura subgemma** Liston, 2006. Russia: **EP** (N). – Europe (WE, EE, NE).
- Euura sylvestris** (Cameron, 1884) [Nematus] (*Nematus silvester* Dalla Torre, 1894; *N. fastosus ponojense* Hellén, 1948; *N. fastosus punctiscuta* Hellén, 1948). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (C, NC), "Siberia" (Zhelochovtsev, Zinovjev, 1995), **FE** (SA). – Europe (WE, EE, NE).
- Euura temporalis** (Hellén, 1970) [Amauronematus]. Russia: **EP** (N, C), **FE** (KA). – Europe (NE).
- Euura tenuiserra** (Lindqvist, 1944) [Amauronematus]. Russia: **EP** (N, NW). – Europe (WE, NE).
- Euura testaceipes** (Brischke, 1883) [Cryptocampus] (*E. cynips* Newman, 1837). Host plant: *Salix* spp. (Salicaceae); larvae form spindle-shaped galls on the petiole or central vein of leaf. Russia: **EP** (NW, C, CR). – Europe (WE, EE, NE), Armenia.
- Euura tibialis** (Newman, 1837) [Nematus] (*Nematus hortensis* Hartig, 1837; *N. xanthobaptus* Förster, 1854; *N. trilineatus* Norton, 1867; *N. catachloris* Snellen van Vollenhoven, 1869; *N. similaris* Norton, 1880; *N. robiniae* Forbes, 1885; *Pteronus solitarius* Rohwer, 1908). Host plant: *Robinia pseudoacacia* (Fabaceae). Russia: **EP** (C, S). – Europe (WE, EE, SE), N America.
- Euura tillbergi** (Malaise, 1920) [Amauronematus] (*Amauronematus tillbergi foveifrons* Malaise, 1920). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (N), **ES** (IR). – Europe (WE, NE), Kyrgyzstan, Korean Peninsula, N America.
- Euura toeniata** (Serville, 1823) [Nematus] (*Nematus taeniatus* Lepeletier, 1823; *Tenthredo humeralis* Zetterstedt, 1838; *Amauronematus aemulus* Konow, 1895; *A. zetterstedti* Malaise, 1920). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (NW), **ES** (IR, YA). – Europe (WE, EE, NE), Korean Peninsula.
- Euura tranquilla** Vårdal et Prous, 2014 (*Amauronematus pacificus* Malaise, 1931; *A. obscurus* Lindqvist, 1962). Russia: **ES** (Lindqvist, 1962), **FE** (KA). – Europe (NE).
- Euura transversalis** (Lindqvist, 1977) [Pteronidea]. Russia: **ES** (IR).
- Euura trautmanni** (Enslin, 1919) [Amauronematus] (*Amauronematus lateralis* Konow, 1895; *A. cameroni* Perkins, 1929; *A. piliserra* Lindqvist, 1943). Russia: **EP** (N, NW), **ES** (Zhelochovtsev, Zinovjev, 1995). – Europe (WE, EE, NE, SE).
- Euura triandrae** (Benson, 1941) [Pontania]. Host plant: *Salix triandra* (Salicaceae). Russia: **EP** (NW). – Europe (WE, EE, NE), Kazakhstan.

- Euura trochanterata** (Malaise, 1931) [Pteronidea] (*Nematus pieli* Takeuchi, 1938). Russia: **FE** (PR). – China (NE).
- Euura tuberculata** (Benson, 1953) [Nematus]. Host plant: *Salix myrtilloides*, *S. saxatilis*, *S. bebbiana*, *S. integra* (Salicaceae). Russia: **EP** (N, NW), **ES** (YA), **FE** (PR, KU, MG, CH). – Europe (WE, NE).
- Euura tunicata** (Zaddach, 1883) [Nematus] (*Nematus laevis* Brischke, 1883). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (NW, C), **ES** (IR). – Europe (WE, EE, NE), Kyrgyzstan.
- Euura turgaiensis** (Safjanov, 1977) [Nematus]. Russia: **WS** (AL). – Kazakhstan.
- Euura udus** (Holmgren, 1883) [Nematus]. Russia: **EP** (N). – Europe (NE).
- Euura umbrata** (Thomson, 1871) [Nematus] (*Nematus collinus* Cameron, 1882; *Pteronus similis* Forsius, 1911; *Pteronidea verrucosae* Kontuniemi, 1966). Host plant: *Betula* spp. (Betulaceae). Russia: **EP** (NW, E), **ES** (IR). – Europe (WE, EE, NE).
- Euura unigallae** (Vikberg et Zinovjev, 2014) [Pontania]. Host plant: *Salix* spp. (Salicaceae); larvae form galls. Russia: **EP** (NW). – Europe (NE).
- Euura vaginosa** (Konow, 1903) [Pachynematus]. Russia: **ES** (IR). – Europe (NE).
- Euura vaga** (Fabricius, 1781) [Tenthredo] (*Tenthredo flaviventris* Gmelin, 1790; *Nematus leucogaster* Hartig, 1840; *N. hypoleucus* Förster, 1854; *N. micraulius* Förster, 1854; *N. punctipleuris* Thomson, 1863; *N. punctulatus* Thomson, 1863; *N. meridionalis* André, 1880; *N. inconspicuus* W.F. Kirby, 1882; *Pachynematus corticosus* MacGillivray, 1901; *Holcocneme ulbrichti* Enslin, 1910; *Pachynematus roscidus* MacGillivray, 1921). Host plant: *Carex* spp. (Poaceae), *Salix* spp. (Salicaceae). Russia: **EP** (N, NW, C, E, NC), **ES** (IR, YA), **FE** (AM, KA, MG). – Europe (WE, EE, NE), Turkey, Kyrgyzstan, Kazakhstan, Mongolia, China (NW).
- Euura variabilis** (Malaise, 1931) [Amauronematus]. Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (Taege et al., 2006), **FE** (KA, MG).
- Euura varians** (Lindqvist, 1962) [Amauronematus]. Russia: **EP** (Zhelochovtsev, Zinovjev, 1995), **ES** (IR). – Europe (NE).
- Euura variator** (Ruthe, 1859) [Nematus] (*Amauronematus longidens* Lindqvist, 1962). Host plant: *Salix* spp. (Salicaceae). Russia: **FE** (MG). – Europe (WE, EE, NE), N America.
- Euura vastatrix** (Zhelochovtsev, 1935) [Pteronidea]. Host plant: *Ribes* spp. (Grossulariaceae). Russia: **WS** (TK), **ES** (KR, IR), **FE** (AM, KH).
- Euura vastidens** (Lindqvist, 1975) [Amauronematus]. Russia: **ES** (IR).
- Euura venusta** (Brischke, 1883) [Cryptocampus] (*Cryptocampus brevicornis* Zaddach, 1883). Host plant: *Salix* spp. (Salicaceae); larvae in galls on the base of a leaf petiole. Russia: **EP** (NW, C), "Siberia" (Zhelochovtsev, Zinovjev, 1995), **FE** (AM). – Europe (WE, EE, NE), Armenia.
- Euura vesicator** (Bremi-Wolf, 1849) [Nematus] (*Nematus helicinus* Brischke, 1850; *N. leptocerus* Förster, 1854; *N. lugdunensis* Snellen van Vollenhoven, 1871). Host plant: *Salix purpurea*, *S. superpurpurea* (Salicaceae); larvae form galls. Russia: **EP** (NW, C), **ES** (YA). – Europe (WE, EE, NE, SE), Caucasus.
- Euura vicina** (Serville, 1823) [Nematus] (*Nematus vicinus* Lepeletier, 1823; *N. longispinis* Kriechbaumer, 1885; *Holcocneme flavipes* Matsumura, 1912). Host plant: *Salix* spp., *Populus* spp. (Salicaceae), *Betula* spp. (Betulaceae), *Rumex* spp. (Polygonaceae). Russia: **EP** (NW, C, NC), **ES** (IR), **FE** (KH, PR, SA, KA). – Europe (WE, EE, NE, SE), Kyrgyzstan, Kazakhstan, Korean Peninsula, Japan (Hok, Hon), N America.
- Euura viduatinus** (Malaise, 1931) [Amauronematus] (*Amauronematus viduatinus apiciscerra* Malaise, 1931). Russia: **ES** (Zhelochovtsev, Zinovjev, 1995), **FE** (KA). – Europe (NE).
- Euura viduatoides** (Lindqvist, 1960) [Amauronematus]. Host plant: *Salix* spp. (Salicaceae). Russia: **ES** (IR). – Europe (WE, NE).
- Euura viduata** (Zetterstedt, 1838) [Tenthredo] (*Nematus lepidotus* Hartig, 1840; *N. brachyotus* Förster, 1854; *N. luctuosus* Förster, 1854; *N. notatus* Förster, 1854; *N. vagus* Zaddach, 1883; *Amauronematus viduatus lugens* Enslin, 1915; *A. viduatus laetus* Enslin, 1915). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (NW, C), **ES** (IR), **FE** (AM, MG). – Europe (WE, EE, NE, SE), Mongolia, China (NC), N America; introduced into Australia, New Zealand.
- Euura villosa** (Thomson, 1863) [Nematus] (*Nematus arcticus* Holmgren, 1869; *N. brachyacanthus palliditarsus* Cameron, 1876; *N. gelidus* W.F. Kirby, 1882). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (N), **ES** (KR, BR), **FE** (CH). – Europe (NE), N America.
- Euura viminalis** (Linnaeus, 1758) [Cynips] (*Nematus vollenhovenii* Cameron, 1874; *N. interstitialis* Cameron, 1876; *N. nigritarsis* André, 1880; *Pontania ciccum* Konow, 1901; *P. viminalis xanthaspis* Enslin,

1915; *P. hungarica* Enslin, 1918). Host plant: *Salix purpurea*, *S. babylonica* (Salicaceae); larvae form galls. Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (TU, YA), **FE** (AM, PR, SA, KU, KA, MG). – Europe (WE, EE, SE), Armenia, Turkey, Mongolia.

Euura viridis (Stephens, 1835) [Nematus] (*Nematus prasinus* Hartig, 1837; *N. polystipus* Förster, 1854; *N. hypoleucus* Costa, 1859; *N. costae* W.F. Kirby, 1882). Host plant: *Betula* spp. (Betulaceae). Russia: **EP** (N), **ES** (IR, BR), **FE** (SA, MG). – Europe (WE, EE, NE, SE), Kazakhstan.

Euura virilis (Zirngiebl, 1955) [Pontania] (*Pontania rufana* Lacourt, 1973). Host plant: *Salix purpurea* (Salicaceae); larvae form galls. Russia: **EP** (NW). – Europe (WE, EE, NE, SE).

Euura viridissima (Möller, 1882) [Nematus] (*Nematus glutinosae* Cameron, 1882). Host plant: *Alnus* spp. (Betulaceae). Russia: **EP** (N, NW, C, E, S, NC). – Europe (WE, EE, NE, SE).

Euura vittata (Serville, 1823) [Nematus] (*Nematus vittatus* Lepeletier, 1823; *N. melanoleucus* Hartig, 1840; *N. exoletus* Eversmann, 1847; *N. continuus* Eversmann, 1847; *N. scabrilvalvis* Thomson, 1871; *Amauronematus konowi* Enslin, 1915; *A. vittatus sternalis* Enslin, 1915; *A. crispus* Benson, 1948). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (N, NW, C, E), **UR**, **ES** (IR, YA), **FE** (KA, MG). – Europe (WE, EE, NE), Kyrgyzstan.

Euura weiffenbachiella Liston et Vikberg, 2017 (*E. weiffenbachii* Ermolenko, 1988). Host plant: *Salix rosmarinifolia*, *S. repens* (Salicaceae). Russia: **ES** (YA). – Europe (WE, EE, NE).

Euura zaitzevi (Enslin, 1919) [Amauronematus]. – Russia: **EP** (N), **WS** (TM).

HEMICROA Stephens, 1835 (*Leptocerca* Hartig, 1837; *Engages* Gistel, 1848; *Leptocercus* Thomson, 1871; *Varna* Ross, 1937). Type species *Tenthredo australis* Serville, 1823. The larvae live in colonies on the *Alnus*, *Betula* (Betulaceae). The genus is distributed in Holarctic and Oriental region. Number of species: World – 13; Palaearctic – 10; Russia – 4.

Hemicroa australis (Serville, 1823) [Tenthredo] (*Tenthredo alni* Linnaeus, 1767; *T. luctuosa* Hill, 1773; *T. australis* Lepeletier, 1823). Host plant: *Betula* spp., *Alnus* spp. (Betulaceae); larvae live alone. Russia: **EP** (N, NW, C, E), **UR**, **WS** (TK, AL), **ES** (YA). – Europe (WE, EE, NE), Kazakhstan.

Hemicroa crocea (Geoffroy, 1785) [Tenthredo] (*Tenthredo rufa* Panzer, 1799; *Hemicroa stigma*

Stephens, 1835; *Leptocercus nigriceps* Thomson, 1871; *Dineura unicolor* Rudow, 1872; *D. americana* Provancher, 1882; *Nematus ardens* Zaddach, 1883; *Dineura pallida* Ashmead, 1890; *Hemicroa dyari* Rohwer, 1918; *H. orientalis* Rohwer, 1921; *H. washingtonia* Rohwer, 1932). Host plant: *Alnus* spp., *Betula* spp., *Corylus* spp. (Betulaceae), *Salix* spp. (Salicaceae); larvae live in groups. Russia: **EP** (N, NW, C, E, NC), **UR**, **ES** (BR, YA), **FE** (AM, KH, PR, SA). – Europe (WE, EE, NE), China (Wei et al., 2006), India, SE Asia, N America.

Hemicroa haematopygia Zhelochovtsev, 1935. Host plant: *Alnus* spp., *Betula* spp. (Betulaceae). Russia: **ES** (IR, BR, YA), **FE** (AM, PR, SA, MG). – Korean Peninsula.

Hemicroa paramushirensis Takeuchi, 1933. Host plant: *Alnus* spp. (Betulaceae). Russia: **FE** (SA, KU). – Japan (Hon).

HOPLOCAMPA Hartig, 1837 (*Macgillivraya* Ashmead, 1898; *Macgillivrayella* Ashmead, 1900). Type species *Tenthredo brevis* Klug, 1816. The larvae in fruits of Rosaceae. The genus is distributed in Holarctic; two species in Oriental region. Number of species: World – > 40; Palaearctic – > 20; Russia – 11.

Hoplocampa alpina (Zetterstedt, 1838) [Tenthredo] (*Selandria pallida* Newman, 1837). Host plant: *Sorbus* spp. (Rosaceae). Russia: **EP** (N, C), **WS** (TM), **ES** (KR). – Europe (WE, EE, NE), Japan (Hon).

Hoplocampa brevis (Klug, 1816) [Tenthredo] (*Tenthredo fallax* Serville, 1823; *T. fallax* Lepeletier, 1823; *T. pyri* Vallot, 1848). Host plant: *Pyrus* spp., *Malus* spp. (Rosaceae). Russia: **EP** (NC, CR). – Europe (WE, EE, SE), Caucasus, SW Asia; introduced into N America.

Hoplocampa chrysorrhoea (Klug, 1816) [Tenthredo] (*H. chrysorrhoea nigrita* Enslin, 1914). Host plant: *Prunus* spp. (Rosaceae). Russia: **EP** (NC, CR). – Europe (WE, EE), Cyprus.

Hoplocampa crataegi (Klug, 1816) [Tenthredo] (*Tenthredo pallida* Serville, 1823; *T. luteola* Serville, 1823; *T. verticata* Serville, 1823; *T. pallida* Lepeletier, 1823; *T. luteola* Lepeletier, 1823; *T. verticata* Lepeletier, 1823). Host plant: *Crataegus* spp. (Rosaceae). Russia: **EP** (NC). – Europe (WE, EE, NE, SE).

Hoplocampa ephippiata Konow, 1897. Host plant: *Malus* spp. (Rosaceae). Russia: **WS** (AL), **ES** (IR, BR), **FE** (AM, KH, PR).

Hoplocampa flava (Linnaeus, 1760) [Tenthredo] (*Tenthredo ruficapilla* Gmelin, 1790; *T. glaucopis*

Rossi, 1790; *Allantus ferrugineus* Panzer, 1803; *Hylotoma ferruginea* Fabricius, 1804; *H. simplex* Fallén, 1807; *Tenthredo brunnea* Klug, 1816; *Hoplocampa flava dimidiata* Costa, 1894; *Tomostethus testaceus* Niegabitowski, 1899). Host plant: *Prunus* spp. (Rosaceae). Russia: EP (S, NC). – Europe (WE, EE, NE, SE), Caucasus, SW Asia.

Hoplocampa fulvicornis (Panzer, 1801) [Tenthredo] (*Tenthredo rutilicornis* Klug, 1816; *T. haemorrhoidalis* Serville, 1823; *T. haemorrhoidalis* Lepeletier, 1823; *Hoplocampa rutilicornis pleuris* Zirngiebl, 1954; *H. prunicola* Benson, 1968). Host plant: *Prunus* spp. (Rosaceae). Russia: EP (CR). – Europe (WE, EE, NE, SE).

Hoplocampa minuta (Christ, 1791) [Tenthredo] (*Tenthredo hylotomoides* Serville, 1823; *T. parvula* Serville, 1823; *T. hylotomoides* Lepeletier, 1823; *T. parvula* Lepeletier, 1823; *T. turcarum* Vallot, 1848; *Hoplocampa fabricii* W.F. Kirby, 1882; *H. minuta dudai* Gregor, 1942). Host plant: *Prunus* spp. (Rosaceae). Russia: EP (NW, CR). – Europe (WE, EE, NE, SE), Armenia, Syria, Uzbekistan, China (NW).

Hoplocampa phantom Zinovjev, 1993. Host plant: *Sorbus* spp. (Rosaceae). Russia: UR, ES (Zhe-lochovtsev, Zinovjev, 1995), FE (PR).

Hoplocampa plagiata (Klug, 1816) [Tenthredo]. Host plant: *Amelanchier* spp., *Crataegus* spp. (Rosaceae). Russia: EP (S). – Europe (WE, EE, SE), Caucasus.

Hoplocampa testudinea (Klug, 1816) [Tenthredo]. Host plant: *Malus* spp., *Pyrus* spp. (Rosaceae). Russia: EP (C, S, NC, CR). – Europe (WE, EE, NE, SE), SW Asia; introduced into N America.

MEGADINEURA Malaise, 1931 (*Stenomesoneura* Wei, 1998). Type species *Dineura grandis* André, 1882. The genus is distributed in East Asia; one species in Oriental region. Number of species: World and Palaearctic – 5; Russia – 1.

Megadineura grandis (André, 1882) [Dineura]. Russia: ES (IR), FE (PR). – China (NE), Korean Peninsula, Japan (Takeuchi, 1952).

MESONEURA Hartig, 1837 (*Pristis* Brullé, 1846; *Mesonevra* Agassiz, 1848; *Lisconeura* Rohwer, 1908). Type species *Tenthredo opaca* Fabricius, 1775. The larvae feed on *Quercus* (Fagaceae). The genus is distributed in Palaearctic. Number of species: World and Palaearctic – 9; Russia – 3.

Mesoneura koreana Ryu, Kim et Lee, 1991. Host plant: *Quercus mongolica* (Fagaceae). Russia: FE (PR). – Korean Peninsula.

Mesoneura lanigera Benson, 1954. Host plant: *Quercus* spp. (Fagaceae). Russia: EP (NC, CR). – Europe (EE), Cyprus.

Mesoneura opaca (Fabricius, 1775) [Tenthredo] (*Tenthredo verna* Klug, 1816; *T. punctigera* Lepeletier, 1823; *T. punctigera* Serville, 1823; *Selandria biloba* Stephens, 1835; *Dineura pallipes* Hartig, 1837; *D. dorsalis* Förster, 1844; *D. selandriiformis* Cameron, 1875; *Mesoneura opaca nigerrima* Enslin, 1914; *M. opaca lucida* Enslin, 1914; *M. opaca obscuriventris* Enslin, 1914). Host plant: *Quercus* spp. (Fagaceae). Russia: EP (C, S, NC, CR). – Europe (WE, EE, NE, SE), Caucasus.

NEMATINUS Rohwer, 1911. Type species *Nematus fuscipennis* Lepeletier, 1823. The larvae feed on Betulaceae. The genus is distributed in Holarctic. Number of species: World – 26; Palaearctic – 22; Russia – 9.

Nematinus acuminatus (Thomson, 1871) [Nematus] (*Nematus nebulosus* Stein, 1881). Host plant: *Betula* spp., *Corylus* spp., *Alnus* spp. (Betulaceae). Russia: EP (NW, C, E), UR, WS (AL), ES (IR, YA), FE (AM, SA, MG). – Europe (WE, EE, NE, SE), Korean Peninsula, Japan (Hok).

Nematinus aterrimus Vikberg, 1982 (*Pteronidea pontanioides* Malaise, 1931). Russia: FE (SA, KA). – Japan (Hok, Hon).

Nematinus bilineatus (Klug, 1819) [Tenthredo] (*Nematus flavus* Gimmerthal, 1844; *N. luteus trimaculatus* Cameron, 1876; *N. flavicans* W.F. Kirby, 1882; *Nematinus bilineatus virilis* Enslin, 1915; *N. bilineatus muliebris* Enslin, 1915; *N. bilineatus luteiformis* Enslin, 1915). Host plant: *Betula* spp. (Betulaceae). Russia: EP (NW). – Europe (WE, EE, NE, SE), Turkey.

Nematinus caledonicus (Cameron, 1882) [Nematus] (*N. nigrosternatus* Malaise, 1931). Host plant: *Betula* spp. (Betulaceae). Russia: EP (NW, C), ES (IR), FE (PR, KU, KA). – Europe (WE, NE).

Nematinus fuscipennis (Serville, 1823) [Nematus] (*Tenthredo rufa* Schrank, 1802; *T. alnina* Bechstein et Scharfenberg, 1805; *Nematus fuscipennis* Lepeletier, 1823; *N. fumipennis* Stephens, 1835; *N. gravenhurstii* Gimmerthal, 1836). Host plant: *Alnus* spp., *Carpinus* spp. (Betulaceae). Russia: EP (N, NW, C, E, S). – Europe (WE, EE, NE, SE).

Nematinus kamtchaticus Malaise, 1931. Russia: ES (KR), FE (SA, KA, MG, CH).

Nematinus luteus (Panzer, 1803) [Nematus] (*Tenthredo lutea* Fabricius, 1804; *Nematinus willigiae*

- Stein, 1926; *N. willigiae pilosus* Benson, 1958). Host plant: Alnus spp., Corylus spp. (Betulaceae). Russia: **EP** (N, NW, C, E, NC). – Europe (WE, EE, NE), Japan (Hok, Hon).
- Nematinus rubrocaudatus** Takeuchi, 1956. Russia: **FE** (KU). – Japan (Hon).
- Nematinus steini** Blank, 1998 (*Tenthredo alneti* Bechstein et Scharfenberg, 1805). Host plant: Alnus spp. (Betulaceae). Russia: **EP** (NW, C). – Europe (WE, EE, NE).
- NEMATUS** Panzer, 1801 (*Nematus* Jurine, 1801; *Nematus* Jurine, 1807; *Craesus* Leach, 1817; *Hypolaepus* W.F. Kirby, 1882; *Paranematus* Zinovjev, 1978). Type species *Tenthredo lucida* Panzer, 1801. The larvae feed on the deciduous trees and shrubs. The genus is distributed mainly in Holarctic, few species in Oriental region. Number species: World – about 40; Palaearctic – about 35; Russia – 20 (+ 1 ssp.).
- Nematus alniastri** (Scharfenberg, 1805) [*Tenthredo*] (*N. varus* Villaret, 1832; *Craesus varus ligeris* Pic, 1923). Host plant: Alnus spp. (Betulaceae). Russia: **EP** (N, NW, C, E, NC). – Europe (WE, EE, NE, SE), Caucasus, Turkey; introduced into N America.
- Nematus brischkei** Zaddach, 1876. Host plant: Carpinus spp., Corylus spp. (Betulaceae). Russia: **EP** (NW). – Europe (WE, EE).
- Nematus coeruleus** Zinovjev, 1978. Host plant: Lonicera coerulea (Caprifoliaceae). Russia: **EP** (NW).
- Nematus egregius** Kajmuk, 1986. Host plant: Crataegus spp. (Rosaceae). Russia: **ES** (YA).
- Nematus inornatus** (Takeuchi, 1936) [*Holcocneme*]. Host plant: Salix spp. (Salicaceae), Vaccinium spp. (Ericaceae). Russia: **FE** (KU). – Japan (Hon, Sado).
- Nematus jakowleffi** (Konow, 1902) [*Holcocneme*]. Host plant: Salix spp. (Salicaceae), ?Betula spp. (Betulaceae). Russia: **ES** (IR), **FE** (AM).
- Nematus japonicus** (Takeuchi, 1921) [*Craesus*] (*N. nigromaculatus* Malaise, 1931). Host plant: Alnus spp. (Betulaceae). Russia: **FE** (PR, SA, KU, KA). – Korean Peninsula, Japan (Hok, Hon).
- Nematus latipes** Villaret, 1832. Host plant: Betula spp. (Betulaceae). Russia: **EP** (C, E), **ES** (BR, YA). – Europe (WE, EE, NE, SE), Kazakhstan.
- Nematus lonicerae** (Weiffenbach, 1957) [*Pachyne-matus*]. Host plant: Lonicera spp., Symphoricarpos spp. (Caprifoliaceae). Russia: **EP** (Zhelochovtsev, Zinovjev, 1995). – Europe (WE, NE, SE).
- Nematus lucens** (Enslin, 1918) [*Pteronidea*] (*Mesoneura nigrostigmata* Haris, 2001). Russia: **EP** (C). – Europe (EE).
- Nematus lucidus** (Panzer, 1801) [*Tenthredo*] (*N. cinctus* Serville, 1823; *N. cinctus* Lepeletier, 1823; *Holcocneme lucidus rufa* Zirngiebl, 1937). Host plant: Crataegus spp., Prunus spp. (Rosaceae). Russia: **EP** (NC), **ES** (YA), **FE** (KU). – Europe (WE, EE, NE, SE), Cyprus, Japan (Hok, Hon).
- Nematus mandshuricus** Zhelochovtsev et Zinojev, 1994. Host plant: Lonicera spp. (Caprifoliaceae). Russia: **ES** (ZB), **FE** (KH).
- Nematus nigrodorsatus** Malaise, 1931. Russia: **FE** (KA).
- Nematus princeps** Zaddach, 1876. Host plant: Salix spp. (Salicaceae). Russia: **EP** (NW), **UR**, **WS** (AL), **ES** (IR, BR, YA), **FE** (AM, PR, SA, KU, MG). – Europe (WE, EE, NE), Mongolia, Japan (Hok, Hon, Shi).
- Nematus septentrionalis** (Linnaeus, 1758) [*Tenthredo*] (*Tenthredo alni* Linnaeus, 1758; *T. largipes* Retzius, 1783; *Nematus laticrus* Villaret, 1832; *Craesus stephensi* Newman, 1837). Host plant: Betulaceae, Oleaceae, Salicaceae. Russia: **WS** (AL), **ES** (KS). – Europe (WE, EE, NE, SE), Armenia, Turkey.
- Nematus shinoharai** (Beneš, 1990) [*Craesus*]. Host plant: Juglans spp. (Juglandaceae). Russia: **FE** (PR). – Japan (Hok, Hon).
- Nematus tataricus** Zinovjev, 1978. Host plant: Lonicera tatarica, L. xylosteum, Symphoricarpu racemosus (Caprifoliaceae). Russia: **EP** (NW, C), **UR**. – Europe (WE, NE), Kazakhstan.
- Nematus tulunensis** Vikberg, 1972. Host plant: Lonicera spp., Symphoricarpu sp. (Caprifoliaceae). Russia: **EP** (NW, C), **UR**, **ES** (IR, YA). – Europe (NE).
- Nematus wahlbergi tavastiensis** Vikberg, 1972. Host plant: Lonicera spp. (Caprifoliaceae). Russia: **EP** (N). – Europe (NE).
- Nematus wahlbergi wahlbergi** Thomson, 1871 (*N. sardiniensis* Costa, 1886). Host plant: Lonicera spp. (Caprifoliaceae). Russia: **EP** (NW, C). – Europe (WE, EE, NE, SE).
- Nematus yokohamensis** (Konow, 1895) [*Holcocneme*]. Host plant: Lonicera spp. (Caprifoliaceae). Russia: **ES** (IR, BR), **FE** (PR, SA). – China (NE), Korean Peninsula, Japan (Hon).
- PLATYCAMPUS** Schiødte, 1839 (*Leptopus* Hartig, 1837; *Erasminus* Gistel, 1848; *Camponiscus* Newman, 1869). Type species *Nematus hypogastricus* Hartig, 1837. The larvae with strongly flattened body; live on the lower surface of leaves of woody plants. The genus is distributed in East Palaearctic; one species in West Europe, and one species

in Oriental region. Number of species: World and Palaearctic – 8; Russia – 6.

Platycampus amaculatus Verzhutskii, 1966. Host plant: *Betula* spp. (Betulaceae). Russia: **ES** (IR), **FE** (KH, PR).

Platycampus amurensis Zinovjev, 1986. Host plant: *Corylus* spp. (Betulaceae). Russia: **FE** (AM, PR).

Platycampus coryli Zinovjev, 1986. Host plant: *Corylus* spp. (Betulaceae). Russia: **FE** (AM, PR).

Platycampus luridiventris (Fallén, 1808) [Tenthredo] (*Tenthredo alnicola* Bechstein et Scharfenberg, 1805; *Nematus hypogastricus* Hartig, 1837; *N. alnivorus* Hartig, 1840; *N. rufipes* Tischbein, 1846; *Leptopus rufipes* Förster, 1854; *Nematus protensus* Förster, 1854; *Camponiscus healaei* Newman, 1869; *Nematus fennicus* André, 1880; *N. tischbeini* André, 1880; *N. alnicola* Zaddach, 1883; *N. cellularis* Brischke, 1884; *Platycampus luridiventris pleuritica* Enslin, 1915). Host plant: *Alnus* spp., *Corylus* spp., *Betula* spp. (Betulaceae). Russia: **EP** (NW, C), **UR**. – Europe (WE, EE, NE), China (Wei et al., 2006), Japan (Hon).

Platycampus speciosus Zinovjev, 1986. Host plant: *Alnus* spp. (Betulaceae). Russia: **FE** (AM, KH, PR).

Platycampus zhelochovtsevi Zinovjev, 1986. Host plant: *Alnus* spp. (Betulaceae). Russia: **FE** (AM, KH, PR, SA).

PRISTIPHORA Latreille, 1810 (*Diphadnus* Hartig, 1837; *Lygaeonematus* Konow, 1890; *Micronematus* Konow, 1890; *Gymnonychus* Marlatt, 1896; *Neopareophora* MacGillivray, 1908; *Neotomostethus* MacGillivray, 1908; *Dineuridea* Rohwer, 1912; *Sala* Ross, 1937; *Pristola* Ross, 1945; *Lygaeotus* Lindqvist, 1952; *Lygaeophora* Lindqvist, 1952; *Nepionema* Benson, 1960; *Melastola* Wong, 1968; *Sharliphora* Wong, 1969; *Oligonematus* Zhelochovtsev, 1988; *Lygaeotus* Liston, 1993; *Lygaeophora* Liston, 1993; *Pristicampus* Zinovjev, 1993). Type species *Pteronus testaceus* Jurine, 1807. The larvae feed on various plants, mainly on deciduous, rarely on coniferous. The genus is distributed in Holarctic, Oriental and Neotropical regions. Number of species: World – about 220; Palaearctic – about 170; Russia – 89.

Pristiphora abbreviata (Hartig, 1837) [Nematus] (*Gymnonychus californicus* Marlatt, 1896). Host plant: *Pyrus* spp. (Rosaceae). Russia: **EP** (NW, C, S, NC), **FE** (AM, KH, PR). – Europe (WE, EE, NE,

SE), Caucasus, Turkey, Kyrgyzstan; introduced into N America.

Pristiphora abietina (Christ, 1791) [Tenthredo] (*Tenthredo pini* Retzius, 1783; *T. abietina* Christ, 1791; *T. abietum* Hartig, 1834; *Nematus abietinus* Dahlbom, 1835; *N. truncatus* Hartig, 1837). Host plant: *Picea* spp. (Pinaceae). Russia: **EP** (NW, C), **ES** (YA). – Europe (WE, EE, NE, SE), Kyrgyzstan.

Pristiphora acutidens Lindqvist, 1977. Russia: **ES** (IR).

Pristiphora affinis (Lindqvist, 1952) [Lygaeonematus]. Host plant: *Salix* spp. (Salicaceae). Russia: **ES** (KR). – Europe (NE).

Pristiphora albilabris (Boheman, 1852) [Nematus] (*Nematus albilabris* Thomson, 1863; *N. collaris* Stein, 1884). Host plant: *Betula* spp. (Betulaceae). Russia: **EP** (N, E), **ES** (IR). – Europe (WE, NE).

Pristiphora albitibia (Costa, 1859) [Nematus] (*Nematus puncticeps* Thomson, 1863; *N. agilis* Zaddach, 1884; *Pristiphora aterrima* Lindqvist, 1977; *P. nigropuncticeps* Haris, 2002). Host plant: *Vicia* spp. (Fabaceae). Russia: **EP** (NW, NC), **WS** (Zhelochovtsev, Zinovjev, 1995), **ES** (IR, BR, YA). – Europe (WE, EE, NE, SE), Mongolia.

Pristiphora alpestris (Konow, 1903) [Pachynematus] (*Lygaeonematus karvoneni* Lindqvist, 1952). Host plant: *Betula pendula*, *B. pubescens* (Betulaceae). Russia: **EP** (N), **ES** (IR). – Europe (WE, EE, NE).

Pristiphora amphibola (Förster, 1854) [Nematus] (*Nematus laetus* Cameron, 1883; *N. fraternus* Cameron, 1885). Host plant: *Picea* spp. (Pinaceae). Russia: **EP** (NW). – Europe (WE, EE, NE).

Pristiphora angulata Lindqvist, 1974. Host plant: *Spiraea chamaedryfolia*, *S. salicifolia* (Rosaceae). Russia: **EP** (NW). – Europe (NE), Mongolia.

Pristiphora aphantoneura (Förster, 1854) [Nematus] (*Tenthredo fulvipes* Fallén, 1808; *Cryptocampus distinctus* Costa, 1882; *Pristiphora pygmaea* Lindqvist, 1964). Host plant: *Lathyrus pratensis* (Fabaceae). Russia: **EP** (N, NW, C, E, NC), **UR**. – Europe (WE, EE, NE, SE), Caucasus, Turkey, Kyrgyzstan, Kazakhstan, Mongolia, China (NW), Korean Peninsula.

Pristiphora appendiculata (Hartig, 1837) [Nematus] (*P. pallipes* Lepeletier, 1823; *P. pallipes* Serville, 1823; *Nematus flavipes* Dahlbom, 1835; *N. fusicornis* Hartig, 1837; *N. enervis* Herrich-Schäffer, 1840; *N. cathoraticus* Förster, 1854; *N. pallicornis* Norton, 1861; *N. pallicornis labratus* Norton, 1861; *Pristiphora grossulariae* Walsh, 1866; *Nematus peletieri* André, 1880; *N. hypobalius* Zaddach, 1884;

N. pumilus Zaddach, 1884; *N. ghilianii* Costa, 1894; *Pristiphora anivskiensis* Haris, 2006). Host plant: Ribes spp. (Grossulariaceae). Russia: **EP** (N, NW, C, E, S, NC), **ES** (IR, BR), **FE** (KH, PR, SA, MG). – Europe (WE, EE, NE, SE), Turkey, Kazakhstan, Mongolia, China (NC), Korean Peninsula, Japan, N America.

Pristiphora apricoti Zinovjev, 1993. Russia: **FE** (PR). – Korean Peninsula.

Pristiphora armata (Thomson, 1863) [Nematus] (*Nematus crassicornis* Hartig, 1837; *N. crataegi* Brischke, 1883; *N. fletcheri* Cameron, 1884; *N. melanostomus* Zaddach, 1884; *N. ensicornis* Jacobs, 1884; *N. nigricollis* Cameron, 1885). Host plant: Crataegus spp. (Rosaceae). Russia: **EP** (NW, C, S, NC), **FE** (SA, KA). – Europe (WE, EE, NE, SE), Turkey, Kazakhstan.

Pristiphora bicalis (Förster, 1854) [Nematus] (*Nematus conspersus* Zaddach, 1883; *N. pruni* Brischke, 1883; *N. lateralis* Brischke, 1885; *N. postumus* Dalla Torre, 1894). Host plant: Prunus spinosa (Rosaceae). Russia: **EP** (NW, C, NC), **ES** (IR, BR), **FE** (SA). – Europe (WE, EE), Turkmenistan.

Pristiphora borea (Konow, 1904) [Lygaeonematus] (*Nematus astutus* Cameron, 1885; *Pachynematus lapponicus* Enslin, 1916). Host plant: Betula nana (Betulaceae). Russia: **EP** (N), **WS** (TM), **ES** (IR, YA, ZB), **FE** (MG). – Europe (WE, EE, NE), N America.

Pristiphora breadalbanensis (Cameron, 1882) [Nematus] (*Lygaeonematus tromsouensis* Kiær, 1898; *L. corpulentus* Konow, 1904; *L. arcticola* Enslin, 1916). Russia: **EP** (N), **ES** (YA), **FE** (CH). – Europe (WE, NE), N America.

Pristiphora bufo (Brischke, 1883) [Nematus] (*Pristiphora pallidula* Konow, 1902; *Pachynematus crassicauda* Lindqvist, 1964; *Pristiphora laricicola* Verzhutskii, 1966). Host plant: Larix spp. (Pinaceae). Russia: **EP** (NW), **WS** (TM), **ES** (TU, IR, BR, YA), **FE** (MG). – Europe (WE, EE, NE), Mongolia.

Pristiphora carinata (Hartig, 1837) [Nematus] (*Tenthredo pallipes* Fallén, 1808; *Nematus canaliculatus* Hartig, 1840; *N. denudatus* Hartig, 1840; *Pachynematus alticola* Enslin, 1916). Host plant: Vaccinium myrtillus (Ericaceae). Russia: **EP** (N, NW, C, E), **UR**, **WS** (AL), **ES** (KR, IR, YA, ZB), **FE** (KA, MG). – Europe (WE, EE, NE), Kyrgyzstan, Mongolia, China (NC).

Pristiphora chromata Rohwer, 1925. Russia: **FE** (PR).

Pristiphora cincta Newman, 1837 (*Nematus quercus* Hartig, 1837; *Tenthredo borealis* Zetterstedt, 1838; *Pristiphora identidem* Norton, 1867; *P. idiota* Norton, 1867; *P. coloradensis* Marlatt, 1896; *P. hoodi* Marlatt, 1896; *P. seorsa* Konow, 1897; *P. idiotiformis* Rohwer, 1910). Host plant: Vaccinium spp. (Ericaceae), Betula spp. (Betulaceae), Salix spp. (Salicaceae). Russia: **EP** (N, C, NC), **ES** (IR, BR, YA), **FE** (AM, KH, PR, SA, KA, MG). – Europe (WE, EE, NE), Kazakhstan, Mongolia, Korean Peninsula, Japan (Hon), N America.

Pristiphora coactula (Ruthe, 1859) [Nematus] (*Nematus winnipeg* Norton, 1867; *N. alpinus* Thomson, 1871; *N. winnipegensis* W.F. Kirby, 1882; *Lygaeonematus pachyvalvis* Konow, 1904; *Pristiphora dawsoni* Rohwer, 1908). Host plant: Vaccinium uliginosum (Ericaceae). Russia: **EP** (N, NW, C), **WS** (TM, AL), **ES** (TU, KR, IR, BR, YA, ZB), **FE** (MG). – Europe (WE, NE), N America.

Pristiphora compressa (Hartig, 1837) [Nematus]. Host plant: Picea spp. (Pinaceae). Russia: **EP** (NW, C, E), **WS** (AL), **ES** (TU, KR, IR). – Europe (WE, EE, NE), Mongolia, China (NC).

Pristiphora condei Lindqvist, 1955. Host plant: Sorbus aucuparia, Amelanchier spicata (Rosaceae). Russia: **EP** (NW, C), **FE** (PR). – Europe (NE, EE).

Pristiphora confusa Lindqvist, 1955. Host plant: Salix spp. (Salicaceae). Russia: **EP** (N), **FE** (PR). – Europe (WE, EE, NE).

Pristiphora conjugata (Dahlbom, 1835) [Nematus] (*Nematus gonymelas* Stephens, 1835; *N. discipennis* Herrich-Schäffer, 1840; *N. discoidalis* Thomson, 1888; *Pristiphora conjugata ulbrichti* Enslin, 1916; *P. conjugata forsiusi* Enslin, 1916). Host plant: Populus spp., Salix spp. (Salicaceae). Russia: **EP** (N, NW, C, E, S, NC), **ES** (IR, YA), **FE** (AM, PR, KA). – Europe (WE, EE, NE, SE), Armenia, Turkey, Iran, Kyrgyzstan, China (NW), Japan.

Pristiphora dasiphorae (Zinovjev, 1993) [Pristicampus]. Host plant: Potentilla fruticosa (Rosaceae). Russia: **ES** (IR, ZB), **FE** (MG).

Pristiphora decipiens (Enslin, 1916) [Lygaeonematus]. Host plant: Picea spp. (Pinaceae). Russia: **EP** (C), **ES** (YA). – Europe (WE, EE, NE), Kyrgyzstan.

Pristiphora dochmocera (Thomson, 1871) [Nematus] (*Tenthredo flavipes* Zetterstedt, 1838; *Nematus congener* W.F. Kirby, 1882; *Pristiphora thomsoni* Lindqvist, 1953). Russia: **EP** (Taeger et al., 2006), **WS** (Zhelochovtsev, Zinovjev, 1995), **FE** (AM). – Europe (NE), Turkey.

- Pristiphora erichsonii** (Hartig, 1837) [Nematus] (*Nematus leachii* Dahlbom, 1835; *N. notabilis* Cresson, 1880). Host plant: Larix spp. (Pinaceae). Russia: **EP** (NW, C), **ES** (IR, BR, YA, ZB), **FE** (AM, KH, SA, KU). – Europe (WE, EE, NE, SE), China (NE), Japan (Hon); introduced into N America.
- Pristiphora friesei** (Konow, 1904) [Lygaeonematus] (*Pristiphora atrata* Lindqvist, 1975). Host plant: Larix decidua, L. eurolepis (Pinaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TM), **ES** (IR). – Europe (WE, EE, NE, SE).
- Pristiphora frigida** (Boheman, 1865) [Nematus] (*P. adelungi* Konow, 1902; *P. gelida* Wong, 1968). Russia: **EP** (N). – Europe (NE), N America.
- Pristiphora geniculata** (Hartig, 1840) [Nematus] (*Nematus cheilon* Zaddach, 1876). Host plant: Sorbus spp. (Rosaceae). Russia: **EP** (NW, NC), **ES** (YA), **FE** (KA). – Europe (WE, EE, NE), Kyrgyzstan, Mongolia, China (NC); introduced into N America.
- Pristiphora gerula** (Konow, 1904) [Lygaeonematus]. Host plant: Picea spp. (Pinaceae). Russia: **EP** (Taege et al., 2006). – Europe (WE, EE, NE).
- Pristiphora glauca** Benson, 1954 (*Pachynematus laricivorus* Takagi, 1931; *Pristiphora takagii* Wong, 1975). Host plant: Larix spp. (Pinaceae). Russia: **EP** (N), **ES** (IR, BR), **FE** (PR, SA). – Europe (WE, NE), Mongolia, Korean Peninsula, Japan (Hon).
- Pristiphora groenblomi** (Lindqvist, 1952) [Lygaeonematus]. Host plant: Betula spp. (Betulaceae). Russia: **EP** (N, E), **ES** (IR). – Europe (WE, EE, NE), Mongolia, Korean Peninsula, N America.
- Pristiphora insularis** Rohwer, 1910 (*Amauronematus amelanchieris* Takeuchi, 1922; *Pristiphora kamtchatica* Malaise, 1931; *P. luteiventris* Koch, 1989). Host plant: Amelanchier asiatica, Chaenomeles japonica, Rosa spp. (Rosaceae). Russia: **EP** (N), **ES** (IR), **FE** (KA, MG). – Europe (WE, NE, SE), N Africa, Japan (Hon), N America.
- Pristiphora jacutica** Beneš, 2014. Host plant: Picea obovata (Pinaceae). Russia: **ES** (YA).
- Pristiphora jakowlewi** (Jakovlev, 1891) [Lygaeonematus]. Russia: **ES** (IR).
- Pristiphora kuznetzovorum** (Enslin, 1919) [Lygaeonematus]. Russia: **EP** (N), **WS** (TM).
- Pristiphora lanifica** (Zaddach, 1883) [Nematus]. Russia: **EP** (NW). – Europe (WE, NE).
- Pristiphora laricis** (Hartig, 1837) [Nematus] (*Nematus ruficollis* Hartig, 1840; *N. leucocnemis* Förster, 1854; *N. oblongus* Cameron, 1882; *N. laricivorus* Brischke, 1883; *N. rusticanus* Brischke, 1884;

- Pachynematus ravidus** Konow, 1903; *Lygaeonematus paedidus* Konow, 1904; *Pachynematus nigricorpus* Takagi, 1931). Host plant: Larix spp. (Pinaceae). Russia: **EP** (NW), **UR**, **WS** (NS), **ES** (KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, MG). – Europe (WE, EE, NE, SE), Mongolia, China (NE), Korean Peninsula, Japan.
- Pristiphora lativentris** (Thomson, 1871) [Nematus] (*Nematus scoticus* Cameron, 1881; *N. extremus* Holmgren, 1883; *Pristiphora bucola* Kincaid, 1900; *Lygaeonematus alpicola* Konow, 1904; *L. pallipes femoralis* Zirngiebl, 1953). Russia: **EP** (N), **WS** (TM, AL), **ES** (TU, KR, IR), **FE** (MG). – Europe (WE, NE), Mongolia, N America.
- Pristiphora leucopodia** (Hartig, 1837) [Nematus] (*Nematus nitens* Borries, 1896; *Pachynematus sagittatus* Konow, 1903; *Nematus piceae* Zhelochovtsev, 1988; *Pristiphora hoverlaensis* Haris, 2001). Host plant: Picea spp. (Pinaceae). Russia: **EP** (NW, C), **UR**, **WS** (Zhelochovtsev, Zinovjev, 1995), **ES** (YA). – Europe (WE, EE, NE).
- Pristiphora leucopus** (Hellén, 1948) [Nematus] (*Nematus vitreipennis* Eversmann, 1864). Host plant: Tilia spp. (Tiliaceae). Russia: **EP** (NW, C, E), **UR**. – Europe (WE, EE, NE).
- Pristiphora longicornis** (Malaise, 1931) [Pachyne-
matus]. Russia: **FE** (KA).
- Pristiphora luteipes** Lindqvist, 1955. Host plant: Salix spp. (Salicaceae). Russia: **EP** (N, C). – Europe (WE, EE, NE, SE).
- Pristiphora maesta** (Zaddach, 1876) [Nematus] (*Nematus brevicornis* Thomson, 1863; *N. parvicornis* W.F. Kirby, 1882; *Pachynematus insularis* Malaise, 1921). Host plant: Malus spp. (Rosaceae). Russia: **EP** (NW, C, NC), **ES** (ZB), **FE** (PR, KA). – Europe (WE, EE, NE), Caucasus.
- Pristiphora malaisei** (Lindqvist, 1952) [Lygaeonematus] (*Mesoneura arctica* Lindqvist, 1959; *Pachynematus incisus* Lindqvist, 1970; *P. intermedius* Verzhutskii, 1974; *Pristiphora mongololaricis* Haris, 2003). Host plant: Potentilla fruticose, ?Comarum palustre (Rosaceae). Russia: **ES** (IR), **FE** (MG). – Europe (WE, NE), Mongolia.
- Pristiphora melanocarpa** (Hartig, 1840) [Nematus] (*Nematus funerulus* Costa, 1859; *N. wuestneii* Stein, 1885; *Pristiphora ortingia* Kincaid, 1900). Host plant: Betula spp. (Betulaceae). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (Zhelochovtsev, Zinovjev, 1995), **ES** (Zhelochovtsev, Zinovjev, 1995), **FE** (KH,

PR, SA). – Europe (WE, EE, NE, SE), Kyrgyzstan, China (NW), Korean Peninsula, N America.

Pristiphora micronematica Malaise, 1931 (*Lygaeonematus leucostoma* Lindqvist, 1952). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (NW), **ES** (Zhelochovtsev, 1988), **FE** (KA). – Europe (WE, NE), N America.

Pristiphora mollis (Hartig, 1837) [Nematus] (*Tenthredo bipunctata* Gmelin, 1790; *Nematus mollis* Hartig, 1837; *Tenthredo lapponica* Zetterstedt, 1838; *Nematus whitei* Cameron, 1878; *Pachymenatus orarius* Kincaid, 1900; *P. kontkaneni* Lindqvist, 1960). Host plant: *Vaccinium myrtillus*, *V. uliginosum* (Ericaceae). Russia: **EP** (N, NW, C, E), **ES** (IR, BR, YA), **FE** (AM, MG, CH). – Europe (WE, EE, NE), Mongolia, Korean Peninsula, N America.

Pristiphora monogyniae (Hartig, 1840) [Nematus] (*Nematus pullus* Förster, 1854; *N. filicornis* Thomson, 1863; *N. hibernicus* Cameron, 1878; *N. nanus* Zaddach, 1884; *N. serotinus* Zaddach, 1884; *N. catulus* Zaddach, 1884; *N. ludens* Costa, 1894). Host plant: *Prunus* spinose, *P. domestica* (Rosaceae). Russia: **EP** (NW, NC). – Europe (WE, EE, NE, SE), Caucasus.

Pristiphora nestor (Zhelochovtsev, 1981) [Nematus]. Russia: **ES** (IR), **FE** (MG).

Pristiphora nigella (Förster, 1854) [Nematus] (*Tenthredo ambigua* Fallén, 1808; *Nematus occultus* Förster, 1854; *N. furvescens* Cameron, 1876; *N. obscurus* Zaddach, 1884; *N. xanthomus* Zaddach, 1884; *N. obscurior* Dalla Torre, 1894). Host plant: *Picea abies* (Pinaceae). Russia: **EP** (NW, C, NC), **ES** (YA). – Europe (WE, EE, NE), Turkey.

Pristiphora nigricans (Eversmann, 1847) [Nematus] (*Nematus anderschi* Zaddach, 1876; *Pristiphora inocreata* Konow, 1902; *P. discolor* Lindqvist, 1975). Host plant: *Sanguisorba officinalis* (Rosaceae). Russia: **EP** (NW, C, E), **ES** (IR). – Europe (WE, EE), Kazakhstan.

Pristiphora nigriceps (Hartig, 1840) [Nematus] (*Nematus bistriatus* Thomson, 1871). Host plant: *Picea abies* (Pinaceae). Russia: **EP** (Zhelochovtsev, 1995), **ES** (YA), **FE** (PR). – Europe (WE, EE, NE).

Pristiphora nigromongolica Haris, 2002. Russia: **ES** (IR). – Mongolia.

Pristiphora pallida (Konow, 1904) [*Lygaeonematus*] (*Lygaeonematus stecki* Nägeli, 1936). Host plant: *Picea abies* (Pinaceae). Russia: "Siberia" (Zhelochovtsev, Zinovjev, 1995). – Europe (WE, EE, NE).

Pristiphora pallidiventris (Fallén, 1808) [Tenthredo] (*Nematus ephippiger* Hartig, 1840; *N. flavigomus* Tischbein, 1846; *N. gemellus* Förster, 1854; *N. marshalli* Cameron, 1875; *N. cirrhostomus* Zaddach, 1883; *Pristiphora pallidiventris denudata* Konow, 1902; *P. zella* Rohwer, 1909; *P. pallicoxa* Rohwer, 1910; *P. xanthotrichela* Rohwer, 1913; *P. pallidiventris haemorrhoidalis* Enslin, 1916; *P. pallidiventris stigmatica* Enslin, 1916; *P. ostiaria* MacGillivray, 1920; *P. pallidiventris megalpina* Lacourt, 1987). Host plant: *Filipendula* spp., *Geum* spp., *Rubus* spp. (Rosaceae). Russia: **EP** (NW, C, E, S, NC), **UR**, **ES** (YA), **FE** (AM, PR, SA, KU). – Europe (WE, EE, NE, SE), Georgia, Armenia, Turkey, Kazakhstan, Mongolia, China (NW), Korean Peninsula, Japan (Hon), N America.

Pristiphora paralella (Hartig, 1840) [Nematus] (*Lygaeonematus tenuicornis* Lindqvist, 1955). Host plant: *Picea* sp. (Pinaceae). Russia: **EP** (NW), **ES** (IR, YA), **FE** (PR). – Europe (WE, EE, NE), Korean Peninsula.

Pristiphora parva (Hartig, 1837) [Nematus] (*Nematus nigricornis* Zaddach, 1883; *N. germanicus* Dalla Torre, 1894; *Lygaeonematus ambiguus* flavater Enslin, 1916). Host plant: *Picea abies* (Pinaceae). Russia: **EP** (NW). – Europe (WE, EE, NE).

Pristiphora pseudodecipliens Beneš et Kříštek, 1976. Host plant: *Picea* spp. (Pinaceae). Russia: **EP** (Zhelochovtsev, Zinovjev, 1995). – Europe (WE, EE, NE).

Pristiphora pseudogeniculata Lindqvist, 1969. Host plant: *Prunus padus* (Rosaceae). Russia: **FE** (KA). – Europe (NE).

Pristiphora pseudomelanocarpa Haris, 2006. Russia: **FE** (KU).

Pristiphora punctifrons (Thomson, 1871) [Nematus] (*Nematus platyceros* Zaddach, 1884; *Pristiphora viridana* Konow, 1902; *Micronematus camtschaticus* Enslin, 1927; *Pristiphora aspericeps* Lindqvist, 1960; *P. flavipicta* Lindqvist, 1975; *P. mongoloxigua* Haris, 2002; *P. flavopleura* Haris, 2002; *P. mongolofausta* Haris, 2003). Host plant: *Rosa* spp. (Rosaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (IR, BR, YA), **FE** (SA, KA, MG). – Europe (WE, EE, NE, SE), Georgia, Central Asia, Kazakhstan, Mongolia, Japan (Hon), N America.

Pristiphora retusa (Thomson, 1871) [Nematus]. Host plant: *Prunus padus* (Rosaceae). Russia: **EP** (NW, C), **ES** (IR, YA), **FE** (KH, PR). – Europe (WE, EE, NE).

- Pristiphora reuteri** (Lindqvist, 1960) [Lygaeonematus]. Host plant: *Salix phyllicifolia* (Salicaceae). Russia: **EP** (N), **UR**, **WS** (TM), **ES** (IR, BR), **FE** (KA, MG). – Europe (NE), N America.
- Pristiphora robusta** (Konow, 1895) [Lygaeonematus]. Host plant: *Picea* spp. (Pinaceae). Russia: **EP** (NW), **ES** (YA). – Europe (EE, NE).
- Pristiphora ruficornis** (Olivier, 1811) [Nematus] (*P. testaceicornis* Serville, 1823; *P. testaceicornis* Lepeletier, 1823; *Nematus fraxini* Hartig, 1837; *N. testaceicornis* Jacobs, 1884; *N. ruficornis integer* Hellén, 1948). Host plant: *Betula pubescens* (Betulaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (Zhelochovtsev, Zinovjev, 1995), **ES** (TU, IR, BR, ZB). – Europe (WE, EE, NE, SE), Armenia, Mongolia.
- Pristiphora rufipes** Serville, 1823 (*P. fusca* Serville, 1823; *Nematus suessionensis* Serville, 1823; *Pristiphora rufipes* Lepeletier, 1823; *P. fusca* Lepeletier, 1823; *Nematus suessionensis* Lepeletier, 1823; *N. selandrioides* Costa, 1859; *N. aquilegiae* Sellen van Vollenhoven, 1866). Host plant: *Aquilegia* spp. (Ranunculaceae). Russia: **EP** (C, S), **WS** (AL), **ES** (KS, KR, YA), **FE** (AM, PR). – Europe (WE, EE, NE, SE), Turkey, Kyrgyzstan; introduced into N America.
- Pristiphora salicivora** (Takeuchi, 1922) [Lygaeonematus]. Host plant: *Salix* spp. (Salicaceae). Russia: **FE** (SA). – Japan (Hon).
- Pristiphora sareptana** Kuznetzov-Ugamskij, 1924 (*P. moravica* Gregor, 1940). Russia: **EP** (C, E, S). – Europe (EE, NE), Kazakhstan.
- Pristiphora saxesenii** (Hartig, 1837) [Nematus] (*P. thalenhorsti* Wong, 1975). Host plant: *Picea* spp. (Pinaceae). Russia: **EP** (NW, C), **WS** (Zhelochovtsev, Zinovjev, 1995), **ES** (YA). – Europe (WE, EE, NE), Kyrgyzstan.
- Pristiphora sermola** Liston, 1993 (*Lygaeonematus variipes* Lindqvist, 1952). Host plant: *Salix* spp. (Salicaceae). Russia: **EP** (NW), **ES** (Zhelochovtsev, Zinovjev, 1995). – Europe (WE, NE).
- Pristiphora similis** Kuznetzov-Ugamskij, 1924. Russia: **EP** (S).
- Pristiphora simplicifrons** Malaise, 1931. Russia: **FE** (PR).
- Pristiphora staudingeri** (Ruthe, 1859) [Nematus] (*P. circularis* Kincaid, 1900; *P. hyperborea* Malaise, 1921; *P. asperlatus* Benson, 1935). Host plant: *Salix herbacea*, *S. phyllicifolia* (Salicaceae). Russia: **EP** (N, NW, C), **ES** (KS, KR, IR, YA, ZB), **FE** (KH, KA, MG). – Europe (WE, EE, NE), Mongolia, China (NC), N America.
- Pristiphora subarctica** (Forsslund, 1936) [Lygaeonematus] (*P. pseudosaxesenii* Lindqvist, 1968). Host plant: *Picea abies* (Pinaceae). Russia: **EP** (C), **ES** (KR). – Europe (WE, EE, NE).
- Pristiphora subbifida** (Thomson, 1871) [Nematus] (*Nematus depressus* Hartig, 1840). Host plant: *Acer campestre* (Sapindaceae). Russia: **EP** (Taeger et al., 2006). – Europe (WE, EE, NE), Cyprus, N America.
- Pristiphora subopaca** Lindqvist, 1955 (*P. coniceps* Lindqvist, 1955; *P. brunniapex* Lindqvist, 1960). Host plant: *Salix caprea*, *S. phyllicifolia* (Salicaceae). Russia: **EP** (NW), **ES** (IR). – Europe (WE, NE).
- Pristiphora tenuiserra** (Lindqvist, 1959) [Microneatus]. Russia: **EP** (Taeger et al., 2006). – Europe (WE, EE, NE).
- Pristiphora testacea** (Jurine, 1807) [Pteronus] (*Tenthredo betulae* Retzius, 1783; *T. betulae* Christ, 1791; *Nematus betulinus* Dahlbom, 1835; *N. betularius* Hartig, 1837; *N. betulae* Hartig, 1837; *N. melanurus* Hartig, 1840; *N. erythrogaster* Thomson, 1871; *N. crassiventris* Cameron, 1878; *N. hartigii* W.F. Kirby, 1882; *N. luteogaster* W.F. Kirby, 1882; *Pristiphora gaunitzi* Lindqvist, 1968). Host plant: *Betula* spp. (Betulaceae). Russia: **EP** (C). – Europe (WE, EE, NE), China (Wei et al., 2006).
- Pristiphora tetrica** (Zaddach, 1883) [Nematus] (*Nematus velatus* Zaddach, 1883; *Pristiphora nievesi* Haris, 2004). Host plant: *Acer* spp. (Sapindaceae). Russia: **EP** (NC). – Europe (WE, SE), N Africa.
- Pristiphora thalictri** (Kriechbaumer, 1884) [Nematus] (*P. henschi* Konow, 1902; *Diphadnus thalictri* Takeuchi, 1922). Host plant: *Thalictrum aquilegifolium*, *T. minus* (Ranunculaceae). Russia: **EP** (NW, C), **ES** (Zhelochovtsev, Zinovjev, 1995), **FE** (KU). – Europe (WE, EE, NE, SE), Kazakhstan, Mongolia, Korean Peninsula, Japan (Hon).
- Pristiphora thalictrovora** Lindqvist, 1962 (*P. thalictrocola* Lindqvist, 1974; *P. rufiventris* Lindqvist, 1974). Host plant: *Thalictrum flavum*, *T. simplex* (Ranunculaceae). Russia: **EP** (Taeger et al., 2006), **ES** (IR). – Europe (NE).
- Pristiphora trochanterica** (Lindqvist, 1952) [Lygaeonematus]. Russia: **EP** (N), **WS** (TM), **ES** (KR). – Europe (NE).
- Pristiphora truncatiserra** Lindqvist, 1975. Russia: **ES** (IR, BR).

Pristiphora werzhutskii Lindqvist, 1970. Host plant: Spiraea spp. (Rosaceae). Russia: **ES** (IR).

Pristiphora wesmaeli (Tischbein, 1853) [Nematus] (*Nematus solea* Snellen van Vollenhoven, 1870). Host plant: Larix spp. (Pinaceae). Russia: **EP** (NW), **WS** (OM), **ES** (KR, IR, YA). – Europe (WE, EE, NE), China (NE), Korean Peninsula, Japan (Hon).

PSEUDODINEURA Konow, 1885 (*Pelmatopus* Hartig, 1837; *Phyllopais* Hering, 1934). Type species *Tenthredo parvula* Klug, 1816. The larvae live in mines on leaves of Ranunculaceae. The genus is distributed in Holarctic. Number of species: World – 12; Palaearctic – 9; Russia – 6.

Pseudodineura clematidis (Hering, 1932) [*Pelmatopus*]. Host plant: Clematis spp., ?Atragene spp. (Ranunculaceae). Russia: **UR** (Taeger et al., 2006). – Europe (WE, EE).

Pseudodineura enslini (Hering, 1923) [*Pelmatopus*]. Host plant: Trollius spp. (Ranunculaceae). Russia: **EP** (C), **ES** (KR, IR). – Europe (WE, EE, NE).

Pseudodineura fuscula (Klug, 1816) [*Tenthredo*] (*Dineura despecta* Hartig, 1837; *Dolerus minutus* Hartig, 1837; *Dineura simulans* Cameron, 1877). Host plant: Ranunculus spp. (Ranunculaceae). Russia: **EP** (N, NW, C, E, S, NC), **ES** (IR). – Europe (WE, EE, NE, SE), Armenia, Kazakhstan; introduced into N America.

Pseudodineura heringi (Enslin, 1921) [*Permatopus*]. Host plant: Anemone spp. (Ranunculaceae). Russia: **EP** (C), **ES** (IR). – Europe (WE, EE, NE).

Pseudodineura mentiens (Thomson, 1871) [Blennocampa] (*Dineura hepaticae* Brischke, 1883; *Pelmatopus mentiens konowi* Enslin, 1921). Host plant: Hepatica nobilis (Ranunculaceae). Russia: **EP** (NW, C, E, NC), **UR**, **FE** (KH, PR). – Europe (WE, EE, NE).

Pseudodineura parvula (Klug, 1816) [*Tenthredo*] (*Tenthredo minuta* Lepetier et Serville, 1828). Host plant: Pulsatilla spp. (Ranunculaceae). Russia: **EP** (C), **ES** (ZB). – Europe (WE, EE).

STAURONEMATUS Benson, 1953 (*Stauronema* Benson, 1948). Type species *Nematus platycerus* Hartig, 1840. The larvae live in mines on leaves. The genus is distributed in Palaearctic. Number of species: one species.

Stauronematus platycerus (Hartig, 1840) [Nematus] (*Nematus vallator* Snellen van Vollenhoven, 1858;

N. cebrionicornis Costa, 1859; *N. callicerus* Thomson, 1863). Host plant: Populus spp., Salix alopechroa (Salicaceae). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **ES** (IR, YA), **FE** (KH). – Europe (WE, EE, NE, SE), Caucasus, Turkey, Iran, Kyrgyzstan, Kazakhstan, China (CC), Korean Peninsula, Japan (Hok, Hon).

Unplaced Nematinae

Amauronematus nigridorsis Jakovlev, 1891. Russia: **ES** (IR).

Cryptocampus fuscus Zaddach, 1883. Russia: **EP** (NW).

Nematus cinereus Eversmann, 1847. Russia: **EP** (E).

Nematus konowi (Jakovlev, 1891) [*Pteronus*]. Russia: **EP** (NC).

Nematus contractus Eversmann, 1847. Russia: **EP** (S).

Nematus fruticum Eversmann, 1847. Russia: **EP** (E).

Nematus schewyrewi (Jakovlev, 1891) [*Pteronus*]. Russia: **EP** (NC).

Nematus (Pteronidea) pyramidalis Hellén, 1948. Russia: **ES** (KR).

Pteronus irkutensis Jakovlev, 1891. Russia: **ES** (IR).

Tenthredo (Selandria) umbrosa Eversmann, 1847. Russia: **EP** (E).

Subfamily SELANDRIINAE

ALPHASTROMBOCEROS Kuznetsov-Ugamskij, 1928 (*Parastromboceros* Takeuchi, 1941). Type species *Strongylogaster konowi* Jakovlev, 1891. The larvae on herbaceous plants. The genus is distributed in Palaearctic and Oriental region. Number of species: World – 5; Palaearctic – 3; Russia – 1.

Alphastromboceros konowi (Jakovlev, 1891) [Strongylogaster] (*Stromboceros filicis* Malaise, 1931). Host plant: Urtica spp. (Urticaceae). Russia: **EP** (C), **FE** (KH, PR, SA). – Europe (WE, EE), China (NE, SE), Korean Peninsula, Japan (Hok, Hon, Kyu).

ANEUGMENUS Hartig, 1837 (*Colposelandria* Enslin, 1912; *Atoposelandria* Enslin, 1913; *Polyselandria* MacGillivray, 1914; *Selandropha* Zirngiebl, 1956). Type species *Tenthredo coronata* Klug, 1818. The larvae on ferns. The genus is distributed in Holarctic, Oriental and Neotropical regions. Number of species: World – 32; Palaearctic – 15; Russia – 8.

Aneugmenus carinifrons (Malaise, 1931) [Selandria] (*Selandria planiceps* Malaise, 1931). Host plant: ?*Lysimachia* spp. (Primulaceae). Russia: **FE** (AM, PR). – China (SE), Korean Peninsula, Japan (Hon, Kyu, Sado, Tsushima).

Aneugmenus coronatus (Klug, 1818) [Tenthredo] (*Selandria analis* Thomson, 1871; *S. cereipes* Snellen van Vollenhoven, 1873; *S. bimaculata* Cobelli, 1892; *S. ogloblini* Dovnar-Zapolskij, 1930). Host plant: *Athyrium* spp., *Dryopteris* spp. (Dryopteridaceae), *Pteridium aquilinum* (Dennstaedtiaceae). Russia: **EP** (NW, C, NC), **WS** (AL), **FE** (AM, KH, PR). – Europe (WE, EE, NE, SE), Georgia, China (NE), Korean Peninsula, Japan (Niu, Wei, 2013).

Aneugmenus fuerstenbergensis (Konow, 1885) [Selandria]. Host plant: *Pteridium aquilinum* (Dennstaedtiaceae). Russia: **EP** (N, C, E). – Europe (WE, SE, EE, NE).

Aneugmenus japonicus Rohwer, 1910 (*A. gratus* Zhelochovtsev, 1951). Host plant: *Pteridium aquelinum* (Dennstaedtiaceae). Russia: **FE** (SA, KU). – China (NC, CC, SE), Korean Peninsula, Japan (Hon, Shi, Kyu).

Aneugmenus kiotonis Takeuchi, 1929. Host plant: *Pteridium aquelinum* (Dennstaedtiaceae). Russia: **FE** (SA, KU). – Japan (Hon, Shi, Kyu, Sado).

Aneugmenus oertzeni (Konow, 1887) [Selandria]. Russia: **EP** (C). – Europe (SE), SW Asia.

Aneugmenus padi (Linnaeus, 1760) [Tenthredo] (*Tenthredo stramineipes* Klug, 1816; *T. albipes* Serville, 1823; *T. albipes* Lepeletier, 1823; *Selandria rufitarsis* Brullé, 1832; *Allantus laticinctus* Brullé, 1832; *Tenthredo cerasorum* Dahlbom, 1835; *Selandria vollenhoveni* Gribodo, 1881; *S. urbis* Ross, 1930). Host plant: *Pteridium aquilinum* (Dennstaedtiaceae). Russia: **EP** (NW, S, NC). – Europe (WE, EE, NE, SE), N Africa, Georgia, Turkey, Iran; introduced into the N America.

Aneugmenus temporalis (Thomson, 1871) [Selandria]. Host plant: *Pteridium aquilinum* (Dennstaedtiaceae). Russia: **EP** (NW, C, E), **UR**, **FE** (AM, KH, PR, SA). – Europe (WE, EE, NE, SE), China (NC, NE, CC), Korean Peninsula, Japan (Niu, Wei, 2013).

BIRKA Malaise, 1944 (*Lineobirka* Wei et Nie, 1997). Type species *Tenthredo cinereipes* Klug, 1816. The genus is distributed in Palaearctic and Oriental region. Number of species: World – 17; Palaearctic – 11; Russia – 4 from 2 subgenera.

Birka (Birka) annulitarsis (Thomson, 1870) [Selandria]. Host plant: *Rubus* spp. (Rosaceae). Russia:

EP (E), **UR**, **WS** (KM, AL), **ES** (YA). – Europe (WE, EE, NE).

Birka (Birka) catellata (Konow, 1900) [Selandria]. Russia: **WS** (KM), **ES** (KR, IR), **FE** (AM).

Birka (Birka) cinereipes (Klug, 1816) [Tenthredo] (*Tenthredo aperta* Hartig, 1837; *Selandria foief-rorns* Thomson, 1870). Host plant: *Myosotis* spp. (Boraginaceae). Russia: **EP** (NW, C, E), **UR**, **WS** (TM, NS, AL), **FE** (KA). – Europe (WE, EE, NE, SE), Caucasus, Turkey.

Birka (Lineobirka) koreana (Takeuchi, 1941) [Nesoselandria] (*B. pallipes* Zhelochovtsev, 1951). Host plant: *Pyrus* spp. (Rosaceae). Russia: **FE** (AM, PR). – China (NE), Korean Peninsula.

BRACHYTHOPS Curtis, 1839 (*Coryna* Lepeletier et Serville, 1828; *Paraselandria* Ashmead, 1898; *Selandridea* Rohwer, 1911; *Pseudoselandria* MacGillivray, 1914). Type species *Brachythops seminigra* Curtis, 1839. The larvae free-living, on *Carex* (Cyperaceae). The genus is distributed in Holarctic. Number of species: World and Palaearctic – 4; Russia – 3.

Brachythops flavens (Klug, 1816) [Tenthredo] (*Tenthredo scapularis* Lepeletier et Serville, 1828; *Phyllostoma puella* Fallén, 1829; *Brachythops seminigra* Curtis, 1839; *Selandria flavescens* Thomson, 1870; *S. decolorata* Cresson, 1880; *Selandridea vanduzeei* Rohwer, 1911; *Selandria flavens flavor* Enslin, 1913; *Pseudoselandria oxalata* MacGillivray, 1914; *Selandria flavens antennalis* Zirngiebl, 1954). Host plant: *Carex* spp. (Cyperaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **ES** (IR, BR), **FE** (AM, KH). – Europe (WE, EE, NE), Armenia, Kazakhstan, China (Wei et al., 2006), N America.

Brachythops nitens (Zhelochovtsev, 1951) [Selandria]. Host plant: *Carex* spp. (Cyperaceae). Russia: **FE** (KH, PR).

Brachythops wuestneii (Konow, 1885) [Selandria] (*Selandria flavistigma* Grönblom, 1939). Host plant: *Carex* spp. (Cyperaceae). Russia: **EP** (N, NW), **WS** (AL), **ES** (IR, BR), **FE** (KH). – Europe (WE, EE, NE), N America.

BUSARBIDEA Rohwer, 1915 (*Canoniades* Forsius, 1929). Type species *Busarbidea himalaiensis* Rohwer, 1915. The genus is distributed in East Palaearctic and Oriental region. Number of species: World – 19; Palaearctic – 8; Russia – 1.

Busarbidea ussuriensis (Malaise, 1931) [Canoniades]. Russia: **FE** (PR).

DOLERUS Panzer, 1801 (*Dolerus* Jurine, 1801; *Dosytheus* Leach, 1817; *Dositheus* Agassiz, 1848; *Loderus* Konow, 1890; *Hamatodentiden* Zirngiebl, 1930; *Multidentiden* Zirngiebl, 1930; *Nodulodentiden* Zirngiebl, 1930; *Pratensisformis* Ermolenko, 1975; *Gonagerformis* Ermolenko, 1975; *Nigerformis* Ermolenko, 1975; *Achaetopriion* Goulet, 1986; *Dicrodolerus* Goulet, 1986; *Neodolerus* Goulet, 1986; *Oncodolerus* Goulet, 1986; *Cyperolerus* Zhelochovtsev, 1988; *Poodolerus* Zhelochovtsev, 1988; *Juncilerus* Zhelochovtsev, 1988; *Equidolerus* Taeger et Blank, 1996). Type species *Tenthredo pedestris* Panzer, 1801. The larvae feed mainly on Poaceae and Equisetaceae, less often on Juncaceae and Cyperaceae. The genus is distributed in Holarctic and Oriental region. Number of species: World – about 225; Palaearctic – about 155; Russia – 75 (+ 8 ssp.) from 9 subgenera.

Dolerus (Achaetopriion) ferrugatus Serville, 1823 (*D. ferrugatus* Lepeletier, 1823; *D. brevicornis* Thomson, 1871; *D. thomsoni* Konow, 1884; *D. thomsoni miricolor* Konow, 1887). Host plant: *Juncus* spp. (Juncaceae). Russia: **EP** (C, E). – Europe (WE, EE, NE, SE), China (Wei et al., 2006).

Dolerus (Achaetopriion) madidus (Klug, 1818) [Tenthredo] (*Tenthredo lateritia* Klug, 1818; *Dolerus chappelli* Cameron, 1877; *D. lamprechti* Konow, 1884; *D. madidus monozonus* Enslin, 1913). Host plant: *Juncus* spp. (Juncaceae). Russia: **EP** (NW, C, E, CR), **WS** (NS). – Europe (WE, EE, NE, SE), Korean Peninsula.

Dolerus (Achaetopriion) pachycerus Hartig, 1837 (*D. taeniatus* Zaddach, 1859; *D. tinctipennis* Cameron, 1881; *D. mocsaryi* Konow, 1895). Host plant: *Juncus* spp. (Juncaceae). Russia: **EP** (N, NW, C, E), **WS** (Zhelochovtsev, 1988), **ES** (BR, YA). – Europe (WE, EE, NE, SE), Kazakhstan.

Dolerus (Achaetopriion) triplicatus (Klug, 1818) [Tenthredo] (*Tenthredo tremula* Klug, 1818; *Dolerus trimaculatus* Lepeletier, 1823; *D. dimidiatus* Lepeletier, 1823; *D. trimaculatus* Serville, 1823; *D. dimidiatus* Serville, 1823; *D. lugubris* Gimmenthal, 1834; *D. steini* Konow, 1885; *D. tremulus aemula* Konow, 1902). Host plant: *Juncus* spp. (Juncaceae). Russia: **EP** (C, S, NC, CR). – Europe (WE, EE, NE, SE), Georgia, Armenia, Turkey.

Dolerus (Cyperolerus) anticus anticus (Klug, 1818) [Tenthredo]. Host plant: *Carex* spp., *Eleocharis palustris* (Cyperaceae). Russia: **EP** (C, E), **UR**, **WS**

(NS), **ES** (Zhelochovtsev, Zinovjev, 1996). – Europe (WE, EE, NE, SE), Caucasus, Turkey, Kazakhstan.

Dolerus (Dicrodolerus) apricus (Norton, 1861) [*Dosytheus*] (*D. luctatus* MacGillivray, 1908; *D. femurru-brum* Rohwer, 1909; *Loderus acerbus* MacGillivray, 1923; *L. alticinctus* MacGillivray, 1923; *L. ancisus* MacGillivray, 1923). Russia: **FE** (MG). – China (Wei et al., 2006), N America.

Dolerus (Dicrodolerus) genucinctus Zaddach, 1859 (*D. annulipes* Thomson, 1871; *Loderus genucinctus tegularis*). Host plant: *Equisetum* spp. (Equisetaceae). Russia: **EP** (NW, C, E), **UR**, **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (IR), **FE** (AM, KH, SA, KU, KA, MG). – Europe (WE, EE, NE), Mongolia, China (NE), Japan (Hok, Hon).

Dolerus (Dicrodolerus) mongolicus (Muche, 1965) [*Loderus*. Host plant: *Equisetum* spp. (Equisetaceae). Russia: **ES** (ZB). – Mongolia, China (NC).

Dolerus (Dicrodolerus) vestigialis vestigialis (Klug, 1818) [Tenthredo]. Host plant: *Equisetum* spp. (Equisetaceae). Russia: **EP** (NW, C, E, NC), **WS** (KM), **ES** (IR, BR). – Europe (WE, EE, NE, SE), Armenia, Turkey, Iran.

Dolerus (Dolerus) aericeps Thomson, 1871 (*D. bajulus* Serville, 1823; *D. bajulus* Lepeletier, 1823; *D. mandibularis* Konow, 1884; *D. aericeps rufipes* Konow, 1888; *D. aericeps erythropus* Enslin, 1913; *D. aericeps theresae* Pic, 1948; *D. aericeps guttatus* Zirngiebl, 1954). Host plant: *Equisetum* spp. (Equisetaceae). Russia: **EP** (N, NW, C, E), **UR**, **WS** (KM, AL), **ES** (TU, IR, BR, YA), **FE** (PR). – Europe (WE, EE, NE, SE), Caucasus, Turkey, Kazakhstan.

Dolerus (Dolerus) armillatus Konow, 1896 (*D. austerus* Muche, 1965). Host plant: Poaceae. Russia: **WS** (AL), **ES** (IR, BR), **FE** (AM, KH, PR). – Mongolia, China (SW, SE), Korean Peninsula, Japan (Hok).

Dolerus (Dolerus) bimaculatus (Geoffroy, 1785) [Tenthredo] (*Tenthredo tristis* Fabricius, 1804; *Dolerus saxatilis* Hartig, 1837). Host plant: *Equisetum* spp. (Equisetaceae). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (AL), **ES** (TU, IR, BR, YA), **FE** (AM, KH, KA, MG). – Europe (WE, EE, NE, SE), Mongolia.

Dolerus (Dolerus) cothurnatus Serville, 1823 (*Tenthredo palustris* Klug, 1818; *Dolerus cothurnatus* Lepeletier, 1823; *Dosytheus junci* Stephens, 1835; *Dolerus busaei* Snellen van Vollenhoven, 1858). Host plant: *Equisetum* spp. (Equisetaceae). Russia: **EP** (N, NW, C, E, S, NC), **ES** (IR), **FE** (MG). – Europe (WE, EE, NE, SE), Mongolia, Korean Peninsula.

Dolerus (Dolerus) elderi elderi Kincaid, 1900 (*D. pusillus* Jakovlev, 1891; *D. cohaesus* MacGillivray, 1908; *D. nutricius* MacGillivray, 1923; *Loderus acriculus* MacGillivray, 1923; *Dolerus nyctelius* MacGillivray, 1923; *D. negotiosus* MacGillivray, 1923; *D. nemorosus* MacGillivray, 1923; *D. pullulus* Zhelochovtsev, 1928; *D. elderi auraneus* Ross, 1931; *D. elderi rubicanus* Ross, 1931; *D. elderi melanus* Ross, 1931; *D. pullulus coloratus* Hellén, 1956; *D. orchonensis* Muche, 1965). Host plant: Equisetum spp. (Equisetaceae). Russia: **EP** (N), **WS** (TM, AL), **ES** (KS, KR, IR, BR, YA), **FE** (KA). – Europe (NE), Tadzhikistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NW, NC), N America.

Dolerus (Dolerus) germanicus aterrimus Zhe-lochovtsev, 1935. Host plant: Equisetum spp. (Equisetaceae). Russia: **FE** (KU). – Japan (Hok).

Dolerus (Dolerus) germanicus claripennis Zhe-lochovtsev, 1935. Host plant: Equisetum spp. (Equisetaceae). Russia: **FE** (KA, MG).

Dolerus (Dolerus) germanicus fuscipennis (Stephens, 1835) [Dosytheus] (*D. arcticus* Thomson, 1871). Host plant: Equisetum spp. (Equisetaceae). Russia: **FE** (KA). – Europe (NE).

Dolerus (Dolerus) germanicus germanicus (Fabricius, 1775) [Tenthredo] (*Tenthredo hortorum* O.F. Müller, 1776; *T. melanocephala* Fourcroy, 1785; *T. eglanteriae* Fabricius, 1793; *T. pedestris* Panzer, 1801; *T. subfusca* Lepeletier, 1823; *Dosytheus xanthopus* Stephens, 1835; *Tenthredo quadricincta* Zetterstedt, 1838; *Dolerus fennicus* André, 1880; *D. rufipes* Grädl, 1881; *D. pratensis nigripes* Konow, 1884; *D. pratensis mediater* Enslin, 1913; *D. pratensis laterater* Enslin, 1913; *D. pratensis terminater* Enslin, 1913; *D. pratensis totus* Malaise, 1931; *D. germanicus berlandi* Pic, 1948; *D. germanicus buyssoni* Pic, 1948; *D. pratensis major* Zirngiebl, 1954). Host plant: Equisetum spp. (Equisetaceae). Russia: **EP** (NW, C), **WS** (NS, KM, AL), **ES** (IR), **FE** (PR, KA). – Europe (WE, EE, NE, SE).

Dolerus (Dolerus) germanicus meridianus Zhe-lochovtsev, 1928. Host plant: Equisetum spp. (Equisetaceae). Russia: **EP** (NC). – Georgia, Armenia.

Dolerus (Dolerus) germanicus orientalis Zhelochovtsev, 1935. Host plant: Equisetum spp. (Equisetaceae). Russia: **FE** (AM, KH, PR, MG). – China (NE), Korean Peninsula.

Dolerus (Dolerus) germanicus sibiricus Zhelochovtsev, 1935. Host plant: Equisetum spp. (Equisetaceae).

Russia: **EP** (N), **WS** (TM), **ES** (KS, KR, IR, BR, ZB), **FE** (MG). – Mongolia.

Dolerus (Dolerus) germanicus subsolanus Zhe-lochovtsev, 1928. Host plant: Equisetum spp. (Equisetaceae). Russia: **UR**, **WS** (OM, AL). – Kazakhstan.

Dolerus (Dolerus) klutchianus Malaise, 1931. Russia: **FE** (KA).

Dolerus (Dolerus) manticatus Konow, 1907. Host plant: Equisetum spp. (Equisetaceae). Russia: **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (KR, IR), **FE** (AM). – Mongolia, China (NC, SW).

Dolerus (Dolerus) nigriceps Konow, 1891. Russia: **EP** (Lacourt, 1999). – Caucasus, Turkey, Iran.

Dolerus (Dolerus) subarcticus Hellén, 1956 (*D. willoughbyi* Benson, 1956). Host plant: Equisetum spp. (Equisetaceae). Russia: **EP** (N, C), **WS** (NS), **ES** (KR, YA). – Europe (NE), Mongolia.

Dolerus (Dolerus) variegatus Jakovlev, 1891 (*D. patricius* Muche, 1965). Host plant: Equisetum spp. (Equisetaceae). Russia: **WS** (NS, AL), **ES** (IR, BR, ZB), **FE** (AM, MG). – Europe (WE), Mongolia, China (NC).

Dolerus (Dolerus) yukonensis Norton, 1872 (*D. scoticus* Cameron, 1881; *D. lateralis* Konow, 1895; *D. articola* Kier, 1898; *D. nuntius* MacGillivray, 1923; *D. nundius* MacGillivray, 1923; *D. scoticus lenensis* Hellén, 1956; *D. thargitai* Zombori, 1994). Host plant: Equisetum spp. (Equisetaceae). Russia: **EP** (N, NW, C, E), **UR**, **WS** (KM), **ES** (IR, BR, YA), **FE** (AM, SA, KU, KA, MG). – Europe (WE, EE, NE), Mongolia, China (NC), N America.

Dolerus (Equidolerus) gessneri André, 1880 (*D. labiosus* Konow, 1897; *D. kamtchaticus* Malaise, 1931; *D. gessneri rufomaculata* Hellén, 1956). Host plant: Equisetum spp. (Equisetaceae). Russia: **EP** (NW, C), **WS** (TM), **ES** (KS, KR, IR, BR, ZB), **FE** (AM, SA, KA, MG). – Europe (WE, EE, NE), China (Wei et al., 2006), Korean Peninsula, Japan (Hok, Hon), N America.

Dolerus (Equidolerus) konowi glacialis Goulet, 1986. Host plant: Equisetum spp. (Equisetaceae). Russia: **FE** (KU, KA, CH). – N America (Alaska).

Dolerus (Equidolerus) pratensis (Linnaeus, 1758) [Tenthredo] (*Tenthredo dubia* Klug, 1818; *T. timida* Klug, 1818; *T. deserta* Klug, 1818; *Dosytheus hyalinialis* Stephens, 1835; *Dolerus dubius niger* Enslin, 1909; *D. dubius atratus* Enslin, 1913; *D. pratensis camtschatcalis* Enslin, 1918; *D. variator* Enslin, 1927; *D. variator mesonotalis* Enslin, 1927; *D. variator mesopleuralis* Enslin, 1927). Host plant: Equisetum

- spp. (Equisetaceae). Russia: **EP** (N, NW, C, E), **UR**, **WS** (NS), **ES** (IR, BR, YA), **FE** (AM, KA). – Europe (WE, EE, NE), Turkey, Mongolia, China (NC), N America; introduced into Tropical Africa.
- Dolerus (Equidolerus) pseudoanticus** Malaise, 1931. Russia: **FE** (KA).
- Dolerus (Equidolerus) rhodogaster** Zhelochovtsev, 1935. Host plant: Equisetum spp. (Equisetaceae). Russia: **ES** (IR, BR), **FE** (AM, PR). – China (NE).
- Dolerus (Equidolerus) subfasciatus** F. Smith, 1874 (*D. picinus* Marlatt, 1898). Host plant: Equisetum spp. (Equisetaceae). Russia: **FE** (KU). – Japan (Hon, Shi).
- Dolerus (Loderus) gilvipes albifrons** (Norton, 1861) [*Dosytheus*] (*D. minusculus* MacGillivray, 1908; *Loderus accuratus* MacGillivray, 1923). Russia: **FE** (CH). – China (NE), N America.
- Dolerus (Loderus) gilvipes gilvipes** (Klug, 1818) [Tenthredo] (*D. tenebrosus* Eversmann, 1847; *Loderus ornatulus* Konow, 1897; *L. ornatulus plagiatus* Konow, 1897; *L. gilvipes camtschatcalis* Enslin, 1927). Russia: **EP** (N, E), **UR**, **ES** (IR, YA), **FE** (AM, PR, KA, MG). – Europe (WE, NE), Mongolia, China (NE, NC), Korean Peninsula, Japan (Goulet, 1986).
- Dolerus (Loderus) pratorum** (Fallén, 1808) [Tenthredo] (*Tenthredo equiseti* Klug, 1818; *Loderus pratorum maculifrons* Hellén, 1956). Host plant: Equisetum spp. (Equisetaceae). Russia: **EP** (N, NW, C, E, NC), **WS** (NS, KM), **ES** (IR, BR, YA), **FE** (AM). – Europe (WE, EE, NE), Kazakhstan, China (NE).
- Dolerus (Neodolerus) anatolii** Heidemaa et Zinovjev, 2004. Russia: **FE** (PR). – Korean Peninsula.
- Dolerus (Oncodolerus) eversmanni eversmanni** W.F. Kirby, 1882 (*Tenthredo palmata* Klug, 1818; *Dolerus fumosus* Eversmann, 1847; *D. trigeminus* Konow, 1884). Host plant: Equisetum spp. (Equisetaceae). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (KM), **ES** (IR, BR, YA), **FE** (AM, KH, MG). – Europe (WE, EE, NE), Georgia, Armenia, Turkey, Mongolia, China (NE).
- Dolerus (Poodolerus) aeneus** Hartig, 1837 (*D. longicornis* Zaddach, 1859; *D. incertus* Zaddach, 1859; *D. elongatus* Thomson, 1871; *D. aeneus zaddachi* Konow, 1886). Host plant: Poa spp., Hordeum vulgare, Triticum spp. (Poaceae), Carex spp. (Cyperaceae). Russia: **EP** (N, NW, C, E, S, NC), **WS** (AL), **ES** (IR), **FE** (KU, KA, MG). – Europe (WE, EE, NE, SE), Armenia, Turkey, Mongolia.
- Dolerus (Poodolerus) albertii** Muche, 1969. Russia: **EP** (NC).
- Dolerus (Poodolerus) anthracinus** (Klug, 1818) [Tenthredo] (*Tenthredo mutilata* Klug, 1818; *Dolerus atricapillus* Hartig, 1837; *D. violaceus* Pigeot, 1918; *D. violaceus obscura* Pigeot, 1918). Host plant: Hordeum vulgare, Poa spp. (Poaceae). Russia: **EP** (N, NW, C, E, S). – Europe (WE, EE, NE, SE).
- Dolerus (Poodolerus) asper** Zaddach, 1859 (*D. oblongus* Cameron, 1882; *D. asper megapteroides* Muche, 1964). Host plant: Deschampsia spp. (Poaceae), Carex spp. (Cyperaceae). Russia: **EP** (N, NW, C, E, S, NC), **ES** (IR, BR, YA), **FE** (AM, MG). – Europe (EE, NE, SE), Armenia, Turkey, Mongolia, China (NC).
- Dolerus (Poodolerus) blanki** Liston, 1995 (*Tenthredo opaca thoracica* Fallén, 1808; *T. thoracica* Klug, 1818). Russia: **EP** (C). – Europe (WE, EE, NE, SE), China (Wei et al., 2006).
- Dolerus (Poodolerus) brevicornis** Zaddach, 1859 (*D. tectus* MacGillivray, 1914). Russia: **EP** (N, NW). – Europe (WE, EE, NE), N America.
- Dolerus (Poodolerus) ciliatus** Konow, 1891 (*D. quasi-ciliatus* Kerenskij, 1926). Russia: **EP** (S, CR). – Europe (EE), Armenia, Turkey, Kazakhstan.
- Dolerus (Poodolerus) coracinus** (Klug, 1818) [Tenthredo]. Russia: **EP** (C). – Europe (WE, EE, NE), Korean Peninsula.
- Dolerus (Poodolerus) dathei** Muche, 1973. Russia: **EP** (NC).
- Dolerus (Poodolerus) derzavini** Malaise, 1931. Russia: **FE** (KA).
- Dolerus (Poodolerus) docilus** Benson, 1956. Russia: **EP** (C, E), **UR**, **WS** (AL). – Kyrgyzstan, Kazakhstan.
- Dolerus (Poodolerus) fumosus** Stephens, 1835 (*D. brachygaster* Hartig, 1837; *D. gracilis* Zaddach, 1859; *D. brevis* Zaddach, 1859; *D. rarus* Zaddach, 1859). Russia: **EP** (NW). – Europe (WE, EE, NE).
- Dolerus (Poodolerus) gibbosus** Hartig, 1837 (*D. planatus* Hartig, 1837). Russia: **EP** (NW, C), **WS** (Zhelochovtsev, Zinovjev, 1996), **FE** (KH, SA). – Europe (WE, EE, NE), Armenia.
- Dolerus (Poodolerus) gonager** (Fabricius, 1781) [Tenthredo] (*Tenthredo erythrogora* Schrank, 1781; *T. geniculata* Geoffroy, 1785; *Dolerus femoratus* Eversmann, 1847; *D. magnicornis* Eversmann, 1847; *D. lucidus* Zirngiebl, 1930; *D. gonager flavitibia* Hellén, 1956). Host plant: Agrostis spp., Festuca spp., Poa spp., Triticum spp., etc. (Poaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (KM, AL),

- ES** (KS). – Europe (WE, EE, NE, SE), Armenia, Azerbaijan, Turkmenistan.
- Dolerus (Poodolerus) haematodes** (Schrank, 1781) [Tenthredo] (*Tenthredo collaris* Donovan, 1808; *Dolerus coeruleascens* Hartig, 1837; *D. micans* Zaddach, 1859; *D. haematodes muliebris* Enslin, 1909; *D. haematodes rufatus* Enslin, 1913). Host plant: Poaceae, Cyperaceae, Juncaceae. Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TM), **ES** (TU, YA, ZB). – Europe (WE, EE, NE, SE), Armenia, Turkey.
- Dolerus (Poodolerus) harwoodi** Benson, 1947. Russia: **EP** (N, NW, C, E, S), **UR**, **WS** (NS), **ES** (IR, BR), **FE** (AM, KH). – Europe (WE, EE, NE).
- Dolerus (Poodolerus) kokujewi** Konow, 1902. Russia: **EP** (C, NC). – Caucasus, SW Asia, Iran.
- Dolerus (Poodolerus) liogaster liogaster** Thomson, 1871 (*D. annulatus* Stein, 1894; *D. liogaster rufo-notatus* Enslin, 1913; *D. liogaster modestus* Dubois, 1920). Host plant: Dactylis spp., Festuca spp., Poa spp. (Poaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**. – Europe (WE, EE, NE), Kazakhstan.
- Dolerus (Poodolerus) liogaster schneideri** Kiær, 1898. Host plant: Poaceae. Russia: **EP** (N), **UR**, **WS** (TM, KM, AL), **ES** (IR, BR). – Europe (NE).
- Dolerus (Poodolerus) melanopterus tauricus** Zhelochovtsev, 1928. Russia: **EP** (CR). – Europe (EE).
- Dolerus (Poodolerus) niger** (Linnaeus, 1767) [Tenthredo] (*Tenthredo angusta* Gmelin, 1790). Host plant: Avena spp., Hordeum spp., Secale spp., Triticum spp., etc. (Poaceae). Russia: **EP** (NW, C, E), **WS** (NS, KM). – Europe (WE, EE, NE, SE).
- Dolerus (Poodolerus) nigratus** (O.F. Müller, 1776) [Tenthredo] (*Tenthredo canescens* Gmelin, 1790; *Dolerus cenchris* Hartig, 1837; *D. leucobasis* Hartig, 1837; *D. fissus* Hartig, 1837; *D. hartigii* Scholtz, 1848; *D. analis* Konow, 1890). Host plant: Triticum spp., Festuca spp., Holcus spp., Poa spp., etc. (Poaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **WS** (KM), **ES** (BR). – Europe (WE, EE, NE, SE), Turkey, China (Wei et al., 2006).
- Dolerus (Poodolerus) nitens** Zaddach, 1859 (*D. coruscans* Konow, 1890; *D. wanda* Ross, 1935). Host plant: Festuca spp., Poa spp., Deschampsia spp. (Poaceae). Russia: **EP** (NW, C, E, NC), **UR**, **WS** (NS, KM), **ES** (Zhelochovtsev, Zinovjev, 1996). – Europe (WE, EE, NE, SE); introduced into N America.
- Dolerus (Poodolerus) nonultimus** Zhelochovtsev, 1941. Russia: **EP** (NC). – Georgia, Armenia, Turkey.
- Dolerus (Poodolerus) picipes** (Klug, 1818) [Tenthredo] (*D. brevitarsus* Hartig, 1837; *D. leucopterus* Zaddach, 1859; *D. ochroneurus* Förster, 1860; *D. intermedius* Cameron, 1881; *D. raddatzi* Konow, 1884; *D. leucopterus sulcatus* Konow, 1884). Host plant: Agrostis spp., Festuca sp. (Poaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (NS). – Europe (WE, EE, NE, SE), Armenia, Kazakhstan.
- Dolerus (Poodolerus) puncticollis** Thomson, 1871 (*D. croaticus* Konow, 1890). Host plant: Poa spp., Dactylis spp. (Poaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**. – Europe (WE, EE, NE, SE), N Africa, Armenia, SW Asia, Iran, Turkmenistan.
- Dolerus (Poodolerus) sanguinicollis** (Klug, 1818) [Tenthredo] (*D. lucens* André, 1880; *D. sanguinicollis reichertii* Konow, 1894; *D. sanguinicollis confusus* Enslin, 1913). Host plant: Poaceae. Russia: **EP** (N, NW, C, E), **UR**, **WS** (KM). – Europe (WE, EE, NE, SE).
- Dolerus (Poodolerus) schmidti** Konow, 1884. Russia: **EP** (N). – Europe (WE, NE).
- Dolerus (Poodolerus) stygius** Förster, 1860 (*D. megapterus* Cameron, 1881; *D. carinatus* Konow, 1884; *D. crassus* Konow, 1884; *D. megalopterus* Schulz, 1906; *D. eutropis* Speiser, 1908). Host plant: Carex spp. (Cyperaceae). Russia: **EP** (N, C, E, S), **UR**, **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (YA). – Europe (WE, EE, NE, SE), Turkey.
- Dolerus (Poodolerus) subalatus** Kerenskij, 1926. Russia: **EP** (C, S, NC). – Europe (EE).
- Dolerus (Poodolerus) vernalis** Ermolenko, 1964. Russia: **EP** (CR). – Europe (WE, EE, SE).
- Dolerus (Poodolerus) vulneratus** Mocsáry, 1878. Russia: **EP** (NW), **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (Zhelochovtsev, Zinovjev, 1996), **FE** (AM). – Europe (NE), China (NE), Korean Peninsula.
- Dolerus (Poodolerus) zhelochovtsevi** Heidemaa et Viitasaari, 2009. Russia: **EP** (NW, C), **WS** (TM). – Europe (WE, EE, NE).
- Dolerus coreanus** Takeuchi, 1927 (*D. merops* Zhelochovtsev, 1928). Russia: **FE** (PR). – China (NE), Korean Peninsula, Japan (Haris, 2000).
- Dolerus ephippiatus** F. Smith, 1874. Host plant: Poaceae (including Hordeum, Triticum, Secale), Equisetum spp. (Equisetaceae). Russia: **FE** (SA, KU). – China (SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Dolerus harukawai** Waterston, 1926. Host plant: Juncus spp. (Juncaceae). Russia: **FE** (PR). – China (NE), Japan (Hon, Kyu).
- Dolerus hyrcanus** Benson, 1968. Russia: **EP** (NC).

Dolerus jesoenensis Matsumura, 1912. Russia: **FE** (SA). – Japan (Hok).

Dolerus lewisii Cameron, 1887. Host plant: *Hordeum* spp., *Triticum* spp. (Poaceae). Russia: **FE** (PR). – Korean Peninsula, Japan (Hon).

Dolerus novograblenovi Malaise, 1931. Russia: **FE** (PR, KA, MG). – Korean Peninsula.

Dolerus okeanskajensis Rohwer, 1925. Russia: **FE** (PR). – Korean Peninsula.

Dolerus sachalinensis Takeuchi, 1936. Russia: **FE** (SA).

Unplaced Dolerus

Dolerus carbonarius Zaddach, 1859. Russia: **EP** (NW).

Dolerus zaddachi W.F. Kirby, 1882 (*D. fumosus* Zaddach, 1859). Russia: **EP** (NW).

NEOSTROMBOCEROS Rohwer, 1912 (*Stypoza* Endrelein, 1920). Type species *Stromboceros metallica* Rohwer, 1912. The larvae feed on the ferns. The genus is distributed in East Palaearctic, Oriental and Australasian regions. Number of species: World – about 135; Palaearctic – about 70; Russia – 2.

Neostromboceros gracilis Takeuchi, 1941. Russia: **FE** (KU). – Japan (Hok, Hon, Shi).

Neostromboceros nipponicus Takeuchi, 1941. Russia: **FE** (KU). – China (SC), Japan (Hon, Shi, Kyu).

NESOSELANDRIA Rohwer, 1910 (*Neobusarbia* Takeuchi, 1928; *Melisandra* Benson, 1939). Type species *Paraselandria imitatrix* Ashmead, 1905. The larvae feed on the ferns and herbaceous plants. The genus is distributed in Palaearctic and Oriental region, two species in Neotropical region; one species introduced into N America. Number of species: World – about 85; Palaearctic – 25; Russia – 2.

Nesoselandria morio (Fabricius, 1781) [Tenthredo] (*Tenthredo ulmi* Schrank, 1802; *T. tristis* Serville, 1823; *T. tristis* Lepeletier, 1823; *Emphytus infuscatus* Eversmann, 1847; *Selandria fabricii* Konow, 1885; *Aneugmenus brunneus* Magretti, 1886). Host plant: Ditrichaceae, Dicranaceae, Rosaceae, Boraginaceae, etc. Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (KM, AL), **ES** (IR, BR), **FE** (AM, KH, PR, KU). – Europe (WE, EE, NE, SE), Georgia, Armenia, Turkey, Kazakhstan, China (SE), Korean Peninsula, Japan (Hon); introduced into N America.

Nesoselandria nipponica Takeuchi, 1929. Host plant: Polypodiophyta. Russia: **FE** (KU). – China (SE), Japan (Hon, Shi, Kyu).

POPPIA Konow, 1904. Type species *Poppia athalioides* Konow, 1904. The larvae feed on the ferns. The genus is distributed in East Asia. Number of species: World and Palaearctic – 2; Russia – 1.

Poppia athalioides Konow, 1904. Host plant: Polypodiophyta, ?*Tilia* sp. (Tiliaceae). Russia: **EP** (NC), **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (YA, ZB), **FE** (AM, KH). – China (NE), Japan (Sado).

PSEUDOHEMITAXONUS Conde, 1932. Type species *Pseudohemitaxonus exsectus* Conde, 1932. The genus is distributed in Palaearctic and Oriental region. Number of species: World – 4; Palaearctic – 3; Russia – 1.

Pseudohemitaxonus parvus Naito, 1969. Russia: **FE** (KU). – Japan (Hok, Hon).

ROCALIA Takeuci, 1952. Type species *Rocalia longipennis* Takeuchi, 1952. The genus is distributed in Palaearctic and Oriental region. Number of species: World – 14; Palaearctic – 3; Russia – 1.

Rocalia longipennis Takeuchi, 1952. Host plant: *Dryopteris* spp. (Dryopteridaceae). Russia: **FE** (KU). – Europe (NE), Japan (Hok, Hon).

SELANDRIA Leach, 1817 (*Cubitella* Malaise, 1944). Type species *Tenthredo serva* Fabricius, 1793. The larvae feed on Poaceae, Juncaceae, Cyperaceae. The genus is distributed in Palaearctic, Oriental and Neotropical regions. Number of species: World – 8; Palaearctic – 4; Russia – 3.

Selandria konoi Takeuchi, 1936. Russia: **FE** (SA). – Japan (Hon).

Selandria melanosterna (Serville, 1823) [Tenthredo] (*Tenthredo melanosterna* Lepeletier, 1823; *Selandria sixii* Snellen van Vollenhoven, 1858; *S. grandis* Zaddach, 1859; *S. interstitialis* Thomson, 1871). Host plant: *Carex* spp., *Scirpus* spp. (Cyperaceae), *Juncus* spp. (Juncaceae). Russia: **EP** (NW, C, E), **ES** (IR). – Europe (WE, EE, NE).

Selandria serva (Fabricius, 1793) [Tenthredo] (*Hylotoma serva mascula* Fallén, 1807; *Tenthredo socia* Klug, 1816; *T. lepida* Serville, 1823; *T. lepida* Lepeletier, 1823; *Selandria dorsalis* Stephens, 1835; *S. excisa* Konow, 1885; *S. serva interstitialis* Konow, 1885; *S. serva fuscitarsis* Benson, 1954; *S. serva punctatus* Zirngiebl, 1956). Host plant: Poaceae, Cyperaceae, Juncaceae. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, TK, NS, AL), **ES** (KR,

IR), **FE** (KH, PR). – Europe (WE, EE, NE, SE), Georgia, Armenia, Turkey, Iran, Kazakhstan, Mongolia, China (NE), Korean Peninsula.

STROMBOCEROS Konow, 1885 (*Strombocerina* Malaise, 1942; *Arbusia* Malaise, 1944). Type species *Tenthredo delicatulus* Fallén, 1808. The larvae on the ferns; pupate inside stems and branches. The genus is distributed in Holarctic, Oriental and Neotropical regions. Number of species: World – 55; Palaearctic – 2; Russia – 2.

Stromboceros delicatulus (Fallén, 1808) [Tenthredo] (*Tenthredo eborina* Klug, 1817; *Synairema alpina* Bremi-Wolf, 1849; *Selandria phthisica* Snellen van Vollenhoven, 1869; *S. virescens* Rudow, 1871; *Tenthredo seesana* Rudow, 1871; *Strongylogaster viridis* Schmiedeknecht, 1881; *Stromboceros delicatulus albiceps* Takeuchi, 1941). Host plant: Dryopteridaceae, Dennstaedtiaceae, Onocleaceae, Polypodiaceae. Russia: **EP** (N, NW, C, E), **UR**, **WS** (NS, KM, AL), **ES** (KR, IR, BR), **FE** (PR, SA, KU, KA). – Europe (WE, EE, NE), China (CC), Korean Peninsula, Japan (Hok, Hon).

Stromboceros koebelei Rohwer, 1910 (*Strongylogaster compressus* Matsumura, 1912; *Stromboceros japonicus* Malaise, 1931). Host plant: *Polystichum* spp. (Dryopteridaceae). Russia: **UR**, **FE** (PR, SA, KU). – Korean Peninsula, Japan (Hok, Hon).

STRONGYLOGASTER Dahlbom, 1835 (*Pseudotaxonus* Costa, 1894; *Polystichophagus* Ashmead, 1898; *Prototaxonus* Rohwer, 1910). Type species *Tenthredo multifasciata* Geoffroy, 1785. The larvae feed on the ferns. The genus is distributed in Holarctic, Oriental and Neotropical regions. Number of species: World – 55; Palaearctic – 25; Russia – 12.

Strongylogaster baikalensis Naito, 1990. Host plant: *Dryopteris* spp. (Dryopteridaceae), *Matteuccia* spp. (Onocleaceae). Russia: **ES** (IR), **FE** (KH). – Europe (EE).

Strongylogaster empriaeformis (Malaise, 1931) [*Thrinax*]. Host plant: Adiantaceae, Dryopteridaceae, Dennstaedtiaceae, Onocleaceae. Russia: **UR**, **WS** (KM), **FE** (PR, KA). – Japan (Hon).

Strongylogaster filicis (Klug, 1817) [Tenthredo] (*Tenthredo carinata* Klug, 1817; *T. atricornis* Stephens, 1835; *Sciopteryx apicalis* Matsumura, 1912). Host plant: *Pteridium aquilinum* (Dennstaedtiaceae). Russia: **EP** (NW, C, E), **UR**, **WS** (AL), **ES** (KR, IR), **FE** (AM, KH, SA). – Europe (WE, EE, NE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Sado).

Strongylogaster macula (Klug, 1817) [Tenthredo] (*Thrinax intermedia* Konow, 1885). Host plant: Dryopteridaceae, *Pteridium aquilinum* (Dennstaedtiaceae). Russia: **EP** (C, E, NC), **UR**, **ES** (IR, BR, ZB), **FE** (PR, KA). – Europe (WE, EE, NE), Armenia, China (NE, SE), Korean Peninsula, Japan (Hok, Hon); introduced into N America.

Strongylogaster mixta (Klug, 1817) [Tenthredo] (*S. femoralis* Cameron, 1875; *Selandria japonica* Matsumura, 1912; *Thrinax contigua orientalis* Malaise, 1931). Host plant: Adiantaceae, Dryopteridaceae, Dennstaedtiaceae. Russia: **EP** (NW, C, E), **UR**, **WS** (KM), **ES** (IR), **FE** (KU, KA). – Europe (WE, EE, NE), Japan (Hok, Hon).

Strongylogaster moiwana Matsumura, 1912. Host plant: *Comogramme* spp. (Adiantaceae), *Osmunda* spp. (Osmundaceae). Russia: **FE** (KU). – Japan (Hok, Hon, Shi).

Strongylogaster multifasciata (Geoffroy, 1785) [Tenthredo] (*Tenthredo lineata* Christ, 1791; *T. linearis* Klug, 1817; *T. alternans* Serville, 1823; *T. alternans* Lepeletier, 1823; *Strongylogaster iridipennis* F. Smith, 1874; *S. caucasicus* Schaposchnikov, 1885; *S. cretensis* Konow, 1887; *S. annularis* Matsumura, 1912; *S. lineata cypria* Benson, 1954). Host plant: Dennstaedtiaceae, Dryopteridaceae. Russia: **EP** (NW, C, E, NC), **UR**, **WS** (TM, NS, AL), **ES** (IR), **FE** (AM, PR, SA, KU). – Europe (WE, EE, NE, SE), Georgia, SW Asia, Iran, China (SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Sado).

Strongylogaster rubra Naito, 1980. Russia: **ES** (IR), **FE** (KU). – Korean Peninsula, Japan (Hok, Hon).

Strongylogaster secunda (Takeuchi, 1929) [Pseudotaxonus] (*Pseudotaxonus tertius* Conde, 1935; *P. secundus humeralis* Takeuchi, 1955). Host plant: *Osmunda japonica* (Osmundaceae). Russia: **FE** (PR, KU). – China (NE, SE), Japan (Hok, Hon, Shi, Kyu).

Strongylogaster struthiopteridis (Malaise, 1931) [*Thrinax*] (*Thrinax maura* Zhelochovtsev, 1988). Host plant: *Athyrium* spp., *Dryopteris* spp. (Dryopteridaceae), *Pteridium aquilinum* (Dennstaedtiaceae). Russia: **UR**, **WS** (KM, AL), **ES** (IR), **FE** (AM, PR, KA). – Europe (NE).

Strongylogaster verzhutskii Naito, 1990. Russia: **ES** (IR).

Strongylogaster xanthocera (Stephens, 1835) [Tenthredo] (*S. geniculata* Thomson, 1871; *S. desbrochersi* Konow, 1892; *S. desbrochersi lepticus* Konow, 1902). Host plant: *Dryopteris* spp., *Polystichum* spp., *Athyrium* spp. (Dryopteridaceae), *Pteridium*

aquinum (Dennstaedtiaceae). Russia: **EP** (NW, C), **WS** (AL), **ES** (IR), **FE** (AM, KH, PR). – Europe (WE, EE, NE, SE), China (NE, SE), Japan (Hon).

THRINAX Konow, 1885 (*Hemitaxonus* Ashmead, 1898; *Epitaxonus* MacGillivray, 1908; *Sahlbergia* Forsius, 1910; *Trearea* Malaise, 1947; *Canonarea* Malaise, 1947). Type species *Thrinax contigua* Konow, 1885. The larvae feed on the ferns. The genus is distributed in Holarctic and Oriental region. Number of species: World – 27; Palaearctic – 20; Russia – 7.

Thrinax angustata (Zhelochovtsev, 1951) [*Hemitaxonus*]. Russia: **FE** (KH, PR). – Korean Peninsula.

Thrinax athyrii (Naito, 1971) [*Hemitaxonus*]. Host plant: *Athyrium otophorum* (Dryopteridaceae). Russia: **FE** (KU). – Japan (Hok, Hon, Shi, Kyu).

Thrinax contigua Konow, 1885 (*Sahlbergia struthiopteridis* Forsius, 1910). Host plant: *Matteuccia* spp. (Onocleaceae). Russia: **EP** (NW), **ES** (IR), **FE** (PR, KU, KA). – Europe (NE), Japan (Hok, Hon).

Thrinax formosana (Takeuchi, 1928) [*Hemitaxonus*]. Russia: **FE** (PR). – China (SE).

Thrinax kamtchatica (Malaise, 1931) [*Hemitaxonus*]. Russia: **FE** (AM, PR, KA).

Thrinax paucipunctatus (Malaise, 1931) [*Hemitaxonus*]. Host plant: *Dryopteris* spp. (Dryopteridaceae). Russia: **ES** (IR), **FE** (KH, PR, KA). – Korean Peninsula, Japan (Hok, Hon).

Thrinax takeuchii (Naito, 1971) [*Hemitaxonus*]. Russia: **FE** (KU). – Japan (Hok, Hon).

Subfamily TENTHREDININAE

AGLAOSTIGMA W.F. Kirby, 1882 (*Laurentia* Costa, 1890; *Bivena* MacGillivray, 1894; *Homoeoneura* Ashmead, 1898; *Neopus* Viereck, 1910; *Macrophyopsis* Enslin, 1912; *Kincaidia* MacGillivray, 1914; *Astochus* MacGillivray, 1914; *Paralloma* Malaise, 1933; *Neurosiobla* Conde, 1935; *Laurentina* Malaise, 1937; *Stigmatosona* Malaise, 1945). Type species *Aglaostigma eburneiguttatum* W.F. Kirby, 1882. The larvae on herbaceous plants. The genus is distributed in Holarctic and Oriental region. Number of species: World – about 55; Palaearctic – 43; Russia – 15 from 5 subgenera.

Aglaostigma (Aglaostigma) amoorense (Cameron, 1876) [Tenthredo] (*Tenthredo amurensis* W.F. Kirby, 1882). Host plant: *Veratrum* spp. (Melanthiaceae).

Russia: **WS** (NS, KM, AL), **ES** (KR, IR, BR), **FE** (AM, KH, PR). – China (NC, NE), Korean Peninsula, Japan (Hon).

Aglaostigma (Aglaostigma) gibbosum (Fallén, 1808) [Tenthredo] (*Perineura alpina* Thomson, 1871; *Aglaostigma eburneiguttatum* W.F. Kirby, 1882; *Tenthredopsis novograblenovi* Malaise, 1931). Host plant: *Filipendula* spp. (Rosaceae). Russia: **EP** (N), **UR**, **ES** (IR), **FE** (PR, KA). – Europe (WE, EE, NE, SE).

Aglaostigma (Astochus) aucupariae (Klug, 1817) [Tenthredo] (*Tenthredo juvenilis* Serville, 1823; *T. juvenilis* Lepeletier, 1823; *Allantus collaris* Dietrich, 1868; *Laurentia craverii* Costa, 1890; *Macrophyia laticarpus* Kriechbaumer, 1891; *Aglaostigma aucupariae lacteore* Benson, 1968). Host plant: *Galium* spp. (Rubiaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (NS, KM), **ES** (IR, BR), **FE** (PR, KU). – Europe (WE, EE, NE), Georgia, Armenia, Turkey, China (Wei et al., 2006).

Aglaostigma (Astochus) fulvipes fulvipes (Scopoli, 1763) [Tenthredo]. Host plant: *Galium* spp. (Rubiaceae). Russia: **EP** (N, NW, C, E, S, NC), **WS** (KM, AL), **ES** (IR). – Europe (WE, EE, NE, SE), Georgia, Turkey, Central Asia, Kazakhstan.

Aglaostigma (Bivena) langei (Konow, 1894) [Rhoggastera] (*A. langei eichleri* Muche, 1975). Host plant: *Epilobium* spp. (Onagraceae). Russia: **EP** (NC). – Europe (WE, EE), Armenia.

Aglaostigma (Macrophyopsis) carbo (Malaise, 1931) [Macrophyopsis]. Russia: **FE** (PR).

Aglaostigma (Macrophyopsis) grosseseratum (Malaise, 1931) [Macrophyopsis]. Russia: **FE** (PR).

Aglaostigma (Macrophyopsis) nebulosum (André, 1881) [Macrophyia] (*Tenthredopsis schmiedeknechti* Konow, 1892). Host plant: *Filipendula* spp. (Rosaceae), *Impatiens* spp. (Balsaminaceae). Russia: **EP** (C, NC), **ES** (IR), **FE** (AM, PR, SA, KU). – Europe (WE, NE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Aglaostigma (Macrophyopsis) occipitosum (Malaise, 1931) [Macrophyopsis]. Host plant: *Angelica polymorpha* (Apiaceae), *Petasites japonicus* (Asteraceae). Russia: **FE** (PR). – China (SE), Korean Peninsula, Japan (Hon, Shi, Kyu, Sado).

Aglaostigma (Macrophyopsis) ochraceum (Malaise, 1931) [Macrophyopsis]. Russia: **FE** (PR).

Aglaostigma (Macrophyopsis) tricolor (Malaise, 1931) [Macrophyopsis]. Russia: **FE** (PR).

Aglaostigma (Neurosiobla) malaisei (Conde, 1935) [Neurosiobla]. Russia: **FE** (PR).

Aglaostigma (Neurosiobla) sachalinense (Conde, 1935) [Neurosiobla] (*A. togashii* Zombori, 1973). Russia: **ES** (Zhelochovtsev, Zinovjev, 1996), **FE** (PR, SA). – Japan (Zinovjev, 1994).

Aglaostigma (Neurosiobla) tertium Zinovjev, 1994. Russia: **FE** (KH, PR).

Aglaostigma sapporonicum (Matsumura, 1912) [Macrophyta] (*Dolerus biguttatus* Matsumura, 1912). Russia: **FE** (PR). – Japan (Hok, Hon).

ARMITARSUS Malaise, 1931. Type species *Armitarsus punctifemoratus* Malaise, 1931. The genus is distributed in East Asia. Number of species: World and Palaearctic – 7; Russia – 4.

Armitarsus albipes Malaise, 1931. Russia: **FE** (AM, PR).

Armitarsus minutus Zhelochovtsev, 1935. Russia: **FE** (AM, PR). – Korean Peninsula, Japan (Hok).

Armitarsus punctifemoratus Malaise, 1931. Host plant: *Alnus* spp. (Betulaceae). Russia: **FE** (PR, SA). – Korean Peninsula, Japan (Hok, Hon).

Armitarsus semirufus Takeuchi, 1936. Russia: **FE** (SA). – Japan (Hok, Hon).

CONASPIDIA Konow, 1898. Type species *Conaspidea sikkimensis* Konow, 1898. The genus is distributed in East Asia and Oriental region. Number of species: World – 23; Palaearctic – 13; Russia – 3.

Conaspidea guttata (Matsumura, 1912) [Eriocampa] (*C. trifasciata* Malaise, 1931; *C. murotai* Togashi, 1984; *C. ishikawai* Togashi, 2005). Host plant: *Kalopanax septemlobus* (Araliaceae). Russia: **FE** (SA, KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Conaspidea hyalina Sato, 1933. Russia: **FE** (KU). – Japan (Hok, Hon, Shi, Kyu).

Conaspidea singularis (Malaise, 1931) [Eriocampa] (*Macrophya flavipes* F. Smith, 1874). Russia: **FE** (KH, PR). – Japan (Hok, Hon).

CORYMBAS Konow, 1903 (*Siobloides* Takeuchi, 1919). Type species *Corymbas koreana* Konow, 1903. The genus is distributed in East Asia and Oriental region. Number of species: World – 7; Palaearctic – 5; Russia – 2.

Corymbas aperta (Takeuchi, 1919) [Siobloides]. Host plant: *Rhododendron* spp. (Ericaceae). Russia: **FE** (KU). – Japan (Hok, Hon, Shi, Kyu).

Corymbas koreana Konow, 1903. Russia: **FE** (PR). – China (SE), Korean Peninsula.

LAGIDINA Malaise, 1945. Type species *Macrophya irritans* F. Smith, 1874. The genus is distributed in East Palaearctic and Oriental region. Number of species: World – 8; Palaearctic – 6; Russia – 1.

Lagidina platycerus platycerus (Marlatt, 1898) [Tenthredo] (*Tenthredo platyceros* Konow, 1899). Host plant: *Iris laevigata* (Iridaceae), *Viola* spp. (Violaceae). Russia: **FE** (KU). – Japan (Hon, Shi, Kyu, Tsushima).

MACROPHYA Dahlbom, 1835 (*Zalagium* Rohwer, 1912; *Pseudomacrophya* Enslin, 1913; *Paramacrophya* Forsius, 1918). Type species *Tenthredo montana* Scopoli, 1763. The genus is distributed in Holarctic and Oriental region. Number of species: World – about 260; Palaearctic – about 160; Russia – 46 (+ 1 ssp.) from 2 subgenera.

Macrophya (Macrophya) albicincta (Schrank, 1776) [Tenthredo] (*Tenthredo albipes* Geoffroy, 1785; *T. albipalpis* Schrank, 1802; *T. luctuosa* Lepetier, 1823; *T. lugubris* Serville, 1823; *T. luctuosa* Serville, 1823; *Macrophya leucopoda* Palma, 1861; *Tenthredo magnicornis* Eversmann, 1864; *Macrophya melanostoma* Rudow, 1871; *M. albicincta decipiens* Konow, 1884; *Perineura cripiae* De-Stefani, 1885; *Macrophya albicincta candidata* Enslin, 1918; *M. albicincta agnani* Pic, 1948; *M. albicincta berlandi* Pic, 1948). Host plant: *Valeriana* spp. (Caprifoliaceae), *Sambucus* spp., *Viburnum* spp. (Adoxaceae). Russia: **EP** (C, S, NC). – Europe (WE, EE, NE, SE), N Africa, Georgia, Armenia, Turkey, Iran.

Macrophya (Macrophya) albipuncta (Fallén, 1808) [Tenthredo] (*Tenthredo arbustorum* Pollich, 1781; *T. liciata* Eversmann, 1847; *Macrophya nivosa* Costa, 1894). Host plant: *Geranium* spp. (Geraniaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (KM, AL), **ES** (KS). – Europe (WE, EE, NE, SE).

Macrophya (Macrophya) annulata (Geoffroy, 1785) [Tenthredo] (*Tenthredo dorsigera* Rossi, 1790; *T. similis* Spinola, 1808; *T. neglecta* Klug, 1817; *Allantus dejectus* Norton, 1860; *Macrophya neglecta nigra* Konow, 1894; *M. annulata theresae* Pic, 1918). Host plant: *Potentilla* spp., *Rosa* spp., *Rubus* sp. (Rosaceae), *Sambucus* spp. (Adoxaceae), *Origanum* spp. (Lamiaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (NS), **ES** (IR, BR, YA), **FE** (AM, PR). – Europe (WE, EE, NE, SE), Armenia, Turkey, Iran, Central Asia.

Macrophya (Macrophya) annulicornis Konow, 1904.

Host plant: *Sambucus* spp. (Adoxaceae). Russia: **UR, WS (AL), ES (IR, BR, ZB), FE (AM, SA)**. – Korean Peninsula.

Macrophya (Macrophya) annulitibia Takeuchi, 1933.

Russia: **FE (PR, SA, KU)**. – China (NE, NC, CC), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Macrophya (Macrophya) apicalis F. Smith, 1874.

Host plant: *Sambucus racemosa*, *S. sieboldiana* (Adoxaceae). Russia: **FE (KU)**. – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Macrophya (Macrophya) blanda (Fabricius, 1775)

[*Tenthredo*] (*Tenthredo ligustrina* Geoffroy, 1785; *T. cylindrica* Panzer, 1799; *T. albilabris* Klug, 1817; *T. lacrymosa* Lepeletier, 1823; *T. lacrymosa* Serville, 1823; *T. cognata* Fallén, 1829; *T. nyctea* Fischer von Waldheim, 1843; *Macrophya blanda brevicornis* Gradl, 1878; *M. albolapidaria* Kuznetzov-Ugamskij, 1927; *Tenthredo reductenotata* Pic, 1928). Host plant: *Rubus* spp., *Fragaria* spp. (Rosaceae). Russia: **EP (C, E, S, NC, CR), UR, WS** (Zhelochovtsev, Zinovjev, 1996). – Europe (WE, EE, NE, SE), Georgia, Armenia, Turkey, Kyrgyzstan, Kazakhstan.

Macrophya (Macrophya) carbonaria F. Smith, 1874.

Host plant: *Sambucus racemosa*, *S. sieboldiana* (Adoxaceae). Russia: **FE (SA, KU)**. – China (CC, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Awa-shima, Awaji-shima, Tsushima).

Macrophya (Macrophya) carinthiaca (Klug, 1817)

[*Tenthredo*]. Host plant: *Geranium* spp. (Geraniaceae). Russia: **EP (N, C), UR, WS (AL)**. – Europe (WE, EE, NE).

Macrophya (Macrophya) caucasica (Mocsáry, 1880)

[*Allantus*] (*M. prasinipes* Konow, 1891). Russia: **EP (NC)**. – Armenia.

Macrophya (Macrophya) chrysura (Klug, 1817)

[*Tenthredo*] (*M. albimacula* Mocsáry, 1881; *M. palidilabris* Costa, 1890). Host plant: *Daucus carota* (Apiaceae). Russia: **EP (S)**. – Europe (WE, EE, SE).

Macrophya (Macrophya) convexiscutellaris Muche, 1969. Russia: **EP (NC)**.**Macrophya (Macrophya) crassula** (Klug, 1817)

[*Tenthredo*] (*Tenthredo maculosa* Serville, 1823; *T. maculosa* Lepeletier, 1823; *Macrophya klugii* Snellen van Vollenhoven, 1869; *M. cora* W.F. Kirby, 1886). Host plant: *Sambucus* spp. (Adoxaceae). Russia: **EP (NC)**. – Europe (WE, EE, SE), Georgia, Armenia, Turkey.

Macrophya (Macrophya) diversipes (Schrank, 1782)

[*Tenthredo*] (*Tenthredo haematopus* Panzer, 1801;

T. ocreata Panzer, 1804; *T. rubripes* Drapiez, 1820; *T. corallipes* Eversmann, 1847; *Macrophya flavipes* Tischbein, 1852; *Tenthredo halensis* Aichinger, 1870; *Macrophya haemotopus immaculiventris* Costa, 1871; *M. eximia* Mocsáry, 1877; *M. caucasica* André, 1881; *M. rubripes* André, 1881; *M. saundersi* W.F. Kirby, 1886; *M. sanguinipes* Mocsáry, 1891; *M. dalmatina* Gasperini, 1891; *M. diversipes passerinii* Ghigi, 1905; *M. diversipes feminina* Enslin, 1913; *M. diversipes masculina* Enslin, 1913; *M. diversipes maculiventris* Enslin, 1913; *M. diversipes nigritarsis* Enslin, 1913). Host plant: *Fragaria* spp., *Rubus* spp. (Rosaceae). Russia: **EP (C, E, S, NC), UR, ES (TU)**. – Europe (WE, EE, SE), Georgia, Armenia, Turkey, Iran, Kyrgyzstan, Kazakhstan.

Macrophya (Macrophya) duodecimpunctata duodecimpunctata (Linnaeus, 1758) [*Tenthredo*] (*Tenthredo signata* Scopoli, 1763; *T. labiata* Geoffroy, 1785; *T. fera* Scopoli, 1786; *T. melanoleuca* Gmelin, 1790; *T. lugubris* Drapiez, 1820; *T. idriensis* Lepeletier, 1823; *T. albamacula* Lepeletier, 1823; *T. albamacula* Serville, 1823; *T. dolens* Eversmann, 1847; *T. curvipes* Gimmerthal, 1847; *Macrophya novemguttata* Costa, 1859; *M. luridicarpa* Costa, 1894; *M. duodecimpunctata nigrina* Konow, 1898). Host plant: Poaceae, *Carex* spp. (Cyperaceae). Russia: **EP (NW, C, E, S, NC), UR, WS (NS, AL)**. – Europe (WE, EE, NE, SE), N Africa, Georgia, Armenia, Turkey.

Macrophya (Macrophya) duodecimpunctata sodalitia Mocsáry, 1909. Host plant: *Carex* spp. (Cyperaceae). Russia: **FE (AM, KH, PR, SA)**. – Korean Peninsula, Japan (Hok, Hon).

Macrophya (Macrophya) erythrocnema Costa, 1859

(*Tenthredo femoralis* Eversmann, 1864). Russia: **EP (NW, C, E), UR**. – Europe (WE, EE, NE, SE), Armenia, SW Asia.

Macrophya (Macrophya) hamata caucasicola Muche, 1969. Russia: **EP (NC)**.

Macrophya (Macrophya) imitator Takeuchi, 1937.

Host plant: *Aster* spp. (Asteraceae). Russia: **FE (AM, PR, SA, KU)**. – Korean Peninsula, Japan (Hok, Hon).

Macrophya (Macrophya) infumata Rohwer, 1925

(*M. scutellata* Kuznetzov-Ugamskij, 1927). Host plant: *Sambucus racemosa* (Adoxaceae). Russia: **EP (E), UR, WS (KM, AL), ES (TU, IR), FE (AM, PR, SA, KU)**. – China (NE), Korean Peninsula, Japan (Hok, Hon).

Macrophya (Macrophya) kongosana Takeuchi, 1937.

Russia: **FE (PR)**. – Korean Peninsula.

Macrophya (Macrophya) koreana Takeuchi, 1937.

Host plant: *Isodon effusus* (Lamiaceae). Russia: **FE** (AM, PR). – China (NC, CC), Korean Peninsula, Japan (Hon).

Macrophya (Macrophya) maculitibia Takeuchi, 1933. Host plant: *Petasites japonicus*, *Parasenecio hastatus* (Asteraceae). Russia: **ES** (IR), **FE** (SA, KU). – Korean Peninsula, Japan (Hok, Hon, Shi).**Macrophya (Macrophya) militaris** (Klug, 1817) [Tenthredo] (*Tenthredo schaefferi* Serville, 1823; *T. schaefferi* Lepeletier, 1823; *Macrophya lepeletieri* Costa, 1859; *M. militaris cabrerae* Konow, 1896; *M. militaris nigriscutis* Enslin, 1913; *M. militaris falsa* Pic, 1928). Host plant: *Rubus* spp. (Rosaceae), *Origanum* spp. (Lamiaceae). Russia: **EP** (S, NC). – Europe (WE, EE, SE), Turkey.**Macrophya (Macrophya) montana montana** (Scopoli, 1763) [Tenthredo] (*Tenthredo trifasciata* Geoffroy, 1785; *T. sulphurata* Gmelin, 1790; *T. melanochra* Gmelin, 1790; *T. tricincta* Christ, 1791; *T. notata* Panzer, 1799; *T. albimana* Lepeletier, 1823; *T. albimana* Serville, 1823; *T. laserpitii* Lepeletier, 1823; *Macrophya rustica scutellaris* Enslin, 1913; *M. rustica pleuralis* Enslin, 1913; *M. rustica martialis* Pic, 1925; *M. rustica luteonotata* Pic, 1925; *M. rustica kabyliana* Pic, 1929). Host plant: *Rubus* spp. (Rosaceae), *Alnus* spp. (Betulaceae). Russia: **EP** (NW, C, E, S, NC, CR), **UR**. – Europe (WE, EE, NE, SE), N Africa, Georgia, Armenia, Turkey, Iran, Turkmenistan.**Macrophya (Macrophya) nemesis** Muche, 1969. Russia: **EP** (NC).**Macrophya (Macrophya) ottomana** Mocsáry, 1881. Russia: **EP** (NC). – Armenia, Turkey.**Macrophya (Macrophya) postica** (Brullé, 1832) [Tenthredo] (*M. ratzeburgii* Tischbein, 1852; *M. histrionica* Snellen van Vollenhoven, 1878; *M. postica nigripileuris* Enslin, 1913; *M. postica luteomaculata* Pic, 1918). Host plant: *Fragaria* spp., *Rubus* spp. (Rosaceae). Russia: **EP** (NC, CR). – Europe (EE, SE), Georgia, Armenia, Turkey.**Macrophya (Macrophya) potanini** Jakovlev, 1891. Russia: **FE** (AM). – China (NC).**Macrophya (Macrophya) recognata** Zombori, 1979 (*M. cognata* Mocsáry, 1881). Russia: **EP** (NC, CR). – Europe (WE, EE, SE), Georgia, Armenia.**Macrophya (Macrophya) ribis** (Schrank, 1781) [Tenthredo] (*Tenthredo leucopus* Gmelin, 1790; *T. exalbida* Gmelin, 1790; *Macrophya bertolini* Cobelli, 1890; *M. ribis morvandica* Pic, 1948). Host plant:

Sambucus spp. (Adoxaceae). Russia: **EP** (NC). – Europe (WE, EE, NE, SE).

Macrophya (Macrophya) rufipes (Linnaeus, 1758) [Tenthredo] (*Tenthredo pavida* Fabricius, 1775; *T. dumetorum* Geoffroy, 1785; *T. multicolor* Geoffroy, 1785; *T. flavifasciata* Christ, 1791; *T. rufipes* Christ, 1791; *T. strigosa* Fabricius, 1798; *T. citreipes* Lepeletier, 1823; *T. citreipes* Serville, 1823; *Allantus ione* Newman, 1837; *Macrophya rufipes orientalis* Mocsáry, 1891; *M. rufipes muliebris* Enslin, 1913; *M. rufipes castiliensis* Enslin, 1914; *M. rufipes reductenotata* Pic, 1929; *M. rufipes diversereducta* Pic, 1929). Host plant: *Vitis* spp. (Vitaceae), *Agromonia* spp. (Rosaceae). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, EE, NE, SE), Armenia, Turkey, Turkmenistan.**Macrophya (Macrophya) sanguinolenta** (Gmelin, 1790) [Tenthredo] (*Tenthredo quadrimaculata* Fabricius, 1787; *T. picea* Gmelin, 1790; *Allantus sambuci* Panzer, 1803; *Tenthredo comma* Fallén, 1808; *Macrophya trochanterica* Costa, 1858; *Tenthredo poecilopus* Aichinger, 1870; *Macrophya quadrimaculata nigrifrons* Jakovlev, 1886; *M. quadrimaculata nigra* Jakovlev, 1888; *M. sanguinolenta borealis* Forsius, 1918; *M. sanguinolenta albitarsis* Enslin, 1918). Host plant: *Veronica* spp. (Plantaginaceae), *Senecia* spp. (Asteraceae), *Galeopsis* spp. (Lamiaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **ES** (IR), **FE** (AM, PR, SA, KU). – Europe (WE, EE, NE, SE), Georgia, Armenia, Turkey, Kazakhstan, Mongolia, China (NE, NC), Korean Peninsula.**Macrophya (Macrophya) sibirica** Forsius, 1918 (*M. sibiricola* Forsius, 1925). Host plant: *Sambucus* spp. (Adoxaceae). Russia: **WS** (AL), **ES** (IR), **FE** (KH, PR). – China (NE, CC), Korean Peninsula.**Macrophya (Macrophya) superba** Tischbein, 1852 (*Tenthredo erythropus* Brullé, 1832; *Macrophya erythropus croatica* Korlević, 1890; *M. flavipennis* Kriechbaumer, 1891; *M. erythropus fluminensis* Strobl, 1901; *M. superba nigricans* Enslin, 1913). Host plant: *Rubus* spp., *Fragaria* spp. (Rosaceae), *Euphorbia* (Euphorbiaceae). Russia: **EP** (NC). – Europe (EE, SE), Armenia, SW Asia.**Macrophya (Macrophya) teutona** (Panzer, 1799) [Tenthredo] (*Tenthredo aureatensis* Schrank, 1802; *Macrophya marginata* Mocsáry, 1881; *M. klugi* Konow, 1894). Host plant: *Euphorbia* spp. (Euphorbiaceae). Russia: **EP** (NC), **UR** (S). – Europe (WE, EE, SE), Turkey.

Macrophyia (Macrophyia) vacillans Malaise, 1931
(*M. vacillans punctifrons* Malaise, 1931). Host plant:
Ligustrum spp., Syringa spp. (Oleaceae). Russia: **FE**
(PR). – Korean Peninsula.

Macrophyia (Pseudomacrophyia) albitarsis Mocsáry,
1909. Russia: **FE** (KH). – Korean Peninsula.

Macrophyia (Pseudomacrophyia) crassuliformis
Forsius, 1925 (*M. brevilabris* Malaise, 1931; *M. brevi-
labris nigroscutellata* Malaise, 1931). Host plant:
?Ligustrum obtusifolium (Oleaceae). Russia: **FE**
(KH, PR). – China (Shinohara, 2015), Korean Pen-
insula, Japan (Hon, Shi).

Macrophyia (Pseudomacrophyia) dibowskii André,
1881. Russia: “Sibiria” (André, 1881), **FE** (AM).

Macrophyia (Pseudomacrophyia) minutissima Take-
uchi, 1937. Russia: **FE** (PR). – Korean Peninsula.

Macrophyia (Pseudomacrophyia) punctumalbum
(Linnaeus, 1767) [Tenthredo] (*Tenthredo erythropus*
Schrank, 1776; *T. punctum* Fabricius, 1781; *T. stel-
lata* Geoffroy, 1785). Host plant: Ligustrum spp.,
Fraxinus spp. (Oleaceae), Crataegus spp. (Rosaceae),
Quercus spp. (Fagaceae). Russia: **EP** (NW, C, S, NC,
CR). – Europe (WE, EE, NE, SE), Georgia, Armenia.

Macrophyia brunnipes André, 1881 (*M. bruneipes*
Dalla Torre, 1894). Russia: “Sibiria” (André, 1881).

Macrophyia coxalis (Motschulsky, 1866) [Dolerus]
(*M. ignava* F. Smith, 1874; *Emphytus japonicus*
W.F. Kirby, 1882; *Macrophyia discreta* Forsius,
1925). Host plant: Rosa spp. (Rosaceae). Russia:
FE (KU). – China (Wei et al., 2006), Korean Pen-
insula, Japan (Hok, Hon, Shi, Kyu, Sado, Awaji-
shima, Oki).

Macrophyia esakii (Takeuchi, 1923) [Pachyprotas-
sis] (*M. exilis* Takeuchi, 1933). Russia: **FE** (SA). –
Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Macrophyia timida F. Smith, 1874 (*M. nigra* Marlatt,
1898; *M. femorata* Marlatt, 1898; *M. nigrita* Enslin,
1910; *M. mikagei* Togashi, 2005). Host plant: Ligus-
trum spp., Syringa vulgaris (Oleaceae). Russia: **FE**
(PR). – China (Shinohara, 2015), Korean Penin-
sula, Japan (Hok, Hon, Shi, Kyu, Sado, Awaji-shima,
Tsushima).

Macrophyia tristis André, 1881. Russia: “Sibiria”
(André, 1881), **FE** (AM).

PACHYPROTASIS Hartig, 1837 (*Lithracia* Cam-
eron, 1902). Type species *Tenthredo rapae*
Linnaeus, 1767. The genus is distributed in Hol-
arctic, Oriental and Neotropical regions. Number

of species: World – about 200; Palaearctic – about
165; Russia – 17.

Pachyprotasis antennata (Klug, 1817) [Tenthredo]
(*P. antennata exannulata* Malaise, 1931). Host plant:
Senecio spp. (Asteraceae), Filipendula spp. (Rosa-
ceae), Fraxinus spp. (Oleaceae), Atropa spp. (Solana-
ceae), etc. Russia: **EP** (N, C, E), **UR**, **WS** (NS, KM,
AL), **ES** (IR, BR), **FE** (AM, PR, SA). – Europe (WE,
EE, NE, SE), Mongolia, China (NE, NC), Korean
Peninsula, Japan (Hok, Hon, Kyu, Sado).

Pachyprotasis elegans Takeuchi, 1956. Russia: **FE**
(KU). – Japan (Hok).

Pachyprotasis erratica erratica F. Smith, 1874 (*M. fla-
voventralis* Matsumura, 1912). Russia: **FE** (SA,
KU). – China (SE), Japan (Hok, Hon, Shi).

Pachyprotasis glabrata Malaise, 1931. Russia: **FE**
(PR). – Korean Peninsula.

Pachyprotasis laeviceps Malaise, 1931. Russia: **FE**
(PR). – Korean Peninsula.

Pachyprotasis limitaris Malaise, 1931. Russia: **FE**
(PR).

Pachyprotasis lineicoxis Malaise, 1931. Russia: **WS**
(Zhelochovtsev, Zinovjev, 1996), **FE** (PR). – China
(SE), Japan.

Pachyprotasis longicornis Jakovlev, 1891 (*Macro-
phyia nigrolineata* Matsumura, 1912; *Pachyprotasis
longicornis kuriharum* Malaise, 1931). Russia: **FE**
(SA, KU). – China (NC), Korean Peninsula, Japan
(Hok, Hon).

Pachyprotasis nigra Stroganova, 1978. Russia: **FE**
(CH).

Pachyprotasis nigronotata Kriechbaumer, 1874
(*P. formosa* Schmiedeknecht, 1881; *P. viridis* Bris-
chke, 1883). Host plant: Plantago spp. (Planta-
ginaceae), Mentha spp. (Lamiaceae), Geum spp.,
Alchemilla spp. (Rosaceae), etc. Russia: **EP** (NW,
C), **ES** (IR), **FE** (AM, KH, PR). – Europe (WE, EE,
NE), China (SE), Korean Peninsula.

Pachyprotasis pedatoria (Malaise, 1931) [*Macro-
phyia*]. Russia: **FE** (PR).

Pachyprotasis rapae (Linnaeus, 1767) [Tenthredo]
(*Tenthredo leucomelas* Strøm, 1768; *T. scripta*
Gmelin, 1790; *T. luctuosa* Schrank, 1802; *T. floricola*
Gravenhorst, 1807; *T. nigrofasciata* Eschscholtz,
1822; *T. duplex* Serville, 1823; *T. duplex* Lepeletier,
1823; *Macrophyia omega* Norton, 1867; *Synairema
americana* Provancher, 1885; *Rhogogaster sayi*
Rohwer, 1908; *Macrophyia obnata* MacGillivray,
1923; *Pachyprotasis rapae nigrosternum* Koch,

1984). Host plant: *Solidago* spp. (Asteraceae), *Scrophularia* spp. (Scrophulariaceae), *Fraxinus* spp. (Oleaceae), etc. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (NS, KM, AL), **ES** (BR, YA, ZB), **FE** (AM, PR, SA, KU, KA, MG). – Europe (WE, EE, NE, SE), Armenia, Turkey, Mongolia, China (Wei et al., 2006), Korean Peninsula, Japan (Hok, Hon), India, N America, Mexico.

Pachyprotasis sejensis sejensis Inomata, 1984. Russia: **FE** (KU). – Japan (Hon).

Pachyprotasis simulans (Klug, 1817) [Tenthredo] (*P. laevicollis* Thomson, 1871; *P. tenuis* Rudow, 1871). Host plant: *Scrophularia* spp. (Scrophulariaceae), *Solidago* spp., *Senecio* spp. (Asteraceae). Russia: **EP** (NW, C), **FE** (PR). – Europe (WE, EE, NE), Mongolia, China (NC).

Pachyprotasis variegata (Fallén, 1808) [Tenthredo] (*P. variegata tenebrosa* Malaise, 1931). Host plant: *Solanum tuberosum* (Solanaceae), *Digitalis* spp. (Plantaginaceae), *Leontodon* spp., *Leucanthemum* spp. (Asteraceae), etc. Russia: **EP** (N, C, E), **UR**, **ES** (BR), **FE** (AM, PR, SA, MG). – Europe (WE, EE, NE), Central Asia, Mongolia, China (NE), Korean Peninsula, Japan.

Pachyprotasis vicaria Malaise, 1931. Russia: **FE** (PR). – Japan (Hok).

Pachyprotasis zukaensis Inomata, 1970. Russia: **FE** (KU). – Japan (Hon).

PERINEURA Hartig, 1837 (*Synairema* Hartig, 1837; *Synaerema* Agassiz, 1848; *Perinevra* Agassiz, 1848; *Synaerema* Schulz, 1906). Type species *Allantus rubi* Panzer, 1803. The genus is distributed in East Asia; one species in the west of Palaearctic and one species in Oriental region. Number of species: World – 11; Palaearctic – 10; Russia – 2.

Perineura japonica Malaise, 1931. Russia: **FE** (PR, SA). – Japan (Hok, Hon).

Perineura rubi (Panzer, 1803) [*Allantus*] (*Tenthredo lividiventris* Fallén, 1808; *T. elegantula* Fallén, 1808; *T. delicatula* Klug, 1817). Host plant: *Rubus* spp. (Rosaceae). Russia: **EP** (NW, C, E). – Europe (WE, EE, NE).

RHOGOGASTER Konow, 1884 (*Rhogogastera* Konow, 1885; *Cytisodaster* Lacourt, 1997). Type species *Tenthredo viridis* Linnaeus, 1758. The larvae on the flowering plants. The genus is distributed in Holarctic and Oriental region. Number

of species: World – 38; Palaearctic – about 35; Russia – 13.

Rhogogaster chlorosoma (Benson, 1943) [Tenthredo]. Host plant: Dennstaedtiaceae, Salicaceae, Betulaceae, Rosaceae. Russia: **EP** (C, E), **UR**, **WS** (AL), **ES** (IR, YA), **FE** (SA, KU). – Europe (WE, EE, NE, SE), Turkey, Mongolia, Japan.

Rhogogaster convergens Malaise, 1931. Russia: **ES** (IR), **FE** (AM, KH, PR). – China (NE), Korean Peninsula, Japan (Hok).

Rhogogaster coryli Muche, 1973. Host plant: *Corylus* spp. (Betulaceae). Russia: **EP** (NC).

Rhogogaster genistae Benson, 1947 (*Rh. genistae viridifrons* Muche, 1973). Host plant: *Genista* spp., *Sarothamnus* spp. (Fabaceae). Russia: **EP** (NC). – Europe (WE, EE, NE, SE), N Africa, Turkey.

Rhogogaster magniserrula Viitasaari, 2015. Russia: **EP** (N). – Europe (NE).

Rhogogaster nigriventris Malaise, 1931 (*Tenthredo varipes* W.F. Kirby, 1882). Host plant: *Geranium* spp. (Geraniaceae), *Rosa* spp. (Rosaceae). Russia: **ES** (IR), **FE** (AM, KH, PR, SA). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Rhogogaster picta (Klug, 1817) [Tenthredo] (*Tenthredo breviuscula* Costa, 1859). Host plant: *Sarothamnus* spp., *Genista* spp., *Cytisus* spp. (Fabaceae). Russia: **EP** (C, E, S, NC), **ES** (TU, ZB). – Europe (WE, EE, NE, SE), N Africa, Georgia, Syria, Kazakhstan, Mongolia.

Rhogogaster polaris Lindqvist, 1964. Russia: **EP** (N), **ES** (IR, BR), **FE** (MG). – Europe (WE, NE).

Rhogogaster punctulata (Klug, 1817) [Tenthredo]. Host plant: Salicaceae, Betulaceae, Rosaceae, Oleaceae. Russia: **EP** (N, NW, C), **UR**, **WS** (NS), **ES** (KR, IR, BR, ZB). – Europe (WE, EE, NE, SE), Turkey, Mongolia, China (NC).

Rhogogaster scalaris (Klug, 1817) [Tenthredo] (*Rh. viridis* auct., nec Linnaeus; *Tenthredo pictipes* Förster, 1850; *Rhogogaster viridis melanonota* Enslin, 1912; *Rh. viridis nigroscutellata* Forsius, 1918; *Rh. viridis lapponica* Enslin, 1918; *Rh. chlorosoma podkumokensis* Muche, 1973; *Tenthredo carpatica* Zhelochovtsev et Zinovjev, 1988). Host plant: polyphage. Russia: **EP** (C, NC), **UR**, **WS** (KM, AL), **ES** (TU, BR), **FE** (KA). – Europe (WE, EE, NE).

Rhogogaster sibirica Enslin, 1912 (*Rh. kudianus* Rohwer, 1925). Russia: **ES** (IR), **FE** (PR).

Rhogogaster tsherepanovi Stroganova, 1988. Russia: **EP** (C), **WS** (NS), **ES** (KR).

Rhogogaster viridis (Linnaeus, 1758) [Tenthredo] (*Tenthredo dryas* Benson, 1943; *Rhogogaster similis* Lindqvist, 1959). Host plant: Salicaceae, Betulaceae, Onagraceae, Fagaceae, Rhamnaceae, Rosaceae, Fabaceae. Russia: **EP** (N, NW, C, E), **WS** (NS, AL), **ES** (IR, BR, YA, ZB), **FE** (PR, KU, KA, MG). – Europe (WE, EE, NE), Kazakhstan, Mongolia, China (NC), Korean Peninsula, SE Asia.

SCIAPTERYX Stephens, 1835 (*Eniscia* Thomson, 1870; *Sciopteryx* Cameron, 1882; *Sciapterina* Ushinskij, 1940). Type species *Tenthredo costalis* Fabricius, 1775. The genus is distributed in Palaearctic. Number of species: World and Palaearctic – 16; Russia – 9.

Sciapteryx caucasica Dovnar-Zapolskij, 1930. Russia: **EP** (NC). – Turkey.

Sciapteryx circassica Dovnar-Zapolskij, 1930. Russia: **EP** (NC). – Georgia, Armenia, Turkey.

Sciapteryx consobrina (Klug, 1816) [Tenthredo] (*Eriocampa marginata* Puls, 1870; *Allantus consobrinus zwickoviensis* Schlechtendal, 1873). Host plant: Adoxa spp. (Adoxaceae), Anemone spp., Ranunculus spp. (Ranunculaceae). Russia: **EP** (NW, C, S, NC). – Europe (WE, EE, NE, SE), Georgia.

Sciapteryx costalis costalis (Fabricius, 1775) [Tenthredo]. Host plant: Ranunculus spp. (Ranunculaceae). Russia: **EP** (C). – Europe (WE, EE, NE, SE).

Sciapteryx dovnari Ushinskij, 1940. Russia: **EP** (NC). – Georgia, Armenia.

Sciapteryx laeta Konow, 1891 (*S. laeta antennata* Muche, 1973). Host plant: Ranunculus spp. (Ranunculaceae). Russia: **EP** (NC). – Georgia, Turkey.

Sciapteryx montana Dovnar-Zapolskij, 1930. Russia: **EP** (NC).

Sciapteryx semenowi Jakovlev, 1886. Host plant: Euclidium spp., Capsella spp. (Brassicaceae). Russia: **EP** (S, CR). – Europe (EE).

Sciapteryx verticalis Muche, 1973. Russia: **EP** (NC).

SIOBLA Cameron, 1877 (*Encarsioneura* Konow, 1890; *Megasiobia* Dovnar-Zapolskij, 1930). Type species *Siobla mooreana* Cameron, 1877. The genus is distributed in Palaearctic and Oriental region. Number of species: World – about 75; Palaearctic – about 55; Russia – 12.

Siobla ferox (F. Smith, 1874) [Macrophyia] (*S. grandis* Matsumura, 1912). Host plant: Balsaminaceae, Polygonaceae, Amaranthaceae. Russia: **FE**

(PR). – China (SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Tsushima).

Siobla frigida (Mocsáry, 1909) [Encarsioneura]. Russia: **FE** (KH). – China (NE, SE).

Siobla grossa Malaise, 1931. Russia: **FE** (AM, PR).

Siobla jucunda (Mocsáry, 1909) [Encarsioneura]. Russia: **FE** (KH, PR, SA, KU). – Japan (Hok, Hon, Shi).

Siobla malaisei Mallach, 1933. Russia: **FE** (PR). – China (NE, CC).

Siobla ruficornis (Gimmerthal, 1834) [Allantus] (*Eriocampa ruficornis* Cameron, 1876; *Siobla sibirica* Malaise, 1931; *S. sibirica bergmani* Malaise, 1931; *S. ruficornis albicornis* Malaise, 1931). Host plant: Epilobium angustifolium (Onagraceae). Russia: **EP** (NW, C, E), **UR**, **WS** (NS, AL), **ES** (IR, BR, YA), **FE** (KH, PR, SA, KU, KA). – Europe (NE), China (NE), Korean Peninsula, Japan (Hon).

Siobla similis (Mocsáry, 1909) [Encarsioneura]. Russia: **FE** (KH, SA).

Siobla sturmii sturmii (Klug, 1817) [Tenthredo] (*Macrophya castanea* Jakovlev, 1888; *M. pacifica* F. Smith, 1874). Host plant: Impatiens spp. (Balsaminaceae), Rubus spp., Rosa spp., Prunus spp. (Rosaceae). Russia: **EP** (NW, C), **WS** (NS), **ES** (YA), **FE** (AM, PR, SA, KU). – Europe (WE, EE, NE), Korean Peninsula, Japan (Hok, Hon, Kyu).

Siobla takeuchii Shinohara, Wei et Niu, 2013. Russia: **FE** (KU). – Japan (Hok, Hon, Kyu).

Siobla venusta venusta (Konow, 1903) [Ercarsioneura]. Russia: **FE** (PR). – China (CC), Korean Peninsula.

Siobla villosa Malaise, 1931. Russia: **FE** (PR). – China (CC).

Siobla zenaida Dovnar-Zapolskij, 1930. Russia: **FE** (KH). – China (NE, CC), Korean Peninsula.

TENTHREDO Linnaeus, 1758 (*Allantus* Jurine, 1801; *Parastatis* W.F. Kirby, 1881; *Dipteromorpha* W.F. Kirby, 1882; *Labidia* Provancher, 1886; *Jermakia* Jakovlev, 1891; *Ischyroceraea* Kiær, 1898; *Rethrax* Cameron, 1899; *Fethalia* Cameron, 1902; *Peus* Konow, 1903; *Clydostomus* Konow, 1908; *Tenthredella* Rohwer, 1910; *Tenthredina* Rohwer, 1910; *Zamacrophyia* Rohwer, 1912; *Metallopeus* Malaise, 1934; *Jakovleviella* Malaise, 1937; *Adungia* Malaise, 1945; *Ebba* Malaise, 1945; *Propodea* Malaise, 1945; *Elinora* Benson, 1946; *Eurogaster* Zirngiebl, 1953; *Cuneala* Zirngiebl, 1956; *Elinopsis* Lacourt, 1986; *Cephaledo* Zhelochovtsev, 1988; *Maculedo* Zhelochovtsev, 1988; *Olivacedo* Zhelochovtsev,

1988; *Temuledo* Zhelochovtsev, 1988; *Zonuledo* Zhelochovtsev, 1988; *Murciana* Lacourt, 1988; *Absentia* Togashi, 1990; *Casipteryx* Taeger, 1991; *Paratenthredo* Taeger, 1991; *Dorhettenyx* Lacourt, 1997; *Endotethryx* Lacourt, 1997; *Sainiella* Lacourt, 1997; *Blankia* Lacourt, 1998). Type species *Tenthredo scrophulariae* Linnaeus, 1758. The larvae on various plants; often polyphages. Adults often found on the inflorescences of Apiaceae, Spiraea, and others; females of many species are predators. The genus is distributed in Holarctic, Oriental and Neotropical regions. Number of species: World – > 900; Palaearctic – about 655; Russia – 158 (+ 13 ssp.) from 14 subgenera.

Tenthredo (Cephaledo) bifasciata diversipennis (Mocsáry, 1891) [Allantus]. Host plant: Asteraceae. Russia: EP (NC). – Georgia, Armenia, SW Asia.

Tenthredo (Cephaledo) bifasciata rossii (Panzer, 1803) [Allantus] (*T. bifasciata* Klug, 1817; *T. scissa* Klug, 1819; *T. rossii pigeoti* Pic, 1948; *T. rossica laeviceps* Hellén, 1955). Host plant: Sonchus spp., Hieracium spp. (Asteraceae). Russia: EP (NW, C). – Europe (WE, EE, NE).

Tenthredo (Cephaledo) bifasciata violacea (André, 1881) [Allantus] (*T. interjecta* Kuznetzov-Ugamskij, 1924). Host plant: Asteraceae. Russia: EP (C, E, S), WS (AL). – Europe (EE, SE), Kazakhstan.

Tenthredo (Cephaledo) caucasica caucasica Eversmann, 1847 (*Allantus moestus* Mocsáry, 1883). Russia: EP (S, NC). – Georgia, Armenia, SW Asia.

Tenthredo (Cephaledo) costata Klug, 1817 (*T. sareptana* Eversmann, 1847; *Allantus faustus* W.F. Kirby, 1882; *A. subcostatus* Jakovlev, 1888; *A. graecus* Konow, 1888; *A. parnasius* Konow, 1888; *A. violaceipennis* Costa, 1890; *A. kiefferi* Konow, 1899; *A. kiettereri ciliix* Enslin, 1910; *Tenthredo kiefferi lugubrata* Enslin, 1914; *Allantus costatus obscurus* Zirngiebl, 1937). Host plant: Chondrilla spp. (Asteraceae), Euphorbia spp. (Euphorbiaceae). Russia: EP (S, NC). – Europe (EE, SE), Georgia, Armenia, SW Asia, Iran.

Tenthredo (Cephaledo) excellens (Konow, 1886) [Allantus] (*T. annulata* Klug, 1817; *Allantus persa* Konow, 1888; *A. persa mandibularis* Enslin, 1910; *Tenthredo persa caja* Enslin, 1912). Russia: EP (S, NC), UR. – Europe (EE), Georgia, Armenia, Turkey, Iran, Turkmenistan.

Tenthredo (Cephaledo) neobesa Zombori, 1980 (*T. zonata* Fallén, 1808; *Allantus obesus* Mocsáry, 1880; *Tenthredo pseudorossii* Taeger, 1985). Host

plant: Hieracium spp., Sonchus spp. (Asteraceae). Russia: EP (E), UR, WS (NS, AL), ES (IR). – Europe (WE, EE, NE), Georgia.

Tenthredo (Cephaledo) segmentaria Fabricius, 1798 (*Allantus steckii* Konow, 1888; *A. stecki trimarginata* Pic, 1933; *A. stecki milliati* Pic, 1933). Russia: ES (TU, IR). – Europe (WE, EE).

Tenthredo (Cephaledo) violascens (Konow, 1886) [Allantus]. EP (NC). – Georgia, Armenia, Turkey.

Tenthredo (Dorhettenyx) jozana (Matsumura, 1912) [*Tenthredopsis*] (*T. fuscata* Enslin, 1920). Russia: WS (AL), FE (PR, SA, KU). – China (SE), Korean Peninsula, Japan (Hok, Hon, Shi).

Tenthredo (Elinora) aulica Enslin, 1912 (*Allantus nigritarsis* Konow, 1886). Russia: EP (NC).

Tenthredo (Elinora) caspia (André, 1881) [Allantus]. Russia: EP (S). – Europe (EE).

Tenthredo (Elinora) dahlii Klug, 1817 (*T. villosa* Brullé, 1832; *T. unicincta* Brullé, 1832; *Allantus xanthorius* Kriechbaumer, 1869; *Macrophyia hartigii* W.F. Kirby, 1882). Russia: EP (S, NC, CR), ES (BR, ZB). – Europe (EE, SE), Uzbekistan.

Tenthredo (Elinora) flaveola Gmelin, 1790 (*T. flavipes* Geoffroy, 1785; *T. rubiginosa* Gmelin, 1790; *T. dispar* Klug, 1817; *T. grata* Lepeletier, 1823; *T. grata* Serville, 1823; *Allantus rufocingulatus* Tischbein, 1852; *A. orientalis* Kriechbaumer, 1869; *A. galicensis* Niegabitowski, 1899; *A. flaveola delagrangei* Pic, 1925). Host plant: Bupleurum spp. (Apiaceae), Raphanus spp., Brassica spp., Sinapis spp. (Brassicaceae). Russia: EP (S, NC, CR), WS (Zhelochovtsev, Zinovjev, 1996). – Europe (WE, EE, NE), Armenia.

Tenthredo (Elinora) koehleri Klug, 1817 (*Allantus funereus* Palma, 1861; *A. koehleri scutellaris* Gregor, 1941). Host plant: Cardamine spp. (Brassicaceae), Geranium sanguineum, Robertianum spp. (Geraniaceae). Russia: EP (NC). – Europe (WE, EE, SE), Turkey.

Tenthredo (Elinora) longipes (Konow, 1886) [Allantus] (*Allantus shestoperovi* Ushinskij, 1936; *Cuneala tricolor* Zirngiebl, 1956). Russia: EP (NC). – Georgia, Armenia, Iran, Turkmenistan.

Tenthredo (Elinora) ornata (André, 1881) [Allantus] (*Allantus caucasicus* Mocsáry, 1880; *A. discolor* Konow, 1890; *A. mocsaryi* Enslin, 1910; *Tenthredo baja* Enslin, 1912; *T. camilla* Enslin, 1912). Russia: EP (NC). – Armenia, Turkey.

Tenthredo (Elinora) persica (André, 1882) [Allantus] (*T. coniensis* Enslin, 1914; *Allantus kareli* Muche,

- 1962; *A. kareli maculatus* Muche, 1962). Russia: **EP** (Lacourt, 1999). – Caucasus, Turkey, Iran.
- Tenthredo (Elinora) pubescens** (André, 1881) [*Allantus*]. Russia: **EP** (NC).
- Tenthredo (Elinora) radoszkowskii** (André, 1881) [*Allantus*] (*Allantus atratus* André, 1883; *A. confinis* Konow, 1886; *A. parviceps* Konow, 1898). Host plant: *Geranium* spp. (Geraniaceae). Russia: **EP** (S, NC). – Caucasus, Turkey, Iran.
- Tenthredo (Elinora) sabariensis** (Mocsáry, 1880) [*Allantus*] (*Allantus breviventris* Cameron, 1887; *A. morawitzi* Jakovlev, 1888; *A. sabariensis moravica* Gregor, 1941). Russia: **EP** (C, E, NC), **UR**. – Europe (EE).
- Tenthredo (Endotethryx) adusta** Motschulsky, 1866 (*T. fulva* Klug, 1817; *T. eversmanni* Ballion, 1869; *T. erratica* F. Smith, 1874; *T. fulva pallidiventris* Jakovlev, 1886; *Allantus moiwasamus* Matsumura, 1912; *Tenthredo urgaensis* Muche, 1965). Imago on *Heraclium* spp. and others Apiaceae. Russia: **EP** (C, E), **UR**, **WS** (NS, AL), **ES** (IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU). – Europe (EE, NE), Mongolia, China (NE), Korean Peninsula, Japan (Hok, Hon, Shi).
- Tenthredo (Endotethryx) campestris** Linnaeus, 1758 (*T. flava* Poda, 1761; *T. poecilechroa* Schrank, 1781; *T. flauicornis* Fabricius, 1781; *T. luteicornis* Fabricius, 1787; *T. mucronata* Gmelin, 1790; *T. flavida* Christ, 1791; *T. umbellatarum* Panzer, 1806; *Tenthredella flavigornis temporalis* Enslin, 1912; *T. flavigornis pallior* Forsius, 1918). Host plant: *Aegopodium* spp. (Apiaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (NS, AL). – Europe (WE, EE, NE), Turkey.
- Tenthredo (Endotethryx) crassa** Scopoli, 1763 (*T. impura* Scopoli, 1763; *T. albicornis* Fabricius, 1781; *T. dealbata* Gmelin, 1790; *T. albata* Turton, 1802; *T. nebulosa* Serville, 1823; *T. nebulosa* Lepeletier, 1823). Host plant: ?*Archangelica* spp. (Apiaceae). Russia: **EP** (NW, C, E), **UR**, **WS** (NS), **ES** (KR). – Europe (WE, EE, NE, SE), Georgia, Kazakhstan.
- Tenthredo (Eurogaster) aaliensis** (Strand, 1898) [Pachyprotasis] (*Rhogogaster arctica* Kiær, 1898; *Tenthredo microps* Konow, 1903; *Rhogogaster kaszabi* Zombori, 1973). Imago on Apiaceae. Russia: **UR**, **ES** (IR, BR, ZB), **FE** (AM, MG). – Europe (WE, NE), Kazakhstan, Mongolia.
- Tenthredo (Eurogaster) convergenata** (Takeuchi, 1955) [Tenthredella]. Russia: **FE** (KU). – Japan (Hok, Hon, Shi).
- Tenthredo (Eurogaster) dealbata** Mocsáry, 1909 (*Tenthredella deaurata* Enslin, 1912). Russia: **WS** (AL), **ES** (IR), **FE** (AM, KH, PR). – Korean Peninsula.
- Tenthredo (Eurogaster) maculiger dioctrioides** (Jakovlev, 1891) [*Allantus*] (*Allantus rupico* Konow, 1908). Imago on Apiaceae, *Ranunculus* spp. (Ranunculaceae). Russia: **FE** (PR). – China (NC, SC), Nepal, India (NE).
- Tenthredo (Eurogaster) maculiger maculiger** (Jakovlev, 1891) [*Allantus*] (*Allantus incinctus* Konow, 1908; *Tenthredo maculigera camtschatica* Forsius, 1928). Imago on Apiaceae, *Ranunculus* spp. (Ranunculaceae). Russia: **WS** (AL), **ES** (TU, IR, BR), **FE** (KA, MG). – Mongolia, China (SW).
- Tenthredo (Eurogaster) mesomela** Linnaeus, 1758 (*T. annularis* Villers, 1789; *T. marginata* Christ, 1791; *T. explanata* Rudow, 1871; *T. bernardi* Konow, 1888; *Tenthredella mesomelas mesopleuralis* Enslin, 1912; *T. mesomelas atramentaria* Enslin, 1912; *Tenthredo mesomelas gigas* Malaise, 1931; *Tenthredella mesomelas reducta* Pic, 1940; *T. mesomelas flavoscutellaris* Pic, 1940; *T. mesomelas mediorufescens* Pic, 1940; *Tenthredo kaszabi* Muche, 1965). Host plant: *Polygonum* spp. (Polyfonaceae), *Heracleum* spp. (Apiaceae), *Ranunculus* spp. (Ranunculaceae), etc. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TK, NS, KM, AL), **ES** (KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KA, MG). – Europe (WE, EE, NE, SE), Armenia, Turkey, Mongolia, China (NE), Korean Peninsula, Japan (Hok, Hon, Shi).
- Tenthredo (Eurogaster) mioceras** (Enslin, 1912) [Tenthredella] (*T. mioceras* Benson, 1943; *T. mesomelas montana* Pasteels, 1946). Host plant: *Ranunculus* spp. (Ranunculaceae), *Heracleum* spp. (Asteraceae), *Dryopteris* spp. (Dryopteridaceae), etc. Russia: **EP** (NW, C, E), **UR**, **WS** (NS, AL), **ES** (BR). – Europe (WE, EE, SE), Kazakhstan, China (Wei et al., 2006).
- Tenthredo (Eurogaster) nitidiceps** (Takeuchi, 1955) [Tenthredella]. Russia: **FE** (KU). – Japan (Hok, Hon).
- Tenthredo (Eurogaster) obsoleta** Klug, 1817 (*T. arctica* Thomson, 1871). Host plant: *Plantago* spp. (Plantaginaceae). Russia: **EP** (C, E), **UR**, **WS** (KM, AL), **ES** (KR, IR, BR), **FE** (AM, PR). – Europe (WE, EE, NE, SE), China (Wei et al., 2006).
- Tenthredo (Eurogaster) opaciceps** (Takeuchi, 1955) [Tenthredella]. Russia: **FE** (SA, KU). – Japan (Hok, Hon).

Tenthredo (Eurogaster) sibiricola (Forsius, 1918)
[Tenthredella]. Russia: **FE** (KA).

Tenthredo (Eurogaster) stulta Jakovlev, 1891 (*Rhogogaster opacella* Mocsáry, 1909; *Tenthredella stulta bernardiformis* Enslin, 1927; *T. stulta temporalis* Enslin, 1927; *Tenthredo elisabethae* Muche, 1965). Host plant: probably polyphage. Russia: **WS** (AL), **ES** (IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KA, MG). – Mongolia, China (NC), Korean Peninsula, Japan (Hok, Hon, Kyu).

Tenthredo (Eurogaster) tenuivaginata (Takeuchi, 1955) [Tenthredella]. Russia: **FE** (SA, KU). – Japan (Hok).

Tenthredo (Maculedo) cinctipleuris (Enslin, 1910) [Allantus]. Russia: **EP** (NC). – Georgia, Armenia, Turkey, Iran.

Tenthredo (Maculedo) maculata Geoffroy, 1785 (*T. unifasciata* Geoffroy, 1785; *T. zonata* Panzer, 1799; *T. succincta* Donovan, 1808; *T. equestris* Panzer, 1809; *T. latizona* Lepeletier, 1823; *T. latizona* Serville, 1823; *T. semseyi* Mocsáry, 1883; *T. andrei* Mocsáry, 1883; *Tenthredella maculata atricollaris* Enslin, 1912; *Tenthredo maculata mediobinotata* Pic, 1926; *T. maculata atroapicalis* Pic, 1926; *T. maculata quittardi* Pic, 1926; *T. maculata wagneri* Zirngiebl, 1940; *T. maculata collaris* Zirngiebl, 1940). Host plant: *Brachypodium* spp., *Dactylis* spp. (Poaceae). Russia: **EP** (C, S). – Europe (WE, EE, SE), Armenia, Turkey.

Tenthredo (Maculedo) trabeata Klug, 1817 (*T. palustris* Klug, 1817; *Perineura fulvitarsis* André, 1881; *Tenthredo atra nigrifemur* Zilahi Kiss, 1927; *T. marinovi* Vassilev, 1978). Host plant: Asteraceae, *Impatiens* spp. (Balsaminaceae). Russia: **EP** (N, C). – Europe (WE, EE, NE, SE).

Tenthredo (Maculedo) vestita André, 1881 (*T. caspica* Mocsáry, 1883; *T. laeta* Konow, 1886; *Allantus limbiferus* Mocsáry, 1891; *Tenthredella celsia* Enslin, 1912; *Tenthredo vestita stria* Enslin, 1920; *T. vestita striata* Enslin, 1920; *T. vestita strigata* Enslin, 1920). Russia: **EP** (NC). – Azerbaijan.

Tenthredo (Olivacedo) eduardi (Forsius, 1919) [Tenthredella] (*Tenthredella enslini* Forsius, 1918; *T. viridans* Benson, 1930; *Tenthredo vivida* Malaise, 1931; *T. pappi* Zombori, 1978). Russia: **FE** (AM, PR). – China (Wei et al., 2006), Korean Peninsula, Japan (Hon, Kyu).

Tenthredo (Olivacedo) fuscoterminata Marlatt, 1898. Russia: **FE** (AM, KH, PR). – China (NE, SE), Korean Peninsula, Japan (Hon, Shi, Kyu).

Tenthredo (Olivacedo) nigrolateralis Malaise, 1931. Russia: **FE** (PR).

Tenthredo (Olivacedo) olivacea olivacea Klug, 1817 (*T. dissimilans* Kincaid, 1900; *Tenthredella olivacea nigrovittata* Enslin, 1912; *Rhogogastera reliqua* MacGillivray, 1919; *Rh. respersus* MacGillivray, 1923; *Rhogogaster viridis montana* Betrem, 1933; *Tenthredo olivacea sekeraei* Gregor, 1941; *T. olivacea bohemica* Gregor, 1941). Host plant: *Ranunculus* spp. (Ranunculaceae), *Plantago* spp. (Plantaginaceae), *Rubus* spp. (Rosaceae), and others. Russia: **EP** (N, NW, C, E), **UR**, **WS** (NS, AL), **ES** (IR, BR, YA, ZB), **FE** (KA, MG). – Europe (WE, EE, NE), N Africa, Mongolia, China (NC), Korean Peninsula, N America.

Tenthredo (Olivacedo) olivacea takedae Matsumura, 1912. Host plant: *Ranunculus* spp. (Ranunculaceae), *Plantago* spp. (Plantaginaceae). Russia: **FE** (AM, PR, SA, KU, KA). – Korean Peninsula, Japan (Hok, Hon).

Tenthredo (Olivacedo) sedankiana Malaise, 1931. Russia: **FE** (PR). – China (NE).

Tenthredo (Olivacedo) subolivacea (Takeuchi, 1955) [Tenthredella]. Russia: **FE** (SA, KU). – Japan (Hok, Hon).

Tenthredo (Olivacedo) tamanukii (Takeuchi, 1936) [Tenthredella]. Russia: **FE** (SA, KU). – Japan (Hok, Hon).

Tenthredo (Olivacedo) viridatrix nippon Shino-hara, 1994. Russia: **FE** (SA, KU). – Japan (Hok, Hon, Shi).

Tenthredo (Olivacedo) viridatrix viridatrix Malaise, 1931. Russia: **FE** (AM, KH, PR). – Korean Peninsula.

Tenthredo (Paratenthredo) monozonus (Kriechbaumer, 1869) [Allantus] (*Allantus dialeucus* Konow, 1898; *A. jugalis* Konow, 1898; *Tenthredo unicincta* Hellén, 1963). Russia: **EP** (NC, CR). – Europe (EE, SE), Caucasus, SW Asia.

Tenthredo (Paratenthredo) shaposhnikovi (Dovnar-Zapolskij, 1930) [Allantus]. Russia: **EP** (NC).

Tenthredo (Paratenthredo) talyshensis Zhelochovtsev, 1988. Russia: **EP** (NC). – Azerbaijan.

Tenthredo (Propodea) fentoni W.F. Kirby, 1882 (*Jermakia spinifera* Mocsáry, 1909; *Tenthredopsis hakiensis* Matsumura, 1912). Host plant: *Euptelea polyandra* (Eupteleaceae). Russia: **FE** (KH, KU). – China (NE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

- Tenthredo (Propodea) rufonotalis** Mallach, 1936
(Dipteromorpha ussuriensis Malaise, 1931). Russia: **FE** (PR). – China (NE, CC, SE), Japan (Hon).
- Tenthredo (Temuledo) brachycera** (Mocsáry, 1909)
 [Allantus] (*Allantus brachycerus guttator* Malaise, 1934; *Tenthredo daetekensis* Togashi, 1976). Russia: **FE** (AM, KH, PR). – China (NE, CC, SE), Korean Peninsula, Japan (Hon).
- Tenthredo (Temuledo) finschi finschi** W.F. Kirby, 1882 (*Allantus kohli* Konow, 1895; *Rhogogaster aerescens* Mocsáry, 1909). Imago on Apiaceae, Spiraea spp. (Rosaceae). Russia: **WS** (KM, AL), **ES** (IR, BR, YA), **FE** (AM, KH, PR, KA, MG). – China (NC, SW), Korean Peninsula, India, SE Asia.
- Tenthredo (Temuledo) finschi seguro** Takeuchi, 1956.
 Imago on Apiaceae, Spiraea spp. (Rosaceae). Russia: **FE** (SA, KU). – Japan (Hok, Hon, Shi, Sado).
- Tenthredo (Temuledo) flavipectus** (Matsumura, 1912) [Allantus] (*Allantus nigripectus* Matsumura, 1912; *Tenthredo xanthomelaena* Enslin, 1920). Russia: **FE** (KU). – Japan (Hok, Hon, Shi, Kyu).
- Tenthredo (Temuledo) hokkaidonis** (Malaise, 1931) [Allantus]. Russia: **FE** (KU). – Japan (Hok, Hon, Shi).
- Tenthredo (Temuledo) japonica** (Mocsáry, 1909)
 [Allantus] (*T. carpinata* Enslin, 1912). Russia: **FE** (KU). – China (Wei et al., 2006), Japan (Hon, Shi, Kyu).
- Tenthredo (Temuledo) kongauana** Rohwer, 1925
 (*T. uchidae* Takeuchi, 1927). Russia: **FE** (PR). – China (Wei et al., 2006), Korean Peninsula, Japan (Hon).
- Tenthredo (Temuledo) kudiana** Rohwer, 1925. Russia: **FE** (AM, PR).
- Tenthredo (Temuledo) nigropicta** (F. Smith, 1874) [Macrophyia] (*T. picta* Motschulsky, 1866; *Rhogogaster nipponica* Rohwer, 1910). Host plant: Pleioblastus spp. (Poaceae). Russia: **FE** (KU). – China (CC, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Tenthredo (Temuledo) notomelas** Enslin, 1920. Russia: **FE** (SA). – Japan (Hok, Hon).
- Tenthredo (Temuledo) ouralensis** (André, 1881)
 [Allantus] (*Allantus uralensis* Dalla Torre, 1894; *A. glesicornis* Konow, 1899; *Tenthredo soyoensis* Togashi, 1973). Russia: **ES** (IR, BR), **FE** (AM, PR). – Turkey, Korean Peninsula.
- Tenthredo (Temuledo) sapporensis** (Matsumura, 1912) [Allantus] (*Allantus leucosternus* Malaise, 1934). Russia: **FE** (SA, KU). – China (NE, NC), Japan (Hok).

- Tenthredo (Temuledo) scutellata** (Mocsáry, 1909)
 [Allantus] (*Allantus temuloides* Zhelochovtsev, 1935). Russia: **WS** (NS), **FE** (AM, KH, PR).
- Tenthredo (Temuledo) temula** Scopoli, 1763 (*T. bicincta* Linnaeus, 1767; *T. semicincta* Schrank, 1781; *T. amoena* Marquet, 1879; *Allantus irkutensis* Jakovlev, 1888; *Tenthredella temula uberior* Enslin, 1912; *T. temula pauperior* Enslin, 1914; *Tenthredo temula xanthaspis* Enslin, 1920; *T. celtica* Benson, 1953). Host plant: *Origanum* spp. (Lamiaceae), *Ligustrum* spp. (Oleaceae). Russia: **EP** (NW, C, E, S, NC), **UR**, **WS** (NS, AL), **ES** (IR, BR), **FE** (AM, PR). – Europe (WE, EE, NE, SE), Georgia, Armenia, Kazakhstan.
- Tenthredo (Temuledo) ussuriensis ussuriensis** (Mocsáry, 1909) [Allantus] (*Allantus andis* André, 1881; *Tenthredo abjecta* Enslin, 1912). Host plant: *Abelia* spp. (Caprifoliaceae). Russia: **FE** (AM, KH, PR). – China (NE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Tenthredo (Tenthredella) alboannulata** (Takeuchi, 1933) [Tenthredella] (*Tenthredella alboannulata kanoi* Takeuchi, 1936). Russia: **FE** (KU). – Japan (Hon, Shi).
- Tenthredo (Tenthredella) albopicta** Puls, 1870 (*T. nigroclypeata* Mocsáry, 1909). Russia: **EP** (NC). – Georgia, Armenia, SW Asia.
- Tenthredo (Tenthredella) amurica** Dalla Torre, 1894 (*T. amurensis* Konow, 1891; *Tenthredella ussuriensis* Forsius, 1918; *T. fagi nigerrima* Forsius, 1918; *Tenthredo konowi* Enslin, 1920). Host plant: *Petasites japonicus* (Asteraceae). Russia: **EP** (NW), **ES** (BR), **FE** (AM, PR, SA). – Europe (WE, NE).
- Tenthredo (Tenthredella) atra** Linnaeus, 1758 (*T. rufipes* Poda, 1761; *T. fuscipes* Gmelin, 1790; *T. obscura* Gmelin, 1790; *T. plebeja* Klug, 1817; *T. dispar* Klug, 1817; *T. relicta* Lepeletier, 1823; *T. scopolii* Lepeletier, 1823; *T. rejecta* Dalla Torre, 1882; *T. scotica* Cameron, 1882; *Tenthredella atra pseudomandibularis* Enslin, 1918; *T. atra pseudoscotica* Enslin, 1918; *T. atra schirmeri* Enslin, 1918; *T. atra orbitalis* Forsius, 1918; *Tenthredo atra transigens* Enslin, 1920; *T. atra nigrifemur* Endre, 1927; *T. atra vagabundica* Malaise, 1931; *T. atra masculina* Gregor, 1941). Host plant: *Mentha* spp. (Lamiaceae), *Plantago* spp. (Plantaginaceae), *Ranunculus* spp. (Ranunculaceae), etc. Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (NS, AL), **ES** (IR, BR, YA), **FE** (KH, PR, KA, MG). – Europe (WE, EE, NE, SE), Kyrgyzstan, Kazakhstan, Mongolia.
- Tenthredo (Tenthredella) atrofasciata** Zhelochovtsev, 1939. Russia: **FE** (AM).

- Tenthredo (Tenthredella) balteata** Klug, 1817
(T. soror Zetterstedt, 1838; *Tenthredella balteata albimacula* Enslin, 1912; *Allantus surensis* Pic, 1939; *Tenthredo tschinggischanensis* Muche, 1965). Host plant: Hypericaceae, Rosaceae, Salicaceae, ferns and others. Russia: **EP** (N, NC), **UR**, **WS** (AL), **ES** (TU, KR, IR, BR), **FE** (KA, MG). – Europe (WE, EE, NE, SE), N Africa, Georgia, Armenia, SW Asia, Iran, Central Asia, Kazakhstan, Mongolia.
- Tenthredo (Tenthredella) basizonata** Malaise, 1938.
 Russia: **FE** (KU). – Japan (Hon, Shi, Kyu).
- Tenthredo (Tenthredella) bipunctula bipunctula** Klug, 1817. Host plant: *Senecio* spp. (Asteraceae). Russia: **EP** (NW), **WS** (NS). – Europe (WE, EE, NE, SE).
- Tenthredo (Tenthredella) bipunctula malaisei** (Takeuchi, 1933) [Tenthredella]. Host plant: *Senecio* spp. (Asteraceae). Russia: **FE** (SA, KU). – Japan (Hon).
- Tenthredo (Tenthredella) borea** Enslin, 1919 (*T. borea* Enslin, 1920; *T. stubbei* Koch, 1984). Russia: **EP** (N), **WS** (TM). – Mongolia.
- Tenthredo (Tenthredella) camtschatalis** (Enslin, 1927) [Tenthredella]. Russia: **FE** (KA).
- Tenthredo (Tenthredella) cockerelli** (Rohwer, 1925) [Tenthredella] (*Tenthredella bituberculata* Takeuchi, 1940). Russia: **FE** (AM, PR). – China (NE, NC, CC), Korean Peninsula.
- Tenthredo (Tenthredella) colon** Klug, 1817 (*Allantus obscurus* Panzer, 1803; *Tenthredo rudowi* André, 1881; *T. nigricollis* W.F. Kirby, 1882; *T. punctulata* Konow, 1887; *T. semicornis* Harrington, 1889; *T. unifasciata* Mocsáry, 1909; *T. colon nigriventris* Enslin, 1912; *Tenthredella cucullata* Enslin, 1912). Host plant: Dennstaedtiaceae, Ranunculaceae, Geraniaceae, Onagraceae, Salicaceae, etc. Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (KM, AL), **ES** (IR, BR, YA), **FE** (AM, KH, PR, SA, KU, KA, MG). – Europe (WE, EE, NE, SE), Caucasus, SW Asia, Kyrgyzstan, Kazakhstan, Mongolia, Korean Peninsula, Japan (Hok, Hon, Sado), N America.
- Tenthredo (Tenthredella) contusa** (Enslin, 1912) [Tenthredella] (*T. lateralis* Mocsáry, 1909; *T. trialbata* Malaise, 1931). Russia: **FE** (AM, KH, PR). – Japan (Hon, Shi), SE Asia.
- Tenthredo (Tenthredella) crenata** (Enslin, 1912) [Tenthredella] (*T. fulvicornis* Mocsáry, 1909; *T. fulviventralis* Mocsáry, 1909). Russia: **ES** (BR), **FE** (AM, KH, SA). – Mongolia.
- Tenthredo (Tenthredella) cunyi** Konow, 1886 (*T. chyzeri* Mocsáry, 1891; *Tenthredella cunyi pleuritica*

- Enslin, 1912; *T. cunyi alboplagiata* Enslin, 1912). Russia: **ES** (IR, BR). – Europe (WE, EE).
- Tenthredo (Tenthredella) decens** Zhelochovtsev, 1939. Russia: **EP** (NW), **ES** (IR, BR, ZB), **FE** (KA).
- Tenthredo (Tenthredella) duplicata** Enslin, 1920 (*T. bimaculata* Konow, 1886). Russia: **EP** (NC). – Georgia, Armenia.
- Tenthredo (Tenthredella) eburata** Konow, 1900 (*Tenthredella rubrocaudata* Takeuchi, 1936). Russia: **EP** (NW, C, E), **UR**, **WS** (NS), **ES** (IR, BR), **FE** (PR). – Europe (NE), China (NE, CC), Korean Peninsula.
- Tenthredo (Tenthredella) eburneifrons** W.F. Kirby, 1882 (*T. lichtwardti* Konow, 1899). Russia: **WS** (AL), **ES** (BR, ZB), **FE** (AM, PR). – Europe (WE, NE), Kazakhstan, Korean Peninsula.
- Tenthredo (Tenthredella) erudita** (Kuznetzov-Ugamskij, 1927) [Tenthredella]. Russia: **FE** (PR).
- Tenthredo (Tenthredella) fagi** Panzer, 1798 (*T. pellicula* Klug, 1817; *T. facigera* Konow, 1899; *Tenthredella fagi melanaspis* Enslin, 1912). Host plant: *Pteridium aquilinum* (Dennstaedtiaceae), *Corylus* spp. (Betulaceae), *Sorbus* spp. (Rosaceae), *Angelica* spp. (Asteraceae). Russia: **EP** (NW, C, E), **UR**, **WS** (NS, AL), **ES** (IR, BR, YA), **FE** (AM, PR, SA, KU). – Europe (WE, EE, NE), Armenia, China (Wei et al., 2006), Korean Peninsula, Japan (Hok, Hon).
- Tenthredo (Tenthredella) ferruginea** Schrank, 1776 (*T. ferruginea* O.F. Müller, 1776; *T. rufipennis* Fabricius, 1793; *T. rufiventris* Panzer, 1799; *T. conspicua* Klug, 1817; *T. schrankii* Lepeletier, 1823; *Allantus laticinctus* Stephens, 1835; *Tenthredo rubecula* Eversmann, 1847; *T. prospera* Erichson, 1851; *T. gymandromorpha* Rudow, 1871; *T. leucocera* Mocsáry, 1909; *T. minuta* Mocsáry, 1909; *Tenthredella dentina* Enslin, 1912; *T. ferruginea leucaspis* Enslin, 1912; *Tenthredo ferruginea rufisternis* Enslin, 1920; *Tenthredella ferruginea frontalis* Pic, 1940; *Perineura sawai* Togashi, 2007). Host plant: Salicaceae, Betulaceae, Rosaceae, ferns, etc. Russia: **EP** (NW, C, E, NC), **UR**, **WS** (AL), **ES** (BR, YA), **FE** (AM, KH, PR, SA, KU, KA, MG). – Europe (WE, EE, NE), Armenia, Turkey, China (NE, CC, SE), Korean Peninsula, Japan (Hok, Hon, Sado), SE Asia.
- Tenthredo (Tenthredella) gifui** Marlatt, 1898 (*T. gifuensis* Konow, 1905; *T. filamentosa* Enslin, 1920). Host plant: *Stellaria* spp. (Caryophyllaceae). Russia: **FE** (SA). – Japan (Hok, Hon, Shi, Kyu).
- Tenthredo (Tenthredella) kurilensis** (Takeuchi, 1931) [Tenthredella]. Russia: **WS** (AL), **FE** (KU). – Japan (Hok, Hon).

- Tenthredo (Tenthredella) leleji** Taeger, Wei et Shino-hara, 2016. Russia: FE (SA). – Japan (Hok, Hon).
- Tenthredo (Tenthredella) livida** Linnaeus, 1758 (*T. dubia* Strøm, 1768; *T. lurida* Ph.L. Müller, 1775; *T. pellucida* O.F. Müller, 1776; *T. bipunctata* O.F. Müller, 1776; *T. annularis* Schrank, 1776; *T. albicornis* Geoffroy, 1785; *T. maura* Fabricius, 1793; *T. carpini* Panzer, 1799; *Allantus aterrimus* Stephens, 1835; *Tenthredella livida clara* Enslin, 1912; *T. livida rubripes* Forsius, 1918; *T. livida rubeola* Forsius, 1918). Host plant: Oleaceae, Lamiaceae, Salicaceae, Caprifoliaceae, Onagraceae, Rosaceae, Betulaceae, Plantaginaceae, etc. Russia: EP (NW, C, E, S, NC), UR, WS (Zhelochovtsev, 1988), ES (KR), FE (AM, KA). – Europe (WE, EE, NE, SE), Georgia, Armenia, Turkey.
- Tenthredo (Tenthredella) luteipennis** Eversmann, 1847 (*T. fallax* Mocsáry, 1880; *T. mocsaryi* André, 1882; *T. menetriesii* W.F. Kirby, 1882; *Tenthredella luteipennis obscuraria* Enslin, 1913). Russia: EP (S, NC). – Georgia, Armenia, SW Asia, Tadzhikistan.
- Tenthredo (Tenthredella) mandibularis** Fabricius, 1804 (*T. rufipes* Gmelin, 1790). Host plant: *Tussilago farfara*, *Petasites* spp. (Asteraceae). Russia: EP (C, S), WS (NS). – Europe (WE, EE, NE, SE), Caucasus.
- Tenthredo (Tenthredella) moniliata** Klug, 1817 (*T. fuscicornis* Eschscholtz, 1822; *Allantus bimaculatus* Gimmerthal, 1836; *Tenthredo poecila* Eversmann, 1847; *T. flavidabris* Gimmerthal, 1847; *T. scita* Erichson, 1851; *T. lachlaniana* Cameron, 1878; *T. poecilopus* Mocsáry, 1880; *T. lachlaniana rubripes* Jakovlev, 1888; *T. lachlaniana immaculata* Jakovlev, 1888; *T. ruthena* Jakovlev, 1891; *T. forsi* Konow, 1908; *T. ruficornis* Mocsáry, 1909; *Tenthredella fuscicornis norvegica* Forsius, 1918; *T. moniliata flavoscutellata* Forsius, 1918; *Tenthredo moniliata immaculosa* Enslin, 1920; *T. moniliata paria* Enslin, 1920; *T. moniliata menyanthidis* Torka, 1929). Host plant: Menyanthaceae, Ranunculaceae, Asteraceae, Rosaceae, Betulaceae, etc. Russia: EP (NW, C, E), UR, WS (AL), ES (IR, BR, YA, ZB), FE (AM, KH, PR, SA, KU, KA, MG). – Europe (WE, EE, NE), Armenia, Mongolia, China (NE, NC, NE, SE), Korean Peninsula.
- Tenthredo (Tenthredella) oryssoides** Jakovlev, 1888 (*T. discophora* Konow, 1896; *T. discophora pullata* Konow, 1898; *T. bigemmis* Konow, 1903; *T. discophora rosea* Enslin, 1913). Russia: EP (NC). – Georgia, Armenia, Turkey.
- Tenthredo (Tenthredella) peninsulae** Malaise, 1931. Russia: ES (YA), FE (AM, KA).
- Tenthredo (Tenthredella) procera** Klug, 1817 (*T. griseocephala* Costa, 1894). Host plant: *Sympytum* spp. (Boraginaceae), *Petasites* spp. (Asteraceae). Russia: EP (C). – Europe (WE, EE).
- Tenthredo (Tenthredella) purpurea** Puls, 1870 (*T. koenigi* Konow, 1896). Russia: EP (NC). – Georgia, Armenia, SW Asia.
- Tenthredo (Tenthredella) rubricoxis** (Enslin, 1912) [*Tenthredella*] (*T. rufipes* Klug, 1817). Host plant: *Senecio* spp. (Asteraceae). Russia: EP (N), WS (TM, NS, KM), FE (PR, MG). – Europe (WE, EE).
- Tenthredo (Tenthredella) serradifera** Conde, 1935. Russia: FE (SA).
- Tenthredo (Tenthredella) silensis** Costa, 1859 (*T. limbata* Klug, 1817; *T. quadridens* Strobl, 1896; *T. limbata nigrifemur* Conde, 1934; *T. oraria* Zombori, 1973). Host plant: Lamiaceae, Ranunculaceae, Rosaceae, Liliaceae, etc. Russia: EP (NW, C, E), UR, WS (AL), ES (TU, IR, BR, YA), FE (KH, AM, SA, KU, KA, MG). – Europe (WE, EE, NE), Mongolia, Korean Peninsula, Japan (Hok, Hon, Sado).
- Tenthredo (Tenthredella) sobrina** Eversmann, 1847 (*Allantus semenowi* Jakovlev, 1891; *Tenthredo semenovi pleuronotata* Forsius, 1934). Russia: EP (C, E, S, NC), UR. – Europe (EE), Kazakhstan.
- Tenthredo (Tenthredella) solitaria solitaria** Scopoli, 1763 (*T. coryli* Panzer, 1799; *T. intermedia* Klug, 1817; *T. coryli albicolon* Klug, 1817; *T. icterica* Costa, 1894; *T. solitaria seeboldi* Konow, 1897; *Tenthredella solitaria rufoterminalata* Enslin, 1912; *T. solitaria rheensis* Koornneef, 1935; *Tenthredo saxonica* Muche, 1968; *T. solitaria vitoshensis* Vassilev, 1973). Host plant: *Euphorbia* spp. (Euphorbiaceae). Russia: EP (NC, C). – Europe (WE, EE, SE), Armenia.
- Tenthredo (Tenthredella) solitaria xylota** (Jakovlev, 1891) [*Allantus*]. Host plant: *Euphorbia* spp. (Euphorbiaceae). Russia: WS (Zhelochovtsev, Zinovjev, 1996), ES (IR, YA), FE (PR). – Kyrgyzstan.
- Tenthredo (Tenthredella) stigma** (Forsius, 1918) [*Tenthredella*] (*T. amplicapitata* Zombori, 1974). Russia: FE (AM, SA). – Korean Peninsula.
- Tenthredo (Tenthredella) velox nigrofemorata** (Takeuchi, 1955) [*Tenthredella*]. Host plant: *Geranium* spp. (Geraniaceae). Russia: FE (SA, KU). – Korean Peninsula, Japan (Hok, Hon).
- Tenthredo (Tenthredella) velox velox** Fabricius, 1798 (*T. biguttata* Hartig, 1837; *T. leucostoma* Rudeow, 1871; *T. gracilenta* Mocsáry, 1879; *T. sylvia*

- W.F. Kirby, 1882; *T. velox sibirica* Konow, 1903; *T. velox bisignata* Konow, 1908; *Tenthredella velox colonoides* Forsius, 1918; *Tenthredo velox alpina* Zirngiebl, 1937; *T. velox nigripleuris* Zirngiebl, 1937). Host plant: Polygonum spp. (Polygonaceae), Alnus spp. (Betulaceae), Salix spp. (Salicaceae). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (NS, KM, AL), **ES** (IR, BR, YA), **FE** (AM, KH, PR, MG). – Europe (WE, EE, NE, SE), Mongolia.
- Tenthredo (Tenthredella) versuta** Mocsáry, 1909 (*Tenthredella hakonensis* Rohwer, 1910). Russia: **FE** (SA). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Tsushima).
- Tenthredo (Tenthredina) cylindrica** (Rohwer, 1911) [*Tenthredina*] (*Conospidia umbrosa* Matsumura, 1912). Russia: **FE** (PR). – China (SE), Japan (Hok, Hon, Shi, Kyu).
- Tenthredo (Tenthredo) alaskana** (Enslin, 1910) [Allantus] (*Allantus heraclei* Kincaid, 1900). Russia: **ES** (YA). – N America.
- Tenthredo (Tenthredo) albiventris albiventris** (Mocsáry, 1880) [Allantus] (*Allantus trivittatus* André, 1881; *A. kussariensis* Konow, 1888; *A. lederi* Konow, 1890; *Tenthredo albiventris hostia* Enslin, 1912; *T. pallidula* Benson, 1930; *T. kerkesta* Dovnar-Zapolskij, 1930). Imago on Conium spp. (Asteraceae). Russia: **EP** (NC). – Europe (WE, SE), Georgia, Armenia, Turkey.
- Tenthredo (Tenthredo) arctica** (Thomson, 1870) [Eniscia] (*Ischyroceraea hyperborea* Kiær, 1898). Host plant: Trifolium repens (Fabaceae); imago on Apiaceae and others. Russia: **EP** (N), **WS** (TM), **ES** (KR, BR, YA), **FE** (CH). – Europe (NE), N America.
- Tenthredo (Tenthredo) arcuata** Förster, 1771 (*Allantus arcuatus alpigena* Heller et Dalla Torre, 1883; *A. clypealis* Konow, 1888). Host plant: Trifolium spp. (Fabaceae). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (TK, NS), **ES** (KR, YA, ZB). – Europe (WE, EE, NE, SE), Armenia, Turkey, Iran, Mongolia, China (Wei et al., 2006).
- Tenthredo (Tenthredo) arcuatoides** Muche, 1965. Russia: **ES** (IR, BR, YA, ZB), **FE** (AM, KH, PR).
- Tenthredo (Tenthredo) brevicornis** (Konow, 1886) [Allantus] (*T. cingulata* Fabricius, 1793; *Allantus arcuatus nitidior* Konow, 1888; *Tenthredo aegra* Enslin, 1912; *Allantus arcuatus vitoensis* Gregor, 1932; *Tenthredo acerrima* Benson, 1952). Host plant: Lotus corniculatus (Fabaceae). Russia: **EP** (C, NC), **ES** (IR, BR), **FE** (PR, KA). – Europe (WE, EE, SE), Georgia, Armenia, Turkey, Mongolia, China (NC).
- Tenthredo (Tenthredo) calvaria** Enslin, 1912 (*Allantus lateralis* Mocsáry, 1909). Russia: **WS** (NS), **ES** (BR, ZB), **FE** (AM, KH, PR). – Mongolia.
- Tenthredo (Tenthredo) devia** (Konow, 1900) [Allantus]. Host plant: polyphage. Russia: **EP** (N), **WS** (TM, AL), **ES** (TU, KR, BR, YA, ZB), **FE** (KU, KA, MG). – Europe (NE), Mongolia, Korean Peninsula, Japan (Hok, Hon), N America.
- Tenthredo (Tenthredo) erasina** Malaise, 1945 (*T. flavipleuris* Muche, 1986). Russia: **ES** (ZB), **FE** (AM, PR). – Mongolia, China (NE).
- Tenthredo (Tenthredo) fuscocostalis** Rohwer, 1925. Russia: **FE** (AM, KH, PR, SA). – Korean Peninsula.
- Tenthredo (Tenthredo) jacutensis** (Konow, 1897) [Allantus]. Imago on various flowers. Russia: **EP** (N), **WS** (TM), **ES** (KR, BR, YA, ZB), **FE** (AM, KH, PR, SA, MG). – Mongolia, Japan (Hok).
- Tenthredo (Tenthredo) korabica** Taeger, 1985. Host plant: polyphage. Russia: **EP** (NC), “Siberia” (Lacourt, 1999). – Europe (WE, EE), Turkey, Mongolia.
- Tenthredo (Tenthredo) liturata** (Mocsáry, 1886) [Allantus] (*Allantus reitteri* Konow, 1886; *Tenthredo reitteri* egredia Enslin, 1914). Russia: **EP** (NC). – Georgia, Armenia, Turkey.
- Tenthredo (Tenthredo) luteocincta** Eversmann, 1847 (*T. luteocincta virago* Enslin, 1913). Russia: **EP** (S). – Georgia, Armenia, Turkey.
- Tenthredo (Tenthredo) marginella fuscipennis** (Konow, 1886) [Allantus] (*T. marginella nigrior* Enslin, 1912). Russia: **EP** (NC). – Armenia.
- Tenthredo (Tenthredo) marginella marginella** Fabricius, 1793 (*T. sigma* Schrank, 1802; *T. succincta* Lepeletier, 1823; *T. succincta* Serville, 1823; *Allantus decipiens* Förster, 1844; *A. heraclei* Rudow, 1872; *A. omissoides* Jakovlev, 1891; *A. amaeonus sabandus* Pic, 1925; *A. marginellus nigroscutellatus* Zirngiebl, 1942; *A. marginellus melanomerus* Zirngiebl, 1942). Host plant: Mentha spp. (Lamiaceae), Plantago spp. (Plantaginaceae), etc. Russia: **EP** (C, E), **UR**, **WS** (TK, NS), **ES** (KR). – Europe (WE, EE, NE, SE), Kyrgyzstan, Kazakhstan.
- Tenthredo (Tenthredo) mongolica** (Jakovlev, 1891) [Allantus] (*T. coreana* Takeuchi, 1927; *T. coreana nigripes* Takeuchi, 1927; *T. erasa* Malaise, 1945). Russia: **WS** (AL), **ES** (TU, BR, ZB), **FE** (AM, KH, PR). – Mongolia, China (NE, NC, CC), Korean Peninsula.
- Tenthredo (Tenthredo) nigripleuris** (Enslin, 1910) [Allantus] (*T. nivalis* Lacourt, 1985). Russia: **EP**

- (N), WS (TM), ES (YA), FE (KA). – Europe (WE, EE), Japan (Hok, Hon).
- Tenthredo (Tenthredo) notha notha** Klug, 1817 [*Allantus perkinsi* Morice, 1919; *A. arcuatus niger* Zirngiebl, 1937]. Host plant: *Trifolium* spp., *Vicia* spp. (Fabaceae). Russia: EP (N, C, E), UR, WS (TK, NS, KM, AL), ES (KR, IR, ZB). – Europe (WE, EE, NE, SE), Kazakhstan, China (Wei et al., 2006).
- Tenthredo (Tenthredo) notha titania** Benson, 1959. Host plant: *Trifolium* spp., *Vicia* spp. (Fabaceae). Russia: EP (NC). – Caucasus, Turkey, Iran.
- Tenthredo (Tenthredo) omissa** (Förster, 1844) [*Allantus*] (*T. omissa melanoceraea* Enslin, 1912; *T. omissa melanomeros* Enslin, 1912; *T. omissa schirmeri* Enslin, 1918). Host plant: *Plantago* spp. (Plantaginaceae). Russia: EP (NW, C, E, S), WS (TM, OM, TK, NS, KM), ES (KR, IR), FE (AM, PR). – Europe (WE, EE, NE, SE), Turkey, Kazakhstan.
- Tenthredo (Tenthredo) pamyreensis** Jakovlev, 1888. Russia: ES (KS). – Tadzhikistan, Kyrgyzstan, Kazakhstan.
- Tenthredo (Tenthredo) propinquata** Klug, 1817. Russia: EP (CR). – Europe (WE, EE, SE), Caucasus, Turkey.
- Tenthredo (Tenthredo) schaefferi** Klug, 1817 (*Allanthus costalis* Costa, 1858). Host plant: *Trifolium* spp., ?*Vicia* spp. (Fabaceae). Russia: EP (E), UR, WS (TK, NS, AL), ES (BR, ZB), FE (MG). – Europe (WE, EE, NE, SE).
- Tenthredo (Tenthredo) scrophulariae** Linnaeus, 1758 (*Allantus scrophulariae joannis* Pic, 1926; *A. scrophulariae repartitus* Pic, 1926; *A. scrophulariae morvandicus* Pic, 1926; *A. scrophulariae branensis* Pic, 1926). Host plant: *Scrophularia* spp., *Verbascum* spp. (Scrophulariaceae). Russia: EP (NW, C, E, NC). – Europe (WE, EE, NE, SE), Armenia, Turkey.
- Tenthredo (Tenthredo) sulphuripes muchei** Koch, 1986. Russia: ES (IR, BR). – Mongolia, China (SW).
- Tenthredo (Tenthredo) sulphuripes sulphuripes** (Kriechbaumer, 1869) [*Allantus*] (*T. beaumonti* Benson, 1950). Host plant: *Bupleurum* spp. (Asteraceae). Russia: EP (NC). – Europe (WE, EE, SE).
- Tenthredo (Tenthredo) vespa** Retzius, 1783 (*T. viminalis* Schrank, 1796; *T. sexannulata* Schrank, 1802; *T. vespiformis* Latreille, 1804; *T. 3 cincta* Fabricius, 1804; *T. 4-cincta* Fallén, 1808; *Allantus multifasciatus* Rudow, 1872; *A. multicinctus* Rudow, 1872; *A. nigrilabris* Frivaldszky, 1877; *Tenthredo vespa stigmatica* Enslin, 1912; *T. vespa omnivinccta* Enslin, 1912; *T. vespa vittozi* Pic, 1933; *Allantus vespa mequignoni* Pic, 1940; *A. vespa monetierensis* Pic, 1940; *Tenthredo vespa inaffectata* Muche, 1965). Host plant: *Spiraea* spp. (Rosaceae), *Lonicera* spp. (Caprifoliaceae), *Viburnum* spp. (Adoxaceae), *Syringa* spp. (Oleaceae), etc. Russia: EP (NW, C, E, NC), UR, WS (NS), ES (IR, BR). – Europe (WE, EE, NE, SE), Georgia, Armenia, Turkey, Mongolia, China (Wei et al., 2006).
- Tenthredo (Zonuledo) amoena** Gravenhorst, 1807 (*T. 2 cincta* Fabricius, 1798; *T. bicincta* Panzer, 1804; *T. cingulum* Klug, 1817; *Allantus quinquecinctus* Gimmerthal, 1834; *A. inversus* Costa, 1894; *A. amoenus lateniger* Pic, 1925; *A. amoenus seminiger* Pic, 1925; *A. amoenus breviniger* Pic, 1925; *A. amoenus lateoluteus* Pic, 1925; *A. amoenus digoniensis* Pic, 1927; *A. amoenus mediomaculatus* Pic, 1927; *A. amoenus multiluteus* Pic, 1927; *A. amena robustior* Pic, 1940). Host plant: *Hypericum maculatum*, *H. perforatum* (Hypericaceae). Russia: EP (NW, C, E). – Europe (WE, EE, NE, SE).
- Tenthredo (Zonuledo) flavipennis** Brullé, 1832 (*Allantus frivaldszkyi* Mocsáry, 1879; *A. laetus* Konow, 1891; *A. luminosus* Konow, 1899). Russia: EP (S, NC). – Europe (EE, SE), Caucasus, SW Asia.
- Tenthredo (Zonuledo) zonula** Klug, 1817 (*T. fasciata* Scopoli, 1763; *T. bifasciata* Geoffroy, 1785; *T. bicincta flava* Christ, 1791; *Allantus similis* Mocsáry, 1880; *A. calcaratus* André, 1881; *A. scutellaris* Konow, 1898). Host plant: *Hypericum perforatum* (Hypericaceae). Russia: EP (NW, C, E, S, NC). – Europe (WE, EE, NE, SE), N Africa, Armenia, SW Asia, Iran, Central Asia.
- Tenthredo abdominalis** (Marsumura, 1912) [Tenthredopsis] (*T. sortitor* Malaise, 1938). Russia: FE (SA, KU). – China (SE), Korean Peninsula, Japan (Hok, Hon), India, Sikkim.
- Tenthredo atroclypeata** Enslin, 1920. Russia: ES (IR).
- Tenthredo atra obscurata** Jakovlev, 1888. Russia: ES (IR).
- Tenthredo caligator** Eversmann, 1847 (*T. nigritarsis* Puls, 1870; *T. morawitzi* Jakovlev, 1888). Russia: EP (S, NC). – Georgia, Armenia, SW Asia.
- Tenthredo cingulifer** (Konow, 1899) [*Allantus*] (*Allantus almasyanus* Mocsáry, 1909). Russia: WS (AL), ES (TU). – Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NW).
- Tenthredo emphytiformis** Malaise, 1931. Russia: FE (KH, PR, KU, KA). – China (Wei et al., 2006), Japan (Hok, Hon, Shi), SE Asia.
- Tenthredo erebia** (Kuznetsov-Ugamskij, 1927) [Tenthredella]. Russia: FE (PR).

- Tenthredo flavomandibulata** (Matsumura, 1912) [Tenthredopsis]. Russia: **FE** (SA, KU). – Korean Peninsula, Japan (Hok, Hon).
- Tenthredo gracilis** (Forsius, 1918) [Tenthredella]. Russia: **FE** (PR).
- Tenthredo intercincta** (Malaise, 1934) [Allantus]. Russia: **FE** (PR). – China (NC).
- Tenthredo kotenkoi** Ermolenko, 1992. Russia: **ES** (ZB).
- Tenthredo longipennis** (Matsumura, 1912) [Allantus] (*T. grandiceps* Malaise, 1938). Russia: **FE** (SA, KU). – Japan (Hok, Hon, Shi).
- Tenthredo mitsuhashii** (Matsumura, 1912) [Macrophyia]. Russia: **FE** (SA, KU). – Japan (Hok, Hon).
- Tenthredo omega** (Takeuchi, 1936) [Tenthredella]. Russia: **FE** (SA, KU). – Japan (Hok, Hon, Shi).
- Tenthredo provida** (F. Smith, 1878) [Allantus] (*T. providus concolor* Malaise, 1945; *T. providus fletcheri* Malaise, 1945; *T. providus altaica* Stroganova, 1980). Russia: **WS** (AL). – China (NW), India (NW).
- Tenthredo providens** F. Smith, 1874 (*T. providentia* Enslin, 1920). Host plant: Oenanthe spp., Cryptotaenia spp., Peucedanum spp., Angelica spp. (Apiaceae). Russia: **FE** (KU). – Japan (Hok, Hon, Shi, Kyu, Sado, Awashima).
- Tenthredo pseudolivacea** Malaise, 1931. Russia: **FE** (PR).
- Tenthredo sibirica** (Kriechbaumer, 1869) [Allantus] (*T. spectabilis* Mocsáry, 1878; *Allantus cephalotes* Jakovlev, 1888; *Jermakia japonica* Rohwer, 1910; *Allantus bicinctus* Matsumura, 1912; *A. bistrigatus* Mallach, 1936). Host plant: Dasiphora spp. (Rosaceae), Ranunculus spp. (Ranunculaceae). Russia: **WS** (AL), **ES** (IR, ZB), **FE** (AM, PR, SA, KU). – Mongolia, China (NC, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu), SE Asia.
- Tenthredo tenuipennis** Malaise, 1931. Russia: **FE** (KH, PR). – China (NE, CC), Korean Peninsula.
- Tenthredo tumida tumida** (Mocsáry, 1909) [Allantus] (*Tenthredella excellens* Kuznetzov-Ugamskij, 1927). Russia: **FE** (AM, KH, PR). – China (NE, SE), Korean Peninsula.
- Tenthredo variicolor** Malaise, 1945. Russia: **FE** (PR). – China (SW), India.
- Tenthredo xanthotarsus** Cameron, 1876. Russia: **WS** (AL), **FE** (AM, SA, KU). – Japan (Hok, Hon, Shi).
- Tenthredo yezoensis** Kumamoto, 1987. Russia: **FE** (SA). – Japan (Hok).

TENTHREDOPSIS Costa, 1859 (*Ebolia* Costa, 1859; *Thomsonia* Konow, 1884; *Eutenthredopsis* Enslin,

1913). Type species *Tenthredo tessellata* Klug, 1817. The larvae on Poaceae, less often on Cyperaceae, Juncaceae, Acoraceae. The genus is distributed in Palaearctic, mainly in west part; two species in Oriental region. Number of species: World and Palaearctic – about 60; Russia – 30 (+ 1 ssp.).

- Tenthredopsis annuligera** (Eversmann, 1847) [Tenthredo] (*Tenthredo albopunctata* Tischbein, 1852; *T. picticornis* Mocsáry, 1880; *T. basimacula* Mocsáry, 1880; *Tenthredopsis casia* Konow, 1898; *T. triforis* Konow, 1898; *T. quadrannulata* Konow, 1898). Host plant: Poaceae. Russia: **EP** (S, CR). – Europe (EE, SE).
- Tenthredopsis auriculata** (Thomson, 1870) [Perineura] (*Tenthredo sachalinensis* Matsumura, 1911; *Tenthredopsis camtschaticalis* Enslin, 1927; *T. camtschaticalis lucens* Enslin, 1927; *T. auriculata camtschatica* Forsius, 1928; *T. sachalinensis nigrolineatus* Takeuchi, 1936). Host plant: Calamagrostis spp. (Poaceae). Russia: **EP** (N, NW, E), **UR**, **WS** (TM), **ES** (IR, BR), **FE** (AM, KH, PR, SA, KA). – Europe (NE).
- Tenthredopsis carinata carinata** Malaise, 1931. Host plant: Poaceae. Russia: **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (TU, IR), **FE** (KA). – Europe (NE).
- Tenthredopsis carinata japonica** Takeuchi, 1956. Host plant: Poaceae. Russia: **FE** (SA, KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Tenthredopsis coquebertii** (Klug, 1817) [Tenthredo] (*Tenthredo labiata* Serville, 1823; *T. labiata* Lepeletier, 1823; *T. rufimana* Spinola, 1843; *Perineura lusitanica* André, 1881; *Tenthredopsis nigricollis* Cameron, 1881; *T. conjungens* Kriechbaumer, 1884; *T. limbilabris* Kriechbaumer, 1884; *Thomsonia braunsi* Konow, 1884; *Tenthredopsis opulenta* Konow, 1887; *T. coqueberti brunnescens* Enslin, 1918; *T. coqueberti nigrata* Gregor, 1941). Host plant: Poaceae. Russia: **ES** (BR). – Europe (WE, EE, SE), China (Wei et al., 2006).
- Tenthredopsis discrepans** Konow, 1890. Russia: **EP** (NC). – Georgia, Armenia.
- Tenthredopsis festiva** Konow, 1890. Russia: **EP** (NC). – Armenia, Azerbaijan, Turkey.
- Tenthredopsis friesei** (Konow, 1884) [Thomsonia] (*Thomsonia laticeps* Konow, 1884; *Tenthredopsis korlevici* Konow, 1887; *T. arrogans* Konow, 1890; *T. nigroscutellata* Konow, 1904; *T. arrogans erythrocoele* Enslin, 1913). Host plant: Calamagrostis spp., Holcus spp. (Poaceae), Juncus spp. (Juncaceae).

- Russia: **EP** (NW, C, E, S, NC), **UR**, **WS** (TM, KM), **ES** (IR). – Europe (WE, EE, NE, SE), Kazakhstan.
- Tenthredopsis gracilis** Jakovlev, 1891. Russia: **ES** (KR).
- Tenthredopsis hungarica** (Klug, 1817) [Tenthredo] (*T. sareptana* Konow, 1894; *T. hungarica atratilobis* Enslin, 1913). Host plant: Poaceae. Russia: **EP** (C, E, S, NC), **UR**, **WS** (KM). – Europe (EE), Kazakhstan, Mongolia.
- Tenthredopsis jakowleffi** Konow, 1896. Russia: **ES** (IR, BR).
- Tenthredopsis kokuewi** Jakovlev, 1891. Russia: **EP** (C).
- Tenthredopsis lactiflua** (Klug, 1817) [Tenthredo] (*T. lactiflua plaga* Enslin, 1913; *T. lactiflua liturata* Enslin, 1913). Host plant: Poaceae. Russia: **EP** (S), **WS** (Zhelochovtsev, Zinovjev, 1996), **ES** (TU, IR). – Europe (WE, EE), Georgia, Kazakhstan.
- Tenthredopsis languida** (Erichson, 1851) [Tenthredo]. Russia: **ES** (IR), **FE** (KH).
- Tenthredopsis ligata** Konow, 1903. Russia: **EP** (NC, CR). – Europe (EE).
- Tenthredopsis litterata** (Geoffroy, 1785) [Tenthredo] (*Tenthredo carbonaria* Linnaeus, 1767; *T. cordata* Geoffroy, 1785; *T. thoracica* Geoffroy, 1785; *T. varia* Gmelin, 1790; *T. flavipes* Christ, 1791; *T. cruciata* Christ, 1791; *T. dimidiata* Fabricius, 1804; *T. rubiginosa* Drapiez, 1819; *T. microcephala* Lepeletier, 1823; *T. microcephala* Serville, 1823; *T. caliginosa* Stephens, 1835; *T. femoralis* Stephens, 1835; *T. analis* Stephens, 1835; *T. orbitalis* Dietrich, 1868; *Tenthredopsis nigriceps* Cameron, 1881; *T. nigronotatus* Cameron, 1881; *Thomsonia thomsoni* Konow, 1884; *Tenthredopsis thomsoni concolor* Konow, 1887; *T. thomsoni nigripes* Konow, 1890; *T. pallida* Konow, 1896; *T. litterata bicolor* Enslin, 1913; *T. litterata melaena* Enslin, 1918; *T. litterata variana* Enslin, 1918; *T. litterata subcarpathica* Gregor, 1927; *T. carbonaria medionotata* Pic, 1948). Host plant: Dactylis spp., Agrostis spp., Calamagrostis spp. (Poaceae). Russia: **EP** (NW, C, E, S, NC), **UR**, **WS** (OM). – Europe (WE, EE, NE, SE), N Africa, Georgia, Turkey.
- Tenthredopsis moscovita** (André, 1881) [Perineura]. Russia: **EP** (C).
- Tenthredopsis nassata** (Linnaeus, 1767) [Tenthredo] (*Tenthredo alneti* Schrank, 1781; *T. perlata* Geoffroy, 1785; *T. apicalis* Geoffroy, 1785; *T. subulata* Gmelin, 1790; *T. melanorhoa* Gmelin, 1790; *T. napata* Turton, 1802; *Allantus tiliae* Panzer, 1803; *Tenthredo ambigua* Klug, 1817; *T. instabilis* Klug, 1817; *T. fulviceps* Stephens, 1835; *T. tristis* Stephens, 1835; *Perineura brevispina* Thomson, 1870; *Tenthredopsis lividiventris* Cameron, 1881; *T. saundersi* Cameron, 1881; *T. inornatus* Cameron, 1881; *T. albomaculatus* Cameron, 1881; *T. dorsivittatus* Cameron, 1881; *Perineura scutellaris flavoguttata* Magretti, 1882; *Thomsonia josephi* Konow, 1884; *Th. obscura* Konow, 1884; *Th. raddatzi* Konow, 1884; *Th. elegans* Konow, 1884; *Tenthredopsis gibberosa* Konow, 1887; *T. fenestrata* Konow, 1890; *T. dorsalis biguttata* Konow, 1890; *T. dorsalis diluta* Konow, 1890; *T. raddatzi indocilis* Konow, 1890; *T. raddatzi dorsata* Konow, 1890; *T. raddatzi vittata* Konow, 1890; *T. raddatzi maura* Konow, 1890; *T. raddatzi inornata* Konow, 1890; *T. raddatzi sagmaria* Konow, 1890; *T. nassata rufata* Konow, 1890; *T. austriaca obscurata* Konow, 1890; *T. elegans nigronotata* Konow, 1890; *T. dorsalis tirolensis* Konow, 1892; *T. konowi* Strobl, 1896; *T. nassata pleurosternalis* Enslin, 1913; *T. parvula atripleuris* Enslin, 1913; *T. parvula atrilobis* Enslin, 1913; *T. parvula atrifemoribus* Enslin, 1913; *T. parvula rubriventris* Enslin, 1913; *T. parvula atramentaria* Enslin, 1913; *T. austriaca rufofemorata* Enslin, 1913; *T. austriaca albata* Enslin, 1913; *T. austriaca candida* Enslin, 1913; *T. nassata metapleuris* Enslin, 1913; *T. nassata trichroma* Enslin, 1913; *T. tristior* Morice, 1914; *T. inornata melanaspis* Enslin, 1918; *T. nassata nigerrima* Endre, 1927; *T. parvula nigribasis* Zirngiebl, 1937; *T. fenestrata quadripunctata* Gregor, 1941; *T. dubia scutellaris* Gregor, 1941; *T. nassata martialis* Pic, 1948; *T. nassata buyssonii* Pic, 1948; *T. virgineus* Muche, 1965; *T. coqueberti ulanbatorensis* Muche, 1965). Host plant: Poaceae, Cyperaceae. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (NS), **ES** (BR, ZB), **FE** (AM, PR, SA, KU, MG). – Europe (WE, EE, NE, SE), Georgia, Armenia, Turkey, Central Asia, Mongolia, China (CC), Japan.
- Tenthredopsis nigella** Konow, 1891. Russia: **EP** (NC). – Europe (SE), Caucasus, Turkey.
- Tenthredopsis nigrescens** Konow, 1897. Russia: **EP** (Lacourt, 1999). – Armenia.
- Tenthredopsis ornata** (Serville, 1823) [Tenthredo] (*Tenthredo neglecta* Serville, 1823; *T. ornata* Lepeletier, 1823; *T. neglecta* Lepeletier, 1823; *Perineura excisa* Thomson, 1870; *Thomsonia excisa binotata* Konow, 1884; *Tenthredopsis excisa atriscutis* Enslin, 1918). Host plant: Poaceae. Russia: **EP** (N, NW, C, E, S, NC), **UR**. – Europe (WE, EE, NE, SE), Armenia.
- Tenthredopsis ornatrix** Konow, 1890. Russia: **EP** (NC). – Europe (EE), Armenia.

Tenthredopsis rufa Konow, 1890. Russia: **EP** (NC). – Georgia, Azerbaijan.

Tenthredopsis scutellaris (Fabricius, 1804) [Tenthredo] (*Tenthredo spreta* Serville, 1823; *T. dorsalis* Serville, 1823; *T. spreta* Lepeletier, 1823; *T. dorsalis* Lepeletier, 1823; *Tenthredopsis flavomaculatus* Cameron, 1881; *T. picticeps* Cameron, 1881; *Perineura scutellaris baldensis* Heller et Dalla Torre, 1883; *Tenthredopsis albipleuris* Konow, 1886; *T. puncticollis* Konow, 1890; *T. austriaca* Konow, 1890; *T. dubia* Konow, 1890; *T. parvula* Konow, 1890; *T. franki* Konow, 1890; *T. thornleyi* Konow, 1899; *T. puncticollis mediatra* Enslin, 1913; *T. flavomaculata centronigra* Enslin, 1913; *T. nassata intermedia* Hellén, 1943). Host plant: Poaceae. Russia: **EP** (NC). – Europe (WE, EE, NE, SE).

Tenthredopsis sordida (Klug, 1817) [Tenthredo] (*T. sordida atriventris* Enslin, 1913; *T. sordida pleuritica* Enslin, 1913). Host plant: Arrhenatherum spp., Lolium spp., Calamagrostis spp., Dactylis spp. (Poaceae), Carex spp. (Cyperaceae). Russia: **EP** (S). – Europe (WE, EE, NE, SE), Turkey.

Tenthredopsis sororia Konow, 1898. Russia: **EP** (NC). – Armenia, Azerbaijan.

Tenthredopsis stigma (Fabricius, 1798) [Tenthredo] (*Tenthredo dorsalis* Spinola, 1808; *T. histrio* Klug, 1817; *Tenthredopsis stigma genualis* Konow, 1890; *T. churchevillei* Konow, 1897; *T. stigma albonotata* Pigeot, 1917). Host plant: Elymus hispidus (Poaceae). Russia: **EP** (NW, C, E, S, NC), **UR**. – Europe (WE, EE, NE, SE).

Tenthredopsis tarsata (Fabricius, 1804) [Tenthredo] (*T. opacipleuris* Stein, 1884; *Thomsonia hilleckeii* Konow, 1884; *Tenthredopsis wuestneii* Stein, 1884). Host plant: Brachypodium sylvaticum (Poaceae). Russia: **EP** (C, S). – Europe (WE, EE, NE, SE).

Tenthredopsis tessellata (Klug, 1817) [Tenthredo] (*Tenthredo ischiadica* Eversmann, 1847;

T. ischiadica Eversmann, 1864; *Perineura cylindrica* Rudow, 1871; *Tenthredopsis tessellata alboplagiata* Konow, 1890; *T. albata* Konow, 1904; *T. tessellata nigratipleuris* Enslin, 1913; *T. tessellata nigratilobis* Enslin, 1913; *T. tessellata nigratiscutis* Enslin, 1913). Host plant: Deschampsia spp., Dactylis spp., Lolium spp. (Poaceae), Aira spp. (Acoraceae). Russia: **EP** (C, E, S), **UR**, **WS** (TK, NS). – Europe (WE, EE, SE), Armenia, Turkey, Iran, Kazakhstan, Mongolia.

Tenthredopsis tischbeinii (Frivaldszky, 1877) [Tenthredo] (*Tenthredo hungarica* Tischbein, 1852; *Perineura konowi* Lethierry, 1887; *Tenthredopsis fischbeini exannulata* Strobl, 1901; *T. pisinna* Konow, 1903; *T. andrei atronotata* Pic, 1948). Host plant: Poaceae. Russia: **EP** (C, S), **WS** (AL), **ES** (TU, IR, BR), **FE** (AM, MG). – Europe (WE, EE, SE), Kazakhstan, Mongolia.

Tenthredopsis viridis Zhelochovtsev, 1941. Russia: **EP** (NC). – Armenia.

TIANMUTHREDO Wei, 1997. Type species *Tianmuthredo nigrodorsata* Wei, 1997. The genus is distributed in Oriental and Palaearctic regions. Number of species: World – 2; Palaearctic and Russia – 1.

Tianmuthredo davei Taeger et Kramp, 2017. Russia: **FE** (PR).

USSURINUS Malaise, 1931. Type species *Ussurinus insignicornis* Malaise, 1931. The genus is distributed in Palaearctic. Only two species.

Ussurinus insignicornis Malaise, 1931. Russia: **FE** (PR).

Ussurinus nobilis Saarinen, 1945. Russia: **EP** (N). – Europe (NE).

INFRAORDER SIRICOMORPHA
SUPERFAMILY PAMPHILIOIDEA

Head closed ventrally, the clypeus being folded back underneath and joined by its lateral arms to the post-genae behind; the result of this is a capsule with 4 holes in it. Mesosternum triangular in shape, clearly defined by sutures and with a presternal bridge in front. Abdomen strongly flattened dorsiventrally.

Male genitalia orthandrious (not twisted). Female sawsheath not exserted strongly. Larva oligopodous.

The superfamily is distributed in North hemisphere, mainly in Temperate zone. Number of taxa: World – 2 families, 2 subfamilies, 11 genera, more than 330 species; Palaearctic – 2/2/11/about 240; Russia – 2/2/10/111 (+ 3 ssp.).

8. FAMILY PAMPHILIIDAE

Very flat and broad species, 7-17 mm long, sun-loving and very fast on the wing. Antennae long, thread-like and many-segmented (18-24 segments). Fore wings with vein *Sc* free and *2r* present. At least the middle and hind tibiae have preapical spines.

The larvae live solitarily or socially. The larvae of the subfamily Cephalciinae feed on conifers, live in a web; the larvae of the subfamily Pamphiliinae feed on angiosperms, live in a web, or in a tube of a rolled leaf held by silk.

The family is distributed in Holarctic, Oriental and Neotropical regions, mainly in Temperate zone. Number of taxa: World – 2 subfamilies, 10 genera, about 300 species; Palaearctic – 2/10/>200; Russia – 2/9/99 (+ 3 ssp.).

References. Dalla Torre, 1894; Konow, 1903; Dovnar-Zapolskij, 1930; Gussakovskij, 1935; Malaise, 1943; Zhelochovtsev, 1968, 1976, 1988; Ermolenko, 1969, 1973, 1976, 1984; Beneš, 1972a,b, 1974, 1976a,b, 2014b; Verzhutskii, 1974, 1981; Zhelochovtsev, Prochorova, 1976; Stroganova, 1978b; Shinohara, 1979, 1980, 1987, 1988a,b, 1992a, 1993, 1995a, 1997a,b, 1998, 2000, 2001a,b, 2002b, 2003, 2005a; Koch, 1982; Achterberg, Aarsten, 1986; Shinohara, Byun, 1993, 1996; Zhelochovtsev, Zinovjev, 1995; Shinohara, Zinovjev, 1996; Shinohara, Lee, 1997; Shinohara, Hara, 2000; Xiao, 2000; Viitasaari, 2002; Shinohara, Zombori, 2003; Shinohara, Vasilenko, 2005; Vasilenko, Dolgov, 2005; Liston et al., 2006, 2014a; Piotrowski et al., 2006; Shinohara, Zhou, 2006; Taeger et al., 2006, 2010; Wei

et al., 2006; Shinohara, Lelej, 2007; Shinohara, Taeger, 2007; Roller, Haris, 2008; Gurov, Battisti, 2009; Schedl, 2009; Sundukov, 2009, 2011b, 2013, 2015; Sundukov, Lelej, 2009, 2012; Hara, 2010; Vasilenko, 2009b, 2010b; Taeger, Blank, 2011; Macek, 2012; Vasilenko, Korshunov, 2012; Ahn et al., 2013; Kulagin, Simonenkova, 2014; Sokolov, 2014; Beneš, Holuša, 2015; Kostyunin, 2015; Popov, 2015; Noblecourt, 2016; Hara, Shinohara, 2017; Shinohara, Kato, 2017.

Subfamily CEPHALCINAE

ACANTHOLYDA A. Costa, 1894 (*Acanthocnema* Costa, 1859; *Itycorsia* Konow, 1897). Type species *Tenthredo erythrocephala* Linnaeus, 1758. The larvae feed on the Pinus, Larix, Picea, Abies, Tsuga, Pseudotsuga (Pinaceae); singly or in groups weave nests from the web. The genus is distributed in Holarctic, Oriental and Neotropical regions; mainly in Holarctic. Number of species: World – > 70; Palaearctic – about 30; Russia – 8 (+ 1 ssp.) from 2 subgenera.

Acantholyda (Acantholyda) erythrocephala (Linnaeus, 1758) [Tenthredo] (*A. erythrocephala vittata* Enslin, 1918; *A. grangeoni* Riou, 1999). Host plant: Pinus spp. (Pinaceae); live in groups, often harm. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TK, NS, KM), **ES** (KR, IR, BR), **FE** (AM). – Europe (WE, EE, NE, SE), Kazakhstan, China (NE, NC), Korean Peninsula, N America.

Acantholyda (Acantholyda) flaviceps (Retzius, 1783) [Tenthredo] (*Lyda cyanea* Klug, 1808). Host plant: Pinus spp. (Pinaceae). Russia: **EP** (NW, C, E), **UR**, **WS** (KM, AL), **ES** (KR). – Europe (WE, EE, NE, SE).

Acantholyda (Itycorsia) aglaia aglaia Zhelochovtsev, 1968 (*A. angarica* Ermolenko, 1969). Host plant: Picea spp. (Pinaceae). Russia: **WS** (KM), **ES** (KR, IR), **FE** (KH, PR). – Japan (Hon).

Acantholyda (Itycorsia) aglaia yezoensis Shinohara et Hara, 2000. Host plant: Picea spp. (Pinaceae). Russia: **FE** (KU). – Japan (Hok).

Acantholyda (Itycorsia) hieroglyphica (Christ, 1791) [Tenthredo] (*Lyda bimaculata* Taschenberg, 1861). Host plant: Pinus spp. (Pinaceae); live alone. Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (NS, KM), **ES** (IR). – Europe (WE, EE, NE, SE), Caucasus, Kazakhstan.

Acantholyda (Itycorsia) laricis (Giraud, 1861) [*Lyda*] (*A. parvula* Zirngiebl, 1937). Host plant: Larix spp.

- (Pinaceae); often harm. Russia: **EP** (N, C), **UR**, **WS** (TK, AL), **ES** (IR, YA), **FE** (AM). – Europe (WE, EE, SE), China (NC), Japan (Hon).
- Acantholyda (Itycorsia) parki** Shinohara et Byun, 1996. Host plant: *Pinus koraiensis* (Pinaceae). Russia: **FE** (PR). – Korean Peninsula.
- Acantholyda (Itycorsia) pirica** Shinohara, 2000. Host plant: ?*Pinus pumila* (Pinaceae). Russia: **FE** (SA). – Japan (Hok).
- Acantholyda (Itycorsia) posticalis pinivora** Enslin, 1918 (*Tenthredo stellata* Christ, 1791; *T. pratensis* Fabricius, 1793; *Acantholida nemoralis magnini* Pic, 1944). Host plant: *Pinus* spp. (Pinaceae); often harm. Russia: **EP** (C), **UR**, **WS** (TK, NS, KM, AL), **ES** (YA, ZB). – Europe (WE, EE, NE, SE), Kazakhstan, Mongolia, China (NE, SE).
- CAENOLYDA** Konow, 1897 (*Dictyolyda* Semenov, 1934). Type species *Tenthredo reticulata* Linnaeus, 1758. The larvae feed on the *Pinus* spp. (Pinaceae). The genus is distributed in West Palaearctic. Number of species: World and Palaearctic – 2; Russia – 1.
- Caenolyda reticulata** (Linnaeus, 1758) [Tenthredo] (*Cephaleia clarkii* Jurine, 1807). Host plant: *Pinus* spp. (Pinaceae). Russia: **EP** (NW, C, E). – Europe (WE, EE, NE, SE).
- CEPHALCIA** Panzer, 1803 (*Cephaleia* Jurine, 1807; *Liolyda* Ashmead, 1898). Type species *Cephaleia arvensis* Panzer, 1803. The larvae feed on the *Pinus*, *Larix*, *Picea*, *Abies*, *Cedrus* (Pinaceae); singly or in groups weave nests from the web. The genus is distributed in Holarctic and Oriental region, mainly in Holarctic. Number of species: World – 43; Palaearctic – 30; Russia – 13 (+ 1 ssp.).
- Cephalcia abietis** (Linnaeus, 1758) [Tenthredo] (*Lyda hypothrofica* Hartig, 1834; *Cephaleia testacea* Gimmerthal, 1836; *Lyda klugii* Hartig, 1837; *L. scutellaris* Thomson, 1871; *L. semiflava* Matsumura, 1912; *Cephaleia mimica* Pic, 1948). Host plant: *Picea* spp. (Pinaceae); live in groups, harm. Russia: **EP** (N, NW, C, E), **UR**, **WS** (KM), **ES** (KR, BR, YA), **FE** (KH, PR, SA). – Europe (WE, EE, NE, SE), Mongolia, China (NE, NC), Japan (Hok).
- Cephalcia alashanica** (Gussakovskij, 1935) [Cephaleia] (*C. alashanica europaea* Beneš, 1976). Russia: **EP** (C). – Europe (WE, EE, NE), Mongolia, China (NC).
- Cephalcia alpina** (Klug, 1808) [Lyda] (*Lyda fallenii* Dalman, 1823; *L. abietina* Hartig, 1837; *L. annulata* Hartig, 1837). Host plant: *Picea* spp., *Larix* spp. (Pinaceae); live alone. Russia: **EP** (N, C), **WS** (TM, AL), **ES** (KR, IR, YA), **FE** (SA). – Europe (WE, EE, NE, SE), N Africa, China (NE), Korean Peninsula, Japan (Hok, Hon).
- Cephalcia arvensis** Panzer, 1803 (*Tenthredo signata* Fabricius, 1781; *Lyda irrorata* Dahlbom, 1835; *L. saxicola* Hartig, 1837; *L. adusta* Dietrich, 1868; *L. irrorata* Thomson, 1871). Host plant: *Picea* spp. (Pinaceae); live alone or in small groups. Russia: **EP** (N, NW, C, E), **UR**, **WS** (NS), **ES** (KR, IR, YA), **FE** (PR, SA, KU). – Europe (WE, EE, NE, SE), Mongolia, China (NE, NC).
- Cephalcia erythrogaster** (Hartig, 1837) [Lyda]. Host plant: *Picea* spp. (Pinaceae); live alone. Russia: **EP** (NW, C). – Europe (WE, EE, NE), China (Wei et al., 2006).
- Cephalcia hartigii** (Bremi-Wolf, 1849) [Lyda]. Host plant: *Abies* spp., *Picea* spp. (Pinaceae). Russia: **EP** (NC). – Europe (WE, EE, SE), N Africa, Georgia.
- Cephalcia intermedia** Hellén, 1948. Russia: **EP** (N). – Europe (NE).
- Cephalcia koebelei** (Rohwer, 1910) [Cephaleia] (*C. baikalica* Verzhutskii, 1973). Host plant: *Larix leptolepis* (Pinaceae). Russia: **ES** (IR, YA). – Japan (Hok, Hon).
- Cephalcia lariciphila japonica** Shinohara, 1997. Host plant: *Larix leptolepis* (Pinaceae). Russia: **FE** (KU). – Japan (Hok, Hon).
- Cephalcia lariciphila lariciphila** (Wachtl, 1898) [Cephaleia]. Host plant: *Larix* spp. (Pinaceae). Russia: **EP** (N, C), **WS** (KM, AL), **ES** (TU, KR, IR, BR, YA), **FE** (PR). – Europe (WE, EE, NE, SE), China (NE, CC), Korean Peninsula.
- Cephalcia nigricoxae** (Matsumura, 1912) [Lyda]. Host plant: *Picea jezoensis* (Pinaceae); harmful to spruce forests in Hokkaido. Russia: **FE** (KU). – Japan (Hok).
- Cephalcia pallidula** (Gussakovskij, 1935) [Cephaleia] (*C. flavistigma* Lindqvist, 1962). Host plant: *Picea* spp. (Pinaceae). Russia: **EP** (N), **ES** (IR, YA), **FE** (PR, SA, KU). – Europe (NE), China (NE).
- Cephalcia sachalinensis** (Matsumura, 1911) [Lyda]. Host plant: *Picea* spp. (Pinaceae). Russia: **FE** (SA).
- Cephalcia variegata** (Takeuchi, 1930) [Cephaleia]. Host plant: *Pinus pumila* (Pinaceae). Russia: **FE** (SA, KU). – Japan (Hok, Hon).

Subfamily PAMPHILIINAE

CHRYSOLYDA Shinohara, 2002. Type species *Pamphilus leucocephalus* Takeuchi, 1938.

The genus is distributed in East Asia. Number of species: one species.

Chrysolyda leucocephala (Takeuchi, 1938) [Pamphilius]. Russia: **FE** (PR). – Korean Peninsula, Japan (Hok, Hon, Kyu).

KELIDOPTERA Konow, 1897. Type species *Lyda maculipennis* J.P.E.F. Stein, 1876. The genus is distributed in West Palaearctic. Number of species: one species.

Kelidoptera maculipennis (J.P.E.F. Stein, 1876) [Lyda]. Russia: **EP** (E). – Turkey.

NEUROTOMA Konow, 1897 (*Gongylocorsia* Konow, 1897). Type species *Tenthredo flaviventris* Retzius, 1783. The larvae feed mainly on the tree and shrub Rosaceae and Quercus (Fagaceae); single or in groups of leaves feeding, weave a web. The genus is distributed in Holarctic and Oriental region, mainly in Holarctic. Number of species: World – 22; Palaearctic – 13; Russia – 7.

Neurotoma atrata Takeuchi, 1930. Host plant: *Quercus acutissima* Fagaceae). Russia: **FE** (PR). – China (SE), Korean Peninsula, Japan (Hon).

Neurotoma fausta (Klug, 1808) [Lyda]. Russia: **EP** (CR). – Europe (WE, EE, SE).

Neurotoma iridescents (André, 1882) [Lyda] (*Cephalcia nigrocoerulea* Rohwer, 1910; *Neurotoma sorbi* Forsius, 1911; *Lyda nigricans* Matsumura, 1912; *Neurotoma nakayamai* Sugihara, 1933; *Pamphilius sajanicus* Stroganova, 1978). Host plant: *Prunus* spp., *Sorbus* spp., *Crataegus chlorosarca* (Rosaceae); live in groups. Russia: **EP** (N, NW), **ES** (KR, BR, YA), **FE** (PR, KU). – Europe (WE, NE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Neurotoma mandibularis (Zaddach, 1866) [Lyda] (*Lyda parisiensis* Giraud, 1870). Host plant: *Quercus* spp. (Fagaceae). Russia: **EP** (CR). – Europe (WE, EE, SE).

Neurotoma nemoralis (Linnaeus, 1758) [Tenthredo] (*Tenthredo punctata* Fabricius, 1798; *Psen caprifolia* Schrank, 1802; *Lyda lucorum* Fallén, 1808; *L. drupacearum* Nördlinger, 1855; *L. maculifrons* Snellen van Vollenhoven, 1858). Host plant: *Prunus* spp. (Rosaceae); live in groups. Russia: **EP** (NW, C, E, S, NC), **UR**. – Europe (WE, EE, NE, SE), Kazakhstan.

Neurotoma saltuum (Linnaeus, 1758) [Tenthredo] (*Tenthredo flaviventris* Retzius, 1783; *Psen pyri*

Schrank, 1802; *Lyda albifrons* Fallén, 1808; *L. clypeata* Klug, 1808; *Tenthredo lutescens* Panzer, 1809; *Pamphilius dimidiatus* Latreille, 1812; *Lyda fasciata* Curtis, 1831; *L. fasciatipennis* Costa, 1864). Host plant: *Prunus* spp., *Malus* spp., *Crataegus* spp. and others Rosaceae; live in groups in a nest of cobwebs, sometimes harm. Russia: **EP** (C, S, NC, CR), **ES** (IR). – Europe (WE, EE, NE, SE), Armenia, Turkey, Korean Peninsula.

Neurotoma sibirica Gussakovskij, 1935 (*N. nemoralis obsoleta* Takeuchi, 1936). Host plant: *Sorbaria sorbifolia* (Rosaceae). Russia: **ES** (IR, YA, ZB), **FE** (AM, KH, PR, SA, KU). – China (NE), Korean Peninsula, Japan (Hok).

ONYCHOLYDA Takeuchi, 1938. Type species *Pamphilius viriditibialis* Takeuchi, 1930. The larvae feed on the *Rubus*, *Filipendula*, *Agrimonia* (Rosaceae), *Cornus* (Cornaceae); live alone and in groups, weave a web. The genus is distributed in Holarctic and Oriental region, mainly in Holarctic. Number of species: World – about 40; Palaearctic – 26; Russia – 11.

Onycholyda armata (Maa, 1949) [Pamphilius]. Russia: **WS** (TK), **FE** (KH, PR). – China (NE, SE), Korean Peninsula.

Onycholyda kumamotonis (Matsumura, 1912) [Lyda]. Host plant: *Filipendula kamtschatica* (Rosaceae). Russia: **FE** (SA, KU). – Japan (Hok, Hon, Kyu).

Onycholyda lucida (Rohwer, 1910) [Pamphilius]. Host plant: *Rubus* spp. (Rosaceae). Russia: **ES** (YA). – Japan (Hon, Shi, Kyu).

Onycholyda minomalis (Takeuchi, 1930) [Pamphilius]. Host plant: *Rubus parvifolius* (Rosaceae). Russia: without regions (Taeger, Blank, 2011). – Japan (Hok, Hon, Shi, Kyu).

Onycholyda nigroclypeata Shinohara, 1987. Russia: **FE** (KH, PR). – China (NE), Korean Peninsula.

Onycholyda rufofasciata (Norton, 1869) [Lyda] (*Lyda rufocincta* Cresson, 1880). Host plant: ?*Rosa* spp. (Rosaceae). Russia: **ES** (IR). – N America.

Onycholyda sertata (Konow, 1903) [Pamphilius] (*Pamphilius thomsoni* Malaise, 1921). Host plant: *Filipendula* spp. (Rosaceae). Russia: **EP** (NW, C, E), **UR**, **WS** (NS, AL), **ES** (YA), **FE** (AM, KH, PR). – Europe (NE), China (NE), Korean Peninsula.

Onycholyda trigaria (Konow, 1897) [Pamphilius]. Russia: **EP** (NC). – Armenia, Azerbaijan, Iran.

Onycholyda viriditibialis (Takeuchi, 1930) [Pamphilius] (*Pamphilius tsherskii* Gussakovskij, 1935).

- Host plant: *Rubus crataegifolius*, *R. microphyllus* (Rosaceae); live in groups. Russia: **FE** (PR). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Onycholyda yezoensis** Shinohara, 1987. Host plant: *Rubus parvifolius*, *R. idaeus* (Rosaceae). Russia: **FE** (SA, KU). – Japan (Hok).
- Onycholyda zinovjevi** Shinohara, 1987. Russia: **FE** (PR). – Korean Peninsula.
- PAMPHILIUS** Latreille, 1803 (*Lyda* Fabricius, 1804; *Anoplolyda* Costa, 1894; *Bactroceros* Konow, 1897). Type species *Tenthredo sylvatica* Linnaeus, 1758. The larvae feed mainly on Rosaceae and Betulaceae, others on Salicaceae, Sapindaceae, Caprifoliaceae, Fagaceae, Cornaceae and Juglandaceae; mostly monophages; live alone or in groups in rolled leaves. The genus is distributed in Holarctic and Oriental region, mainly in Holarctic. Number of species: World – about 115; Palaearctic – about 95; Russia – 56 (+ 1 ssp.).
- Pamphilius albopictus** (Thomson, 1871) [Lyda] (*Pamphilius altaicus* Gussakovskij, 1935; *P. viridipes* Achterberg et Aartsen, 1986). Host plant: *Prunus* spp. (Rosaceae). Russia: **EP** (N, C), **UR**, **WS** (KM, AL), **ES** (IR, YA), **FE** (KH, PR, KA). – Europe (WE, NE), Korean Peninsula.
- Pamphilius alnicola** Ermolenko, 1973. Host plant: *Alnus hirsuta* (Betulaceae). Russia: **FE** (PR, KU). – Japan (Hok, Hon).
- Pamphilius alnivorus** Shinohara, 2005. Host plant: *Alnus hirsuta* (Betulaceae). Russia: **ES** (YA), **FE** (KH, PR, SA, KU). – Japan (Hok, Hon).
- Pamphilius archiducalis** Konow, 1897. Host plant: *Alnus* spp. (Betulaceae). Russia: **FE** (SA, KU). – Japan (Hok, Hon).
- Pamphilius armeniacus** Shinohara, 1988. Russia: **EP** (NC). – Armenia.
- Pamphilius aucupariae** Vikberg, 1971. Host plant: *Sorbus* spp. (Rosaceae). Russia: **ES** (YA). – Europe (NE).
- Pamphilius balteatus** (Fallén, 1808) [Lyda] (*P. cingulatus* Latreille, 1812; *Lyda suffusa* Hartig, 1837). Host plant: *Rosa* spp., *Prunus* spp., *Spiraea* spp. (Rosaceae); live alone. Russia: **EP** (NW, C, E), **UR**, **WS** (NS), **ES** (IR, YA, ZB), **FE** (PR, SA, KU, KA, MG). – Europe (WE, NE, SE), Korean Peninsula, Japan (Hok).
- Pamphilius betulae** (Linnaeus, 1758) [*Tenthredo*] (*Tenthredo fulva* Retzius, 1783; *Lyda aurita* Klug, 1808). Host plant: *Populus* spp. (Salicaceae); live alone. Russia: **EP** (NW, C, E), **UR**, **WS** (NS, KM, AL), **ES** (IR, YA), **FE** (KH, PR, SA, KU, MG). – Europe (WE, EE, NE, SE), Turkey. **Pamphilius borisi** Beneš, 1972. Russia: **ES** (IR).
- Pamphilius brevicornis brevicornis** Hellén, 1948. Host plant: *Populus* spp. (Salicaceae). Russia: **EP** (N), **WS** (NS, KM), **ES** (IR), **FE** (PR, SA, KA). – Europe (WE, NE), Korean Peninsula.
- Pamphilius caucasicus** Gussakovskij, 1935. Russia: **EP** (NC). – Georgia.
- Pamphilius convexus** Shinohara, 1988. Russia: **FE** (KH, PR). – Korean Peninsula.
- Pamphilius coreanus** Takeuchi, 1938 (*Anoplolyda frontimacula* Malaise, 1943). Russia: **FE** (KH, PR). – Korean Peninsula.
- Pamphilius croceus** Shinohara, 1986. Russia: **FE** (PR). – Korean Peninsula.
- Pamphilius festivus** C. Pesarini et F. Pesarini, 1984. Host plant: *Populus* spp. (Salicaceae). Russia: **EP** (NW, C, S), **UR**, **WS** (NS, KM, AL), **ES** (KS, KR). – Europe (WE, EE, NE, SE).
- Pamphilius fumipennis** (Curtis, 1831) [Lyda] (*Lyda stigma* Stephens, 1835; *L. carpini* Brischke, 1892). Host plant: *Corylus* spp., *Alnus* spp. (Betulaceae). Russia: **EP** (N, NW, C, E). – Europe (WE, EE, NE, SE).
- Pamphilius gyllenhali** (Dahlbom, 1835) [Lyda]. Host plant: *Salix* spp. (Salicaceae); live alone. Russia: **EP** (N, NW, C, E), **UR**. – Europe (WE, NE, SE).
- Pamphilius heecheonparki** Shinohara, 1998. Russia: **ES** (IR, YA), **FE** (KH, PR). – Korean Peninsula.
- Pamphilius hilaris** (Eversmann, 1847) [Lyda]. Host plant: *Rosa* spp. (Rosaceae). Russia: **UR**, **WS** (TK), **FE** (PR, KA). – China (CC), Japan (Hon).
- Pamphilius histrio** Latreille, 1812. Host plant: *Populus* spp. (Salicaceae); live alone. Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (NS), **ES** (YA). – Europe (WE, EE, NE, SE), Kazakhstan, Korean Peninsula, Japan (Hok).
- Pamphilius hortorum** (Klug, 1808) [Lyda] (*P. hortorum bicinctus* Benson, 1945). Host plant: *Rubus* spp. (Rosaceae). Russia: **EP** (N, NW, C, E), **UR**, **WS** (NS, KM, AL), **ES** (IR, YA), **FE** (KH, PR, SA, KU, MG). – Europe (WE, EE, NE, SE), China (NE), Korean Peninsula, Japan (Hok, Hon).
- Pamphilius inanitus** (Villers, 1789) [*Tenthredo*] (*Tenthredo agrestis* Panzer, 1804; *Lyda fallax* Serville, 1823; *L. fallax* Lepeletier, 1823). Host plant: *Rosa* spp. (Rosaceae). Russia: **EP** (NW), **ES** (YA). – Europe (WE, EE, NE, SE).

Pamphilius itoi Shinohara, 1985. Host plant: *Alnus hirsuta* (Betulaceae). Russia: **FE** (PR, KU). – Japan (Hok, Hon).

Pamphilius jucundus (Eversmann, 1847) [Lyda] (*Lyda arbuti* Zaddach, 1866). Host plant: *Fragaria* spp. (Rosaceae). Russia: **EP** (NW, C, E), **UR**, **WS** (NS). – Europe (WE, EE, NE).

Pamphilius kamikochensis Takeuchi, 1930. Host plant: *Prunus ssiori* (Rosaceae). Russia: **FE** (KU). – Japan (Hok, Hon).

Pamphilius kontuniemii Shinohara, 2003. Russia: **EP** (N, NW, NC). – Europe (WE, EE, NE, SE), Georgia.

Pamphilius kyutekparki Shinohara, 1991. Russia: **FE** (PR). – Korean Peninsula.

Pamphilius latifrons (Fallén, 1808) [Lyda] (*Lyda maculosa* Zaddach, 1866). Host plant: *Populus* spp., *Salix* spp. (Salicaceae). Russia: **EP** (N, C), **WS** (KM), **ES** (IR, BR, ZB), **FE** (PR, MG). – Europe (WE, EE, NE, SE), Kazakhstan.

Pamphilius leleji Shinohara et Taeger, 2007. Russia: **FE** (PR).

Pamphilius lethierryi (Konow, 1887) [Lyda]. Russia: **EP** (NC). – Europe (WE, EE, SE), Georgia, Armenia.

Pamphilius lobatus Maa, 1950. Russia: **FE** (PR). – China (NE), Korean Peninsula, Japan (Hok, Hon).

Pamphilius marginatus (Serville, 1823) [Lyda] (*Lyda marginata* Lepeletier, 1823; *L. bicolor* Herrich-Schäffer, 1833; *Tenthredo ratzeburgii* Ratzeburg, 1844). Host plant: *Corylus* spp., *Carpinus* spp., *Ostrya* spp. (Betulaceae). Russia: **EP** (C). – Europe (WE, EE, NE, SE).

Pamphilius masao Shinohara, 2005. Host plant: *Betula ermanii*, *Alnus maximowiczii* (Betulaceae). Russia: **FE** (SA, KU). – Japan (Hok).

Pamphilius maximus Shinohara, 1995. Russia: **FE** (PR). – Korean Peninsula.

Pamphilius montanus pulcher Shinohara, 1988. Host plant: *Sorbus* spp. (Rosaceae). Russia: **ES** (YA). – Korean Peninsula, Japan (Hok).

Pamphilius pallidus Shinohara, 1988. Russia: **FE** (PR).

Pamphilius pallipes (Zetterstedt, 1838) [Lyda] (*Lyda flavipes* Zetterstedt, 1838; *L. variegata* Zaddach, 1866; *Pamphilius pallidipes* Dalla Torre, 1894). Host plant: *Betula* spp., *Alnus* spp. (Betulaceae). Russia: **EP** (N, NW, C, E), **WS** (TK, NS, KM, AL), **ES** (BR, YA), **FE** (AM, KH, PR, KU, KA, MG). – Europe (WE, EE, NE), Iran, Kazakhstan, Mongolia, China (NE, NC), Korean Peninsula.

Pamphilius persicum MacGillivray, 1907. Russia: **EP** (NC). – Armenia, N America.

Pamphilius pictifrons Gussakovskij, 1935. Host plant: ?*Rosa* spp. (Rosaceae). Russia: **ES** (YA), **FE** (MG).

Pamphilius planifrons Beneš, 1976. Russia: **FE** (PR). – Korean Peninsula.

Pamphilius pugnax Konow, 1897. Russia: **EP** (NC). – Azerbaijan.

Pamphilius sapporensis (Matsumura, 1912) [Lyda] (*P. rugosus* Beneš, 1976). Host plant: *Filipendula kamtschatica* (Rosaceae). Russia: **FE** (SA, KU). – Japan (Hok).

Pamphilius stramineipes (Hartig, 1837) [Lyda] (*P. facetus* Konow, 1898; *P. stramineipes medialis* Enslin, 1917). Host plant: *Rosa* spp. (Rosaceae). Russia: **EP** (N, C, E, NC), **UR**, **WS** (TM, OM, NS), **ES** (KR), **FE** (SA, KU). – Europe (WE, EE, NE), Armenia, Japan (Hok).

Pamphilius sulphureipes sulphureipes W.F. Kirby, 1882 (*Anoplolyda viriditibialis aino* Takeuchi, 1936). Host plant: *Rosa* spp. (Rosaceae). **FE** (AM, PR, SA). – Korean Peninsula.

Pamphilius sylvaticus (Linnaeus, 1758) [Tenthredo] (*Tenthredo fulvipes* Retzius, 1783; *Cephalcia nemorum* Panzer, 1803). Host plant: *Sorbus* spp., *Crataegus* spp., *Prunus* spp. (Rosaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (NS, KM), **ES** (YA). – Europe (WE, EE, NE, SE), Turkey.

Pamphilius takeuchii Beneš, 1972 (*P. jucundus* Takeuchi, 1930). Russia: **FE** (PR). – Japan (Hok, Hon, Shi, Kyu).

Pamphilius tricolor Beneš, 1974. Russia: **EP** (N), **WS** (TM), **ES** (TU, KR, ZB), **FE** (KH, PR, KA, MG). – Korean Peninsula, Japan (Hok, Hon).

Pamphilius ussuriensis Shinohara, 1988. Russia: **FE** (PR).

Pamphilius vafer (Linnaeus, 1767) [Tenthredo] (*Tenthredo depressa* Schrank, 1781; *Psen alni* Schrank, 1802). Host plant: *Alnus* spp., *Betula* spp. (Betulaceae). Russia: **EP** (N, NW, C, E), **UR**, **WS** (NS, AL), **ES** (TU, IR, YA). – Europe (WE, EE, NE, SE), Georgia, Kazakhstan.

Pamphilius varius (Serville, 1823) [Lyda] (*Lyda varia* Lepeletier, 1823; *L. infida* Zaddach, 1866; *Pamphilius flavifrons* W.F. Kirby, 1882; *Lyda konowi* Jakovlev, 1891; *Anoplolyda engelhardtii* Dovnar-Zapolskij, 1930). Host plant: *Alnus* spp., *Betula* spp. (Betulaceae). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (NS, KM), **ES** (KR, BR, YA), **FE** (AM, PR, SA, KU, KA). – Europe (WE, EE, NE, SE), Georgia,

- Kazakhstan, Korean Peninsula, Japan (Hok, Hon, Kyu).
- Pamphilius venustus** (F. Smith, 1874) [Lyda]. Russia: FE (Shinohara, 1988b). – Japan (Hok, Hon, Shi).
- Pamphilius virescens** Malaise, 1931. Russia: ES (IR), FE (KA). – Korean Peninsula, Japan (Hok).
- Pamphilius viridulus** Shinohara, 2001. Host plant: Prunus spp., Crataegus chlorosarca (Rosaceae). Russia: FE (PR).
- Pamphilius volatilis** (F. Smith, 1874) [Lyda]. Host plant: Prunus spp., Crataegus chlorosarca (Rosaceae). Russia: FE (PR). – Korean Peninsula, Japan (Hok, Hon, Kyu).
- Pamphilius zhelochovtsevi zhelochovtsevi** Beneš, 1974. Host plant: Rosa spp. (Rosaceae). Russia: FE (KH, PR). – Korean Peninsula.

Pamphilius zhelochovtsevi nipponicus Shinohara, 1993. Host plant: Rosa spp. (Rosaceae). Russia: FE (KU). – Japan (Hon).

Pamphilius zinovjevi Shinohara, 1988. Russia: FE (AM, PR). – Korean Peninsula.

PSEUDOCEPHALEIA Zirngiebl, 1937. Type species *Pseudocephaleia brachycercus* Zirngiebl, 1937. The genus is distributed in Palaearctic. Number of species: World and Palaearctic – 2; Russia – 1.

Pseudocephaleia praeteritorum (Semenov, 1934) [Dictyolyda] (*P. brachycercus* Zirngiebl, 1937; *Cephalcia pseudoreticulata* Zombori, 1967). Russia: WS (AL). – Europe (EE, SE), Caucasus, Turkey, China (Wei et al., 2006).

9. FAMILY MEGALODONTESIDAE

Flat-bodied flower-haunting species, very fast on the wing. Characterized by their many-segmented flabellate antennae, with flattened prolongations from the apices of the flagellar segments. Body very broad and flat, pamphiloid in form, with wasp-like colouring and yellowish-brown marked wings. The larvae live socially in webs on herbaceous plants – Apiaceae and Rutaceae.

The family is distributed in the Palaearctic and Oriental regions, but was rich and diverse group in the Mesozoic. Number of taxa: World and Palaearctic – 1 genus, 37 species; Russia – 1/12.

R e f e r e n c e s. Jakovlev, 1888; Gussakovskij, 1935; Maa, 1949b; Zombori, 1971; Togashi, 1973; Verzhutskii, 1973, 1974, 1981; Ermolenko, 1984; Zhelochovtsev, 1988; Zhelochovtsev, Zinovjev, 1995; Taeger, 1998, 2002; Vasilenko, Dolgov, 2005; Liston et al., 2006; Taeger et al., 2006, 2010; Wei et al., 2006; Roller, Haris, 2008; Schedl, 2009; Sundukov, Lelej, 2009, 2012; Taeger, Blank, 2011; Kostyunin, 2015; Noblecourt, 2016; Wang et al., 2016.

MEGALODONTES Latreille, 1803 (*Tarpa* Fabricius, 1804; *Megalodus* Rafinesque, 1815; *Melanopus* Konow, 1897; *Tristactus* Konow, 1897; *Rhipidioceros* Konow, 1897; *Forficulotarpa* Pic, 1918; *Tristactoides* Chevin, 1985). Type species *Tenthredo cephalotes* Fabricius, 1781. The larvae feed on the Apiaceae and Rutaceae; live in nests from cobwebs. The genus is distributed in Palaearctic and Oriental region. Number of species: World and Palaearctic – 37; Russia – 12.

Megalodontes cephalotes (Fabricius, 1781) [*Tenthredo*] (*Tarpa klugii* Leach, 1817; *T. spissicornis* Klug, 1824; *T. fuscipennis* Herrich-Schäffer, 1840). Host plant: Laserpitium spp., Libanotis spp., Peucedanum spp., Seseli libanotis (Apiaceae). Russia: **EP** (N, C, E), **UR**. – Europe (WE, EE, SE), Turkey.

Megalodontes eversmanni (Freymuth, 1870) [*Tarpa*] (*Tarpa loewii* J.P.E.F. Stein, 1876; *T. multicincta* Mocsáry, 1891). Russia: **EP** (NC). – Europe (SE), Caucasus, Turkey, Iran.

Megalodontes fabricii (Leach, 1817) [*Tarpa*] (*Tarpa megacephala* Klug, 1824; *T. redimita* Klug, 1824). Russia: **EP** (C, S, E, NC, CR), **UR**. – Europe (WE, EE, SE), Turkey.

Megalodontes flabellicornis (Germar, 1825) [*Tarpa*] (*Tarpa exornata* Zaddach, 1866; *T. coronata* Zaddach, 1866; *T. albicincta* J.P.E.F. Stein, 1876; *T. speciosa* Mocsáry, 1877). Russia: **EP** (C, NC). – Europe (EE, SE), Georgia, Turkey.

Megalodontes flavicornis (Klug, 1824) [*Tarpa*]. Russia: **EP** (C, S, NC, CR). – Europe (WE, EE, SE), Caucasus, Turkey, Iran, Turkmenistan.

Megalodontes interruptus Gussakovskij, 1935 (*M. antoniae* Zombori, 1971). Russia: **ES** (IR). – Mongolia, China (NE).

Megalodontes medius Konow, 1897. Russia: **EP** (S, NC, CR). – Europe (EE), Azerbaijan, Turkey.

Megalodontes plagicephalus (Fabricius, 1804) [*Tarpa*] (*Tarpa flabellata* Eversmann, 1847; *T. plagicephala latialis* Manzone, 1893; *Megalodontes aquilus* Konow, 1902; *M. nigritegulis* Konow, 1904; *M. xanthocerus* Gussakovskij, 1935). Host plant: Peucedanum alsaticum, Seseli libanotis (Apiaceae). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (OM, NS, AL). – Europe (WE, EE, SE), Georgia, Armenia, Kazakhstan.

Megalodontes quinquecinctus (Klug, 1824) [*Tarpa*] (*M. nitidus* Maa, 1949; *M. tsunekii* Togashi, 1973; *M. apakensis* Togashi, 1973). Russia: **EP** (CR), **ES** (BR, ZB). – Mongolia, China (NE, NC).

Megalodontes scythicus Zhelochovtsev, 1988 (*M. eversmanni* Gussakovskij, 1935). Russia: **EP** (S, NC, CR), **UR**. – Europe (EE), Kazakhstan.

Megalodontes spiraeae (Klug, 1824) [*Tarpa*] (*Tarpa pectinicornis* Klug, 1824; *T. ruthena* Jakovlev, 1888; *T. borealis* Jakovlev, 1891; *Megalodontes siberiensis* Rohwer, 1925; *M. coreensis* Takeuchi, 1927). Host plant: Sphallerocarpus gracilis (Apiaceae), Spiraea spp. (Rosaceae). Russia: **UR**, **WS** (NS, AL), **ES** (KS, KR, IR, BR, ZB), **FE** (AM, KH, PR). – Mongolia, China (NE, NC, SW), Korean Peninsula.

Megalodontes thor Taeger, 2002. Host plant: Peucedanum oreoselinum (Apiaceae). Russia: **EP** (Taeger, 2002), **UR**, **WS** (AL). – Europe (WE, EE), Kazakhstan.

SUPERFAMILY CEPHOIDEA

One recent family: Cephidae.

10. FAMILY CEPHIDAE

Slender insects, with a long cylindrical or laterally compressed body. Slow on the wing.

Head with hypostomal bridge separating oral cavity from occipital foramen. Antenna long and thread-like or slightly clavate, 16-30-segmented. Hind margin of pronotum almost straight, being nearly the shortest distance between the tegulae; mesonotum short, never extending in front of the tegulae; cenchri absent. Tibiae usually with preapical spines on the hind and middle legs; fore tibia with only one modified apical spur. Fore wings with the vein M joining Rs after Rs has left $Sc+R$. Abdomen constricted slightly at the apex of the 1st segment. Female ovipositor exserted and clearly visible from above. Male often with some of the apical sternites fringed or with patches of modified setae; male genitalia orthandrous, parameres continuous with parameral plates, devoid of muscles and apical cupping discs, and with aedeagus tubular.

The larvae internal borers in stems or twigs of herbaceous plants, shrubs and trees; lead a secretive way of life.

The family is distributed worldwide. Number of taxa: World – 2 subfamilies, 24 genera, about 165 species; Palaearctic – 1/16/>120; Russia – 1/10/57.

R e f e r e n c e s. Dalla Torre, 1894; Konow, 1897; Kokujev, 1910; Dovnar-Zapskij, 1926a, 1928, 1931a; Enslin, 1927; Gussakovskij, 1935; Takeuchi, 1955; Er-molenko, 1971b, 1984; Muche, 1971a, 1975b; Stroganova, 1976, 1981, 1982, 1987; Zhelochovtsev, Prochorova, 1976; Verzhutskii, 1981; Zhelochovtsev, 1988; Liston, 1995; Yang, 1995; Zhelochovtsev, Zinovjev, 1995; Ivie, Zinovjev, 1996; Wei, Nie, 1997a; Naito et al., 1998; Shinohara, 1999; Ivie, 2001; Smith, Shinohara, 2002; Chen et al., 2004; Taeger, 2005; Liston et al., 2006, 2014a; Pavlusenko, 2006; Taeger et al., 2006, 2010; Wei et al., 2006; Vasilenko, 2007, 2010b, 2011b; Roller, Haris, 2008; Humala, Polevoi, 2009; Schedl, 2009; Wei, Smith, 2010; Wu, 2010; Blank, Taeger, 2011; Taeger, Blank, 2011; Macek, 2012; Sundukov, Lelej, 2012; Vasilenko, Korshunov, 2012; Beneš, 2014b; Kostyunin, Efimov, 2014; Liston, Prous, 2014; Beneš, Holuša, 2015; Kostyunin, 2015; Niu et al., 2015; Sundukov, 2015, 2017; Lesieur et al., 2016; Noblecourt, 2016.

Subfamily CEPHINAE

CAENOCEPHUS Konow, 1896. Type species

Caenocephus jakowleffi Konow, 1896. The genus is distributed in Holarctic and Oriental region. Number of species: World – 5; Palaearctic – 3; Russia – 2.

Caenocephalus lunulatus (Strobl, 1895) [Cephus] (*C. jakowleffi* Konow, 1896). Russia: **ES** (IR). – Europe (WE, EE).

Caenocephalus xanthopus Shinohara, 1999. Russia: **FE** (PR). – Korean Peninsula, Japan (Hon).

CALAMEUTA Konow, 1896 (*Monoplopus* Konow, 1896; *Haplocephalus* Benson, 1935; *Ephemerocephalus* Zhelochovtsev, 1968). Type species *Cephus filiformis* Eversmann, 1847. The larvae feed on the Poaceae. The genus is distributed in Holarctic and Oriental region. Number of species: World – 22; Palaearctic – 20; Russia – 10 from 2 subgenera.

Calameuta (Calameuta) bicornes Stroganova, 1976. Russia: **WS** (AL).

Calameuta (Calameuta) filiformis (Eversmann, 1847) [Cephus] (*Cephus elongatus* Snellen van Vollenhoven, 1858; *C. arundinis* Giraud, 1863; *C. marginatus* Kawall, 1864; *C. erberi* Damianiitsch, 1866; *C. vagabundus* Mocsáry, 1886; *C. infernalis* Dovnar-Zapskij, 1926; *Calameuta atrata* Dovnar-Zapskij, 1931; *C. rugosa* Dovnar-Zapskij, 1931; *C. amurensis* Gussakovskij, 1935). Host plant: Phragmites spp., Arrenatherum spp., Elytrigia spp., etc. (Poaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (NS, KM, AL), **ES** (IR), **FE** (KH, PR). – Europe (WE, EE, NE, SE), N Africa, Caucasus, Asia Minor, Iran, Kyrgyzstan, Kazakhstan, China (NW).

Calameuta (Calameuta) filum (Gussakovskij, 1935) [Cephus]. Host plant: Poaceae. Russia: **EP** (C, E, S, NC), **UR**, **WS** (NS, KM, AL), **ES** (KS, KR, IR). – Europe (WE, EE, NE, SE), Uzbekistan, Kazakhstan.

Clameuta (Calameuta) gracilicornis (Konow, 1896) [Cephus]. Russia: **EP** (NC). – Armenia, Azerbaijan.

Calameuta (Calameuta) haemorrhoidalis (Fabricius, 1781) [Tenthredo] (*Astatus analis* Klug, 1803; *A. floralis* Klug, 1803; *Trachelus haemorrhoidalis* Jurine, 1807; *Cephus atripes* Stephens, 1835; *C. quadriguttatus* Costa, 1882; *C. variabilis* Mocsáry, 1886; *C. quadriguttulatus* Dalla Torre, 1894; *C. quadrisignatus* Costa, 1894; *C. haemorrhoidalis signifer* Konow, 1896; *C. pseudotabidus* Kokujev, 1910; *C. diversipes* Ghigi, 1915; *Trachelus syriacus*

Pic, 1917). Host plant: Poaceae. Russia: **EP** (S, NC, CR). – Europe (WE, EE, SE), Caucasus, Asia Minor, Turkmenistan.

Calameuta (Calameuta) idolon (Rossi, 1794) [*Ichneumon*] (*Cephus mittrei* Guérin, 1844; *C. bellieri* Sichel, 1860; *C. variegatus* J.P.E.F. Stein, 1876; *Monoplopus apicicornis* Pic, 1916). Host plant: species of Poaceae. Russia: **EP** (CR). – Europe (EE, SE), N Africa, Caucasus, Asia Minor.

Calameuta (Calameuta) pallipes (Klug, 1803) [*Astata*] (*Cephus phthiscus* Fabricius, 1804; *C. pusillus* Stephens, 1835; *C. immaculatus* Stephens, 1835; *C. cultarius* Hartig, 1837; *C. pallidipes* Dalla Torre, 1894). Host plant: *Phleum* spp. (Poaceae). Russia: **EP** (NW, C, S), **UR**, **ES** (KS). – Europe (WE, EE, NE), Asia Minor, Iran, Kazakhstan.

Calameuta (Calameuta) punctata (Klug, 1803) [*Astata*] (*Cephus pravei* Dovnar-Zapskij, 1926). Russia: **EP** (NC). – Europe (SE), Georgia, Turkmenistan.

Calameuta (Calameuta) sachalinensis (Takeuchi, 1938) [*Cephus*]. Russia: **FE** (SA). – China (SW).

Calameuta (Ephemerocephus) kuzini Zhelochovtsev, 1968. Russia: **WS** (AL). – Kazakhstan.

CEPHUS Latreille, 1803 (*Astatus* Jurine, 1801; *Peronistilus* Ghigi, 1905; *Pironistilomorphus* Pic, 1916; *Fossulocephus* Pic, 1917; *Pseudocephus* Dovnar-Zapskij, 1931). Type species *Sirex pygmaeus* Linnaeus, 1767. The larvae feed on the Poaceae. The genus is distributed in Holarctic and Oriental region. Number of species: World – > 40; Palaearctic – about 40; Russia – 20.

Cephus brachycercus Thomson, 1871 (*C. punctulatus* Konow, 1896; *C. brachycercus tibialis* Dovnar-Zapskij, 1926). Host plant: Poaceae. Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (NS, KM, AL), **ES** (IR), **FE** (PR, SA, KU). – Europe (WE, EE, NE, SE), N Africa, Turkey, Uzbekistan, Kyrgyzstan, Kazakhstan, China (SW), Japan (Hon).

Cephus camtschatcalis Enslin, 1927. Russia: **FE** (KA).

Cephus daghestanicus (Dovnar-Zapskij, 1931) [*Pseudocephus*]. Russia: **EP** (NC).

Cephus excisus (Dovnar-Zapskij, 1931) [*Pseudocephus*]. Russia: **EP** (NC).

Cephus filippovi Zhelochovtsev, 1968. Russia: **ES** (ZB).

Cephus fumipennis Eversmann, 1847 (*C. carbonarius* Jakovlev, 1891). Host plant: *Agropyron* spp., ?*Phalaris* spp. (Poaceae). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (TK, NS, KM, AL). – Europe (WE, EE, NE, SE), N Africa, Caucasus, Turkey, Syria,

Tajikistan, Kazakhstan (SE), Mongolia, China (NW, CC, SW).

Cephus gracilis Costa, 1860 (*C. gracilis laterimaculatus* Strobl, 1901). Host plant: species of Poaceae. Russia: **EP** (S). – Europe (SE).

Cephus hyalinatus Konow, 1897. Host plant: species of Poaceae. Russia: **WS** (KM, AL), **ES** (KS, TU, KR, IR, BR), **FE** (AM, KH, PR, SA, KU, KA). – Mongolia, Korean Peninsula, Japan (Hok).

Cephus infuscatus Thomson, 1871 (*C. infuscatus* André, 1881; *C. andreae* W.F. Kirby, 1882). Host plant: *Phalaris arundinacea* (Poaceae). Russia: **EP** (NW, C). – Europe (WE, EE, NE), China (NE, CC).

Cephus nigrinus Thomson, 1871. Host plant: *Milium* spp., *Poa* spp. (Poaceae). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (NS, KM, AL), **ES** (IR). – Europe (WE, EE, NE), Turkey, Syria, Iran, Kazakhstan.

Cephus nigripennis (Takeuchi, 1927) [*Eumetabolus*] (*Monoplopus japonicus* Forsius, 1928; *Cephus graminis* Maa, 1944). Host plant: *Elytrigia* spp., *Agropyron* spp. (Poaceae). Russia: **ES** (IR), **FE** (PR). – China (NE, CC, SE), Korean Peninsula, Japan (Hon, Shi, Tsushima).

Cephus parvus (Dovnar-Zapskij, 1931) [*Pseudocephus*]. Russia: **EP** (NC).

Cephus pseudopilosulus Dovnar-Zapskij, 1926. Russia: **EP** (NC), **WS** (AL).

Cephus pulcher Tischbein, 1852 (*C. luteomarginatus* Giraud, 1857). Host plant: *Agropyron* spp. and others (Poaceae). Russia: **EP** (S, NC, CR), **WS** (NS, KM, AL). – Europe (EE, SE), Turkey, Uzbekistan, Kazakhstan.

Cephus pygmeus (Linnaeus, 1767) [*Sirex*] (*Tenthredo polygona* Gmelin, 1790; *Banchus viridator* Fabricius, 1804; *Cephus subcylindricus* Gravenhorst, 1807; *C. leskii* Lepeletier, 1823; *C. flavisternum* Costa, 1882; *C. clypearis* Costa, 1894; *C. pygmaeus palaestinus* Pic, 1918; *C. tanaiticus* Dovnar-Zapskij, 1926). Host plant: *Agropyron* spp., *Avena* spp., *Phleum* spp., *Triticum* spp., etc. (Poaceae). Russia: **EP** (C, NC, CR), **UR**, **WS** (NS, KM, AL). – Europe (WE, EE, NE, SE), N Africa, Caucasus, SW Asia, Iran, Central Asia, China (CC); introduced into N America.

Cephus rjabovi Dovnar-Zapskij, 1926. Russia: **EP** (S, NC).

Cephus sareptanus Dovnar-Zapskij, 1928. Russia: **EP** (S). – Kazakhstan.

Cephus spinipes (Panzer, 1800) [*Banchus*] (*C. cultratus* Eversmann, 1847; *C. pilosulus* Thomson, 1871;

C. pumilus André, 1881; *C. similis* Mocsáry, 1886; *C. affinis* Kokujev, 1910; *C. exilis* Kokujev, 1910; *C. pilosulus incompletus* Hellén, 1935). Host plant: Phleum spp., Dactylis spp. (Poaceae). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (NS, KM, AL). – Europe (WE, EE, NE, SE), N Africa, Armenia, Kyrgyzstan, Kazakhstan, China (NW).

Cephus tshettui Stroganova, 1981. Russia: **WS** (AL). **Cephus zahaikevitshi** (Ermolenko, 1971) [Pseudocephus]. Russia: **EP** (CR).

CHARACOPYGUS Konow, 1899. Type species *Characopygus moricei* Konow, 1899. The larvae feed on the Asteraceae. The genus is distributed in West Palaearctic. Number of species: World and Palaearctic – 4; Russia – 2.

Characopygus modestus Dovnar-Zapolskij, 1931. Russia: **UR**, **WS** (KM), **ES** (IR).

Characopygus scythicus Dovnar-Zapolskij, 1931. Host plant: Centaurea spp. (Asteraceae). Russia: **EP** (S), **UR**. – Europe (EE), Armenia, Turkey.

JANUS Stephens, 1829 (*Phylloecus* Newman, 1838; *Ephippionotus* A. Costa, 1860). Type species *Janus connectens* Stephens, 1829. The larvae in the branches of trees and shrubs of Rosaceae, Salicaceae, Fagaceae, Adoxaceae. The genus is distributed in Holarctic and Oriental region. Number of species: World – 26; Palaearctic – 13; Russia – 4.

Janus compressus (Fabricius, 1793) [Sirex] (*Cephus flaviventris* Förster, 1844; *Ephippionotus luteiventris* Costa, 1860; *Phylloecus eburneus* André, 1881; *Cephus foersteri* André, 1881; *C. balearicus* Kriechbaumer, 1894; *Janus foersteri ulmi* Pic, 1918). Host plant: Pyrus spp., Crataegus spp., Malus spp. (Rosaceae). Russia: **EP** (NC). – Europe (WE, EE, SE), N Africa, Turkey.

Janus cynosbati (Linnaeus, 1758) [Tenthredo] (*Lyda geoffroyi* Lepeletier, 1823; *Cephus femoratus* Curtis, 1830; *Phylloecus faunus* Newman, 1838; *Ephippionotus cephalotes* Costa, 1860; *Cephus melanarius* Mocsáry, 1891; *C. niger* Brischke, 1892). Host plant: Quercus pubescens, Q. robur (Fagaceae). Russia: **EP** (NW, C). – Europe (WE, EE, NE, SE), Turkey.

Janus formosus (Zhelochovtsev, 1935) [Cephus] (*J. orientalis* Gussakovskij, 1935). Russia: **WS** (NS, AL), **FE** (PR). – China (NE).

Janus luteipes (Lepeletier, 1823) [Cephus] (*J. connectens* Stephens, 1829; *J. bifrons* Newman, 1838; *Cephus emichi* Mocsáry, 1886; *Janus luteipes diversipes*

Pic, 1918). Host plant: Salix spp., Populus spp. (Salicaceae), ?Viburnum spp. (Adoxaceae). Russia: **EP** (NW, C), **FE** (PR). – Europe (WE, EE, NE, SE), N Africa, Georgia.

PACHYCEPHUS J.P.E.F. Stein, 1876. Type species *Pachycephus smyrnensis* J.P.E.F. Stein, 1876. The genus is distributed in West Palaearctic. Number of species: World and Palaearctic – 2; Russia – 2.

Pachycephus cruentatus cruentatus (Eversmann, 1847) [Cephus] (*P. caspicus* Dovnar-Zapolskij, 1931). Russia: **EP** (S, NC, CR). – Europe (EE), Caucasus, Turkey.

Pachycephus smyrnensis smyrnensis J.P.E.F. Stein, 1876 (*P. aeneovarius* Kohl, 1905; *P. brevis* Ghigi, 1915; *Spatulocephus sanctus* Pic, 1916). Russia: **EP** (NC). – Europe (WE, SE), N Africa, Armenia, Azerbaijan, SW Asia.

PHYLLOECUS Newman, 1838 (*Hartigia* Schiødte, 1839; *Cerobactrus* Costa, 1860; *Macrocephus* Schlechtendal, 1878; *Cephosoma* Gradl, 1881; *Adirus* Konow, 1899; *Paradirus* Dovnar-Zapolskij, 1931; *Hissarocephus* Gussakovskij, 1945). Type species *Phylloecus faunus* Newman, 1838. The larvae in the branches of Rosaceae. The genus is distributed in Holarctic, Oriental and Neotropical regions; mainly in Palaearctic. Number of species: World – about 30; Palaearctic – > 20; Russia – 9.

Phylloecus agilis (F. Smith, 1874) [Cephus] (*Hartigia draconis* Maa, 1944; *H. draconis bipunctata* Maa, 1944; *H. draconis collaris* Maa, 1944). Russia: **FE** (PR, SA). – China (CC, SE), Japan (Hon, Shi, Kyu).

Phylloecus coreanus (Takeuchi, 1938) [Hartigia]. Russia: **FE** (PR). – Korean Peninsula.

Phylloecus etorofensis (Takeuchi, 1955) [Hartigia]. Host plant: Rosa spp. (Rosaceae). Russia: **EP** (C, E), **WS** (NS, KM), **ES** (TU, YA), **FE** (KU). – Europe (NE).

Phylloecus linearis (Schrank, 1781) [Tenthredo] (*Cephus quinquefasciatus* Stephens, 1835; *C. major* Eversmann, 1847; *Macrocephus agrimoniae* Goury, 1912). Host plant: Agrimonia spp. (Rosaceae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (NS, KM, AL), **ES** (IR, ZB). – Europe (WE, EE, NE, SE), N Africa, Georgia, Turkey, Kyrgyzstan, Kazakhstan.

Phylloecus niger (M. Harris, 1779) [Sirex] (*Astatas satyrus* Panzer, 1801; *Cerobactrus major* Costa,

1860; *Cephus brachypterus* Damianitsch, 1866; *C. glabellifer* Thomson, 1871; *Cephosoma syringae* Grall, 1881; *Phylloecus cruciatus* Costa, 1894). Host plant: Rubus spp. and others (Rosaceae). Russia: **EP** (S, NC, CR), **UR**, **WS** (NS, KM). – Europe (WE, EE, SE), N Africa, Armenia, Turkey, Syria, Iran, Kazakhstan, China (NC, CC).

Phylloecus pyrrha (Zhelochovtsev, 1968) [Hartigia]. Russia: **ES** (ZB).

Phylloecus sibiricola Jakovlev, 1891 (*P. cylindricus* Jakovlev, 1891; *Hartigia affinis* Dovnar-Zapolskij, 1931). Host plant: Rosa spp. and others (Rosaceae). Russia: **WS** (TK, NS, KM, AL), **ES** (IR), **FE** (AM, PR). – Kazakhstan, Mongolia.

Phylloecus viator (F. Smith, 1874) [Cephus] (*Macrocephus japonicus* Ghigi, 1915). Host plant: Rubus spp., Rosa spp. (Rosaceae). Russia: **FE** (PR). – China (NE, NC, SE), Korean Peninsula, Japan (Hok, Hon).

Phylloecus xanthostoma (Eversmann, 1847) [Cephus] (*Cerobactrus facialis* Costa, 1864; *Macrocephus ulmariae* Schlechtendal, 1878; *Phyllaecus giraudi* Schlechtendal, 1880; *Hartigia semenovi* Dovnar-Zapolskij, 1931; *H. jakovlevi* Dovnar-Zapolskij, 1931). Host plant: Filipendula spp. (Rosaceae). Russia: **EP** (S, E), **UR**. – Europe (WE, EE, NE), N Africa, Turkey.

STENOCEPHUS Shinohara, 1999. Type species *Stenocephus oncogaster* Shinohara, 1999. The genus is distributed in Palaearctic and Oriental region. Number of species: World – 2; Palaearctic – 1; Russia – 1.

Stenocephus oncogaster Shinohara, 1999. Russia: **FE** (KU). – China (CC), Korean Peninsula, Japan (Hok, Hon).

SYRISTA Konow, 1896 (*Neosyrista* Benson, 1935). Type species *Cephus parreyssi* Spinola, 1843. The larvae in the branches of Rosaceae. The genus is distributed in Palaearctic and Oriental region. Number of species: World – 3; Palaearctic – 2; Russia – 2.

Syrista parreyssii (Spinola, 1843) [Cephus] (*Cephus orientalis* Tischbein, 1852; *C. spectabilis*

J.P.E.F. Stein, 1876; *Macrocephus robustus* Mocsáry, 1883; *Cephus parreyssi rufiventris* Jakovlev, 1888). Host plant: Rosa spp. (Rosaceae). Russia: **EP** (NC). – Europe (SE), Caucasus, SW Asia, Iran.

Syrista similis Mocsáry, 1904 (*Cephus subrufa* Matsamura, 1912; *Neosyrista japonica* Benson, 1935). Host plant: Rosa spp. (Rosaceae). Russia: **FE** (KU). – China (NC, CC), Japan (Hok, Hon, Shi, Kyu).

TRACHELUS Jurine, 1807 (*Astatus* Panzer, 1801; *Cepha* Billberg, 1820; *Ateuchopus* Konow, 1896; *Eumetabolus* Schulz, 1906; *Trachelastatus* Morice et Durrant, 1915; *Microcephus* Benson, 1935; *Neateuchopus* Benson, 1935). Type species *Sirex tabidus* Fabricius, 1775. The genus is distributed in Palaearctic; introduced into the Nearctic. Number of species: World and Palaearctic – 8; Russia – 5.

Trachelus libanensis (André, 1881) [Cephus] (*Cephus nigritarsis* André, 1881; *Ateuchopus armenius* Konow, 1896). Russia: **EP** (NC). – Georgia, Armenia, SW Asia.

Trachelus pallipes Stroganova, 1976. Russia: **UR**. – Kazakhstan (NW).

Trachelus tabidus (Fabricius, 1775) [Sirex] (*Sirex macilentus* Fabricius, 1793; *Cephus mandibularis* Serville, 1823; *C. nigritus* Serville, 1823; *C. mandibularis* Lepeletier, 1823; *C. nigritus* Lepeletier, 1823; *C. vittatus* Costa, 1875; *Calameuta johnsonii* Ashmead, 1903). Host plant: Triticum spp., Secale cereale, Hordeum spp., Avena spp. (Poaceae); dangerous pest. Russia: **EP** (S, NC, CR). – Europe (WE, EE, SE), N Africa, Caucasus, SW Asia; introduced into N America.

Trachelus tigris (Benson, 1935) [Neateuchopus]. Russia: **EP** (S).

Trachelus troglodyta (Fabricius, 1787) [Sirex] (*Cephus mocsaryi* W.F. Kirby, 1882; *Astatus tenuicornis* Konow, 1902). Host plant: Secale cereale (Poaceae); dangerous pest. Russia: **EP** (NW, C, E, S, NC). – Europe (WE, EE, NE, SE), N Africa, Georgia, SW Asia, Kazakhstan.

SUPERFAMILY SIRICOIDEA

Head with hypostomal bridge separating oral cavity from occipital foramen behind; labrum spatulate. Pronotum short medially and emarginate behind; mesosternum triangular, usually with presternal bridge in front, though the sutures are usually obsolete. Male genitalia orthandrious; parameres articulating on the parameral plates and moved by special muscles, and with apical cupping discs. Female ovipositor strongly exserted. Larva of reduced oligopod type. The superfamily is distributed worldwide. Number of taxa: World – 2 recent families, 4 subfamilies, 40 genera, about 260 species; Palaearctic – 2/3/21/about 105; Russia – 2/3/9/41 (+ 1 ssp.).

11. FAMILY SIRICIDAE

Large insects, generally over 14 mm, with exserted ovipositors, and differing from all other Hymenoptera in their minute tegulae. Mouthparts with maxillary palp 1-segmented; labial palp 2- to 3-segmented, the last segment enlarged and bearing a large apical sensory cup, and the 1st segment not enlarged. Antennae setaceous, 17–30-segmented (4–8-segmented in *Teredon* Norton), with a long curved 1st segment at least as long as the 3rd. Neck short. Pronotum strongly emarginate behind. Mesonotum not divided transversely, but with a lateral lobe divided off each side of the scutum. Fore wings with the membrane corrugated at the apex and with a large apical appendicular cell. Tibiae without preapical spines, and front tibia with only one apical spur. Abdomen cylindrical and contracted at base of 1st segment, which is medially divided; last segment with a horn-like projection, called the cornus. The larvae are white with only vestigial thoracic legs. They live in the wood of coniferous and deciduous trees; generation is usually biennial. Many species are technical pests of wood.

The family is distributed worldwide. Number of taxa: World – 2 subfamilies, 11 genera, > 130 species; Palaearctic – 2/7/75; Russia – 2/5/20 (+ 1 ssp.).

R e f e r e n c e s. Jakovlev, 1900; Semenov, 1921; Gussakovskij, 1935; Semenov, Gussakovskij, 1935; Maa, 1949a; Benson, 1965a; Stroganova, 1968; Verzhutskii, 1981; Khomentovskij, 1983; Zhelochovtsev, 1988; Zhelezovtsev, Zinovjev, 1992, 1996; Blank, 1996; Vasilenko, 2004, 2011b; Liston et al., 2006, 2014a; Taeger et al., 2006, 2010; Wei et al., 2006; Lelej, Taeger, 2007a; Roller, Haris, 2008; Schedl, 2009; Taeger, Blank, 2011;

Schiff et al., 2012; Sundukov, Lelej, 2012; Vasilenko, Korshunov, 2012; Ahn et al., 2013; Beneš, 2014b; Goulet et al., 2015; Kostyunin, 2015; Sundukov, 2015; Noblecourt, 2016.

Subfamily SIRICINAE

SIREX Linnaeus, 1760 (*Neuropachys* Heer, 1858; *Urocerites* Heer, 1867; *Paururus* Konow, 1896). Type species *Ichneumon juvencus* Linnaeus, 1758. The larvae feed on coniferous wood of Pinaceae. The genus is distributed in Holarctic and Oriental region; introduced into the Neotropical, Ephiopian and Australian regions. Number of species: World – 28; Palaearctic – 15; Russia – 6.

Sirex atricornis Kjellander, 1945. Russia: **WS** (NS). – Europe (NE).

Sirex cyaneus dux (Semenov, 1921) [*Paururus*]. Host plant: ?*Abies nordmanniana* (Pinaceae). Russia: **EP** (Benson, 1965a). – Europe (EE), Georgia, Armenia, Israel.

Sirex ermak (Semenov, 1921) [*Paururus*]. Host plant: *Pinus* spp., *Picea* spp., *Abies* spp., *Larix* spp. (Pinaceae). Russia: **EP** (C), **WS** (TK, NS, KM, AL), **ES** (KS, TU, IR, BR, YA, ZB), **FE** (AM, PR, SA, KU, KA). – Kazakhstan, Mongolia, China (NE).

Sirex juvencus juvencus (Linnaeus, 1758) [*Ichneumon*] (*Urocerus feisthameli* Brullé, 1832; *Sirex dubia* W.F. Kirby, 1882; *S. australis* W.F. Kirby, 1882; *S. neglectus* W.F. Kirby, 1882; *S. hirsutus* W.F. Kirby, 1882; *S. leseleuci* Tournier, 1890; *Paururus juvencus virago* Enslin, 1918; *P. sucineiceps* Koornneef, 1935; *P. juvencus maculata* Hellén, 1935). Host plant: *Pinus* spp., *Picea* spp., *Abies* spp., *Larix* spp. (Pinaceae). Russia: **EP** (N, C, E, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (TU, KR, IR, BR), **FE** (KH, PR, SA, KA). – Europe (WE, EE, NE, SE), Georgia, Mongolia, China (NE, CC, SE), Japan (Hok, Hon), India; introduced into N America and Australia.

Sirex mongolorum (Semenov et Gussakovskij, 1935) [*Paururus*]. Host plant: *Abies* spp., *Pinus* spp., *Picea* spp. (Pinaceae). Russia: **ES** (BR, ZB). – Mongolia.

Sirex noctilio Fabricius, 1793 (*S. melanocerus* Thomson, 1871; *Paururus atlantidis* Ghigi, 1909). Host plant: *Pinus* spp., *Picea* spp., *Abies* spp., *Larix* spp. (Pinaceae). Russia: **EP** (N, NW, C, E, NC, CR), **UR**, **WS** (NS, KM, AL), **ES** (TU, KR, IR, BR, YA), **FE** (AM, KH, MG). – Europe (WE, EE, NE, SE), N

Africa, Turkey, Kazakhstan, Mongolia; introduced into N and S America, Australia, New Zealand, S Africa.

UROKERUS Geoffroy, 1762 (*Xanthosirex* Semenov, 1921; *Eosirex* Piton, 1940). Type species *Ichneumon gigas* Linnaeus, 1758. The larvae mainly feed on coniferous wood of Pinaceae. The genus is distributed in Holarctic and Oriental region; one species introduced into the Neotropical region. Number of species: World – 33; Palaearctic – 28; Russia – 6 (+ 1 ssp.).

Urocerus antennatus (Marlatt, 1898) [Sirex] (*Sirex ogumae* Matsumura, 1911; *S. antennatus immaculatus* Zirngiebl, 1937). Host plant: *Abies* spp., *Pinus* spp., *Picea* spp. (Pinaceae), *Cryptomeria* spp. (Cupressaceae). Russia: **ES** (ZB), **FE** (PR, SA, KU). – China (NE), Korean Peninsula, Japan (Hok, Hon, Shi).

Urocerus argonautarum (Semenov, 1921) [Sirex]. Host plant: *Abies nordmanniana*, *Picea* spp. (Pinaceae). Russia: **EP** (NC). – Georgia, Turkey.

Urocerus fantoma (Fabricius, 1781) [Sirex] (*Sirex tardigradus* Cederhjelm, 1798; *Xanthosirex phantasma* Semenov, 1921; *Sirex augur pallida* Ermolenko, 1957). Host plant: *Pinus* spp., *Picea* spp., *Abies* spp., *Larix* spp. (Pinaceae). Russia: **EP** (NW, C), **WS** (TM, KM), **ES** (KR, IR, ZB), **FE** (PR). – Europe (WE, EE, NE, SE), Kazakhstan, Mongolia.

Urocerus gigas gigas (Linnaeus, 1758) [*Ichneumon*] (*Sirex marisca* Linnaeus, 1760; *S. hungaricus* Christ, 1791; *S. psylluis* Fabricius, 1793; *Urocerus lefebvre* Guérin, 1833; *Sirex faustus* Costa, 1894; *Urocerus gigas taiganus* Benson, 1943; *U. gigas luteogaster* Berland, 1947; *Sirex gigas montana* Ermolenko, 1957). Host plant: *Pinus* spp., *Picea* spp., *Abies* spp. and others (Pinaceae), *Populus* spp., *Salix* spp. (Salicaceae), *Betula* spp. (Betulaceae), etc. Russia: **EP** (N, NW, C, S, NC, CR), **UR**, **WS** (TM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, SA, KA, MG, CH). – Europe (WE, EE, NE, SE), SW Asia, Kazakhstan, Mongolia, China (NC, NW), Korean Peninsula; introduced into S America.

Urocerus gigas orientalis Maa, 1949. Host plant: *Abies* spp., *Picea* spp. and others (Pinaceae). Russia: **ES** (ZB), **FE** (SA, KA). – China (CC), Korean Peninsula, Japan (Hok).

Urocerus sah (Mocsáry, 1881) [Sirex]. Host plant: *Juniperus* spp. (Cupressaceae). Russia: **EP** (NC,

CR). – Europe (EE, NE), Azerbaijan, Turkey, Afghanistan, Kazakhstan; introduced into N America.

Urocerus yasushii (Yano, 1917) [Sirex] (*Sirex umbra* Semenov, 1921; *Urocerus sachalinensis* Conde, 1935). Host plant: *Picea* spp. (Pinaceae). Russia: **ES** (ZB), **FE** (KH, PR, SA, KU). – China (NE), Korean Peninsula, Japan (Hon).

XOANON Semenov, 1921. Type species *Xoanon mysta* Semenov, 1921. The genus is distributed in East Asia and Oriental region. Number of species: World – 2; Palaearctic – 1; Russia – 1.

Xoanon matsumurae (Rohwer, 1910) [Sirex] (*X. mysta* Semenov, 1921). Host plant: *Pinus* spp., *Picea* spp., *Abies* spp. (Pinaceae). Russia: **ES** (BR, ZB), **FE** (PR, SA, KU). – China (NE, NC), Korean Peninsula, Japan (Hok, Hon, Shi).

Subfamily TREMICINAE

TREMEX Jurine, 1807 (*Xyloterus* Hartig, 1837; *Xyloecematum* Heyden, 1868). Type species *Sirex fuscicornis* Fabricius, 1787. The larvae feed the wood of deciduous trees. The genus is distributed in Holarctic and Oriental regions; one species was introduced into the Neotropical region. Number of species: World – 33; Palaearctic – 21; Russia – 5.

Tremex alchymista Mocsáry, 1886. Russia: **EP** (NC). – Europe (WE, EE).

Tremex apicalis Matsumura, 1912 (*T. propheta* Semenov, 1921). Host plant: *Acer* spp. (Sapindaceae), *Prunus* spp. (Rosaceae). Russia: **ES** (ZB), **FE** (SA). – China (NE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Tremex fuscicornis (Fabricius, 1787) [Sirex] (*Sirex struthiocamelus* Villers, 1789; *S. camelogigas* Christ, 1791). Host plant: *Betula* spp. (Betulaceae), *Populus* spp. (Salicaceae), *Robinia* spp. (Fabaceae), *Acer* spp. (Sapindaceae), *Ulmus* spp. (Ulmaceae). Russia: **EP** (NW, C, E, S, NC, CR), **WS** (TM, TK, NS, KM, AL), **ES** (BR, ZB), **FE** (AM, PR, SA, KU). – Europe (WE, EE, NE, SE), Armenia, Kazakhstan, China (NE, NC, SW, SE), Korean Peninsula, Japan (Hok, Hon); introduced into S America.

Tremex magus magus (Fabricius, 1787) [Sirex] (*Sirex nigrita* Fabricius, 1787). Host plant: *Betula* spp. (Betulaceae), *Populus* spp. (Salicaceae). Russia: **EP** (C, E, S), **UR**, **WS** (KM, AL). – Europe (WE, EE, SE), Kazakhstan.

Tremex satanas Semenov, 1921. Russia: **ES** (BR), **FE** (PR).

XERIS Costa, 1894 (*Neoxeris* Saini et Singh, 1987).

Type species *Ichneumon spectrum* Linnaeus, 1758. The larvae feed on coniferous wood (Pinaceae, Cupressaceae). The genus is distributed in Holarctic, Oriental and Neotropical regions. Number of species: World – 16; Palaearctic – 8; Russia – 2.

Xeris malaisei Maa, 1949. Host plant: *Abies* spp. (Pinaceae), *Cryptomeria* spp. (Cupressaceae). Russia:

FE (PR). – China (NE), Korean Peninsula, Japan (Hok, Hon).

Xeris spectrum (Linnaeus, 1758) [Ichneumon] (*Sirex nanus* O.F. Müller, 1776; *S. emarginatus* Fabricius, 1793). Host plant: *Pinus* spp., *Picea* spp., *Abies* spp., *Larix* spp. (Pinaceae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TM, NS, KM, AL), **ES** (TU, KR, BR, ZB), **FE** (AM, PR, SA, KU). – Europe (WE, EE, NE, SE), Georgia, SW Asia, Kazakhstan, Japan (Hok, Hon, Shi, Kyu).

12. FAMILY XIPHYDRIIDAE

Large or medium insects 7–20 mm long. Antennae setaceous and 13–19 segmented, with a long curved 1st segment at least as long as 3rd. Neck long; cervical sclerites viewed from the side appear longer than broad. Mesonotum with a transverse furrow. Fore wings corrugated apically, and without an appendiculate cell; vein r_2+r_3 present, and base of Rs not angled sharply near its base, so that its basal portion is not in a line with M . Tibiae without preapical spines; front tibia with only one apical spur. Last abdominal segment without an apical horn-like projection.

The larvae white with only vestigial thoracic legs; and they bore in the wood of deciduous trees, mainly Betulaceae, Salicaceae and Ulmaceae.

The family is distributed worldwide, except of Afro-tropical region. Number of taxa: World – 2 subfamilies, 29 genera, about 150 species; Palaearctic – 1/14/50; Russia – 1/4/21.

References. Jakovlev, 1900; Takeuchi, 1938; Gussakovskij, 1947; Maa, 1949a; Stroganova, 1968; Zhelochovtsev, 1968, 1988; Ermolenko, 1979; Verzhutskii, 1981; Zhelochovtsev, Zinovjev, 1996; Togashi, 1998; Liston et al., 2006, 2014a; Taeger et al., 2006, 2010; Wei et al., 2006; Shcherbakov, 2008; Smith, 2008; Humala, Polevoi, 2009; Schedl, 2009; Smith et al., 2011; Smith, Shinohara, 2011; Sundukov, 2011a, 2015; Taeger, Blank, 2011; Vasilenko, 2011b; Sundukov, Lelej, 2012; Vasilenko, Korshunov, 2012; Beneš, 2014b; Kostyunin, 2015; Noblecourt, 2016.

Subfamily XIPHYDRIINAE

EUXIPHYDRIA Semenov et Gussakovskij, 1935.

Type species *Xiphydria potanini* Jakovlev, 1891. The larvae feed the wood of deciduous trees. The genus is distributed in East Asia and Oriental region. Number of species: World – 6; Palaearctic – 4; Russia – 2.

Euxiphydria amphybolia Sundukov, 2012. Host plant: *Phellodendron amurense* (Rutaceae), *Betula mandshurica* (Betulaceae), *Acer ukurunduense* (Sapindaceae). Russia: **FE** (PR, SA, KU).

Euxiphydria potanini (Jakovlev, 1891) [Xiphydria] (*Xiphydria ruficeps* Mocsáry, 1909; *X. ruficeps* Matsumura, 1912; *X. akazui* Matsumura, 1932; *X. maidli* Zirngiebl, 1937; *Euxiphydria subtrifida* Maa, 1944). Host plant: *Acer mono*, *A. mayrpii*

(Sapindaceae), *Betula ermanii* (Betulaceae). Russia: **FE** (KH, PR, SA, KU). – China (NE, NC, SW, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Sado).

PLATYXIPHYDRIA Takeuchi, 1938. Type species *Platyxiphydria tiphiformis* Takeuchi, 1938. The larvae feed the wood of deciduous trees. The genus is distributed in East Asia; one species was introduced into the Oriental region. Number of species: World – 8; Palaearctic – 7; Russia – 1.

Platyxiphydria sikhotealinensis Sundukov, 2012. Host plant: *Tilia amurensis* (Tiliaceae). Russia: **FE** (PR).

XIPHYDRIA Latreille, 1803 (*Urocerus* Jurine, 1801; *Hybonotus* Klug, 1803; *Xiphydria* Panzer, 1806; *Xiphiura* Fallén, 1813; *Xiphydria* Norton, 1869; *Xiphidion* Provancher, 1875; *Konowia* Brauns, 1884; *Pseudoxiphydria* Enslin, 1911; *Apoxiphia* Maa, 1949; *Nasoxiphia* Maa, 1949). Type species *Ichneumon camelus* Linnaeus, 1758. The larvae feed the wood of deciduous trees. The genus is distributed in Holarctic and Oriental regions. Number of species: World – 36; Palaearctic – 22; Russia – 17.

Xiphydria alnivora Matsumura, 1927. Host plant: *Alnus hirsuta* (Betulaceae). Russia: **FE** (PR, KU). – Japan (Hok, Hon).

Xiphydria annulitibia Takeuchi, 1936. Host plant: *Alnus hirsuta* (Betulaceae). Russia: **FE** (SA, KU). – Korean Peninsula, Japan (Hok, Hon, Shi).

Xiphydria betulae (Enslin, 1911) [*Pseudoxiphydria*] (*Pseudoxiphydria markewitschi* Ermolenko, 1960; *Konowia guntionensis* Zombori, 1960). Host plant: *Betula* spp. (Betulaceae). Russia: **EP** (C, E), **UR**, **WS** (NS, KM, AL), **ES** (KR), **FE** (AM, KH). – Europe (WE, EE, NE, SE).

Xiphydria buyssoni Konow, 1903. Host plant: *Alnus hirsuta* (Betulaceae). Russia: **ES** (ZB), **FE** (PR). – Japan (Hon, Shi, Kyu).

Xiphydria camelus (Linnaeus, 1758) [*Ichneumon*] (*X. eborata* Konow, 1899; *X. jozana* Matsumura, 1927; *X. kawakamii* Matsumura, 1927; *X. kuccharonis* Matsumura, 1927). Host plant: *Alnus* spp., *Betula* spp. (Betulaceae), *Populus* spp. (Salicaceae), *Ulmus* spp. (Ulmaceae) and others. Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, KR, IR), **FE** (AM, KH, PR, SA, KU, KA). – Europe (WE, EE, NE, SE), Turkey, Kazakhstan, China (NE, NC), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

- Xiphydria caucasica** Semenov et Gussakovskij, 1935.
Russia: **EP** (NC). – Georgia.
- Xiphydria jakovlevi** Semenov et Gussakovskij, 1935.
Host plant: *Alnus hirsuta* (Betulaceae). Russia: **FE** (PR, KU). – China (NE, CC, SE), Korean Peninsula.
- Xiphydria kastsheevi** Ermolenko, 1979. Russia: **FE** (PR).
- Xiphydria longicollis** (Geoffroy, 1785) [Tenthredo] (*Urocerus annulatus* Jurine, 1807). Host plant: *Quercus* spp. (Fagaceae), *Betula* spp. (Betulaceae), *Acer* spp. (Sapindaceae) and others. Russia: **EP** (C, CR). – Europe (WE, EE, SE).
- Xiphydria megapolitana** (Brauns, 1884) [Konowia].
Host plant: *Betula* spp. (Betulaceae). Russia: **EP** (NW, C), **ES** (BR), **FE** (PR). – Europe (WE, EE, NE, SE).
- Xiphydria ogasawarai** Matsumura, 1927. Host plant:
Acer mayrii, *A. mono* (Sapindaceae). Russia: **FE** (PR, KU). – Japan (Hok, Hon).
- Xiphydria palaearctica** Semenov, 1921 (*X. jezoensis* Matsumura, 1927). Host plant: *Alnus* spp., *Betula* spp. (Betulaceae), *Acer* spp. (Sapindaceae), *Ulmus* spp. (Ulmaceae) and others. Russia: **ES** (IR), **FE** (PR, SA, KU). – China (NE), Korean Peninsula, Japan (Hok, Hon, Sado).
- Xiphydria picta** Konow, 1897 (*Konowia bifurcata* Maa, 1949). Host plant: *Alnus* spp. (Betulaceae). Russia:

- EP** (NW, C, E, S, NC, CR). – Europe (WE, EE, NE, SE), Georgia, Kazakhstan.
- Xiphydria popovi** Semenov et Gussakovskij, 1935. Host plant: *Alnus* spp., *Betula* spp. (Betulaceae). Russia: **ES** (TU, ZB), **FE** (AM, KH, PR). – China (NE).
- Xiphydria prolongata** (Geoffroy, 1785) [Tenthredo] (*Sirex dromedarius* Fabricius, 1787; *Xyphidria fasciata* Lepeletier, 1823). Host plant: *Betula* spp. (Betulaceae), *Salix* spp., *Populus* spp. (Salicaceae), *Ulmus* spp. (Ulmaceae). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (TK, NS, KM, AL). – Europe (WE, EE, NE, SE), Azerbaijan, Turkmenistan, Kazakhstan, Mongolia.
- Xiphydria ribesii** (Zhelochovtsev, 1968) [Euxiphidria]. Host plant: *Ribes* spp. (Grossulariaceae). Russia: **WS** (TK).
- Xiphydria sulcata** Maa, 1949. Russia: **FE** (SA).
- XIPHYDRIOLA** Semenov, 1921. Type species
Xiphydriola amurensis Semenov, 1921. The genus is distributed in East Asia. Number of species: World and Palaearctic – 2; Russia – 1.
- Xiphydriola amurensis** Semenov, 1921. Russia: **FE** (PR). – China (NE), Korean Peninsula.

INFRAORDER ORUSSOMORPHA
SUPERFAMILY ORUSSOIDEA

One family: Orussidae.

13. FAMILY ORUSSIDAE

Very rare insects, characterized by the insetrion of the antennae on the ventral aspect of the head, below the apparent clypeus and below the lower margins od the eyes; the antenna is 11-segmented and setiform in the male, but in the female is only 10-segmented, with the 9th segment unique in form, swollen and longer than any other segment; head behind with a hypostomal bridge separating the oral cavity from the occipital foramen; labrum spatulate. Pronotum strongly emarginate behind; mesosternum triangular without presternal bridge and with sutures obsolete. The wing-venation is reduced mostly to pigmented bands; there is no vein $2r$ in the fore wing and no cross veins $r-m$ or $m-cu$ in the hind wings. Male genitalia orthandrious; parameres continuous with parameral plates, without muscles and whitout apical cupping discs; aedeagus trough-like. Female sawsheath leveled, not exserted.

The Orussidae is a small family of parasitic wasps. Their biology is incompletely known, but the majority of the evidence bearing on the lifestyle of Orussidae indicates that they are idiobiont ectoparasites of wood-boring beetle larvae, primarily Buprestidae and Sircidae and Xiphydriidae also (Vilhelmsen et al., 2013).

The family is distributed worldwide. Number of taxa: World – 2 subfamilies, 17 genera and about 90 species; Palaearctic – 2/5/about 20; Russia – 1/2/4.

References. Conde, 1935; Zhelochovtsev, 1988; Kraus, 1998; Blank et al., 2006; Taeger et al., 2006, 2010; Togashi, 2008; Schedl, 2009; Taeger, Blank, 2011; Zombori, 2013; Beneš, 2014b; Choi et al., 2014; Sundukov, 2014, 2017; Noblecourt, 2016.

Subfamily ORUSSINAE

ORUSSUS Latreille, 1797. Type species *Oryssus coronatus* Fabricius, 1798. The genus is distributed worldwide, except Neotropical region. Number of species: World – 28; Palaearctic – 14; Russia – 3.

Orussus abietinus (Scopoli, 1763) [Sphecius] (*Tenthredo degener* Christ, 1791; *Sirex vespertilio* Fabricius, 1793; *Oryssus coronatus* Fabricius, 1798; *O. albopunctatus* Gimmerthal, 1836; *O. hyalinipennis* Costa, 1860). Parasite of *Buprestis* spp. (Buprestidae). Russia: EP (C, S, NC, CR), UR, ES (IR, ZB). – Europe (WE, EE, NE, SE), Georgia, SW Asia, Iran, Kazakhstan.

Orussus coreanus Takeuchi, 1938. Russia: FE (AM, KH, PR, SA). – China (NE), Korean Peninsula.

Orussus rufipes Tsuneki, 1963. Parasite of *?Xiphydria ogasawarai* Matsumura, 1927 (Xiphydriidae). Russia: FE (KU). – Japan (Hok, Hon).

PSEUDORYSSUS Guiglia, 1954. Type species *Oryssus henschii* Mocsáry, 1910. The genus is distributed in Palaearctic. Number of species: World – 2; Palaearctic – 2; Russia – 1.

Pseudoryssus henschii (Mocsáry, 1910) [Oryssus] (*P. emanuelis* Guiglia, 1956). Russia: EP (Taeger et al., 2006). – Europe (WE, EE, SE), N Africa, NW Asia.

**SUBORDER APOCRITA
ACULEATA**

INFRAORDER VESPOMORPHA

SUPERFAMILY CHRYSIDOIDEA

57. FAMILY DRYINIDAE

A.S. LELEJ AND V.M. LOKTIONOV

Dryinids are relatively uncommon insects whose larvae are parasitoids of Hemiptera (Auchenorrhyncha): planthoppers (Fulgoroidea: Cixiidae, Delphacidae, Dictyopharidae, Issidae, Tropiduchidae), leafhoppers (Membracoidea: Cicadellidae), or treehoppers (Membracoidea: Membracidae). Small (most are 5–8 mm) black insects with a reduced wing venation. Antennae 10-segmented, rising low on face, and not distinctly elbowed. Hind wing with a jugal lobe. The two sexes are often quite different and can be associated only by rearing. Most females are remarkable in that the fore tarsi are developed into chelae used for grasping and holding the hoppers used as hosts. Female dryinids catch a host adult or nymph with their chelae, sting and temporarily paralyze it, and lay eggs between the two mesosomal and metasomal segments. The parasitoid larva feeds internally on the host, almost during most of its development being a part of the body of the larva protruding as a saclike structure. The parasitoid when full grown, leaves the host and spins a silken cocoon nearby. Some species are polyembryonic, with 40 to 60 individuals developing from a single egg. They play important roles in the natural control of some insect pests, and have potential significance as biological control agents.

Cosmopolitan. Number of taxa: World – 15 subfamilies, 50 genera, 1827 species, Palaearctic – 9/24/257, Russia – 5/8/63.

References. Kieffer, 1914; Haupt, 1941, 1944; Ponomarenko, 1965, 1966, 1970, 1972, 1975, 1978, 1979, 1992, 1995, 2000, 2007; Olmi, 1984, 2003, 2004; Lelej, Kurzenko, 2006; Ponomarenko, Olmi, 2007; Guglielmino et al., 2013; Xu et al., 2013; Olmi, Xu, 2015.

Subfamily APHELOPINAE

APHELOPUS Dalman, 1823 (*Antaphelopus* Benoit, 1951; *Gymnaphelopus* Benoit, 1951). Type species *Dryinus atratus* Dahlman, 1823. Parasitoids

of Cicadellidae (subfam. Typhlocybinae). The genus is distributed worldwide. Number of species: World – 78, Palaearctic – 18, Russia – 7.

Aphelopus atratus (Dalman, 1823) [Dryinus] (*A. holomelas* Richards, 1939). Parasitoid of Cicadellidae: Typhlocybinae. Russia: EP (Olmi, Xu, 2015), FE (PR). – Europe (WE, NE, SE, EE), Armenia, Kazakhstan, Japan (Hon).

Aphelopus bennetti Olmi, 2004. Russia: FE (SA). – Japan (Hon, Shi).

Aphelopus camus Richards, 1939 (*A. empoasca* Ponomarenko, 1970). Parasitoid of Cicadellidae: Typhlocybinae. Russia: EP (C), ES (IR), FE (PR). – Europe (WE, NE, SE, EE), Tunisia, Turkey, Japan (Hon).

Aphelopus maetoi Olmi, 1995. Russia: FE (PR, SA). – China (NE), Japan (Hon).

Aphelopus melaleucus (Dalman, 1818) [Gonatopus] (*Ceraphron albipes* Ratzeburg, 1848; *A. albipes* Kieffer, 1905; *A. trisulcatus* Kieffer, 1914). Parasitoid of Cicadellidae: Typhlocybinae. Russia: EP (N, NW, C), ES (IR), FE (SA, KU). – Europe (WE, NE, SE, EE), Turkey, Armenia, Lebanon, China (NC), Japan (Hon).

Aphelopus serratus Richards, 1939. Parasitoid of Cicadellidae: Typhlocybinae. Russia: EP (Olmi, Xu, 2015). – Europe (WE, NE, SE, EE), China (NC), Japan (Hon, Shi).

Aphelopus urbaini Olmi, 2003. Russia: FE (SA, KU). – China (NC), Japan (Hon).

Subfamily ANTEONINAE

ANTEON Jurine, 1807 (*Chelogynus* Haliday, 1838; *Neochelogynus* R. Perkins, 1905; *Paranteon* R. Perkins, 1905; *Prosanteon* R. Perkins, 1905; *Lasianteon* Kieffer, 1913; *Lasianteon* Kieffer, 1913; *Xenanteon* Kieffer, 1913; *Allanteon* Kieffer, 1914; *Hirtanteon* Benoit, 1951). Type species *Anteon jurineanum* Latreille, 1809. Parasitoid of Cicadellidae. The genus is distributed worldwide. Number of species: World – 423, Palaearctic – about 50, Russia – 16.

Anteon arcuatum Kieffer, 1905. Parasitoid of Cicadellidae: Idiocerinae. Russia: EP (C), UR. – Europe (WE, NE, SE, EE), Mongolia, China (NC).

Anteon brachycerum (Dalman, 1823) [Dryinus]. Parasitoid of Cicadellidae: Macropsinae. Russia: EP (C). – Europe (WE, NE, EE), Japan (Hon), Nepal.

Anteon ephippiger (Dalman, 1818) [Gonatopus] (*A. albidicollis* Kieffer, 1905; *Chelogynus albidocollis* (!):

- Ponomarenko, 1978; *A. pyonganensis* Móczár, 1983). Parasitoid of Cicadellidae: Deltcephalinae, Macropsinae. Russia: **EP** (NW, E, CR), **UR**, **ES** (YA), **FE** (PR, SA, KU). – Europe (WE, NE, SE, EE), Morocco, Turkey, Lebanon, Kyrgyzstan, Kazakhstan, Mongolia, China (NE, NC), Korean Peninsula, Japan (Hok, Hon, Kyu).
- Anteon esakii** Yasumatsu, 1960. Parasitoid of Cicadellidae: Ledrinae. Russia: **FE** (PR). – China (NC), Korean Peninsula, Japan (Kyu).
- Anteon exiguum** (Haupt, 1941) [Chelogynus]. Russia: **FE** (PR). – Europe (WE, NE, EE), Korean Peninsula.
- Anteon fasciale** (Thomson, 1860) [Chelogynus]. Russia: **EP** (C). – Europe (WE, NE).
- Anteon flavigorne** (Dalman, 1818) [Gonatopus] (*A. sericeus* Kieffer, 1905; *A. subflavigornis* Haupt, 1941). Parasitoid of Cicadellidae: Deltcephalinae, Idiocerinae, Macropsinae. Russia: **EP** (NW, C, CR), **UR**, **FE** (PR). – Europe (WE, NE, SE, EE), Algeria, Kazakhstan, Mongolia, Korean Peninsula.
- Anteon fulviventre** (Haliday, 1828) [Dryinus] (*Chelogynus fulviventris*: Ponomarenko, 1978). Russia: **FE** (SA). – Europe (WE, NE, SE, EE), Morocco, Algeria, Kazakhstan.
- Anteon gaullei** Kieffer, 1905 (*A. cameroni* Kieffer, 1905; *Antaeon rufulocollis* Chitty, 1908; *Chelogynus rufulocollis*: Ponomarenko, 1978; *Ch. cameroni*: Ponomarenko, 1978). Parasitoid of Cicadellidae. Russia: **EP** (C), **ES** (IR). – Europe (WE, NE, SE, EE), Kyrgyzstan, Kazakhstan, Korean Peninsula.
- Anteon hikense** Olmi, 1995. Russia: **FE** (PR). – Korean Peninsula, Japan (Kyu).
- Anteon infectum** (Haliday, 1837) [Dryinus]. Parasitoid of Cicadellidae: Iassinae. Russia: **EP** (C), **FE** (PR). – Europe (WE, NE, SE, EE), Azerbaijan, Turkey, Japan (Hon).
- Anteon jurineanum** Latreille, 1809 (*Gonatopus brevicornis* Dalman, 1818; *A. brevicorne*: Ponomarenko, 1978). Parasitoid of Cicadellidae: Deltcephalinae, Macropsinae. Russia: **EP** (NW, C), **UR**, **WS** (AL), **FE** (PR, SA, KU, KA). – Europe (WE, NE, SE, EE), Turkey, Jordan, Turkmenistan, Kazakhstan, Mongolia, China (NE, NC, SW), Korean Peninsula, Japan (Hok, Hon, Kyu), Nepal.
- Anteon pubicorne** (Dalman, 1818) [Gonatopus] (*Chelogynus lucidus*: Ponomarenko, 1978; *A. mongolicum* Móczár, 1983). Parasitoid of Cicadellidae: Deltcephalinae, Macropsinae. Russia: **EP** (NW, C, E, S, CR), **UR**, **WS** (AL), **ES** (KR, IR), **FE** (PR, SA, KA, MG). – Europe (WE, NE, SE, EE), Morocco, Turkey, Armenia, Kazakhstan, Mongolia, China (NE, NC, NW), Korean Peninsula, Japan (Hok, Hon).
- Anteon scapulare** (Haliday, 1837) [Dryinus] (*A. carinatus* Kieffer, 1905; *A. longiforceps* Kieffer, 1905; *Chelogynus lanionis* Haupt, 1941). Parasitoid of Cicadellidae: Iassinae. Russia: **FE** (PR). – Europe (WE, EE), Turkey, Japan (Kyu).
- Anteon takenoi** Olmi, 1995. Russia: **FE** (PR). – China (NE, NC, SW), Korean Peninsula, Japan (Kyu).
- Anteon tripartitum** Kieffer, 1905 (*Chelogynus silvaticus* Ponomarenko, 1970). Parasitoid of Cicadellidae. Russia: **EP** (C), **UR**. – Europe (WE, NE, SE, EE), Japan (Hon).
- LONCHODRYINUS** Kieffer, 1905 (*Prenanteon* Kieffer, 1913; *Psilanteon* Kieffer, 1913). Type species *Lonchodryinus tricolor* Kieffer, 1905. Parasitoids of Cicadellidae. The genus is distributed worldwide. Number of species: World – 16, Palaearctic – 4, Russia – 1.
- Lonchodryinus ruficornis** (Dalman, 1818) [Gonatopus] (*Prenanteon semenovi* Ponomarenko, 1970; *P. pektusanense* Móczár, 1983; *P. ruficornis*: Ponomarenko, 1978; *L. kotenkoi* Ponomarenko, 1992). Parasitoid of Cicadellidae (excluding Typhlocybinae). Russia: **EP** (N, C, S), **UR**, **WS** (AL), **ES** (IR, BR, YA, ZB), **FE** (KH, SA, KU, KA, MG). – Europe (WE, NE, SE, EE), Kyrgyzstan, Kazakhstan, Mongolia, China (NE, NC, NW, WP), Korean Peninsula, Japan (Hok, Hon, Shi).

Subfamily BOCCHINAE

MYSTROPHORUS Förster, 1856. Type species *Mystrophorus formicaeformis* Ruthe, 1859. Parasitoids of Cicadellidae: Deltcephalinae. Palaearctic genus with 2 species.

Mystrophorus formicaeformis Ruthe, 1859. Parasitoid of Cicadellidae: Deltcephalinae. Russia: **EP** (C). – Europe (WE, NE, SE, EE).

Mystrophorus apterus Ponomarenko, 2000. Parasitoid of Cicadellidae: Deltcephalinae. Russia: **EP** (C).

Subfamily DRYININAE

DRYINUS Latreille, 1804 (*Campylonyx* Westwood, 1835; *Chelothelius* Reinhard, 1863; *Chlorodryinus* R. Perkins, 1905; *Paradryinus* R. Perkins, 1905;

Mesodryinus Kieffer, 1906; *Plastodryinus* Kieffer, 1906; *Hesperodryinus* R. Perkins, 1907; *Pero-dryinus* R. Perkins, 1907; *Lestodryinus* Kieffer, 1911; *Tridryinus* Kieffer, 1913; *Bocchoides* Benoit, 1953; *Richardsidryinus* Móczár, 1965; *Avodryinus* Ponomarenko, 1981; *Alphadryinus* Olmi, 1984). Type species *Dryinus formicarius* Latreille, 1804 = *Sphex collaris* Linnaeus, 1767. Parasitoids of Acanaloniidae, Cixiidae, Delphacidae, Dictyopharidae, Flatidae, Fulgoridae, Issidae, Lophopidae, Ricanidae, Tropiduchidae. The genus is distributed worldwide. Number of species: World – 293, Palaearctic – 11, Russia – 2.

Dryinus albrechti (Olmi, 1984) [*Richardsidryinus* (*Richardsidryinus albrechti*): Ponomarenko, 1992]. Russia: UR. – Canary Islands.

Dryinus tarraconensis Marshall, 1868 (*D. szepligetii* Kieffer, 1905; *Lestodryinus gregori* Hoffer, 1936; *L. bidens* Haupt, 1937). Parasitoid of Dictyopharidae. Russia: EP (S), FE (PR). – Europe (WE, SE, EE), Morocco, Turkey, Iraq, Tajikistan.

Subfamily GONATOPODINAE

ECHTHRODELPHAX R. Perkins, 1903 (*Pterognatopus* Abdul-Nour, 1976). Type species *Echthrodelpax fairchildii* R. Perkins, 1903. Parasitoids of Delphacidae. The genus is distributed worldwide. Number of species: World – 12, Palaearctic – 2, Russia – 1.

Echthrodelpax tauricus Ponomarenko, 1970. Russia: EP (CR).

GONATOPUS Ljungh, 1810 (*Dicondylus* Haliday, 1830; *Labeo* Haliday, 1833; *Labeola* Haldeman, 1842; *Chalcogonatopus* R. Perkins, 1905; *Eugonatopus* R. Perkins, 1905; *Neogonatopus* R. Perkins, 1905; *Pachygonatopus* R. Perkins, 1905; *Paragonatopus* R. Perkins, 1905; *Pseudogonatopus* R. Perkins, 1905; *Mercetia* Kieffer, 1906; *Platygonatopus* Kieffer, 1906; *Agonatopoides* R. Perkins, 1907; *Agonatopus* R. Perkins, 1907; *Apterodryinus* R. Perkins, 1907; *Cyrtogonatopus* Kieffer, 1907; *Pristogonatopus* Kieffer, 1907; *Digonatopus* Kieffer, 1913; *Trigonatopus* Kieffer, 1913; *Laberius* Kieffer, 1914; *Trichogonatopus* Hellen, 1930; *Laberinus* Ogloblin, 1932; *Metagonatopus* Ogloblin, 1932; *Allogonatopus* Haupt, 1938; *Donisthorpina* Richards, 1939; *Epigonatopoides* Richards, 1939; *Plectrogonatopus*

Richards, 1939; *Tetrodontochelys* Richards, 1939; *Idologonatopus* Ogloblin, 1953; *Madecagonatopus* Benoit, 1953; *Rhynchogonatopus* Benoit, 1953; *Pseudogonatopoides* Williams, 1956; *Cyrtogonatopoides* Ponomarenko, 1966; *Plectrogonatopoides* Ponomarenko, 1975; *Acrodontochelys* Currado, 1976; *Dolichochelys* Olmi et Currado, 1976; *Megagonatopus* Olmi et Currado, 1976; *Neogonatopoides* Abdul-Nour, 1976; *Nogatopus* Ponomarenko, 1979; *Tetradontochelys* J. Perkins, 1976; *Paradicondylus* Olmi, 1986). Type species *Gonatopus formicarius* Ljungh, 1810. Parasitoids of Acanaloniidae, Cicadellidae (except for Typhlocybinae, Idiocerinae, Macropsinae), Delphacidae, Flatidae, Issidae, Lophopidae, Tropiduchidae. The genus is distributed worldwide. Number of species: World – 441, Palaearctic – more than 50, Russia – 20.

Gonatopus baeticus (Ceballos, 1927) [Mercetia] (*Dryinus victorovi* Ponomarenko, 1970). Parasitoid of Dictyopharidae. Russia: EP (E, CR). – Europe (WE, SE, EE), Morocco, Algeria, Turkey, Turkmenistan, Uzbekistan.

Gonatopus bicolor (Haliday, 1828) [Dryinus] (*Dicondylus bicolor*: Ponomarenko, 1978). Russia: EP (C), FE (KU). – Europe (WE, NE, SE, EE), Kazakhstan.

Gonatopus camelinus Kieffer, 1904 (*Pseudogonatopus camelinus*: Ponomarenko, 1992, 1995). Parasitoid of Delphacidae. Russia: WS (KM). – Europe (WE, SE), Morocco, Algeria, Egypt, Turkey, Iran, United Arab Emirates, Tajikistan, Turkmenistan, Uzbekistan, Japan (Kyu).

Gonatopus chersonensis Ponomarenko, 1970. Russia: EP (CR). – Kyrgyzstan, Kazakhstan, Mongolia.

Gonatopus clavipes (Thunberg, 1827) [Gelis] (*G. sepsoides* Westwood, 1833; *Labeo pusillus* Szépligeti, 1901; *G. borealis* Sahlberg, 1910; *G. campestris* Ponomarenko, 1965; *G. rhaensis* Ponomarenko, 1970; *G. sepsoides*: Ponomarenko, 1992, 1995). Parasitoid of Cicadellidae: Deltcephalinae. Russia: EP (N, C, E, S, NC, CR), WS (KM), ES (KR), FE (AM, PR, SA, KU). – Europe (WE, NE, SE, EE), Tunisia, Turkey, Azerbaijan, Iran, Uzbekistan, Kazakhstan, Mongolia, Korean Peninsula, Japan (Hok, Hon).

Gonatopus distinctus Kieffer, 1906 (*G. septemdentatus* J. Sahlberg, 1910; *Pseudogonatopus distinctus*: Ponomarenko, 1978). Parasitoid of Delphacidae. Russia: EP (N, S, CR), FE (KU). – Europe (WE, NE, SE, EE), Mongolia.

Gonatopus distinguendus Kieffer, 1905 (*G. excavatus* Sahlberg, 1910; *Allogonatopus procerus* Haupt,

- 1938; *G. thomsoni* Hellen, 1953; *G. rossicus* Ponomarenko, 1965). Parasitoid of Cicadellidae: Deltocephalinae. Russia: **EP** (NW, C, E, S, NC), **UR**, **ES** (IR), **FE** (PR). – Europe (WE, NE, SE, EE), Morocco, Tunisia, Turkey, Iran, Kyrgyzstan, Kazakhstan, Mongolia, China (NE).
- Gonatopus formicarius** Ljungh, 1810 (*G. formicinus* Krogerus, 1932). Parasitoid of Cicadellidae: Deltocephalinae. Russia: **EP** (C, E, S), **UR**, **FE** (PR). – Europe (WE, NE, SE, EE), Turkey, Uzbekistan, Kazakhstan.
- Gonatopus graecus** Olmi, 1984. Parasitoid of Cicadellidae: Deltocephalinae. Russia: **EP** (S). – Europe (SE).
- Gonatopus helleni** (Raatiainen, 1961) [*Dicondylus*] (*Dicondylus sibiricus* Ponomarenko, 1965; *Pseudogonatopus rufus* Ponomarenko, 1979; *Dicondylus helleni*: Móczár, 1983; *D. dichromus*: Ponomarenko, 1992, 1995). Parasitoid of Delphacidae. Russia: **EP** (S), **ES** (BR). – Europe (WE, NE, SE, EE), Mongolia.
- Gonatopus horvathi** Kieffer, 1906 (*Cyrtogonatopoides borealis* Ponomarenko, 1966). Parasitoid of Cicadellidae: Deltocephalinae. Russia: **UR**, **WS** (KM), **ES** (YA), **FE** (KH, SA). – Europe (WE, NE, SE, EE), Kazakhstan, Mongolia, China (NE).
- Gonatopus lunatus** Klug, 1810 (*Neogonatopus lunatus*: Ponomarenko, 1978). Parasitoid of Cicadellidae: Deltocephalinae. Russia: **EP** (C, E, S, NC), **UR**, **WS** (KM, AL), **ES** (IR, ZB). – Europe (WE, NE, SE, EE), Morocco, Turkey, Israel, Jordan, Lebanon, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, Korean Peninsula.
- Gonatopus nepalensis** (Olmi, 1984) [*Pseudogonatopus*] (*Pseudogonatopus nepalensis*: Ponomarenko, 1992, 1995). Russia: **ES** (ZB), **FE** (PR). – Nepal.
- Gonatopus pallidus** (Ceballos, 1927) [*Dicondylus*] (*Donistorpina pallida*: Ponomarenko, 1992, 1995). Russia: **FE** (PR). – Europe (NE, SE), Algeria.
- Gonatopus pedestris** Dalman, 1818 (*G. leucostomus* Sahlberg, 1910; *Pachygonatopus arnoldii* Ponomarenko, 1966; *Tetrodonthelys pedestris*: Ponomarenko, 1992, 1995). Parasitoid of Cicadellidae: Deltocephalinae. Russia: **EP** (N, C, E, S, CR), **ES** (KR), **FE** (SA). – Europe (WE, NE, SE, EE), China (NE, CC), Japan (Hon, Kyu, Shi).
- Gonatopus planiceps** Kieffer, 1904. Parasitoid of Cicadellidae: Agalliinae. Russia: **EP** (S). – Europe (WE, SE, EE), Morocco, Algeria, Egypt, Turkey, United Arab Emirates, Turkmenistan, Kazakhstan.
- Gonatopus popovi** Ponomarenko, 1965. Parasitoid of Cicadellidae: Deltocephalinae. Russia: **EP** (E).
- Gonatopus solidus** (Haupt, 1938) [*Allogenatopus*] (*Agonatopoides johannae* Currado et Olmi, 1975; *A. solidus*: Ponomarenko, 1992, 1995). Parasitoid of Delphacidae. Russia: **ES** (ZB), **FE** (PR). – Europe (WE, NE, SE, EE).
- Gonatopus spectrum** (Van Vollenhoven, 1874) [*Dryinus*] (*Neogonatopus albidus* auct.). Parasitoid of Cicadellidae. Russia: **EP** (Olmi, Xu, 2015). – Europe (WE, NE, SE, EE), Kyrgyzstan, Kazakhstan.
- Gonatopus striatus** Kieffer, 1905 (*Agonatopoides tau-ricus* Ponomarenko, 1965). Parasitoid of Cicadellidae. Russia: **EP** (CR). – Europe (WE, NE, SE, EE), Turkey, Mongolia.
- HAPLOGONATOPUS** R. Perkins, 1905 (*Monogonatopus* Richards, 1939; *Congodryinus* Benoit, 1950). Type species *Haplogonatopus apicalis* R. Perkins, 1905. Parasitoids of Delphacidae. The genus is distributed worldwide. Number of species: World – 7, Palaearctic – 2, Russia – 1.
- Haplogonatopus oratorius** (Westwood, 1833) [*Gonatopus*] (*H. atratus* Esaki et Hashimoto, 1932; *H. suchovi* Ponomarenko, 1970; *Monogonatopus oratorius*: Ponomarenko, 1978). Parasitoid of Cicadellidae, Delphacidae. Russia: **EP** (E), **WS** (OM), **ES** (KR). – Europe (WE, NE, SE, EE), Turkey, Lebanon, China (NE, NC, NW, SE), Korean Peninsula, Japan (Hon, Kyu, Shi, Ryukyu, Ogasawara), Northern Mariana Islands (Guam).

58. FAMILY EMBOLEMIDAE

S.A. BELOKOBYLSKIJ

Small wasps parasitizing nymphs of plant hoppers (Hemiptera: Auchenorrhyncha) (Olmi et al., 2014a, 2014b). Females of Embolemidae are micropterous, brachypterous or macropterous, but males are always macropterous; third segment of maxillary palp much larger than other palpal segments; antenna 10-segmented, with long (female) or short (male) scape, articulated on two prominent contiguous frontal processes far above the clypeus; in macropterous forms, fore wing with long tubular developed radial and distal abscissa of medial vein, second radiomedial vein usually present; tibial spurs 1, 1, 2. Three extant genera are known in this family: *Embolemus* Westwood, parasitoids of nymphs of Cixiidae, *Ampulicomorpha* Ashmead, parasitoids of nymphs of Achilidae, and *Troglebolemus* Olmi, Mita et Guglielmino, 2014 with unknown biology. Also three fossil genera were described in Embolemidae: *Baissobius* Rasnitsyn, 1975, *Embolemopsis* Olmi, Rasnitsyn et Guglielmino, 2010 and *Cretembolemus* Olmi, Rasnitsyn, Brothers et Guglielmino, 2014.

Number of taxa: World – 3 genera, 39 species, Palaearctic – 3/10, Russia – 2/5.

R e f e r e n c e s . Trjapitzin, 1978; Rasnitsyn, Matveev, 1989; Belokobylskij, 1990, 1995; Olmi, 1996; Belokobylskij, Lelej, 2012; Varrone, Olmi, 2012; Guglielmino, Buckle, 2013; Olmi et al., 2014a,b.

AMPULICOMORPHA Ashmead, 1893. Type species *Ampulicomorpha confusa* Ashmead, 1893. Small genus with worldwide distribution. parasitoids of the nymphs of Achilidae living in rotten logs

and feeding on hyphal sheets of shelf fungi (Olmi et al., 2014a, 2014b). Number of species: World – 25 (including 1 fossil), Palaearctic – 2, Russia – 1.

Ampulicomorpha thauma Rasnitsyn et Matveev, 1989. Russia: EP (S), FE (KU). – Korean Peninsula, ?Japan.

EMBOLEMUS Westwood, 1833 (*Myrmecomorphus* Westwood, 1833; *Polyplanus* Nees, 1834; *Formila* de Romand, 1846; *Pedinomma* Forster, 1856). Type species *Embolemus ruddii* Westwood, 1833. Largest and worldwide distributed embolemid genus, which are most abundant in the tropical areas. The members of this genus are known as parasitoids of nymphs of Cixiidae living in the soil and feeding on roots (Varrone, Olmi, 2012; Olmi et al., 2014a,b). Number of species: World – 34 (including 1 fossil), Palaearctic – 7, Russia – 4.

Embolemus pecki Olmi, 1997. Russia: FE (PR). – Europe (SE), China (NO, NE, SE), Korean Peninsula, Vietnam, Indonesia.

Embolemus sensitivus Xu, Olmi et Guglielmino, 2012 (*E. ruddii* auct.). Russia: FE (AM, PR). – China (NO, SW, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu), Vietnam.

Embolemus ruddii Westwood, 1833 (*Myrmecomorphus rufescens* Westwood, 1833; *Pedinomma rufescens* var. *antennalis* Kieffer, 1906; *Pedinomma rufescens* var. *holochlora* Marshall, 1906; *Pedinomma rufescens* var. *hypochlora* Marshall, 1906; *Polyplanus sickershusanus* Nees, 1834; *Formila chevrolatii* de Romand, 1846). Russia: EP (NW, S), ES (IR). – Europe (WE, SE, NE, EE).

Embolemus tauricus Olmi, Belokobylskij et Guglielmino, 2014. Russia: EP (CR).

59. FAMILY BETHYLIDAE

A.S. LELEJ AND K.I. FADEEV

Small to medium sized, usually dark-colored wasps. Females of many species are wingless and antlike in appearance. In a few species, both winged and wingless forms occur in each sex. These wasps are parasitoids of the larvae of Lepidoptera and Coleoptera. Several species attack moths or beetles that infest grain or flour. A few species will sting people. Widely distributed in tropical and subtropical areas. Number of extant taxa: World – 84 genera, 2340 species, Palaearctic – 20 / 244, Russia – 20 / 34.

References. Kieffer, 1906, 1914; Evans, 1964; Nagy, 1968; Perkins, 1976; Trjapitsin, 1978; Móczár, 1984; Gordh, Móczár, 1990; Gorbatovsky, 1995, 1998; Argaman, 2003; Terayama, 2003a,b; Vargas-Rojas, Terayama, 2006; Macek et al., 2007; Polaszek, 2011; Lelej, 2012a; Humala, Polevoi, 2011, 2015; Kostjukov et al., 2014; Nakonechnaya et al., 2014; Zamprogno, Azevedo, 2014; Paukkunen, Kozlov, 2015; Terayama, 2016; Fadeev, 2017.

Subfamily PRISTOCERINAE

ACREPYRIS Kieffer, 1905 (*Neopristocera* Yasumatsu, 1955). Type species *Epyris reticulatus* Kieffer, 1904 (= *Pristocera armifera* Say, 1828). The genus is distributed in New World, Oriental region and Palaearctic. Number of species: World – 38, Palaearctic – 7, Russia – 1.

Acrepyris masii (Giordani Soika, 1933) [Pristocera]. Russia: EP (NC, CR). – Europe (SE).

PRISTOCERA Klug, 1808 (*Dicrogenium* Stadelmann, 1894, *Neurepyris* Kieffer, 1905; *Kathepyris* Kieffer, 1907; *Mangesia* Kieffer, 1911; *Nomineia* Kieffer, 1911; *Trichelobrachium* Kieffer, 1914; *Neodicrogenium* Benoit, 1957; *Diepyris* Benoit, 1957). Type species *Bethylus depressa* Fabricius, 1805. The genus is distributed in Afrotropical, Palaearctic and Oriental regions. Number of species: World – 124, Palaearctic – 2, Russia – 1.

Pristocera depressa (Fabricius, 1805) [Bethylus] (*Pseudisobrachium roubali* Menozzi, 1925). The parasite of larva of *Agriotes obscurus* L. (Elateridae). Russia: EP (S, CR). – Europe (SE, EE).

PSEUDISOBRACHIUM Kieffer, 1904 (*Monepyris* Kieffer, 1905; *Xestobethylus* Cameron, 1909;

Plutobethylus Kieffer, 1910; *Lyssepyris* Kieffer, 1913; *Xantepyris* Kieffer, 1913; *Parisobrachium* Kieffer, 1914; *Afrisobrachium* Benoit, 1957; *Edapholigon* Ogloblin, 1963). Type species *Pseudisobrachium laticeps* Kieffer, 1904. The females are often found in ant nests, it is suggested that they are parasitoids of beetle larvae. The genus is distributed in worldwide except Australia. Number of species: World – 140, Palaearctic – 9, Russia – 2.

Pseudisobrachium belokobylskyi Gorbatovsky, 1995.

Russia: FE (PR).

Pseudisobrachium subcyanum (Haliday, 1838) [*Epyris*] (*P. carpentieri* Kieffer, 1906). Russia: EP (C, E, S, NC, CR), UR. – Europe (WE, EE).

Subfamily EPYRINAE

Tribe EPYRINI

EPYRIS Westwood, 1832 (*Calyzoa* Westwood in Hope, 1837; *Dolus* Motschulsky, 1863; *Muellerella* de Saussure, 1892; *Homoglenus* Kieffer, 1904; *Paracalyzoa* Cameron, 1909; *Artiepyris* Kieffer, 1913; *Psilepyris* Kieffer, 1913; *Parepyris* Kieffer, 1913; *Pseudocalyoza* Turner, 1915; *Calyzella* Enderlein, 1920). Type species *Epyris niger* Westwood, 1832. The parasitoids of larvae of Tenebrionidae. The genus is distributed worldwide. Number of species: World – 219, Palaearctic – 40, Russia – 6.

Epyris alias Gorbatovsky, 1995. Russia: FE (PR).

Epyris chasanicus Gorbatovsky, 1995. Russia: FE (PR).

Epyris erythrocerus Kieffer, 1906. Russia: EP (S). – Europe (EE), N Africa.

Epyris kurzenkoi Gorbatovsky, 1995. Russia: FE (PR).

Epyris leleji Gorbatovsky, 1995. Russia: FE (PR).

Epyris tardus Kieffer, 1906. Russia: EP (S). – Europe (EE).

HOLEPYRIS Kieffer, 1904 (*Rysepyris* Kieffer, 1906; *Misepyris* Kieffer, 1913; *Parepyris* Brèthes, 1913).

Type species *Holepyris africanus* Kieffer, 1904. The parasitoids of larvae of beetles Cucujidae, Curculionidae, Tenebrionidae and moth Pyralidae. The genus is distributed worldwide. Number of species: World – 102, Palaearctic – 36, Russia – 1.

Holepyris kasparyani Gorbatovsky, 1995. Russia: FE (PR).

LAELIUS Ashmead, 1893 (*Paralaelius* Kieffer, 1905; *Allepyris* Kieffer, 1905). Type species *Laelius*

trogodermatis Ashmead, 1893. The parasitoids of larvae beetle Dermestidae, Bostrichidae, Anobiidae. The genus is distributed in Holarctic. Number of species: World – 23, Palaearctic – 15, Russia – 3.

Laelius antropovi Gorbatovsky, 1995. Russia: **FE** (PR).
Laelius femoralis (Förster, 1860) [*Bethylus*]. Russia: **EP** (C). – Europe (WE, NE).

Laelius foersteri Kieffer, 1914 (*Bethylus rufipes* Förster, 1860, nom. praeocc., nec Say, 1836). Russia: **EP** (N). – Europe (WE, EE).

Tribe SCLERODERMINI

SCLERODERMUS Latreille, 1809 (*Sclerochroa* Förster, 1850; *Scleroderma* Costa, 1864; *Neoscleroderma* Kieffer, 1905). Type species *Sclerodermus domesticus* Latreille, 1809. Number of species: World – 70, Palaearctic – 26, Russia – 1.

Sclerodermus domesticus Klug, 1809 (*Scleroderma fonscolombei* Westwood, 1881). Russia: **EP** (C, S, NC, CR). – Europe (WE, EE).

Tribe CEPHALONOMIINI

CEPHALONOMIA Westwood, 1833 (*Holopedina* Förster, 1850; *Cephaloderma* Hoffer, 1936). Type species *Cephalonomia formiciformis* Westwood, 1833. Number of species: World – 33, Palaearctic – 19, Russia – 2.

Cephalonomia formiciformis Westwood, 1833 (*Holopedina polypori* Förster, 1850). The parasite of the beetles *Cis* Latr. and *Ennearthron* Mellié (Ciidae). Russia: **EP** (CR). – Europe (WE, EE).

Cephalonomia hammi Richards, 1939. Russia: **EP** (N). – Europe (WE, NE).

PARARHABDEPYRIS Gorbatovsky, 1995. Type species *Pararhabdepyris paradoxus* Gorbatovsky, 1995. The genus is distributed in Eastern Palaearctic, Oriental region (Thailand) and Australia. Number of species: World – 3, Palaearctic – 1, Russia – 1.

Pararhabdepyris paradoxus Gorbatovsky, 1995. Russia: **FE** (PR).

PLASTANOXUS Kieffer, 1905 (*Snappania* Hedqvist, 1975). Type species *Anoxus chittendenii* Ashmead, 1893. The parasitoids of larvae beetle Cucujidae. The genus is distributed worldwide. Number of species: World – 10, Palaearctic – 6, Russia – 1.

Plastanoxus evansi Gorbatovsky, 1995. Russia: **FE** (PR).

Subfamily MESITIINAE

CLYTROVORUS Nagy, 1972. Type species *Sclerodermus multilobatus* Costa, 1864. Monotypic Palaearctic genus.

Clytovorus multilobatus Costa, 1864 [*Sclerodermus*] (*Mesitius fuscicornis* Kieffer, 1906; *M. horvathi* Kieffer, 1906; *M. zavadili* Hoffer, 1936; *M. viator* Nagy, 1968). Russia: **EP** (S), **UR**. – Europe (SE, EE).

GERBEKAS Argaman, 2003. Type species *Mesitius carcelii* Westwood, 1874. Monotypic Palaearctic genus.

Gerbekas carcelii (Westwood, 1874) [*Mesitius*] (*M. halidayellus* Westwood, 1874; *M. hungaricus* Kieffer, 1906; *M. gracilis* Kieffer, 1906; *M. nagyi* Móczár, 1969). Russia: **EP** (S). – Europe (WE, SE, EE), Tunisia, Israel, North Kazakhstan.

METRIONOTUS Móczár, 1970. Type species *Metrionous pappi* Móczár, 1970 (= *Mesitius minimus* Kieffer, 1906). Monotypic genus is distributed in Palaearctic and Afrotropical regions.

Metrionotus minimus (Kieffer, 1906) [*Mesitius*] (*Mesitius minor* Benoit, 1968; *Metrionotus pappi* Móczár, 1970). Russia (Argaman, 2003). – Europe (WE, SE), Egypt, Cyprus, Turkey, Syria, Lebanon, Israel, Saudi Arabia, Yemen, Iraq, Afghanistan, South Africa, Namibia.

PARVOCULUS Móczár, 1970. Type species *Parvoculus myrmecophilus* Móczár, 1970 (= *Mesitius indicus* Kleffer, 1905). Number of species: World – 3, Palaearctic – 2, Russia – 1.

Parvoculus indicus Kieffer, 1905 [*Mesitius*] (*Mesitius pyrenaicus* Kleffer, 1906; *M. caspicus* Nagy, 1969, *M. brevispinosus* Benoit, 1968, *Incertosulcus krombeini* Móczár, 1970, *Metrionotus africanus* Móczár, 1970, *M. szelenyi* Móczár, 1970; *M. mocsaryi* Móczár, 1970; *M. yarrowi* Móczár, 1970; *Parvoculus myrmecophilus* Móczár, 1970). Russia: **EP** (S). – Europe (WE), Egypt, Jordan, Israel, Yemen, Iraq, India, Democratic Republic of Congo, South Africa, Botswana, Namibia.

SULCOMESITIUS Móczár, 1970 (*Topcobius* Nagy, 1972). Type species *Sulcomesitus longispinus*

Móczár, 1970. The genus is distributed in Europe, Asia, and Africa. Number of species: World – 54, Palaearctic – 2, Russia – 1.

Sulcomesitus moczari Gorbatovsky, 1995. Russia: **FE** (PR).

UKAYAKOS Argaman, 2003. Type species *Mesitius obscurus* Kieffer, 1906. Monotypic Palaearctic genus.

Ukayakos obscurus (Kieffer, 1906) [Mesitius] (*Metriotonotus bekkeri* Móczár, 1984; *Heterocoelia nikolskiae* Móczár, 1984). Russia: **EP** (S). – Europe (WE), Tajikistan.

Subfamily BETHYLINAE

BETHYLUS Latreille, 1802 (*Perisemus* Förster, 1856; *Episemus* Thomson, 1862; *Anoxus* Thomson, 1862; *Digoniozus* Kieffer, 1905). Type species *Omalus fuscicornis* Jurine, 1807. The parasitoids of Lepidoptera larvae Tortricidae, Gelechiidae, Noctuidae. The genus is distributed in Holarctic. Number of species: World – 34, Palaearctic – 29, Russia – 2.

Bethylus fuscicornis (Jurine, 1807) [Omalus] (*B. syngenesiae* Haliday, 1834; *Perisemus triareolatus* Förster, 1851; *Episemus variabilis* Thomson, 1862, *B. berlandi* Arlé, 1929). The parasitoid of the larvae *Helicoverpa armigera* (Hübner) (Noctuidae). Russia: **EP** (N, NC), **ES** (YA), **FE** (AM, KH, PR, SA, KU, KA, MG). – Europe (WE, NE, EE), N Africa, Japan (Hon, Hon).

Bethylus cephalotes (Förster, 1860) [*Perisemus*]. The parasitoid of the larvae *Helicoverpa armigera* (Hübner) (Noctuidae). Russia: **EP** (NC). – Europe (WE).

GONIOZUS Förster, 1856 (*Parasierola* Cameron, 1883; *Progoniozus* Kieffer, 1905; *Perisierola* Kieffer, 1914). Type species *Bethylus claripennis* Förster, 1851. The parasitoids of Lepidoptera larvae Tortricidae, Gelechiidae, Pyralidae. The genus is distributed worldwide. Number of species: World – 142, Palaearctic – 12, Russia – 4.

Goniozus claripennis (Förster, 1851) [Bethylus] (*B. fuscipennis* Förster, 1851; *G. distigmus* Thomson,

1861; *Goniosus* (!) *flavipennis* (!): Nakonechnaya et al., 2014). The parasitoid of the larvae of *Sparganothis pilleriana* Den. et Schiff., *Eupoecilia ambiguella* Hb. (Tortricidae) and *Helicoverpa armigera* (Hübner) (Noctuidae). Russia: **EP** (N, NC), Siberia (Kieffer, 1914). – Europe (WE, NE, EE), Caucasus, Iran.

Goniozus gallicola (Kieffer, 1905) [Parasierola]. The parasitoid of the larvae *Lobesia botrana* (Den. et Schiff.) (Tortricidae), *Apomyelois ceratoniae* (Zeller), *Ephestia elutella* (Hübner), *Cryptoblabes gnidiella* Mill. (Pyralidae), *Borkhausenia* sp. (Oecophoridae) and *Helicoverpa armigera* (Hübner) (Noctuidae). Russia: **EP** (S, NC). – Europe (WE, SE, EE).

Goniozus japonicus Ashmead, 1904. The parasitoids of Lepidoptera larvae Gelechiidae, Gracillariidae, Pyralidae, Tortricidae, Glyphipterigidae. Russia: **ES** (YA), **FE** (KH, PR). – China (NC, CC, SE), Korean Peninsula, Japan (Hon, Shi, Kyu, Ryu).

Goniozus ussuricus (Gorbatovsky, 1995) [Parasierola] (*Parasierola ussurica*: Gorbatovsky, 1995). Russia: **FE** (PR).

ODONTEPYRIS Kieffer, 1904 (*Trissomalus* Kieffer, 1905; *Prosierola*: Gorbatovsky, 1995). Type species *Odontepyris flavinervus* Kieffer, 1904. The genus is distributed in Old World and Australia. Number of species: World – 26, Palaearctic – 3, Russia – 2.

Odontepyris orientalis (Gorbatovsky, 1995) [Prosierola]. Russia: **FE** (PR).

Odontepyris erucarus (Szelényi, 1958) [Parasierola]. The parasitoid of the larvae *Helicoverpa armigera* (Hübner) (Noctuidae). Russia: **EP** (E, S, NC, CR). – Europe (EE).

SIEROLA Cameron, 1881 (*Lelejola* Gorbatovsky, 1998). Type species *Sierola testaceipes* Cameron, 1881. The genus is distributed in Eastern Palaearctic, Oriental region, Australia, and Hawaii. Number of species: World – 198, Palaearctic – 3, Russia – 1.

Sierola ashmeadi Gorbatovsky, 1995. Russia: **FE** (PR). – Japan (Hon).

60. FAMILY CHRYSIDIDAE – CUCKOO WASPS

P. ROSA, A.S. LELEJ, S.A. BELOKOBILSKIJ,
V.M. LOKTIONOV AND L.A. ZAYTSEVA

The Chrysidae, commonly known as cuckoo wasps, jewel wasps or ruby tail wasps (*осы-блестянки* or *золотые осы*) are a cosmopolitan family which include about 2.500 species in the world (Aguiar et al., 2013), subdivided into 92 genera. The Russian species belong to two subfamilies, Cleptinae and Chrysidae, parasitoids or cleptoparasites of other Hymenoptera (sawflies, solitary wasps and bees) and Lepidoptera (Kimsey, Bohart, 1991). Another subfamily (Amiseginae), not yet recorded in the Palaearctic region with the exception of Japan, is likely to occur in Russian Far East (Kurzenko, Lelej, 2007) and members of this subfamily are known as egg parasitoids of walking sticks (Phasmatidae).

Like in Europe, the highest diversity of cuckoo wasps is found in the southern part of Russia and relatively few species have been observed in its northern part (Nikol'skaya, 1978). In fact, chrysids are usually thermophilous and search for sandy sites, clay brick and stone walls, rocky steppes, deserts and semideserts; but they can be found also in forest margins, clearings with sun-exposed dead tree trunks and stumps and gardens with sun-exposed dead wood, on wooden walls, wood poles and fence posts and all places where their hosts live (Rosa, 2006; Paukkunen et al., 2015). In the highest latitude and on mountains they fly on rocky outcrops and alpine meadows (Rosa, 2006; Paukkunen et al., 2015). Adults can be often observed feeding on flowers as Apiaceae and Euphorbiaceae, but many other flowers are occasionally visited by chrysids (Rosa, 2004).

Despite Russia being the largest country in the world, extending across the entirety of northern Asia and much of eastern Europe and incorporating a wide range of environments and landforms, studies on Chrysidae there are unexpectedly scarce. In fact, the only comprehensive catalogue dates back to Radoszkowski (1866) and the only key, restricted to the European part of the USSR, was compiled by Nikol'skaya (1978); it was mainly based on Linsenmaier (1959) and Móczár (1967) monographs, without adding new data or records. Nikol'skaya (1978) listed 99 species in the European part of USSR, which represent a small percentage compared with the European fauna, numbering about 600 species and subspecies (Linsenmaier, 1959; Soon, Rosa, 2015).

Even though the chrysidid fauna of few regions (the Caucasus, Crimea and Lower Volga) is well known, the species composition in large parts of Russia is almost unknown. Some administrative regions (Provinces and Territories) or Republics are too poorly studied. For example, no chrysids have been recorded yet in the Provinces of Smolensk, Orel and Kaliningrad, even when the faunas of the adjacent countries (Poland, Lithuania and Belarus) are well studied. The Siberian fauna has been recently studied by Rosa et al. (2017b, c), while Rosa et al. (2017d) gave other Russian records from its Far East and the European part.

Number of taxa: World – 95 genera, about 2815 species, Palaearctic – 36/1557, Russia – 23/330.

R e f e r e n c e s. Pallas, 1771; Assmuss, 1862; Radoszkowski, 1866; Dalla Torre, 1892; Uchida, 1927; Kuznetzov-Ugamskij, 1927; Hammer, 1950; Tsuneki, 1947, 1948; 1950, 1953a,b, 1955, 1959, 1963, 1979; Linsenmaier, 1959, 1968, 1987; Móczár, 1967; Semenov, 1967; Nikol'skaya, 1978; Kimsey, Bohart, 1991; Terayama et al., 2005; Rosa, 2004, 2006; Kurzenko, Lelej, 2007; Paukkunen et al., 2014, 2015; Terayama et al., 2016; Rosa et al., 2015; 2017a, b, c, d.

Subfamily CLEPTINAE

Number of taxa: World – 3 genera, 163 species, Palaearctic – 1/92, Russia – 1/15.

CLEPTES Latreille, 1802 (*Holcobleptes* Móczár, 1962; *Leiocleptes* Móczár, 1962; *Zimmermannia* Móczár, 1962; *Melanocleptes* Móczár, 1962; *Chrysocleptes* Móczár, 1962; *Oxycleptes* Móczár, 1962; *Neocleptes* Kimsey, 1981; *Maculosicléptes* Móczár, 1988). Type species *Sphex semiaurata* Linnaeus, 1761. Parasitoids of sawflies *Neodiprion* sp. (Diprionidae), *Hoplocampa brevis* (Klug), *Nematus hispidae* Smith, *N. ribesii* Scopoli, *Pristiphora* sp. (Tenthredinidae). Holarctic genus. Number of species: World – 92, Palaearctic – 71, Russia – 15.

Cleptes aerosus aerosus Förster, 1853. Russia: **EP** (CR). – Europe (SE, WE, EE), Georgia, Armenia, Turkey.

Cleptes dauriensis Móczár, 1997. Russia: **ES** (ZB).

Cleptes femoralis Mocsáry, 1890. Russia: **EP** (E). – Europe (SE), Caucasus, Turkey.

Cleptes halinae Kuznetzov-Ugamskii, 1927 (*C. hyalinae*: Kurzenko, Lelej, 2007; Lelej, Kurzenko, 2012). Russia: **FE** (PR).

Cleptes ignitus (Fabricius, 1787) [Ichneumon] (*C. ignita*: Radoszkowski, 1866; *C. igneta* (!): Radoszkowski, 1880; *C. ignites* (!): Shibaev, 2006). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, W Asia.

Cleptes insidiosus du Buysson, 1891. Russia: **EP** (NC).

Cleptes nitidulus (Fabricius, 1793) [Ichneumon] (*C. fallax* Mocsáry, 1889). Parasitoid of *Nematus* spp. and *Euura* spp. (Tenthredinidae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), Turkey.

Cleptes obsoletus Semenov, 1891. Russia: **EP** (S).

Cleptes primorensis Móczár, 2000. Russia: **FE** (PR).

Cleptes putoni du Buysson, 1886 (*C. saussurei* Mocsáry, 1889; *C. putonii* (!): Dalla Torre, 1892). Russia: **EP** (S). – Europe (WE, SE, EE), Turkey, Jordan.

Cleptes radoszkowskii Mocsáry, 1889. Russia: **EP** (E, NC). – Georgia, Azerbaijan, Uzbekistan.

Cleptes semiauratus (Linnaeus, 1761) [Sphex] (*C. palipes* Lepeletier, 1806). Parasitoid of *Endelomyia aethiops* (Gmelin), *Euura ribesii* (Scopoli) and *Pristiphora incisa* (Lindqvist) (Tenthredinidae). Russia: **EP** (NW, C, E, CR), **UR**. – Europe (WE, NE, SE, EE), Caucasus, Turkey.

Cleptes semicyaneus Tournier, 1879 (*C. elegans* Mocsáry in Mocsáry et Szepligeti, 1901). Russia: **EP** (C, E, S, CR), **ES** (IR). – Europe (WE, EE).

Cleptes splendidus (Fabricius, 1794) [Ichneumon] (*C. consimilis* du Buysson, 1887; *C. caucasicus* Semenov, 1920). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), N Africa, Georgia, Turkey, Iran.

Cleptes striatipleurus Rosa, Forshage, Paukkunen et Soon, 2015 (*C. semiauratus* auct.). Russia: **EP** (C, E, NC, CR), ?**WS**, ?**ES**, ?**FE**. – Europe (SE, EE), ?N America.

Subfamily CHYSIDINAE

Number of taxa: World – 56 genera, 2,488 species, Palaearctic – 31/1,461, Russia – 22/315.

Tribe ELAMPINI

Number of taxa: World – 25 genera, 1706 species, Palaearctic – 14/518, Russia – 11/123.

CHRYSELLAMPUS Semenov, 1932. Type species *Ellampus heros* Semenov, 1892 (*Parellampus* Semenov, 1932). Number of species: World – 11, Palaearctic – 10, Russia – 2.

Chrysellampus duplipunctatus Tsuneki, 1948 (*Philoctetes duplipunctatus*: Kurzenko, Lelej, 2007). Russia: **ES** (ZB), **FE** (KH, PR). – China (NE, NC), Korean Peninsula.

Chrysellampus sculpticollis (Abeille de Perrin, 1878) [*Omalus*]. Parasitoid of *Pemphredon rugifer* (Dahlbom) (Crabronidae). Russia: **EP** (CR). – Europe (WE, SE, EE), Caucasus, Turkmenistan.

COLPOPYGA Semenov, 1954. Type species *Hedychrum flavipes* Eversmann, 1858. Holarctic genus. Number of species: World – 7, Palaearctic – 4, Russia – 1.

Colpopyga flavipes (Eversmann, 1858) [Hedychrum] (*Spintharis pallipes*: Mocsáry, 1889; *Spintharina pallipes*: Zimmermann, 1950). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Middle East.

ELAMPUS Spinola, 1806 (*Ellampus* Agassiz, 1846; *Notozus* Förster, 1853). Type species *Chrysis panzeri* Fabricius, 1804. Nest parasites of Crabronidae. Distributed worldwide except Australian and Oriental regions. Number of species: World – 63, Palaearctic – 41, Russia – 21.

Elampus agnolii Rosa, 2017. Russia: **ES** (TU).

Elampus albipennis (Mocsáry, 1889) [Ellampus] (*E. ambigus* (!): Radoszkowski, 1877; *Notozus viridis* Tournier, 1890, nom. praeocc., nec Cresson, 1865; *E. tournieri* Dalla Torre, 1892). Russia: **EP** (C, E, S, CR), **UR**, **ES** (IR, BR), **FE** (PR). – Europe (EE), Saudi Arabia, UAE, W Asia, Iran, Turkmenistan.

Elampus assamensis (Mocsáry, 1911) [Ellampus]. Russia: **FE** (PR). – India (Assam).

Elampus bidens (Förster, 1853) [*Notozus*] (*E. femoralis* Eversmann, 1858; *Omalus superbus* Abeille de Perrin, 1878; *Notozus superbus* var. *rufescens* du Buysson, 1896; *E. bidens tristis* Tsuneki, 1970). Russia: **EP** (C, E), **WS** (OM), **ES** (IR, BR), **FE** (AM). – Europe (WE, SE, EE), Caucasus, W Asia, Japan (Hok, Hon, Shi, Kyu).

Elampus caeruleus Dahlbom, 1854 (*Omalus viridiventris* Abeille de Perrin, 1878; *E. coeruleus* (!): Becker, 1880). Russia: **EP** (E, S, CR), **UR**, **ES** (KR), **FE** (PR). – Europe (WE, SE, EE), Caucasus, China (NE).

Elampus coloratus Rosa, 2017. Russia: **ES** (TU). – Mongolia.

Elampus constrictus (Förster, 1853) [*Notozus*] (*E. panzeri*: Eversmann, 1858, Trautmann, 1927; *E. productus*: Becker, 1869; *N. productus* var. *vulgaris*

du Buysson, 1892; *N. angustatus*: du Buysson, 1896; *E. spina*: Sahlberg, 1910; *N. spina* ab. *prasina* Hellén, 1920; *N. panzeri*: Trautmann, 1927; *N. ambiguus*: Nikol'skaya, 1978; *Ellampus ambiguus*: Shibaev, 2006; *Ellampus constrictus*: Shibaev, 2006). Parasitoid of *Mimesa* spp. (Crabronidae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (BR, ZB), **FE** (AM, PR). – Europe (WE, NE, SE, EE), N Africa, China (NE, NC).

Elampus eversmanni (Mocsáry, 1889) [Ellampus] (*Elampus ambiguus* Eversmann, 1858, nom. praeocc., nec Dahlbom, 1854; *Notozus eversmanni*: du Buysson, 1892; *Ellampus eversmannii* (!): Dalla Torre, 1892). Russia: **EP** (E, S, NC, CR). – Caucasus, Iran, Central Asia, Kazakhstan.

Elampus foveatus (Mocsáry, 1914) [Ellampus] (*Elampus konowi*: Vinokurov, 2013). Russia: **EP** (NC), **ES** (IR). – Europe (WE, NE, SE, NE).

Elampus pallasi (Semenov, 1967) [Notozus]. Russia: **ES** (IR). – Kazakhstan.

Elampus panzeri (Fabricius, 1804) [Chrysis] (*Ch. scutellaris* Panzer, 1798, nom. praeocc., nec Fabricius, 1794; *Notozus panzeri*: Woldstedt, 1875; *Ellampus olgae* Semenov, 1891; *Notozus scutellaris*: Bischhoff, 1913; *Notozus constrictus*: Trautmann, 1927; *E. scutellaris*: Kurzenko, Lelej, 2007). Parasitoid of *Mimesa bicolor* (Jurine) *M. equestris* (Fabricius) and *M. lutaria* (Fabricius) (Crabronidae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (OM, AL), **ES** (IR, BR, YA, ZB), **FE** (PR). – Europe (WE, NE, SE, EE), Caucasus, W Asia, China (NE).

Elampus petri (Semenov, 1967) [Notozus]. Russia: **EP** (C), **ES** (TU, ZB).

Elampus pliginskii (Semenov, 1967) [Notozus]. Russia: **EP** (CR). – Georgia.

Elampus pyrosomus (Förster, 1853) [Notozus] (*E. chrysotonotus*: Eversmann, 1858). Russia: **EP** (C, E, S, NC), **UR**. – Europe (SE, EE), Caucasus, Turkey.

Elampus retusus (Semenov, 1967) [Notozus]. Russia: **ES** (ZB), **FE** (AM).

Elampus rufitarsis (Tournier, 1879) [Notozus]. Russia: **EP** (S).

Elampus sanzii Gogorza, 1887. Russia: **EP** (C, E, S), **ES** (IR, BR). – Europe (SE), Turkey, Middle East.

Elampus spina (Lepeletier, 1806) [Hedychrum] (*E. productus*: Eversmann, 1858; *Ellampus frivalszkyi*: Mocsáry, 1882). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), North Africa, Georgia, W Asia.

Elampus tauricus (Semenov, 1967) [Notozus]. Russia: **EP** (CR).

Elampus turcmenicus (Linsenmaier, 1968) [Omalus]. Russia: **EP** (S). – Turkmenistan.

Elampus ussuriensis (Semenov, 1967) [Notozus]. Russia: **ES** (YA, ZB), **FE** (PR).

HABA Semenov, 1954. Type species *Holopyga almasyana* Mocsáry, 1911. Palaearctic genus. Number of species: World – 5, Palaearctic – 4, Russia – 1.

Haba almasyana (Mocsáry, 1911) [Holopyga]. Russia: **EP** (S). – Kyrgyzstan.

HEDYCHRIDIUM Abeille de Perrin, 1878 (*Euchrum* Semenov, 1954; *Euchridium* Semenov, 1954; *Zarudnium* Semenov, 1954; *Zarudnidium* Semenov, 1954; *Cyrtachridium* Semenov, 1954). Type species *Hedychrumb minutum* Lepeletier, 1806 [= *Hedychriderium ardens* (Coquebert, 1801)]. Nest parasites of Crabronidae. Distributed worldwide except Australian region. Number of species: World – 290, Palaearctic – 200, Russia – 39.

Hedychridium adventicum Zimmermann, 1962. Russia: **EP** (C), **UR**. – Europe (WE, SE), Turkey.

Hedychridium aheneum (Dahlbom, 1854) [Hedychriderum] (*Hedychrumb callosum* Radoszkowski, 1877; *H. incrassatum*: De Stefani, 1888; *Holopyga ahenea*: Dalla Torre, 1892; *Z. incrassatum* (!): Vinokurov, 2006; *Hedychridium incrassatum*: Vinokurov, 2009). Russia: **EP** (S, NC, CR), **UR**. – Europe (SE, EE), Caucasus, Turkey, Middle East.

Hedychridium ardens ardens (Coquebert, 1801) [Chrysis] (*Hedychrumb minutum*: Eversmann, 1858; *H. integrum*: Assmuss, 1862). Parasitoid of *Diodontus tristis* (Vander Linden), *Oxybelus bipunctatus* Olivier, *Tachysphex nitidus* (Spinola), *T. obscuripennis* (Schenck) and *T. pompiliformis* (Panzer) (Crabronidae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (OM, TK, AL), **ES** (KS, TU, KR, IR, BR, ZB). – Europe (WE, NE, SE, EE), Caucasus.

Hedychridium arenisi Rosa, 2017. Russia: **ES** (TU).

Hedychridium asianum Linsenmaier, 1997 (*H. integrum asianum* Linsenmaier, 1997). Russia: **WS** (AL), **ES** (TU, KR, IR, BR). – Mongolia, China (NW).

Hedychridium belokobylskiji Rosa, 2017. Russia: **ES** (KS, TU, KR).

Hedychridium caputaureum G. Trautmann et W. Trautmann, 1919 (*H. roseum* var. *caputaureum* G. Trautmann et W. Trautmann, 1919). Parasitoid of *Astata minor* Kohl (Crabronidae). Russia: **EP** (NW, NC), **UR**. – Europe (WE, NE, EE), W Asia.

- Hedychridium caucasicum** Trautmann, 1926 (*H. sculpturatum* var. *caucasicum* Trautmann, 1926). Russia: **EP** (NC). – Caucasus, Turkey.
- Hedychridium chloropygum chloropygum** du Buysson, 1888. Russia: **EP** (NC). – Europe (WE, SE).
- Hedychridium coriaceum coriaceum** (Dahlbom, 1854) [Hedychrum] (*Hedychrum coriaceum* (!): Eversmann, 1858; *Holopyga* (*Hedychridium*) *coriacea*: Mocsáry, 1889). Parasitoid of *Lindenius albilibris* (Fabricius) (Crabronidae). Russia: **EP** (N, NW, C, E, NC), **UR**. – Europe (WE, NE, SE, EE), Turkey.
- Hedychridium cupreum** (Dahlbom, 1845) [Hedychrum] (*Chrysis integra*: Nylander, 1859; *Holopyga integra*: Mocsáry, 1889, *Hedychridium integrum*: Hellén, 1920, Vorontsovskij, 1930, Nikol'skaya, 1978, Zvantsov, 1988, Vinokurov, 2006, 2009). Parasitoid of *Dryudella pinguis* (Dahlbom), *D. stigma* (Panzer), *Harpactus lunatus* (Dahlbom) and *H. tumidus* (Panzer) (Crabronidae). Russia: **EP** (N, NW, C, NC), **UR**, **ES** (KS, TU, IR). – Europe (WE, NE, EE), Mongolia, China (NC).
- Hedychridium elegantulum** du Buysson, 1887. Possible parasitoid of *Miscophus bicolor* (Jurine) (Crabronidae). Russia: **EP** (C, NC). – Europe (WE, SE, EE).
- Hedychridium erschovi** (Radoszkowski, 1877) [Hedychrum] (*Euchrum turanum* Semenov, 1954). Russia: **EP** (NC). – Iran, Central Asia, Kazakhstan.
- Hedychridium femoratum** (Dahlbom, 1854) [Hedychrum]. Parasitoid of *Dryudella stigma* (Panzer) (Crabronidae). Russia: **EP** (NC), **UR**, **ES** (KS, TU, KR). – Europe (WE, SE), Turkey.
- Hedychridium gabriellae** Rosa, 2017. Russia: **ES** (TU).
- Hedychridium jucundum** (Mocsáry, 1889) [*Holopyga* (*Holopyga iucunda* Mocsáry, 1889)]. Russia: **EP** (C, S, NC), **UR**. – Europe (WE, SE, EE), Caucasus, Turkey.
- Hedychridium krajniki krajniki** Balthasar, 1946. Russia: **EP** (C), **UR**. – Europe (WE, SE, EE), Caucasus.
- Hedychridium leleji** Rosa, 2017. Russia: **ES** (TU).
- Hedychridium loktionovi** Rosa, 2017. Russia: **WS** (AL).
- Hedychridium longigena** Rosa, 2017. Russia: **ES** (IR).
- Hedychridium lucidiventre** Semenov, 1967. Russia: **EP** (NC). – Kazakhstan.
- Hedychridium mediocrum** Linsenmaier, 1987 (*H. scutellare*: Linsenmaier, 1959; *H. mediocre* Kimsey in Kimsey et Bohart, 1991). Russia: **UR**. – Europe (WE, SE, EE).
- Hedychridium monochroum** du Buysson, 1888 (*Cryptechridium breviceps* Semenov, 1954; *C. pusio* Semenov et Nikol'skaya, 1954). Russia: **EP** (S), **UR**. – Europe (WE, SE, EE), Caucasus, Turkmenistan, Tajikistan, Uzbekistan. From Europe to India.
- Hedychridium moricei** du Buysson, 1904 (*H. flavipes* var. *moricei*: Trautmann, 1927; *Zarudnidium clarum* Semenov, 1967). Russia: **EP** (S). – Europe (SE), Azerbaijan, Turkey, Middle East.
- Hedychridium parkanense** Balthasar, 1946. Russia: **EP** (S). – Europe (SE, EE).
- Hedychridium propodeale** Rosa, 2017. Russia: **ES** (KS, TU, KR).
- Hedychridium proshchalykini** Rosa, 2017. Russia: **ES** (TU).
- Hedychridium purpurascens** (Dahlbom, 1854) [Hedychrum] (*H. purpureascens* (!): Vinokurov, 2007; *H. purpureissens* (!): Vinokurov, 2006). Russia: **EP** (NC), **WS** (TK). – Europe (WE, NE, EE).
- Hedychridium roseum** (Rossi, 1790) [*Chrysis*] (*Chrysis carnea* var. *rosea* Rossi, 1790; *Euchrum roseum*: Nikol'skaya, 1978; (!)*Chrysura simplex*: Leontiev, 2015; *Euchrum roseum nanum*: Vinokurov, 2013). Parasitoid of *Astata boops* (Schrank), *Tachysphex pompiliformis* (Panzer) and possibly *Dryudella stigma* (Panzer), *Tachysphex pectinipes* (Linnaeus), *T. pompiliformis* (Panzer), and *Harpactus tumidus* (Panzer) (Crabronidae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), Abkhazia, Georgia, Azerbaijan, China (NE).
- Hedychridium rossicum** Gussakovskij, 1948 (*H. valesiense* Linsenmaier, 1959). Parasitoid of *Astata boops* (Schrank) (Crabronidae). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), Azerbaijan.
- Hedychridium satunini** Semenov, 1967 (*H. catunini* (!): Vinokurov, 2004). Russia: **EP** (NC). – Georgia, Azerbaijan.
- Hedychridium sculpturatum** (Abeille de Perrin, 1877) [Hedychrum] (*Euchrum sculpturatum*: Nikol'skaya, 1978; Vinokurov, 2006). Russia: **EP** (S, NC), **UR**, **WS** (TK). – Europe (WE, SE, EE), Caucasus.
- Hedychridium scutellare** (Tournier, 1878) [Hedychrum] (*H. mediocre* Linsenmaier, 1959). Russia: **EP** (NC). – Europe (WE, SE).
- Hedychridium sibiricum** Rosa, 2017. Russia: **ES** (TU).
- Hedychridium trossolus** (Semenov, 1954) [*Euchridium*]. Russia: **EP** (NC). – Tajikistan.
- Hedychridium tsunekii** Linsenmaier, 1959. Russia: **FE** (KH). – Korean Peninsula.
- Hedychridium turcicum** Arens, 2010 (*H. valesiense turcicum* Arens, 2010). Russia: **ES** (KR). – Turkey.

Hedychridium uvarovi Semenov, 1967. Russia: **EP** (NC).

Hedychridium zelleri (Dahlbom, 1845) [Hedychrum]. Parasitoid of *Misophus niger* Dahlbom and *M. concolor* Dahlbom (Crabronidae). Russia: **EP** (C, E, NC). – Europe (WE, NE, EE).

HEDYCHRUM Latreille, 1802 (*Cymura* Dahlbom, 1845; *Wollmania* Mocsáry, 1902). Type species *Chrysis lucidula* Fabricius, 1775 [= *Hedychrum nobile* (Scopoli, 1763)]. Nest parasites of *Cerceris* spp. and *Philanthus* spp. (Crabronidae). Number of species: World – 129, Palaearctic – 59, Russia – 14.

Hedychrum alexii Semenov, 1967. Russia: **EP** (NC). – Kazakhstan.

Hedychrum aureicolle Mocsáry, 1889 (*H. aureicolle* (!): Vinokurov, 2006; *H. aureicollis* (!): Vinokurov, 2009). Russia: **EP** (NC). – Europe (SE), Abkhazia, Turkey, Cyprus, Palestine, Iran.

Hedychrum chalybaeum Dahlbom, 1854 (*H. coeruleans*: Mocsáry, 1889; *H. szaboi* Mocsáry, 1889; *H. chalybeatum* (!): Kuznetsova, 1990; *H. cornescens* (!): Blagoveschenskaya, 1994). Parasitoid of *Cerceris interrupta* (Panzer) (Crabronidae). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (AL), **ES** (TU, KR, IR, BR, ZB), **FE** (AM, PR). – Europe (WE, NE, SE), Mongolia, China (NC, NE).

Hedychrum cholodkovskii Semenov, 1967. Russia: **EP** (NC), **UR**. – Kazakhstan.

Hedychrum gerstaeckeri gerstaeckeri Chevrier, 1869 (*H. Marianum* Mocsáry, 1911; *H. gerstaeckeri japonicum*: Linsenmaier, 1959). Parasitoid of *Cerceris rybyensis* (Linnaeus) and *C. ruficornis* (Fabricius) (Crabronidae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (AL, TK), **ES** (KR, YA), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), Georgia, China (SE), Japan.

Hedychrum gerstaeckeri plicatus Kilimnik, 1993 (*H. gerstaeckeri plisatus* (!): Shibaev, 2006). Russia: **EP** (C, NC), **UR**. – Europe (EE).

Hedychrum kozhantshikovi Semenov, 1967. Russia: **EP** (NC). – Kazakhstan.

Hedychrum longicolle Abeille de Perrin, 1877 (*H. collare* Semenov, 1892; *H. nobiliforme* Semenov, 1967; *H. trapezicolle* Semenov, 1967). Parasitoid of Crabronidae. Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (TK), **FE** (PR). – Europe (WE, SE), N Africa, Georgia, W Asia, Turkmenistan, Kazakhstan, China.

Hedychrum niemelai Linsenmaier, 1959 (?*H. semiviolaceum*: Vorontsovij, 1930; *H. aureicolle*: Kurzenko,

Lelej, 2007). Parasitoid of *Cerceris arenaria* (Linnaeus), *C. ruficornis* (Fabricius), *C. rybyensis* (Linnaeus), *C. quadrispasiata* (Panzer), and *C. quinquefasciata* (Rossi) (Crabronidae). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (NS), **ES** (KS, TU, KR, IR, BR, YA), **FE** (AM, PR). – Europe (WE, NE, SE, EE), China (NE).

Hedychrum nobile (Scopoli, 1763) [Sphex] (*H. lucidulum*: Eversmann, 1858, Assmuss, 1862; *H. lucidulum* var. *regium*: Eversmann, 1858; *Chrysis lucidula*: Nylander, 1859; *Ch. regia*: Nylander, 1859; *H. hediculum* (!): Radoszkowski, 1880; (!) *Ch. dichroa*: Leontiev, 2015). Parasitoid of *Cerceris arenaria* (Linnaeus), *C. rybyensis* (Linnaeus), *C. quadrispasiata* (Panzer), *C. quadricincta* (Panzer) (Crabronidae). Larvae of *H. nobile* can also feed on prey (Coleoptera, Curculionidae) stored in the nest. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TK, NS, AL), **ES** (KS, TU, KR, IR, BR, ZB). – Europe (WE, NE, SE, EE), Abkhazia, Caucasus.

Hedychrum rutilans rutilans Dahlbom, 1854 (*H. intermedium*: Nikolskaya, 1978; (!) *Chrysura austriaca*: Leontiev, 2015). Parasitoid of *Philanthus triangulum* Fabricius and *P. coronatus* (Thunberg) (Crabronidae). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **ES** (KR, ZB). – Europe (WE, NE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey.

Hedychrum rutilans ermak Semenov, 1967. Russia: **WS** (AL), **ES** (KS, TU, KR, IR, BR, ZB), **FE** (AM, PR).

Hedychrum simile Mocsáry, 1889 (*H. cyaneum* Radoszkowski, 1889, nom. praeocc., nec Brullé, 1846; *H. simili* (!): Uchida, 1927; *H. simile* f. *pullatum* Tsuneki, 1953; *H. simile aereum* Tsuneki, 1970). Parasitoid of *Cerceris arenaria* (Linnaeus) (Crabronidae). Russia: **FE** (AM, PR). – Mongolia, China (NE, NC), Korean Peninsula, Japan (Hon).

Hedychrum tobiasi Kilimnik, 1993. Russia: **EP** (NC). – Europe (EE).

Hedychrum virens virens Dahlbom, 1854 (*H. grandis* Tournier, 1890; *H. grande*: Dalla Torre, 1892). Parasitoid of *Cerceris tuberculata* de Villers (Crabronidae). Russia: **EP** (C, S, NC, CR), **UR**, **WS** (TK). – Europe (WE, SE), Georgia, Azerbaijan, Turkey, Middle East.

Hedychrum viridilineolatum Kilimnik, 1993. Russia: **FE** (PR). – Europe (EE).

HOLOPYGA Dahlbom, 1854 (*Pseudbedychrom* Abeille de Perrin, 1879; *Oar* Semenov, 1954; *Psacas* Semenov, 1954). Type species *Holopyga amoena*

- Dahlbom, 1845. Parasitoids of Crabronidae. Distributed worldwide, mostly in the Palaearctic. Number of species: World – 110, Palaearctic – 88, Russia – 17.
- Holopyga austrialis** Linsenmaier, 1959. Russia: **EP** (S). – Europe (WE, SE, EE).
- Holopyga amoenula amoenula** Dahlbom, 1845 (*H. amoenella* (!): Vorontsovij, 1930). Russia: **EP** (NC). – Europe (SE), Turkey.
- Holopyga amoenula occidenta** Linsenmaier, 1959. Russia: **ES** (KR). – Europe (WE, SE), Turkey.
- Holopyga chrysonota chrysonota** (Förster, 1853) [Ellampus]. Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (OM, AL), **ES** (KS, TU, KR, IR, YA). – Europe (WE, SE, EE), Caucasus, China (NC).
- Holopyga fervida fervida** (Fabricius, 1781) [Chrysis] (*H. chloroidea*: du Buysson, 1892, *H. curvata*: Bischoff, 1910; Blagoveschenskaya, 1994). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, NE, SE, EE), N Africa, Turkey, Middle East, Iran.
- Holopyga generosa asiatica** Trautmann, 1926 (*H. amoenula*: Mocsáry, 1889; *H. gloriosa* var. *asiatica* Trautmann, 1926; *H. ovata proviridis* Linsenmaier, 1959; *H. gloriosa* var. *viridis*: du Buysson, 1892). Russia: **WS** (AL), **ES** (KS, TU, KR) – Caucasus.
- Holopyga generosa generosa** (Förster, 1853) [Ellampus] (*H. ovata* var. *b*: Eversmann, 1858; *H. ovata*: Assmuss, 1862; *H. gloriosa* var. *ovata*: du Buysson, 1892; *H. amoenula*: Zvantsov, 1988; (!) *Pseudospinolia neglecta*: Leontiev, 2015). Parasitoid of *Astata boops* (Schrank) (Crabronidae). The females lay their eggs in nymphs of Heteroptera before they have been captured and brought to the nest by the host. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TK, NS, AL), **ES** (KS, TU, KR, IR). – Europe (WE, NE, SE, EE), China, Korean Peninsula.
- Holopyga ignicollis** Dahlbom, 1854 (*H. ovata* var. *ignicollis* Dahlbom, 1854; *H. cingulata* Semenov, 1967; *H. aereomaculata* (!): Vinokurov, 2006; *H. cingulata* Semenov, 1967). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (KS, KR, IR). – Europe (WE, SE, EE), Caucasus, Middle East, Kyrgyzstan, Kazakhstan.
- Holopyga inflammata inflammata** (Förster, 1853) [Ellampus]. Russia: **EP** (C). – Europe (WE, NE, SE, EE), N Africa, W Asia.
- Holopyga jurinei** Chevrier, 1862. Russia: **EP** (S, NC). – Europe (WE, SE, EE), N Africa, Turkey, W Asia.
- Holopyga lucida** (Lepeletier, 1806) [Hedychrum]. Russia: **EP** (C, S), **UR**, **ES** (IR) – Europe (WE, SE), Turkey.
- Holopyga medvedevi** Semenov, 1967. Russia: **EP** (NC). – Kazakhstan.
- Holopyga metallica** (Dahlbom, 1854) [Hedychrum] (*H. sibirica* Semenov, 1967). Parasitoid of *Dryudella stigma* (Panzer) (Crabronidae). Russia: **EP** (N, NW), **ES** (IR). – Europe (NE).
- Holopyga minuma** Linsenmaier, 1959. Russia: **EP** (E), **ES** (KS, TU, KR). – Europe (SE, EE), Turkey, Middle East, Iran.
- Holopyga mlokosiewitzii** (Radoszkowski, 1877) [Hedychrum]. Russia: **EP** (NC). – Europe (SE), Caucasus, Turkey, Palestine.
- Holopyga pavlovskii** Semenov et Nikol'skaya, 1954. Russia: **EP** (NC). – Tajikistan.
- Holopyga punctatissima punctatissima** Dahlbom, 1854. Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE), Caucasus, Turkey, Iran.
- Holopyga turkestanica** Mocsáry, 1909 (*H. punctatissima* var. *turkestanica* Mocsáry, 1909; *H. crassepuncta* Semenov, 1954; *H. ovata crassepuncta*: Linsenmaier, 1968). Russia: **EP** (NC). – Caucasus, Turkey, Iran, Turkmenistan, Kazakhstan.
- Holopyga vigora** Linsenmaier, 1959. Russia: **EP** (NC). – Europe (SE), Turkey.
- OMALUS** Panzer, 1801. Type species *Chrysis aenea* Fabricius, 1787. Nest parasites of *Passaloecus* spp., *Pemphredon* spp., *Psenulus* spp. (Crabronidae). Number of species: World – 34, Palaearctic – 26, Russia – 5.
- Omalus aeneus** (Fabricius, 1787) [Chrysis] (*O. aeneus* var. *pygialis* du Buysson, 1887; *Ellampus zarudnyi* Semenov, 1932; *Ellampus aeneus* var. *chevrieri*: Bischoff, 1910). Parasitoids of *Passaloecus corniger* Shuckard, *P. eremita* Kohl, *P. gracilis* (Curtis), *P. singularis* Dahlbom, *P. turionum* Dahlbom, *Pemphredon lethifer* (Shuckard), *P. lugubris* (Fabricius), and *Psenulus pallipes* (Panzer) (Crabronidae). The females lay their eggs in live aphids and do not enter the host nest as in the case of *O. biaccinctus*. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TK, AL), **ES** (KS, TU, KR, IR, YA, ZB), **FE** (AM, KH, PR, SA, KU). – Europe (WE, NE, SE, EE), N Africa, Georgia, Uzbekistan, China (NC, SE), Japan (Hok, Hon, Kyu). Probably accidentally was introduced into N America.
- Omalus berezovskii** (Semenov, 1932) [Ellampus]. Russia: **ES** (KS, TU), **FE** (PR). – China (NC, SW).
- Omalus biaccinctus** (du Buysson, 1892) [Ellampus]. Parasitoids of *Passaloecus turionum* Dahlbom,

P. gracilis (Curtis), and *P. eremita* Kohl (Crabronidae). The females lay their eggs in living aphids at the hunting site of their host, and the egg is brought into the host's nest concealed in the aphid prey. Russia: EP (NW, C, NC, CR). – Europe (WE, NE, SE, EE), W Asia.

Omalus hohlbeki (Semenov, 1932) [Ellampus]. Russia: EP (NC). – Kazakhstan.

Omalus stella (Semenov et Nikol'skaya, 1954) [Ellampus]. Russia: EP (S, NC). – Tajikistan, Kazakhstan.

PHILOCTETES Abeille de Perrin, 1879 (*Diplorrhos* Aaron, 1885; *Dictenulus* Semenov, 1932). Type species *Holopyga cicatrix* Abeille de Perrin, 1879 [= *Philoctetes micans* (Klug, 1835)]. Nest parasites of Crabronidae. Number of species: World – 54, Palaearctic – 48, Russia – 12.

Philoctetes bidentulus (Lepeletier, 1806) [Hedychrum] (*Elampus bidentatus* (!) Eversmann, 1858; *Elampus wesmaeli*: De Stefani, 1888; *Elampus appendicinus*: Dalla Torre, 1892). Parasitoid of *Pseun caliginosus* Westwood (Crabronidae). Russia: EP (C, E, S, NC), UR, ES (KS, IR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, W Asia.

Philoctetes bogdanovii (Radoszkovski, 1877) [Holopyga] (*Elampus rudowi*: Mocsáry, 1889; *E. bagdanovi* (!): du Buysson, 1892; *E. bogdanowii* (!): Bischhoff, 1913; *Ph. bogdanovi* (!): Rosa, 2006). Russia: EP (C, S, CR), UR. – Europe (SE, EE), Azerbaijan, Turkey, W Asia, Iran.

Philoctetes conifer (Semenov, 1932) [Ellampus]. Russia: EP (NC).

Philoctetes cynthiae Rosa, 2017. Russia: ES (TU). – Mongolia.

Philoctetes horvathi (Mocsáry, 1889) [Ellampus] (*Elampus wesmaeli* Mocsáry, 1882, nom. praeocc., nec Chevrier, 1862; *E. horváthii* (!): Dalla Torre, 1892). Russia: EP (S), ES (TU, KR). – Europe (WE, SE, EE), China (NC), Korean Peninsula.

Philoctetes kuznetsovi (Semenov, 1932) [Ellampus]. Russia: EP (E, NC), ES (IR). – Georgia.

Philoctetes lyubae Rosa, 2017. Russia: ES (TU).

Philoctetes mongolicus (du Buysson, 1901) [Ellampus] (*E. horvathi* var. *mongolicus* du Buysson, 1901). Russia: EP (S), ES (KS, TU, YA, ZB). – Central Asia, Mongolia, China (NC).

Philoctetes pylnovi (Semenov, 1932) [Ellampus]. Russia: UR. – Uzbekistan.

Philoctetes sareptanus (Mocsáry, 1889) [Ellampus] (*E. schulthessi* Mocsáry, 1890; *E. schulthessi* var.

subauratus Mocsáry, 1890; *Omalus sareptanus* var. *inflammatus*: Trautmann, 1927). Russia: EP (C, E, S, NC), UR, ES (IR), FE (KH). – Iran.

Philoctetes stackelbergi (Semenov, 1932) [Ellampus]. Russia: WS (AL).

Philoctetes truncatus (Dahlbom, 1831) [Chrysis]. Parasitoid of *Diodontus tristis* (Vander Linden) (Crabronidae). Russia: EP (C, E), UR, WS (TK), ES (TU, KR, IR). – Europe (WE, NE, SE, EE), N Africa.

PROCHRIDUM Linsenmaier, 1968. Type species *Holopyga hirtipes* Mocsáry, 1902. Number of species: World – 5, Palaearctic – 4, Russia – 1.

Prochridium sp. Russia: EP (S). Specimens of this genus have been recently collected in the south of European part and their description is in preparation.

PSEUDOMALUS Ashmead, 1902. Type species *Omalus semicircularis* Aaron, 1885 [= *Pseudomalus janus* (Haldeman, 1844)]. Nest parasites of Crabronidae that prey on aphids: *Passaloecus* spp., *Pemphredon* spp. and *Diodontus*. Number of species: World – 33, Palaearctic – 29, Russia – 10.

Pseudomalus agnolii Rosa, 2017. Russia: EP (C).

Pseudomalus auratus auratus (Linnaeus, 1758) [Sphecius] (*Omalus curvitiventris* Tournier, 1879; *Elampus testaceicornis* du Buysson, 1892; *E. bergiellus* Semenov, 1932). Parasitoid of *Passaloecus corniger* Shuckard, *P. eremita* Kohl, *P. insignis* (Vander Linden), *P. gracilis* (Curtis), *P. monilicornis* Dahlbom, *P. singularis* Dahlbom, *P. turionum* Dahlbom, *Pemphredon inornata* Say, *P. lethifer* (Shuckard), *P. lugens* Dahlbom, *P. lugubris* (Fabricius), *P. rugifer* (Dahlbom), and *Diodontus tristis* (Vander Linden) (Crabronidae). Russia: EP (N, NW, C, E, S, NC, CR), UR, WS (TK, AL), ES (KS, TU, KR, IR, BR), FE (PR, KA). – Europe (WE, SE, NE, EE), N Africa, Caucasus, China (NE, NC), Korean Peninsula, Japan (Hon). Accidentally was introduced into N America.

Pseudomalus auratus viridiventris (Mocsáry, 1890) [Ellampus]. Russia: EP (NC). – Caucasus.

Pseudomalus bergi (Semenov, 1932) [Ellampus]. Russia: EP (NC). – Iran, Kazakhstan.

Pseudomalus cupratus (Mocsáry, 1889) [Ellampus] (*E. auratus* var. *cupratus* Mocsáry, 1889). Russia: EP (C, CR). – Europe (WE, SE, EE).

Pseudomalus grandis (Tsuneki, 1950) [Ellampus]. Russia: FE (PR). – Japan (Hon, Kyu).

Pseudomalus nipponicus (Tsuneki, 1970) [Omalus].

Russia: **ES** (KR), **FE** (PR). – Japan (Hok, Hon).

Pseudomalus punctatus (Uchida, 1927) [Philotetes].

Russia: **ES** (KS, KR, YA), **FE** (AM, PR, SA, KA). – Korean Peninsula, Japan (Hok, Hon, Kyu).

Pseudomalus pusillus pusillus (Fabricius, 1804)

[Chrysis] (*Elampus minutus* Dahlbom, 1854: Becker, 1865; *Omalus pusillum* (!): Becker, 1880; *Ellampus auratus* var. *virescens* Mocsáry, 1889; *E. bordini* Semenov, 1932; *E. ruthenus* Semenov, 1932; *E. pusillus* var. *schniedeknechti*: du Buysson, 1896). Parasitoid of *Passaloecus eremita* Kohl, *P. insignis* (Vander Linden), and *Pemphredon lethifer* (Shuckard) (Crabronidae). Russia: **EP** (N, C, E, S, NC, CR), **UR**, **WS** (OM, AL), **ES** (KS, TU, KR, IR, YA, ZB). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Kazakhstan.

Pseudomalus pusillus semicupreus (Linsenmaier, 1959) [Omalus] (*P. pucillus* (!) *semicupreus*: Vinokurov, 2013). Russia: **EP** (NC). – Europe (SE).**Pseudomalus triangulifer** (Abeille de Perrin, 1877)

[Omalus] (*Ellampus auratus* var. *triangulifer*: du Buysson, 1892). Parasitoid of *Passaloecus insignis* (Vander Linden), *Pemphredon lugubris* (Fabricius), *P. lethifer* (Shuckard), *P. lugens* Dahlbom, *P. montana* Dahlbom, and *P. rugifer* (Dahlbom) (Crabronidae). Russia: **EP** (N, NW, C, NC), **UR**, **WS** (AL), **ES** (KS, TU, IR, YA, ZB), **FE** (KH, PR, KA, MG). – Europe (WE, NE, SE, EE), Caucasus, Turkey, China (NW).

Pseudomalus violaceus (Scopoli, 1763) [Sphex] (*Omalus coeruleus*: Eversmann, 1858; *Ellampus coerulescens*: Mocsáry, 1882; *E. violaceus*: De Stefani, 1888; *O. violaceus*: Trautmann, 1927 et auct.). Parasitoid of *Pemphredon lugubris* (Fabricius), *Passaloecus corniger* Shuckard and *P. eremita* Kohl (Crabronidae). Russia: **EP** (NW, C, E, NC), **UR**, **WS** (TK, AL), **ES** (IR, ZB), **FE** (PR). – Europe (WE, NE, SE, EE), Georgia, Middle East, China (NC).

Tribe CHRYSIDINI

Number of taxa: World – 28 genera, about 1700 species, Palaearctic – 15/934, Russia – 10/189.

CHRYSIDEA Bischoff, 1913. Type species *Chrysis pumila* Klug, 1845. Number of species: World – 22, Palaearctic – 4, Russia – 2.**Chrysidea disclusa disclusa** (Linsenmaier, 1959) [Chrysis] (*Chrysis pumila disclusa* Linsenmaier, 1959). Russia: **EP** (NC). – Europe (WE, SE, EE).**Chrysidea pumila** (Klug, 1845) [Chrysis] (*Chrysis assimilis* Abeille de Perrin, 1879; *Chrysidea pumilla* (!): Vinokurov, 2004; *Chrysidea persica* Radoszkowski, 1881). Russia: **EP** (C, S, NC), **UR**. – Europe (WE, SE, EE), N Africa.

CHRYYSIS Linnaeus, 1761 (*Chrysogona* Förster, 1853; *Dichrysis* Lichtenstein, 1876; *Tetrachrysis* Lichtenstein, 1876; *Hexachrysis* Lichtenstein, 1876; *Cornuchrysis* Baltasar, 1953; *Gonodontochrysis* Semenov, 1954). Type species *Sphex ignita* Linnaeus, 1758. More than 80 species groups are known within *Chrysis* (Kimsey, Bohart, 1991; Linsenmaier, 1999; Tarbinsky, 2002; Rosa, 2005). Palaearctic species are parasitoids of solitary bees (Megachilidae) and solitary wasps (Sphecidae, Crabronidae, Vespidae (Eumeninae, Masarinae)). It is the largest genus in the Chrysididae, worldwide distributed. Number of species: World – about 1100, Palaearctic – 708, Russia – 140.

Chrysis aello Semenov et Nikol'skaya, 1954. Russia: **EP** (NC). – Tajikistan.**Chrysis aestiva** Dahlbom, 1854 (*Ch. moczari*: Vinokurov, 2006). Russia: **EP** (NC). – Europe (SE), Caucasus, Turkey, Palestine, Iran.**Chrysis albanica** Trautmann, 1927 (*Ch. succincta* var. *albanica* Trautmann, 1927). Russia: **EP** (S, NC, CR), **UR**. – Europe (SE, EE), Georgia.**Chrysis ambigua** Radoszkowski, 1891 (*Ch. taczanowskyi* (!) var. *ambigua*: Trautmann, 1927). Russia (Trautmann, 1927). – Europe (SE), Turkey, Palestine, Iran, Turkmenistan.**Chrysis amneris** Balthasar, 1953 (*Ch. arnoldii* Semenov, 1967). Russia: **Southern Russia** (Strumia, 2008). – Sudan, Saudi Arabia, United Arab Emirates, Palestine, Turkmenistan.**Chrysis amurensis** Semenov, 1967. Russia: **WS** (NS, AL), **ES** (KS), **FE** (AM, PR).**Chrysis analis analis** Spinola, 1807 (*Ch. analis* var. *rubescens* Radoszkowski, 1880; *Ch. sybarita* (!) var. *rubescens*: Bischoff, 1910). Parasitoid of *Osmia andrenoides* Spinola (Megachilidae). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), Caucasus.**Chrysis angolensis** Radoszkowski, 1881 (*Ch. fuscipennis*: Mocsáry, 1889; *Ch. fuscipennis murasaki*: Tsuneki, 1953). Parasitoid of *Sceliphron* (Sphecidae) and *Eumenes* (Vespidae: Eumeninae). Russia: **ES** (ZB), **FE** (AM, KH, PR). – China (NE, NC, CC, SW, SE), Korean Peninsula, Japan (Hon, Shi, Kyu, Tsushima, Ryu). Worldwide except Europe (with exception of Cyprus).

- Chrysis angustifrons angustifrons** Abeille de Perrin, 1878 (*Ch. lagodechii* Radoszkowski, 1889; *Ch. pyrrha* Semenov, 1967). Russia: **EP** (NC). – Europe (WE, SE, EE), Georgia.
- Chrysis angustula angustula** Schenck, 1856 (*Ch. ignita* var. *brevidens*: du Buysson, 1895). Parasitoid of *Ancistrocerus trifasciatus* (Müller), *Symmorphus bifasciatus* (Linnaeus), *S. allobrogus* (de Saussure), *S. connexus* (Curtis), and *S. debilitatus* (de Saussure) (Vespidae: Eumeninae). Russia: **EP** (N, NW, NC), **UR**, **ES** (IR). – Europe (WE, NE, SE, EE), Caucasus, W Asia, China (NE, NC).
- Chrysis auriceps** Linsenmaier, 1959. Russia: **UR**. – Europe (SE, EE), Turkey, Middle East.
- Chrysis bergi** Semenov, 1967. Russia: **EP** (NC). – Europe (EE).
- Chrysis bianchii** Semenov, 1892. Russia: **EP** (NC). – Caucasus, Turkmenistan.
- Chrysis bicolor** Lepeletier, 1806 (*Ch. succincta* var. *bicolor*: Mocsáry, 1889; *Ch. succincta* ab. *virideocincta* Hellén, 1920). Parasitoid of *Tachysphex obscuripennis* (Schenck), *T. pompiliformis* (Panzer), and *Dinetus pictus* (Fabricius) (Crabronidae). Russia: **ER** (N, NW, C, E, S, NC, CR), **UR**, **WS** (OM), **ES** (IR, YA), **FE** (AM, KH, PR, SA). – Europe (WE, NE, SE, EE), N Africa, Caucasus.
- Chrysis bilobata** Balthasar, 1953. Russia: **EP** (NC). – Turkey, Middle East.
- Chrysis borealis** Paukkunen, Ødegaard et Soon, 2015. Possible parasitoid of *Ancistrocerus parietum* (Linnaeus) and *A. scoticus* (Curtis) (Vespidae: Eumeninae). Russia: **EP** (N, NW). – Europe (NE).
- Chrysis brevitarsis** Thomson, 1870. Parasitoid of *Discoelius zonalis* Panzer and *D. priesneri* Mader (Vespidae: Zethinae). Russia: **EP** (NC, CR), **ES** (TU). – Europe (WE, NE, SE, EE).
- Chrysis buda** Bohart, 1991. Russia: **ES** (BR). – China (NC, CC).
- Chrysis calimorpha calimorpha** Mocsáry, 1882. Russia: **EP** (C, S, NC). – Europe (WE, SE, EE).
- Chrysis carnifex** Mocsáry, 1889. Russia: **FE** (PR).
- Chrysis caspiensis** Linsenmaier, 1959. Russia: **EP** (S, NC). – Europe (EE).
- Chrysis castigata** Linsenmaier, 1951. Russia: **ES** (TU, BR). – Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan.
- Chrysis caucasicola** Balthasar, 1953 (*Ch. analis* var. *caucasica* Mocsáry, 1912, nom. praeocc., nec Radoszkowski, 1877; *Ch. analis caucasicola* Semenov, 1967, nom. praeocc., nec Balthasar, 1953; *Ch. analis caucasiensis* Linsenmaier, 1959). Russia: **EP** (NC). – Georgia, Azerbaijan.
- Chrysis cerastes** Abeille de Perrin, 1877 (!) *Ch. immaculata*: Leontiev, 2015. Parasitoid of *Syneuodynerus egregius* (Herrich-Schaeffer) (Vespidae: Eumeninae). Russia: **EP** (E, NC, CR), **WS** (AL). – Europe (WE, SE, EE), Azerbaijan, Caucasus.
- Chrysis chinensis** Mocsáry, 1912. Russia: **WS** (AL), **ES** (KS, ZB). – Europe (WE), Mongolia, China (NE, CC).
- Chrysis chosenensis** Tsuneki, 1950 (*Ch. nitidula chosenensis* Tsuneki, 1950; *Ch. chosenensis* f. *clariventris* Tsuneki, 1953; *Ch. chosenensis* (!): Kurzenko, Lelej, 2007). Russia: **FE** (PR). – Korean Peninsula.
- Chrysis chrysoprasina** Förster, 1853. Russia: **EP** (C), **UR**. – Europe (WE, SE, EE), Armenia, W Asia.
- Chrysis chrysostigma** Mocsáry, 1889 (*Ch. ramburi*: Balthasar, 1953 et auct.). Russia: **EP** (C, NC). – Europe (WE, SE, EE), Caucasus, Turkey.
- Chrysis cingulicornis cingulicornis** Förster, 1853. Russia: **EP** (S, CR), **UR**, **ES** (KR). – Europe (WE, SE, EE), Georgia, Caucasus, Central Asia.
- Chrysis cingulicornis pseudopyrrhina** Linsenmaier, 1959. Russia: **EP** (S).
- Chrysis circe** Mocsáry, 1889. Russia: **EP** (NC). – Europe (SE, EE), Caucasus, Turkey.
- Chrysis clarinicollis** Linsenmaier, 1951 (*Ch. ignita* var. *clarinicollis* Linsenmaier, 1951). Russia: **EP** (S), **UR**. – Europe (NE, WE, SE, EE), N Africa.
- Chrysis comparata** Lepeletier, 1806 (*Ch. distinguenda*: Radoszkowski, 1866; *Ch. chevrieri*: Mocsáry, 1882 et auct.; *Ch. chevrieri* var. *orientalis* Mocsáry, 1889). Russia: **EP** (C, S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Caucasus.
- Chrysis comta** Förster, 1853. Russia: **EP** (S, NC, CR), **UR**. – Europe (WE, SE, EE), Caucasus, Turkey, Iran, China (NE, NC).
- Chrysis consanguinea consanguinea** Mocsáry, 1889. Russia: **WS** (AL), **ES** (KS, TU, IR). – Europe (WE, SE, EE), N Africa.
- Chrysis cyaneata** Mocsáry, 1909 (*Ch. uvarovi* Semenov, 1967). Russia: **EP** (NC). – Kazakhstan.
- Chrysis cylindrica** Eversmann, 1858 (*Ch. viridula* var. *integra*: De Stefani, 1888 et auct.; *Ch. viridula* var. *erythromelas* du Buysson, 1895). Parasitoid of *Tropidodynerus interruptus* (Brullé) (Vespidae: Eumenidae). Russia: **EP** (E, S, CR), **UR**, **ES** (YA). – Europe (WE, SE, EE), Caucasus.
- Chrysis daphne** Smith, 1874. Russia: **FE** (PR). – Japan (Hon).

Chrysis daphnis daphnis Mocsáry, 1889. Russia: **EP** (CR). – Europe (WE, SE, EE), N Africa.

Chrysis daphnis syriensis Linsenmaier, 1959. Russia: **EP** (NC). – Turkey, Syria, Palestine, Iran.

Chrysis dauriana Linsenmaier, 1959 (*Ch. mongolica* Semenov, 1967, nom. praeocc., nec Mocsáry, 1914; *Ch. mongoliana* Bohart, 1991). Russia: **WS** (AL), **ES** (TU, KR, BR, ZB), **FE** (PR). – Mongolia.

Chrysis diacantha diacantha Mocsáry, 1889 (*Ch. calypso* Semenov et Nikol'skaya, 1954). Russia: **EP** (NC). – Europe (SE), Caucasus, Middle East, Central Asia.

Chrysis decora Mocsáry, 1887 (*Ch. superba* Radoszkowski, 1877, nom. praeocc., nec Cresson, 1865; *Ch. rutilans* var. *asiatica* Mocsáry, 1889; *Ch. rutilans* var. *mesasiatica* Semenov, 1912). Russia: **EP** (E, S, NC, CR), **UR**, **WS** (TK), **ES** (ZB). – Turkey, Palestine, Iran, Turkmenistan, Kyrgyzstan, Kazakhstan.

Chrysis distincta distincta Mocsáry, 1887 (*Ch. analis* var. *incerta* Radoszkowski, 1880, nom. praeocc., nec Dahlbom, 1854). Russia: **EP** (S, NC, CR). – Georgia, Azerbaijan, Turkey, Cyprus, Palestine, Iran.

Chrysis distincta thalhammeri Mocsáry, 1889 (*Ch. distincta talhameri* (!): Vinokurov, 2013). Russia: **EP** (NC). – Europe (SE, EE), Caucasus, Middle East, Central Asia.

Chrysis elegans Lepeletier, 1806. Russia: **ER** (CR), **UR**. – Europe (WE, SE, EE), Caucasus.

Chrysis equestris Dahlbom, 1854. Parasitoid of *Discoelius dufourii* Lepeletier de Saint Fargeau and *D. zonalis* (Panzer) (Vespidae: Zethinae). Russia: **EP** (C, E), **UR**, **FE** (PR). – Europe (WE, NE, SE, EE).

Chrysis fasciata Olivier, 1791. Russia: **EP** (C, E, CR), **UR**, **WS** (TK), **ES** (IR), **FE** (AM, PR). – Europe (WE, NE, SE, EE), N Africa.

Chrysis fouqueti csikiana Mocsáry, 1912 (*Ch. nitidularia*: Vinokurov, 2006). Russia: **EP** (NC), **WS** (AL). – Kazakhstan.

Chrysis frankenbergeri Balthasar, 1953 (*Ch. frankenbergeri* (!): Vinokurov, 2006). Russia: **EP** (NC). – Europe (SE, EE), Armenia, Azerbaijan, Palestine.

Chrysis frivaldszkyi frivaldszkyi Mocsáry, 1882 (*Ch. frivaldszkyi* (!): Nikolskaya, 1978 *Ch. frivaldskii* (!): Vinokurov, 2006). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (KS, TU, KR, IR). – Europe (WE, SE, EE), Caucasus, Turkey.

Chrysis fulgida Linnaeus, 1761 (*Ch. cruenta* Mocsáry, 1883; *Ch. fulgida* var. *concolor* Mocsáry, 1912). Parasitoid of *Symmorphus allobrogus* (de Saussure), *S. bifasciatus* (Linnaeus), *S. crassicornis* (Panzer)

and *S. murarius* (Linnaeus), and possibly *Ancistrocerus parietum* (Linnaeus) (Vespidae: Eumeninae). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (TM, TK, AL), **ES** (KS, TU, KR, IR, ZB), **FE** (AM, PR, KA). – Europe (WE, NE, SE, EE), Caucasus, China (NE).

Chrysis galloisi (du Buysson, 1908) [Tetrachrysis] (*Ch. rubrifasciata* Tsuneki, 1950; *Ch. galboisi* (!): Vinokurov, 2006). Russia: **ES** (BR), **FE** (PR). – China, Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Chrysis germari germari Wesmael, 1839 (*Ch. nitidula* Germar, 1817, nom. praeocc., nec Fabricius, 1775; *Ch. succinctula*: Radoszkowski, 1866). Russia: **ER** (C, E), **UR**. – Europe (WE, SE, EE), Caucasus.

Chrysis gertabi Radoszkowski, 1891 (*Ch. taczanowskyi* (!) var. *germabi* (!): Trautmann, 1927). **Russia** (Trautmann, 1927). – Turkmenistan, Central Asia.

Chrysis glasunovi Semenov, 1967. Russia: **EP** (S, NC, CR). – Georgia, Caucasus.

Chrysis gracillima gracillima Förster, 1853 (*Ch. gracilima* (!): Vinokurov, 2006). Parasitoid of *Microdynerus exilis* (Herrich-Schäffer), *M. parvulus* (Herrich-Schäffer), *M. nudunensis* (de Saussure) (Vespidae: Eumeninae). Russia: **EP** (C, S, NC), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), N Africa, Middle East, Iran.

Chrysis graelsii graelsii Guérin-Méneville, 1842 (*Ch. analis*: Eversmann, 1858; *Ch. sybarita*: Radoszkowski, 1889; *Ch. chevrieri*: du Buysson, 1895; *C. chevrieri* var. *pusilla* du Buysson, 1895; *Ch. sybarita pekinensis*: Vinokurov, 2006; *Ch. sybarita opaca*: Vinokurov, 2006). Parasitoid of *Euodynerus disconotatus* (Lichtenstein), *E. notatus* (Jurine), and *E. quadrifasciatus* (Fabricius) (Vespidae: Eumeninae). Russia: **EP** (N, C, NC, CR), **UR**, **WS** (NS), **ES** (KS, TU, KR, BR). – Europe (WE, NE, SE, EE), Caucasus, Asia Minor, China (NC).

Chrysis grohmanni krkiana Linsenmaier, 1959. Russia: **EP** (S, NC). – Europe (SE, EE).

Chrysis grumorum Semenov, 1892. Russia: **EP** (NC). – China (WP).

Chrysis ignita ignita (Linnaeus, 1758) [Sphex]. Parasitoid of *Ancistrocerus auctus* (Fabricius), *A. gazella* (Panzer), *A. nigricornis* (Curtis), *A. parietum* (Linnaeus), *A. parietinus* (Linnaeus), *A. scoticus* (Curtis), *A. trifasciatus* (Müller) and possibly *Gymnomerus laevipes* (Shuckard), *Euodynerus posticus* (Herrich-Schäffer), *Allodynerus delphinalis* (Giraud), *Symmorphus bifasciatus* (Linnaeus) (Vespidae: Eumeninae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TK, AL), **ES** (IR, BR, ZB), **FE** (AM, KH),

- PR, SA, KA). – Europe (WE, NE, SE, EE), N Africa, China (NE, NC), Korean Peninsula, India.
- Chrysis illigeri** Wesmael, 1839 (*Ch. succincta* ab. *chrysoprasina*: Hellén, 1920). Parasitoid of *Tachysphex pompiliformis* (Panzer) (Crabronidae). Russia: **EP** (NW, C, S, NC), **UR**, **WS** (AL), **ES** (TU, KR, IR, BR, YA), **FE** (PR). – Europe (WE, NE, SE, EE).
- Chrysis immaculata** du Buysson, 1898. Russia: **EP** (C, NC). – Europe (WE, SE, EE).
- Chrysis impressa** Schenck, 1856 (*Ch. valida*: Humala, Polevoi, 2009). Parasitoid of *Ancistrocerus parietinus* (Linnaeus), *A. claripennis* Thomson, *A. trifasciatus* (Müller) (Vespidae: Eumeninae). Russia: **EP** (N, NW, S, NC, CR). – Europe (WE, NE, SE, EE), Central Asia.
- Chrysis inaequalis inaequalis** Dahlbom, 1845 (*Ch. inaequalis* (!): Vorontsovij, 1930). Parasitoid of *Eumenes coarctatus* (Linnaeus) (Vespidae: Eumeninae). Russia: **EP** (E, S, NC, CR), **UR**, **Siberia**. – Europe (WE, SE, EE), Central Asia, China (NE, NC).
- Chrysis inaequalis sapphirina** Semenov, 1912. Russia: **EP** (S). – Europe (SE), Cyprus, Palestine, Iran, Kyrgyzstan.
- Chrysis inaequipunctata** Bischoff, 1910. Russia: **EP** (E). – Uzbekistan.
- Chrysis inconsueta** Rosa, 2017. Russia: **WS** (AL). – Tajikistan.
- Chrysis indigotea** Dufour et Perris, 1840 (*Ch. zaitzevi* Semenov, 1967; *Ch. indogotea* (!): Bischoff, 1913; *Ch. indigotea* var. *Dagestanica* Mocsáry, 1889; *Ch. ignite* (!) *dagestanica* (!): Vinokurov, 2006). Parasitoid of Eumeninae (Vespidae), possibly *Gymnomerus laevipes* (Shuckard). Russia: **EP** (S, NC, CR), **Siberia**, **FE** (AM). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Central Asia.
- Chrysis insperata** Chevrier, 1870. Russia: **EP** (C, S, NC), **UR**. – Europe (WE, SE, EE), Azerbaijan, Caucasus.
- Chrysis interjecta interjecta** du Buysson, 1894 (*Ch. interjecta hemichlora*: Vinokurov, 2013). Russia: **EP** (NC), **UR**. – Europe (WE, SE, EE), Middle East.
- Chrysis io** Semenov, 1967. Russia: **EP** (CR).
- Chrysis iris** Christ, 1791 (*Ch. nitidula*: Assmuss, 1862 et auct.; *Ch. soluta*: Radoszkowski, 1866). Parasitoid of *Symmorphus allobrogus* (de Saussure), *S. crassicornis* (Panzer), and *S. murarius* (Linnaeus) (Vespidae: Eumeninae). Russia: **EP** (NW, C, E, NC), **UR**, **ES** (KS, TU). – Europe (WE, NE, SE, EE), Caucasus.
- Chrysis japonica** Cameron, 1887 (*Ch. daurica* Mocsáry, 1914; *Ch. japonica* var. *cyanea* Uchida, 1927).
- Russia: **ES** (ZB), **FE** (AM). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Tsushima, Ryu).
- Chrysis jaxartis** Semenov, 1910 (*Ch. sybarita* var. *jaxartis* Semenov, 1910; *Ch. valesiana perezi*: Vinokurov, 2006). Russia: **EP** (NC). – Europe (SE), Turkey, Iran, Kazakhstan.
- Chrysis karafutonis** Tosawa, 1932. Russia: **FE** (SA).
- Chrysis kolazyi** Mocsáry, 1889 (*Tetrachrysis grohmanni* var. *sareptana* Trautmann, 1926; *Ch. kolazi*: Kimsey, Bohart, 1991). Russia: **EP** (S). – Europe (WE, SE).
- Chrysis lanata** Mocsáry, 1912 (*Ch. sapphirus* Semenov, 1967). Russia: **FE** (PR). – Uzbekistan.
- Chrysis lanceolata** Linsenmaier, 1959 (*Ch. lanciolata* (!): Vinokurov, 2013). Possibly Parasitoid of *Diodontus minutus* (Fabricius) and *Miscophus bicolor* Jurine (Crabronidae). Russia: **EP** (S, NC, CR), **UR**. – Europe (SE, EE).
- Chrysis leachii** Shuckard, 1837 (*Ch. leachi* (!): Nikol'skaya, 1978). Russia: **EP** (C, E, S, NC, CR), **WS** (AL). – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Caucasus, Middle East, Iran.
- Chrysis leptomandibularis** Niehuis, 2000. Possibly parasitoid of *Symmorphus debilitatus* (de Saussure) (Vespidae: Eumeninae). Russia: **EP** (N, NW), **UR**. – Europe (WE, NE, SE, EE), Turkey.
- Chrysis longula longula** Abeille de Perrin, 1879 (*Ch. ignita* var. *longula* Abeille de Perrin, 1879; *Ch. longula aeneopaca* Linsenmaier, 1959). Parasitoid of *Ancistrocerus antilope* (Panzer), *Euodynerus posticus* (Herrich-Schäffer), *A. parietinus* (Linnaeus), *Symmorphus crassicornis* (Panzer), and *S. murarius* (Linnaeus) (Vespidae: Eumeninae). Russia: **EP** (N, NW, C, E, NC, CR), **UR**, **WS** (AL), **ES** (IR, BR). – Europe (WE, NE, SE, EE), Central Asia.
- Chrysis longula sublongula** Linsenmaier, 1951. Russia: **EP** (NC). – Europe (WE, NE, SE, EE).
- Chrysis lyda** Rosa, 2017. Russia: **EP** (NC). – Abkhazia.
- Chrysis maderi** Linsenmaier, 1959. Russia: **EP** (NC). – Europe (SE).
- Chrysis marginata marginata** Mocsáry, 1889. Russia: **EP** (NC), **UR**. – Europe (WE, SE, EE), Caucasus, Turkey, Middle East, Iran, Central Asia.
- Chrysis martinella martinella** du Buysson, 1900 (*Ch. satunini*: Vinokurov, 2006). Russia: **EP** (NC). – Europe (SE), Azerbaijan, Turkey.
- Chrysis mediadentata** Linsenmaier, 1951 (*Ch. ignita* var. *mediadentata* Linsenmaier, 1951; (!) *Ch. longula*: Leontiev, 2015). Parasitoid of *Odynerus spinipes* (Linnaeus) and *O. reniformis* (Gmelin) (Vespidae:

- Eumeninae). Russia: **EP** (E, NC), **ES** (IR). – Europe (WE, NE, SE, EE).
- Chrysis mediata mediata** Linsenmaier, 1951 (*Ch. ignita* var. *mediata* Linsenmaier, 1951). Parasitoid of *Odynerus spinipes* Linnaeus, *Ancistrocerus trifasciatus* (Müller) (Vespidae: Eumeninae). Russia: **EP** (NC), **WS** (TK), **ES** (YA), **FE** (PR). – Europe (WE, NE, SE, EE).
- Chrysis mirabilis** Radoszkowski, 1877 (*Ch. sinuosa*: Radoszkowski, 1866; *Ch. verae* Semenov, 1967). Russia: **EP** (CR). – Europe (SE), Georgia, Caucasus, Turkey.
- Chrysis mokrousovi** Rosa, 2017. Russia: **EP** (NC). – Abkhazia.
- Chrysis mutabilis mutabilis** du Buysson, 1887 (*Ch. mutabilis* var. *araxana*: du Buysson, 1895). Russia: **EP** (NC, CR). – Europe (SE), Georgia, Azerbaijan, Cyprus, Turkey, Middle East, Iran, Central Asia.
- Chrysis mysta** du Buysson, 1900. Russia: **Southern Russia** (Linsenmaier, 1969). – Turkey, Middle East, Syria, W Asia, Iran, Kazakhstan.
- Chrysis mysticalis** Linsenmaier, 1959 (*Ch. inaequalis*: Radoszkowski, 1889). Russia: **EP** (S, CR), **WS** (AL), **ES** (TU, KR, ZB). – Europe (WE, SE), Caucasus.
- Chrysis neobule** Semenov, 1954. Russia: **EP** (NC). – Caucasus, Tajikistan, Uzbekistan, Kazakhstan.
- Chrysis nevelskiana** Semenov, 1967. Russia: **FE** (PR).
- Chrysis nox** Semenov, 1954. Russia: **WS** (AL). – Tajikistan, Mongolia.
- Chrysis obtusidens** Dufour et Perris, 1840. Russia: **EP** (C, E, NC, CR), **WS** (TK). – Europe (WE, SE, EE), Caucasus.
- Chrysis pavesii** Rosa, 2017. Russia: **WS** (AL).
- Chrysis perrini** Radoszkowski, 1880 (*Ch. analis* var. β: Eversmann, 1858; *Ch. analis* var. *perrisi* Radoszkowski, 1880; *Ch. analis*: Mocsáry, 1889; *Ch. sybarita* (!) var. *perrini*: Bischoff, 1910). Russia: **EP** (NC), **UR**. – Caucasus.
- Chrysis phryne phryne** Abeille de Perrin, 1878. Russia: **EP** (NC). – Europe (WE, SE, EE).
- Chrysis placida** Mocsáry, 1879. Russia: **EP** (S, NC, CR), **UR**. – Europe (SE, EE), Caucasus, Turkey.
- Chrysis popovi** Semenov, 1954. Russia: **WS** (AL). – Tajikistan.
- Chrysis proauriceps** Rosa, 2017. Russia: **ES** (KR).
- Chrysis pseudobrevitarsis** Linsenmaier, 1951 (*Ch. ignita* var. *pseudobrevitarsis* Linsenmaier, 1951). Parasitoid of *Ancistrocerus antilope* (Panzer), *Euodynerus notatus* (Jurine), *E. quadrifasciatus* (Fabricius) (Vespidae: Eumeninae). Russia: **EP** (N, NW, NC, CR), **WS** (AL), **ES** (KS, TU, KR). – Europe (WE, NE, SE, EE), Mongolia.
- Chrysis pulchella pulchella** Spinola, 1807 (!) *Ch. pyrrhina*: Leontiev, 2015). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), Iraq, Iran, Central Asia.
- Chrysis pushkiniana** Semenov, 1967 (*Ch. graeca* Arens, 2004). Russia: **EP** (CR). – Europe (SE).
- Chrysis ragusae** De Stefani, 1888 (*Ch. taurica* Mocsáry, 1889; *Ch. ragusai* (!): Linsenmaier, 1959). Russia: **EP** (CR). – Europe (WE, SE, EE), Asia Minor.
- Chrysis remota remota** Mocsáry, 1889 (*Ch. margiana*: Vinokurov, 2006). Russia: **EP** (NC). – Caucasus, Turkmenistan, Kyrgyzstan, Central Asia.
- Chrysis rubripygia** Tsuneki, 1950. Russia: **FE** (PR). – Japan (Hok, Hon, Shi, Kyu).
- Chrysis ruddii ruddii** Shuckard, 1837 (*Ch. auripes*: Radoszkowski, 1880; *Ch. ruhdi* (!): Levi et al., 1974). Parasitoid of *Eumenes coronatus* (Panzer), *Ancistrocerus oviventris* (Wesmael), *A. parietum* (Linnaeus), *A. scoticus* (Curtis), *Eumenes* spp., *Odynerus spinipes* (Linnaeus), and *O. reniformis* (Gmelin) (Vespidae: Eumeninae). Russia: **EP** (N, NW, C, NC, CR), **UR**. – Europe (WE, NE, SE, EE), Caucasus, Asia Minor.
- Chrysis rufitarsis** Brullé, 1833. Russia: **EP** (S). – Europe (WE, SE, EE), N Africa, Caucasus, Central Asia.
- Chrysis russica** Linsenmaier, 1959. Russia: **Southern Russia** (Linsenmaier, 1959).
- Chrysis rutilans extranea** Linsenmaier, 1959. Russia: **ES** (KS), **FE** (PR). – China (NE, SW), Japan.
- Chrysis rutilans rutilans** Olivier, 1791 (?) *Ch. splendida* var. *aurotecta*: Bischoff, 1910; *Ch. splendida*: Humala, 1997). Parasitoid of *Gymnomerus laevipes* (Shuckard), *Jucancistrocerus caspicus* (Giordani Soika), *Katamenes flavigularis* (Blüthgen) (Vespidae: Eumeninae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (KS, TU, IR), **FE** (PR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, China, Japan.
- Chrysis rutiliventris rutiliventris** Abeille de Perrin, 1879. Russia: **EP** (NC), **Siberia**. – Europe (WE, SE, EE).
- Chrysis sardarica** Radoszkowski, 1890 (*Ch. sardarica lepidula* Linsenmaier, 1959; *Ch. sardarica perrecta*: Vinokurov, 2006; *Ch. cardarica* (!) *lepidula*: Vinokurov, 2006). Russia: **EP** (NC). – N Africa, Caucasus, Turkey, Middle East, Central Asia.

- Chrysis schencki** Linsenmaier, 1968 (*Ch. ignita schenckiana* Linsenmaier, 1959, nom. praeocc., nec Mocsáry, 1912; *Ch. ignita*: Humala, Polevoi, 2009). Parasitoid of *Ancistrocerus trifasciatus* (Müller), *A. gazella* (Panzer), and *A. nigricornis* (Curtis) (Vespidae: Eumeninae). Russia: **EP** (N, NW, NC), **WS** (TK). – Europe (WE, NE, SE), Central Asia, Japan.
- Chrysis sculpturata** Mocsáry, 1912 (*Ch. ignita* var. *sculpturata* Mocsáry, 1912). Russia: **EP** (NC). – Europe (WE, SE), Caucasus.
- Chrysis scutellaris** Fabricius, 1794. Possibly parasitoid of *Megachile leachella* Curtis and *Pseudanthidium lituratum* (Panzer) (Megachilidae). Russia: **EP** (C, E, S, NC). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey.
- Chrysis serena serena** Radoszkowski, 1891 (*Ch. pyrrhina* Dahlbom, 1845; Eversmann, 1858; *Ch. pyrrhina siciliaca* Linsenmaier, 1959; Vinokurov, 2006; *Ch. pyrrhina rhodesiaca*: Vinokurov, 2006). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), Azerbaijan, Caucasus, Middle East, Syria, Iran, Chia (NE).
- Chrysis sexdentata** Christ, 1791 (*Ch. caucasica* Radoszkowski, 1877; *Ch. micans*: du Buysson, 1896). Parasitoid of *Ancistrocerus parietum* (Linnaeus), *Euodynerus dantici* (Rossi) (Vespidae: Eumeninae). Russia: **EP** (C, S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Caucasus, Iran, Central Asia.
- Chrysis sibirica** Rosa, 2017. Russia: **ES** (TU, BR).
- Chrysis singula** Radoszkowski, 1891. Russia: **EP** (S). – Caucasus, Turkmenistan.
- Chrysis solida** Haupt, 1957 (*Ch. mediata fenniensis* Linsenmaier, 1959; *Ch. scintillans* Valkeila, 1971). Parasitoid of *Ancistrocerus trifasciatus* (Müller), *Euodynerus notatus* (Jurine) and possibly *Symmorphus debilitatus* (de Saussure) (Vespidae: Eumeninae). Russia: **EP** (N, NW, NC), **ES** (KR), **FE** (KA). – Europe (WE, NE, EE), Caucasus, Japan.
- Chrysis sooni** Rosa, 2017. Russia: **FE** (PR).
- Chrysis soror** Dahlbom, 1854 (*Ch. ariadne* Mocsáry, 1889; *Ch. scutellaris* var. *ariadne*: Balthasar, 1954; *Ch. soror calandra* Semenov, 1967). Russia: **EP** (C, E, S, NC), **UR**, **ES** (KR, IR). – Europe (SE, EE), Caucasus, Middle East, Central Asia.
- Chrysis speciosa** Radoszkowski, 1877. Russia: **Western Siberia** (Bischoff, 1913). – Cyprus, Iran, Uzbekistan.
- Chrysis splendidula** Rossi, 1790 (*Ch. tamara* Semenov, 1967; *Ch. cyanopyga*: Eversmann, 1858; *Ch. splendidula abdominalata*: Vinokurov, 2006; *Ch.*

- splendidula chlorisans*: Vinokurov, 2006). Parasitoid of *Eumenes coarctatus* (Linnaeus), *Eumenes mediterraneus* Kriechbaumer and *E. pomiformis* (Fabricius), *Gymnomerus laevipes* (Shuckard), *Symmorphus allobrogus* (de Saussure), *S. laevipes* (Shuckard) (Vespidae: Eumeninae). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, NE, SE, EE), Caucasus, Central Asia, China (NE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Tsushima, Ryu).
- Chrysis subcoriacea** Linsenmaier, 1959. Possibly Parasitoid of *Ancistrocerus trifasciatus* (Müller) (Vespidae: Eumeninae). Russia: **EP** (N, NW, NC). – Europe (WE, NE, SE, EE), Central Asia, Japan.
- Chrysis subsinuata fallax** Marquet, 1882 (*Ch. medioris*: Dalla Torre, 1892; *Ch. subsinuata unfaisciata*: Vinokurov, 2006; *Ch. mochii*: Vinokurov, 2006; *Ch. subsinuata nochii* (!): Vinokurov, 2006; *Ch. subsinuata fallex* (!): Vinokurov, 2006). Russia: **EP** (NC, CR). – Europe (SE, EE), Armenia, Turkey.
- Chrysis succincta** Linnaeus, 1767 (*Ch. siccincta* (!): Buganin et al., 2000). Russia: **EP** (N, C, E, S), **UR**, **ES** (BR, ZB). – Europe (WE, NE, ?SE, EE), Caucasus.
- Chrysis taczanowskii** Radoszkowski, 1877 (*Ch. taczanowskyi* (!): Bischoff, 1910; *Ch. taczanowskyi* (!): Trautmann, 1927; *Ch. znojkoi* Semenov, 1967; *Ch. znoikoi* (!): Vinokurov, 2006). Parasitoid of *Euodynerus disconotatus* (Lichtenstein) (Vespidae: Eumeninae). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), N Africa, Azerbaijan, Caucasus, Palestine, Syria, Iran.
- Chrysis tenella** Mocsáry, 1889 (*Ch. tenella* var. *chalcophana*: du Buysson, 1894; *Ch. tenella* var. *mlokosiewitzii*: du Buysson, 1894). Russia: **EP** (NC). – Caucasus.
- Chrysis terminata** Dahlbom, 1854. Parasitoid of *Ancistrocerus nigricornis* (Curtis) (Vespidae: Eumeninae). Russia: **EP** (S), **UR**. – Europe (WE, NE, SE, EE), Central Asia.
- Chrysis tianshanica** Semenov, 1967. Russia: **EP** (NC). – Central Asia.
- Chrysis tragica** Semenov, 1967. Russia: **EP** (NC).
- Chrysis uljanini sickmanni** Mocsáry, 1892 (*Ch. uljanini*: Bischoff, 1913; *Ch. sarafschana sickmanni*: Tsuneki, 1953; *Ch. uljanini*: Kurzenko, Lelej, 2007 et auct.). Russia: **FE** (AM, KH, PR).
- Chrysis valesiana** Frey-Gessner, 1887 (*Ch. sybarita* var. *valesiana* Frey-Gessner, 1887). Parasitoid of *Tropidodynerus interruptus* (Brullé), *Paravespa rex* (von Schulthess), *Paragymnomerus spiricornis*

(Spinola) (Vespidae: Eumeninae). Russia: EP (S, NC, CR). – Europe (WE, SE, EE).

Chrysis valida Mocsáry, 1912 (*Ch. ignita* var. *valida* Mocsáry, 1912). Russia: EP (C, NC), UR, WS (TK, AL), FE (AM, PR). – Europe (WE, NE, EE), Turkey.

Chrysis verna Dahlbom, 1854. Russia: EP (CR). – Europe (SE, EE), Caucasus, Palestine.

Chrysis vicana Vinokurov, 2010. Russia: EP (NC).

Chrysis vinokurovi Rosa, 2017. Russia: EP (NC).

Chrysis viridula Linnaeus, 1761 (*Ch. bidentata* var. *gemma* du Buysson, 1895). Parasitoid of *Odynerus spinipes* (Linnaeus), *O. reniformis* (Gmelin) and *O. melanocephalus* (Gmelin) (Vespidae: Eumeninae). Russia: EP (N, C, NC, CR), UR, WS (AL), ES (TU, KR, IR), FE (KH, PR). – Europe (WE, NE, SE, EE), China (NE), Korean Peninsula, Japan.

Chrysis westerlundi Trautmann, 1927. Russia: EP (N). – Europe (NE).

Chrysis zetterstedti Dahlbom, 1845 (*C. fasciata*: Banaszak, 1980 et auct.). Possibly parasitoid of *Euodynerus notatus* (Jurine) (Vespidae: Eumeninae). Russia: EP (NW, C, E, S, NC, CR), UR, WS (AL), ES (IR), FE (PR). – Europe (NE, EE).

CHRYWSURA Dahlbom, 1845 (*Olochrysis* Lichtenstein, 1876; *Monochrysis* Lichtenstein, 1876; *Holochrysis* Rye, 1878; *Arctochrysis* Haupt, 1956). Type species *Chrysis austriaca* Fabricius, 1804. Parasitoids of solitary bees (Megachilidae). Mostly Holarctic genus. Number of species: World – 124, Palaearctic – 113, Russia – 20.

Chrysura alticola (Semenov, 1912) [Chrysis]. Russia: WS (AL). – Tajikistan.

Chrysura austriaca (Fabricius, 1804) [Chrysis]. Parasitoid of bees *Hoplitis adunca* (Panzer), *H. anthocopoides* (Schenck), *Osmia parietina* Smith (Megachilidae). Russia: EP (C, E), Siberia (Radoszkowski, 1889). – Europe (WE, NE, SE, EE), N Africa.

Chrysura cuprea (Rossi, 1790) [Chrysis] (*Chrysis coeruleipes*: Radoszkowski, 1866). Parasitoid of *Osmia bicolor* Schrank, *O. versicolor* Latreille, *O. rufohirta* Latreille, *O. spinulosa* Trautmann, *O. andrenoides* Spinola (Megachilidae). Russia: EP (S, NC, CR). – Europe (WE, SE, EE), Caucasus, Abkhazia.

Chrysura desertorum (du Buysson, 1887) [Chrysis] (*Chrysis cyanocoelia*: Mocsáry, 1889). Russia: Southern Russia (Linsenmaier, 1969). – Georgia, Turkey, Israel.

Chrysura dichroa (Dahlbom, 1854) [Chrysis] (*Chrysis baeri* Radoszkowski, 1866). Parasitoid of *Osmia* (Megachilidae), which build nests in empty snail shells: *O. rufohirta* Lepetier de Saint Fargeau, *O. andrenoides* Spinola, *O. aurulenta* Panzer, *O. caerulescens* (Linnaeus), *O. ferruginea* Latreille, *O. spinulosa* (Kirby), *O. versicolor* Latreille. Russia: EP (C, E, NC), UR, WS (TK). – Europe (WE, NE, SE, EE), Caucasus, Central Asia.

Chrysura filiformis (Mocsáry, 1889) [Chrysis]. Russia: EP (NC). – Europe (WE, SE, EE), Caucasus, Turkey.

Chrysura hirsuta (Gerstaecker, 1869) [Chrysis] (*Chrysis bicolor*: Assmuss, 1862; *Ch. osmiæ*: Westerlund, 1892). Parasitoid of solitary bees: *Osmia inermis* (Zetterstedt), *O. nigriventris* (Zetterstedt), *O. parietina* Curtis, *O. uncinata* Gerstaecker, and *Hoplitis tuberculata* (Nylander) (Megachilidae). Russia: EP (N, NW, C), UR, ES (IR, BR, YA). – Europe (WE, NE, SE, EE), Caucasus, China (NE, NC, SW), Korean Peninsula, Japan (Hon, Shi).

Chrysura ignifrons (Brullé, 1833) [Chrysis] (*Chrysis aurifrons*: du Buysson, 1894; *Ch. gnifrons* (!): Vinokurov, 2006). Russia: EP (NC). – Europe (WE, SE), N Africa, Middle East, Central Asia.

Chrysura koma (Tsuneki, 1950) [Chrysis]. Russia: FE (PR). – China, Korean Peninsula.

Chrysura laevigata (Abeille de Perrin, 1879) [Chrysis] (*Chrysis dichroa* var. *loevigata* (!): Trautmann, 1927). Russia: EP (NC), UR. – Europe (WE, SE, EE), N Africa, Azerbaijan, Middle East, Iran.

Chrysis lydiae lydiae (Mocsáry, 1889) [Chrysis]. Russia: EP (NC). – Europe (SE), Turkey, Middle East.

Chrysura nikolaji Rosa, 2017 (*Chrysis medea* Semenov, 1967, nom. praeocc., nec Balthasar, 1953). Russia: EP (NC). – Abkhazia.

Chrysura oraniensis porphyrea (Mocsáry, 1889) [Chrysis] (*Chrysis oraniensis*: du Buysson, 1894). Parasitoid of *Anthidium sticticum* (Fabricius) (Megachilidae). Russia: EP (NC). – Europe (SE), Turkey.

Chrysura pseudodichroa (Linsenmaier, 1959) [Chrysis]. Russia: EP (NC, CR). – Europe (WE, SE, EE), Turkey, Middle East, Iran.

Chrysura purpureifrons (Abeille de Perrin, 1878) [Chrysis] (*Chrysis purpureifrons helleniensis* Linsenmaier, 1968). Russia: EP (C, S, NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Iran, Central Asia.

Chrysura radians (Harris, 1776) [Chrysis] (*Chrysis bicolor*: Radoszkowski, 1880; *Ch. pustulosa*: Mocsáry, 1889). Parasitoid of solitary bees (Megachilidae), which usually nest in cavities in dead wood: *Hoplitis adunca* (Panzer), *H. anthocopoides* (Schenck), *Osmia caerulescens* (Linnaeus), *O. leiana* (Kirby). Russia: **EP** (N, C, E, S, NC), **UR**, **ES** (IR, BR). – Europe (WE, NE, SE, EE), N Africa, Abkhazia, W Asia.

Chrysura refulgens (Spinola, 1806) [Chrysis]. Russia: **EP** (NC). – Europe (WE, SE, EE), N Africa, Caucasus, China (SE).

Chrysura simplex simplex (Dahlbom, 1854) [Chrysis]. Parasitoid of the solitary bee *Osmia caementaria* Gerstaecker (Megachilidae). Russia: **EP** (NC). – Europe (WE, SE, EE), N Africa, Caucasus.

Chrysura trimaculata (Förster, 1853) [Chrysis]. Parasitoid of solitary bees in the genus *Osmia* (Megachilidae). Russia: **ES** (TU). – Europe (WE, SE, EE), Caucasus, Turkey.

Chrysura varicornis (Spinola, 1838) [Chrysis] (*Chrysis sodalis* Mocsáry, 1892). Russia: **Southern Russia** (du Buysson, 1894). – Europe (WE, SE), N Africa, Turkey, Middle East, Central Asia.

EUCHROEUS Latreille, 1809 (*Brugmoia* Radoszkowski, 1877). Type species *Chrysis purpurata* Fabricius, 1775. The generic name *Euchroeus* Latreille, 1809 and the specific name *Chrysis purpurata* Fabricius, 1787 were conserved by IZN (1998). As a result, the generic name *Brugmoia* Radoszkowski, 1877 (used by Kimsey and Bohart (1991)) is a junior synonym of *Euchroeus* Latreille, 1809, and the name *quadrata* is a junior synonym of *purpurata* Fabricius. Number of species: World – 26, Palaearctic – 22, Russia – 6.

Euchroeus limbatus limbatus Dahlbom, 1854. Russia: **EP** (S, NC), **UR**. – Caucasus, Turkey, Central Asia.

Euchroeus mongolicus Tsuneki, 1947. Russia: **ES** (BR). – China (NC).

Euchroeus purpuratus purpuratus (Fabricius, 1787) [Chrysis] (*Euchraeus* (!) *quadratus*: Assmuss, 1862; *E. purpuratus* var. *smaragdinus* Trautmann, 1926; *Brugmoia quadrata*: Kurzenko, Lelej, 2007). Recorded as a possible parasitoid of *Ammophila* Kirby and *Podalonia* Fernald (Sphecidae). Russia: **EP** (C, E, S), **UR**, **WS** (OM, AL), **ES** (KS, KR, BR). – Europe (WE, SE, EE), Central Asia.

Euchroeus purpuratus irradians (Semenov, 1910) [Pseudochrysis]. Russia: **UR**, **ES** (KR). – Kyrgyzstan.

Euchroeus par (Semenov, 1967) [Spinolia]. Russia: **ES** (BR).

Euchroeus rugulosus (Mocsáry, 1909) [Chrysis] (*E. doursi*: Trautmann, 1927; *Brugmoia rugulae* Kimsey, 1991). Russia: **ER** (S, CR), **UR**, **ES** (KR). – Uzbekistan, Kyrgyzstan, Kazakhstan.

Euchroeus victorius Rosa, 2017. Russia: **ES** (KS).

PENTACHRYYSIS Lichtenstein, 1876. Type species *Chrysis amoena* Eversmann, 1857. Number of species: World – 6, Palaearctic – 5, Russia – 1.

Pentachrysis amoena (Eversmann, 1858) [Chrysis] (*Chrysis amöna* (!): Dalla Torre, 1892). Russia: **EP** (S), **WS** (OM). – Central Asia, Mongolia.

PSEUDOCHRYYSIS Semenov, 1891 (*Pseudospinolia* Linsenmaier, 1951; *Neospinolia* Linsenmaier, 1968). Type species *Chrysura humboldti* Dahlbom, 1845. Nest parasitoids of potter wasps (Vespidae: Eumeninae). Number of species: World – 18, Palaearctic – 17, Russia – 7.

Pseudochrysis aureicollis (Abeille de Perrin, 1879) [Chrysis]. Russia: **EP** (NC). – Europe (SE), N Africa, Caucasus, Iran.

Pseudochrysis fahringeri Trautmann, 1926. Russia: **WS** (AL), **ES** (KS, KR). – Europe (WE).

Pseudochrysis gengiskhan Rosa, 2017. Russia: **ES** (TU).

Pseudochrysis humboldti (Dahlbom, 1845) [Chrysis] (*Chrysis cyanura* Dahlbom, 1854; *Ch. fedtchenkoi*: Radoszkowski, 1880). Parasitoid of *Paravespa grandis albida* Blütgen (Vespidae: Eumeninae). Russia: **EP** (E, S, NC, CR), **UR**, **ES** (IR). – Europe (WE, SE, EE), Caucasus, Azerbaijan, Iran, Central Asia, China (NC).

Pseudochrysis incrassata (Spinola, 1838) [Chrysis] (*Euchroeus gratus minutus*: Linsenmaier, 1968). Russia: **EP** (S, NC). – Europe (WE, SE), Caucasus, Turkey.

Pseudochrysis neglecta (Shuckard, 1836) [Chrysis] (*Ch. austriaca* Dahlbom, 1829, nom. praeocc., nec Fabricius, 1804; *Ch. integrella* Dahlbom, 1854; *Ch. tadzhika* Semenov et Nikol'skaya, 1954). Nest parasite of *Odynerus spinipes* (Linnaeus) and *O. reniformis* (Gmelin) and possibly occasionally of *Ancistrocerus parietum* (Linnaeus) and *Gymnomerus laevipes* (Shuckard) (Vespidae: Eumeninae). Russia: **EP** (N, C, E, S, NC, CR), **UR**, **WS** (TK, AL), **ES** (KS, TU, KR, IR), **FE** (KH). – Europe (WE, NE, SE, EE), Turkey, Tajikistan, China (NW), N America.

Pseudochrysis uniformis (Dahlbom, 1854) [Chrysis] (*Chrysis fasciata* Spinola, 1805, nom. praeocc., nec Olivier, 1791). Russia: **EP** (E). – Europe (SE, EE), N Africa, Caucasus, Turkey, Asia, Iran.

SPINOLIA Dahlbom, 1854 (*Polyodontus* Radoszkowski, 1877; *Achrysis* Semenov, 1892). Type species *Spinolia magnifica* Dahlbom, 1854 [= *Spinolia lamprosoma* (Förster, 1853)]. Parasitoids of potter wasps (Vespidae: Eumeninae) and digger wasps (Crabronidae). Number of species: World – 19, Palaearctic – 19, Russia – 6.

Spinolia chalcites (Mocsáry, 1890) [Chrysis]. Russia: **EP** (E, S, NC). – Turkey, Iran.

Spinolia dournovii (Radoszkowski, 1866) [Chrysis] (*Ch. dournovoi* (!): Radoszkowski, 1880). Russia: **EP** (NC). – Europe (SE), N Africa, Turkey, Middle East, Iran.

Spinolia lamprosoma (Förster, 1853) [Chrysis] (*Spinolia* (!) *lamprosoma*: Vorontsovskij, 1930). Russia: **UR**. – Europe (WE, SE), Caucasus, Turkey.

Spinolia morawitzii (Mocsáry, 1889) [Chrysis] (*Ch. morawitzi* (!): Dalla Torre, 1892). Russia: **EP** (S). – Caucasus, Turkey, Turkmenistan.

Spinolia stchurovskyi (Radoszkowski, 1877) [Polyodontus] (*P. stchurovskyi* Radoszkowski, 1877). Russia: **EP** (S), **UR**. – Central Asia.

Spinolia unicolor (Dahlbom, 1831) [Chrysis]. Parasitoid of *Pterocheilus phaleratus* (Panzer) (Vespidae: Eumeninae) and possibly *Bembecinus hungaricus* (Frivaldszky) (Crabronidae). Russia: **EP** (C). – Europe (WE, NE, SE, EE), Mongolia.

SPINTHARINA Semenow, 1892. Type species *Chrysis vagans* Radoszkowski, 1877. Parasitoids of solitary wasps (Masarinae). Number of species: World – 26, Palaearctic – 21, Russia – 2.

Spintharina mocsaryi (Radoszkowski, 1890) [Spintharis] (*Chrysis cuprata*: Radoszkowski, 1880). Russia: **Southern Russia** (Linsenmaier, 1968). – Caucasus, Turkey, Middle East.

Spintharina versicolor (Spinola, 1808) [Chrysis] (*Ch. patriarchalis* Radoszkowski, 1880; *Ch. kuznetzovi* Semenov, 1967). Parasitoid of *Celonites abbreviatus* (Villers) (Vespidae: Masarinae). Russia: **EP** (CR). – Europe (WE, SE), Armenia, Iran.

STILBUM Spinola, 1806. Type species *Chrysis calens* Fabricius, 1781. Nest parasite of solitary

wasps (Vespidae: Eumeninae; Sphecidae) and bees (Megachilidae). Number of species: World – 5, Palaearctic – 3, Russia – 2.

Stilbum calens (Fabricius, 1781) [Chrysis] (*Stilbum nobile*: Mocsáry, 1882). Parasitoid of *Katamenes flavigularis* Blüthgen, *K. arbustorum* (Panzer), *Delta uncuiculatum* (Villers) (Vespidae: Eumeninae), *Sceliphron madraspatanum tubifex* (Latreille) (Sphecidae) and possibly *Chalicodoma parietina* (Geoffroy) (Megachilidae). Russia: **EP** (S, NC, CR), **UR**, **ES** (BR, ZB). – Europe (WE, SE, EE), N Africa, Central Asia, China (NE, NC), Japan (Hon, Shi, Kyu).

Stilbum cyanurum (Forster, 1771) [Chrysis] (*S. splendidum*: Radoszkowski, 1866; *S. amethystinum*: Mocsáry, 1882; *S. pici*: Linsenmaier, 1959). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), Caucasus, China (NE, NC, CC, SE). Worldwide except N and S America.

TRICHRYYSIS Lichtenstein, 1876 (*Alocochrysis* Haupt, 1956). Type species *Sphex cyanea* Linnaeus, 1758. Nest parasites of solitary wasps (Crabronidae and Pompilidae). Number of species: World – 36, Palaearctic – 11, Russia – 3.

Trichrysis cyanea (Linnaeus, 1758) [Sphex]. Parasitoid of *Trypoxylon figulus* Linnaeus, *Pemphredon rugifer* (Dahlbom) (Crabronidae) and *Auplopus carbonarius* (Scopoli) and *Dipogon* sp. (Pompilidae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TK, AL), **ES** (KS, TU, KR, IR, BR, ZB), **FE** (AM, KH, PR, SA, KA). – Europea (WE, SE, NE, EE), N Africa, Central Asia, China (NE, NC), Korean Peninsula, Japan (Hok, Hon).

Trichrysis lusca (Fabricius, 1804) [Chrysis] (*Praestochrysis lusca*: Kurzenko, Lelej, 2007). Russia: **FE** (PR). – China (NC, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu), SE Asia, India, Papua-New Guinea, Australia. Accidentally was introduced into Europe.

Trichrysis pellucida (du Buysson, 1887) [Chrysis] (*Chrysis buyssoni* Mocsáry, 1899, replacement name for secondary homonym *Ch. pellucida* du Buysson, 1887, nec *Brugmoia pellucida* Radoszkowski, 1877, currently *Euchroeus pellucidus* (Radoszkowski, 1877); *T. buyssoni*: Kurzenko, Lelej, 2007). Russia: **FE** (KH, PR, SA). – Turkey, China (NC, CC), Korean Peninsula.

Tribe PARNOPINI

Number of taxa: World – 3 genera, 22 species, Palaearctic – 3/9, Russia – 1/3.

PARNOPES Latreille, 1797. Type species *Chrysis carnea* Fabricius, 1775 [= *Parnopes grandior* (Pallas, 1771)]. Holarctic genus. Parasitoids of digger wasps in the tribe Bembicini (Crabronidae). Number of species: World – 17, Palaearctic – 6, Russia – 3.

Parnopes glasunowi Semenov, 1901. Russia: EP (S). – Central Asia.

Parnopes grandior (Pallas, 1771) [*Chrysis*] (*Parnopes carnea*: Eversmann, 1858). Parasitoid of *Bembix rostrata* (Linnaeus) (Crabronidae). Russia: EP (NW, C, E, S, NC, CR), UR, WS (NS), ES. – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Caucasus, Central Asia.

Parnopes popovii Eversmann, 1857 (*Parnopes popovi* (!): Bischoff, 1913). Russia: EP (C), UR, ES (KS, TU, KR, IR, BR, ZB), FE (AM, PR). – Mongolia, China (NE, NC), Korean Peninsula.

Species erroneously reported for Russia and to be excluded from Russian fauna

The following records were originally described in ex Soviet countries (Kazakhstan, Uzbekistan, etc.), and later reported as “southern Russia” and “Russia” and to be excluded from the Russian fauna, till precise records and identifications are available. We also include species considered erroneously or doubtfully identified.

Cleptes afer Lucas, 1849. Northern African species (Lisenmaier, 1999). Distributional data from Crimea are datable and should be related to *Cl. ignitus* other species subsequently described in the *nitidulus* group.

Cleptes mocsaryi Semenov, 1891. Currently the species is known only on specimens from Montenegro (type locality) and Greece (Arens, 2014).

Hedychridium cupratum (Dahlbom, 1854). Endemic species which is living only on the Alps (Austria, France, Germany, Italy, Switzerland) (Rosa et al., 2017d).

Hedychridium ibericum Lisenmaier, 1959. Endemic of the Iberian peninsula (Mingo, 1994). The record is related to another species in the *H. ardens* species-group.

Hedychridium lampadum Lisenmaier, 1959. It is an unnecessary replacement name for *H. lampas*

(Christ, 1790) (Kimsey, Bohart, 1991). *H. lampadum* Lisenmaier includes females of different species (Rosa, 2006; Arens, 2010).

Hedychridium maculiventre Lisenmaier, 1959. It could be referable to *H. palestinense* Balthasar, 1953 (Arens, 2010) or other species in the *H. roseum* species-group with metallic reflections on the second sternite.

Hedychrum coelestinum Spinola, 1838. African species, also known from Saudi Arabia in the Palaearctic (Lisenmaier, 1999). The record from Russia is very likely a misidentification by Kirchner (1867).

Hedychrum sinicum Semenov, 1967. Described from Sichuan (China) and misidentified by Vinokurov (2006).

Hedychrum gerstaeckeriformes (nom. nudum). Kilimnik (1993) labeled the holotype in ZIN as *Hedychrum gerstaeckeriformes*, but later described this species as *H. tobiasi*. Vinokurov (2006, 2007, 2009, 2013) reported both *H. gerstaeckeriformes* and *H. tobiasi*, therefore one of the two identification is related to an unknown species.

Holopyga gloriosa (Fabricius, 1793) This name has been used for a long time, although mistakenly, till today; in particular many European and Russian species have been considered as varieties of *H. gloriosa*. Previous identifications of Russian chrysids and identified as *H. gloriosa* must be referred to different species, in particular *H. inflammata*, *H. lucida*, *H. chrysonota* and females of *H. jurinei* and *H. ignicollis*. The name *Chrysis gloriosa* Fabricius, 1793 was suppressed by ICZN (1998, Opinion 1906).

Holopyga pulawskii Lisenmaier, 1968. This species described from Turkmenistan: Aşgabat, locality given as southern Russia by Kimsey and Bohart (1991).

Omalus imbecillus (Mocsáry, 1889). Citations of this species from Russia are related to specimens collected in Turkmenistan (Strumia, 2008).

Pseudomalus auratus virescens (Mocsáry, 1889). Móczár (1964) designated the lectotype and synonymised *P. auratus virescens* with *P. auratus* (Linnaeus). The type series includes specimens belonging to different species: *Pseudomalus auratus* (Linnaeus, 1758), *Omalus biaccinctus* du Buysson, 1893 and *Pseudomalus pusillus* (Fabricius, 1804).

Chrysis abeillei Gribodo, 1879. Balthasar's (1953) record of this species from Sarepta is based on Trautmann's (1927) wrong interpretation of *Chrysis abeillei*. The species is known only for Syria, so far.

Chrysis altaica Mocsáry, 1912. The holotype of this species originates from Kazakhstan: Semipalatinsk.

Chrysis angustula gracilis Schenck, 1856. Niehuis (2000), after type examination, synonymised *Chrysis gracilis* Schenck with *C. angustula* Schenck. Vinokurov (2010) listed both *C. angustula* and *C. angustula gracilis* for the same localities and his identification is possibly referable to *C. leptomandibularis* Niehuis, 2000 or *C. corusca* Valkeila, 1971.

Chrysis aurotecta Abeille de Perrin, 1878. It is an endemic species of Corsica and Sardinia which was misidentified by Vinokurov (2006).

Chrysis bidentata Linnaeus, 1767. It is a synonym of *Ch. viridula* Linnaeus, 1761. *Ch. bidentata* sensu Linsenmaier is a misidentification of *C. consanguinea* Mocsáry, 1889 (Linsenmaier, 1997). Specimens erroneously identified as *C. bidentata* usually belong to different species within the *Ch. viridula* species-group.

Chrysis bihamata Spinola, 1838. The record from Russia (Linsenmaier, 1969) is in error and referable to other species in the *C. bihamata* group.

Chrysis distincta exigua Mocsáry, 1889. The record from southern Russia (Linsenmaier, 1959) is related to Western Asian countries.

Chrysis distinguenda Spinola, 1838. It is a *nomen dubium* cited by Eversmann (1858).

Chrysis gribodoi Abeille de Perrin, 1877. The identification by Bischoff (1910) is doubtful and related to other species later described in the *Ch. succincta* species-group.

Chrysis grohmanni grohmanni Dahlbom, 1854. The distribution of this species is limited to western Mediterranean countries. Citations from Southern Russia, Middle East and central Asia are referable to other species or subspecies within the *Ch. succincta* species-group.

Chrysis imperatrix du Buysson, 1887. Dalla Torre (1892) reported “Russia” as type locality for *Chrysis imperatrix*, but the species was described from Turkmenistan and the species is not yet recorded for Russia.

Chrysis ignita bischoffi Linsenmaier, 1959. We examined some specimens identified by Vinokurov (2006) as *C. ignita bischoffi* who belong to different species and different subgroups within the *C. ignita* species-group, all with darkened colouration of metasoma.

Chrysis ignita melaensis Linsenmaier, 1968. It is an endemic subspecies restricted to the Corsica and Sardinia (Linsenmaier, 1987). The specimens

examined (Vinokurov, 2010) belong to different taxa with darkened colouration of the metasoma.

Chrysis mauritanica Trutmann, 1927. This species is known only for NW Africa: Tunisia and Morocco (Linsenmaier, 1959, 1999; Kimsey, Bohart, 1991). Vinokurov’s (2006, 2009) records are referable to other species in the *Ch. leachii* species-group.

Chrysis mediata berberiana Linsenmaier, 1959. This is a NW African species known from Algeria, Morocco and Tunisia (Linsenmaier, 1959, 1999). The identification (Vinokurov, 2010) is referable to other species in the *Ch. ignita* species-group.

Chrysis pseudoignita Linsenmaier, 1987. Species endemic of the Canary islands (Palma, Tenerife) and possibly conspecific with *C. magnidens* Peréz, 1895 also endemic of the Canary Islands. Vinokurov’s identifications (2010) are referable to different species.

Chrysis rutilans rigiana Linsenmaier, 1951. It was found only in a small Swiss Valley (Wallis, Mt. Rigi) so far, and specimens from North Caucasus (Vinokurov, 2006) are related to other species in the *Ch. splendidula* species-group.

Chrysis splendidula decora Mocsáry, 1887. It is the replacement name for *Ch. superba* Radoszkowski, also named *Ch. rutilans mesasiatica* or *Ch. insperata mesasiatica*. It is a small and slender species, not related to *Ch. splendidula*. Vinokurov’s (2004) identification is possibly related to a colour variation of *Ch. splendidula*.

Chrysura dichropis (du Buysson, 1894). Species known from Palestine and Syria. The record of this species from North Caucasus (Vinokurov, 2012) is referable to other species in the *Chrysura dichroa* species-group.

Chrysura fulminatrix (du Buysson, 1888). This species is known only on the holotype, collected in Algeria. The record from North Caucasus (Vinokurov, 2012) is referable to other species in the *Chrysura dichroa* species-group.

Chrysura interdichroa (Linsenmaier, 1959). This species is known only from specimens collected in Palestina, so far. The record of this species from North Caucasus (Vinokurov, 2012) is referable to other species in the *Chrysura dichroa* species-group.

Chrysura prodichroa (Linsenmaier, 1959). This is a synonym of *Chrysura isabellae* (Trautmann, 1926) (*C. radians* species-group). Vinokurov’s (2012) record is related to a species in the *C. dichroa* species-group.

Euchroeus orientis Semenov, 1910. The species described from Xinjiang (West China) and its record from East Russia (Linsenmaier, 1968) is erroneous.

Euchroeus purpuratus consularis du Buysson, 1896.
This subspecies is known only for NW Africa:

Algeria, Morocco and Tunisia (Linsenmaier, 1999). Records from southern Russia (Trautmann, 1927) are doubtful and related to other species later described by Semenov-Tian-Shanskij.

SUPERFAMILY SCOLIOIDEA

61. FAMILY SAPYGIDAE

A.S. LELEJ AND N.V. KURZENKO

Small and medial size wasps. Their larvae are clepto-parasites in the cells of megachilid bees (Megachilidae: *Ashmeadiella* Cockerell, *Osmia* Panzer, *Heriades* Spinola, *Hoplitis* Klug, *Chelostoma* Latreille, *Dianthidium* Cockerell), apid bees (Apidae: *Anthophora* Latreille, *Xylocopa* Latreille), or colletid bees (Colletidae: *Hylaeus* Fabricius). Nearctic *Fedtschenkia anthracina* (Ashmead) is the parasitoid of *Pterocheilus trichogaster* Bohart (Vespidae: Eumeninae). On eclosion from its egg the first instar sapygid larva destroys and consumes the egg of the host; later instars consume the food supplied in the cells, some Nearctic species are ectoparasitoids on host larvae. Includes two recent subfamilies: Fedtschenkiinae (1 genus, 5 species from arid area of Central Asia and North America) and Sapyginae (11 genera, 66 species, distributed worldwide, except Australia).

Number of taxa: World – 12 genera, 66 species, Palaearctic – 6/31, Russia – 5/9.

R e f e r e n c e s . Dalla Torre, 1897; Gussakovskij, 1932; Sugihara, Kim, 1936; Yasumatsu, Sugihara, 1938; Yasumatsu, 1939; Pate, 1947a,b; Tobias, 1965b, 1978; Guiglia, 1969, 1971; Krombein, 1979; Kurzenko, 1986, 1995, 1996, 2004, 2009; Kurzenko, Gusenleitner, 1994; Kurzenko et al., 1995; Matsumoto et al., 2007; Kurzenko, 2012.

Subfamily SAPYGINAE

MONOSAPYGA Pic, 1920. Type species *Sapyga (Monosapyga) theresae* Pic, 1920. Number of species: World – 2, Palaearctic – 2, Russia – 1.

Monosapyga clavicornis (Linnaeus, 1758) [Apis] (*Scolia prisma* Fabricius, 1787; *Masaris crabroniformis* Panzer, 1797; *Sapyga theresae* Pic, 1920; *S. triinterrupta* Pic, 1920; *S. ininterrupta* Pic, 1920). Cleptoparasite of bees *Chelostoma florisomne* (Linnaeus) and *Osmia* Panzer (Megachilidae). Russia: EP (C, CR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Armenia, Azerbaijan, Turkey.

POLOCHRIDIUM Gussakovskij, 1932. Type species *Polochridium eoum* Gussakovskij, 1932. Monotypic genus.

Polochridium eoum Gussakovskij, 1932 (*P. koreanum* Sugihara et Kim, 1936). Russia: FE (PR). – China (NC), Korean Peninsula.

POLOCHRUM Spinola, 1805 (*Dimorpha* Panzer, 1806; *Aclastocera* Foerster, 1855). Type species *Polochrum repandum* Spinola, 1805. Number of species: World – 3, Palaearctic – 2, Russia – 1.

Polochrum repandum Spinola, 1805 (*Aclastocera frivaldszki* Förster, 1855). Cleptoparasite of *Xylocopa violacea* Linnaeus (Apidae). Russia: EP (CR). – Europe (WE, SE, EE), Caucasus, Azerbaijan, Turkey.

SAPYGA Latreille, 1796 (*Hellus* Fabricius, 1804). Type species *Scolia quinquepunctata* Fabricius, 1781. The genus includes four subgenera: *Arthrosapya* Pic, 1920; *Polosapyga* Kurzenko, 1994; *Recotosapyga* Kurzenko, 1986, and *Sapyga* Latreille, 1796. Number of species: World – 28, Palaearctic – 17, Russia – 5.

Sapyga (Sapyga) caucasica schevyrevi Morawitz, 1889. Russia: EP (NC). – Caucasus, Turkey, Iran.

Sapyga (Sapyga) coma Yasumatsu et Sugihara, 1938. Cleptoparasite of *Osmia pedicornis* Cockerell and *O. cornifrons* Radoszkowski (Megachilidae). Russia: FE (PR). – China (CC), Korean Peninsula, Japan (Hon).

Sapyga (Sapyga) morawitzi Turner, 1911 (*S. gussakovskii* Kurzenko, 1986). Russia: South Russia (Gusenleitner, 2013c). – Europe (SE), Caucasus, Turkey, Iran.

Sapyga (Sapyga) quinquepunctata (Fabricius, 1781) [*Scolia quadriguttata* Fabricius, 1781; *Sphex tricolor* Schrank, 1781; *Sirex pacca* Fabricius, 1787; *Scolia sexguttata* Fabricius, 1793; *S. octoguttata* Fabricius, 1793; *Sapyga punctata* Klug, 1803; *Hellus sexpunctatus* Fabricius, 1804; *H. quadriguttatus* Fabricius, 1804; *Sapyga punctum* Lamarck, 1817; *S. varia* Lepeletier et Audinet-Serville, 1825; *S. decipiens* Lepeletier et Audinet-Serville, 1825; *S. quinquepunctata* var. *arvenica* Pic, 1925). Cleptoparasite of *Chalicodoma* Latreille, *Osmia* Panzer, *Heriades* Spinola, *Megachile* Latreille (Megachilidae). Russia: EP (C, S, CR). – Europe (WE, SE, NE, EE), N Africa, Caucasus, Turkey.

Sapyga (Sapyga) similis (Fabricius, 1793) [*Sirex (Sapyga variegata* Dahlbom, 1844; *S. exornata* Gerstäcker, 1861; *S. pedestris* Gerstäcker, 1861; *S. rufipes* A. Costa, 1864). Cleptoparasite of *Osmia* Panzer, *Chelostoma* Latreille (Megachilidae). Russia:

EP (N, NW, C), **WS** (TK), **ES** (IR, BR, YA), **FE** (AM, KH, PR, SA, MG). – Europe (WE, NE, SE, EE). Kazakhstan, Mongolia, China (NC).

SAPYGINA A.Costa, 1887. Type species *Sapyga decemguttata* Jurine, 1807. The genus includes two subgenera: *Sapygina* Costa, 1887 and *Sapygella* Benoit, 1950. Number of species: World – 13, Palaearctic – 4, Russia – 1.

Sapygina decemguttata (Jurine, 1807) [Sapyga] (*Meria millefolii* Lepeletier et Audinet-Serville, 1825; *Sapyga chelostomae* Robineau-Desvoidy, 1836; *Myzine panzeri* Guérin-Méneville, 1837; *Polochrum cylindricum* Schenk, 1857; *Sapyga nigra* Tournier, 1889). Russia: **EP** (CR). – Europe (WE, SE, EE), Turkey, Israel.

62. FAMILY SCOLIIDAE

M.V. MOKROUSOV AND A.S. LELEJ

Large, robust, hairy wasps, mostly 20–30 mm, usually black with a yellow band on the metasoma. Larvae of these wasps are ectoparasitoids of the larvae of scarabaeid beetles (Scarabaeoidea). The adults are commonly found on flowers. The females burrow into the ground to locate a host and when they find the grub, they sting it and paralyze it, then burrow deeper into the soil and construct a cell around the grub. Number of taxa: World – 43(143) genera, 560 species, Palaearctic – 11/100, Russia – 5/20.

R e f e r e n c e s. Pallas, 1771; Eversmann, 1846, 1849; Erichson, 1849; Bischoff, Bradley, 1929; Steinberg, 1962; Bradley, 1972; Hamon, 1993; Argaman, 1996; Osten, 1999, 2005; Terayama, Nagase, 2016; Lelej, Mokrousov, 2017.

Subfamily SCOLIINAE

Tribe CAMPSOMERINI

COLPA Dufour, 1841 (*Trielis* de Saussure, 1863; *Campsoscolia* Betrem, 1933). Type species *Scolia interrupta* Fabricius, 1781 (=*Vespa galbula* Pallas, 1771). The genus includes four subgenera: *Colpa* Dufour, 1841; *Carbonelis* Betrem, 1972; *Crioscolia* Bradley, 1951; *Heterelis* Costa, 1887. Number of species: World – 31, Palaearctic – 12, Russia – 4.

Colpa (Colpa) galbula (Pallas, 1771) [Vespa] (*Scolia sexmaculata* Fabricius, 1781; *S. interrupta* Fabricius, 1781; *Sphex insubrica* Scopoli, 1786; *Scolia sareptana* Eversmann, 1849). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), Algeria, Georgia, Azerbaijan, Cyprus, Turkey, W Kazakhstan.

Colpa (Colpa) klugii (Vander Linden, 1827) [*Scolia*]. Russia: **EP** (CR). – Europe (SE, EE), Turkey, Iran.

Colpa (Crioscolia) tartara (de Saussure, 1880) [Elis] (*Scolia mongolica* F. Morawitz, 1889). Russia: **EP** (NC). – Armenia, Afghanistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, China (NW).

Colpa (Heterelis) quinquecincta (Fabricius, 1793) [*Scolia*] (*Scolia rufiventris* Fischer von Waldheim, 1843, nom. praeocc., nec Fabricius, 1804; *S. grisea* Eversmann, 1849; *S. vetula* Eversmann, 1849). Russia: **EP** (C, S, NC, CR), **UR**, **WS** (AL). – Europe (WE, SE, EE), N Africa, Georgia, Azerbaijan,

Turkey, Israel, Iraq, Iran, W Pakistan, Central Asia, Kazakhstan.

MEGACAMPSOMERIS Betrem, 1928 (*Fiharbuxa* Argaman, 1996). Type species *Tiphia grossa* Fabricius, 1804. Number of species: World – 40, Palaearctic – 6, Russia – 2.

Megacampsomeris grossa matsumurai (Betrem, 1941) [*Campsomeris*]. Russia: **FE** (PR). – Japan (Hon).

Megacampsomeris prismatica (Smith, 1855) [*Scolia*] (*Elis iris* Magretti, 1892; *Scolia simillima* Tullgren, 1904; *M. uchidai* Betrem, 1941). Russia: **FE** (PR). – China (NE, NC, CC, SW, SE), Korean Peninsula, Japan (Hon), SE Asia, India.

Tribe SCOLIINI

CARINOSCOLIA Betrem, 1927 (*Nokbibula* Argaman, 1996). Type species *Scolia opalina* Smith, 1857. Number of species: World – 8, Palaearctic – 2, Russia – 1.

Carinoscolia vittifrons (Sichel in de Saussure et Sichel, 1864) [*Scolia*] (*S. fascinata* Smith, 1873; *Discolia signatifrons* Pérez, 1905; *D. fasciata* Matsuura, 1917). Russia: **FE** (AM, KH, PR). – China (NE, SW, SE), Korean Peninsula, Japan (Hon, Kyu, Yakushima, Ryu).

MEGASCOLIA Betrem, 1928. Type species *Scolia procer* Illiger, 1802. The genus includes two subgenera: *Megascolia* Betrem, 1928 and *Regiscolia* Betrem et Bradley, 1964. Number of species: World – 15, Palaearctic – 3, Russia – 1.

Megascolia (Regiscolia) maculata (Drury, 1773) [*Sphex*] (*Scolia flavifrons* Fabricius, 1775; *S. haemorrhoidalis* Fabricius, 1787). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Abkhazia, Caucasus, Cyprus, Turkey, Syria, Israel, Iran, Turkmenistan, W Kazakhstan.

SCOLIA Fabricius, 1775 (*Lisoca* A. Costa, 1858; *Scolioides* Guiglia et Capra, 1934; *Kokarevta* Argaman, 1996; *Buzatlana* Argaman, 1996; *Orlovinga* Argaman, 1996; *Lisoca* Argaman, 1996; *Pardesiya* Argaman, 1996). Type species *Scolia quadrivittata* Fabricius, 1775 (=*Vespa sexmaculata* O.F. Müller, 1766). The genus includes two subgenera: *Discolia* de Saussure, 1863 and *Scolia* Fabricius, 1775. Number of species: World – more than 200, Palaearctic – 50, Russia – 12.

Scolia (Discolia) hirta (Schrink, 1781) [Apis] (*Discolia alutus* Nagy, 1967; *D. mongolina* Nagy, 1970). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (TM, AL), **ES** (KS, TU, KR, IR, BR, ZB). – Europe (WE, NE, SE, EE), Abkhazia, Caucasus, Cyprus, Turkey, Israel, Iran, Turkmenistan, Uzbekistan, Kazakhstan, Mongolia.

Scolia (Discolia) histrionica (Fabricius, 1787) [Tiphia] (*S. loebischii* Dalla Torre, 1897; *S. palaeanarctica* Gussakovskij, 1932; *S. decorata* auct.). Russia: **FE** (AM, KH, PR, KU). – Mongolia, China (NE, NC, CC, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Yakushima), SE Asia.

Scolia (Discolia) oculata (Matsumura, 1911) [Discolia] (*S. pseudounifasciata* Betrem, 1928). Russia: **FE** (KH, PR). – China (NC, CC, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Tsushima, Yakushima).

Scolia (Discolia) schrenckii Eversmann, 1846. Russia: **WS** (TM, AL), **ES** (IR). – Kazakhstan, Uzbekistan, Kyrgyzstan, Mongolia, China (NW).

Scolia (Discolia) sinensis de Saussure in de Saussure et Sichel, 1864 (*S. indica* Uchida, 1925). Russia: **FE** (KH, PR). – Pakistan, Central Asia, SE Kazakhstan, China (NE, NC, CC, SW, WP, SE), Korean Peninsula, Japan (Hon), India (Jammu and Kashmir).

Scolia (Scolia) fallax fallax Eversmann, 1849, nom. protectum (*Vespa tricolor* Pallas, 1771, nom. oblitum; *S. moreana* Muche, 1962; *S. galbula* auct.). Russia: **EP** (E, S, CR). – Europe (SE, EE), Azerbaijan, Kazakhstan.

Scolia (Scolia) fallax syriacola Betrem, 1935. Russia: **EP** (NC). – Georgia, Azerbaijan, Turkey, Syria, Lebanon, Israel, Iran, Turkmenistan.

Scolia (Scolia) fuciformis (Scopoli, 1786) [Sphe] (*Scolia insubrica* Rossi, 1790, nom. praeocc., nec *Sphex insubrica* Scopoli, 1786; *Scolia amabilis* Eversmann, 1849). Russia: **EP** (C, E, S, NC, CR), **ES** (IR). – Europe (WE, SE, EE), Azerbaijan, Cyprus, Turkey, Israel, Iran, Kazakhstan.

Scolia (Scolia) garrula Erichson in Ménétriés, 1849 (*S. laeta* Eversmann, 1849; *S. menetriesi* de Saussure, 1859). Russia: **EP** (S). – Central Asia, Kazakhstan.

Scolia (Scolia) gussakovskii Steinberg, 1953. Russia: **EP** (S). – Europe (EE), Turkmenistan, Uzbekistan, Kazakhstan.

Scolia (Scolia) potanini F. Morawitz, 1889. Russia: **ES** (BR), **FE** (AM). – Mongolia, China (NE, NC).

Scolia (Scolia) quadricincta quadricincta (Scopoli, 1787) [Sphe] (*Scolia dejeani* Vander Linden, 1829; *S. flaviceps* Eversmann, 1846; *S. galbula* auct.). Russia: **EP** (C, S, NC, CR). – Europe (WE, SE, EE), Georgia, Azerbaijan, Kyrgyzstan, Kazakhstan, W Mongolia.

Scolia (Scolia) quadricincta mangichlakensis Radoszkowski, 1879 (*S. flaviceps flaviceps*: Steinberg, 1962; *S. flaviceps mangichlakensis*: Steinberg, 1962). Russia: **EP** (NC). – Caucasus, Cyprus, Turkey, Israel, Iran, Turkmenistan, Tajikistan, Uzbekistan, S Kazakhstan.

Scolia (Scolia) sexmaculata (O.F. Müller, 1766) [Vespa] (*S. quadripunctata* Fabricius, 1775). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (TK), **ES** (KS, IR). – Europe (WE, SE, EE), N Africa, Abkhazia, Caucasus, Cyprus, Turkey, Syria, Lebanon, Israel, Iran, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan.

63. FAMILY TIPHIIDAE

M.V. MOKROUSOV AND A.S. LELEJ

Ectoparasitoids of beetles larva (Coleoptera). Distributed worldwide, mainly in arid, tropical and subtropical regions. Number of taxa: Word – 7 subfamilies, 120 genera, 2000 species; Palaearctic – 4/17/100; Russia – 3/6/28.

References. Tobias, 1978; Kimsey, 1991; Argaman, 1992, 1994; Agnoli, 2005; Boni Bartalucci, 2010, 2016; Terayama, 2016; Mokrousov, 2018.

Subfamily METHOCHINAE

Ectoparasitoids of subterranean tiger beetles larvae (Cicindelidae). The subfamily is distributed worldwide except Australian region. Number of taxa: Word – 2 genera, 75 species; Palaearctic – 1/10, Russia – 1/2.

METHOCHA Latreille, 1804 (*Methocha* Latreille, 1805; *Metocha* auct.; *Metoca* auct.). Type species *Mutilla articulata* Latreille, 1792. The genus includes three subgenera: *Methocha* Latreille, 1804, *Dryinopsis* Brues, 1910, and *Andreus* Ashmead, 1903. Number of species: World – 66, Palaearctic – 5, Russia – 2.

Methocha (Dryinopsis) picipes (F. Morawitz, 1890) [*Methocha*]. Russia: EP (C, NC), ES (IR, BR, ZB), FE (AM, KH, PR). – Europe (EE), Turkmenistan, Tajikistan, Uzbekistan.

Methocha (Methocha) articulata (Latreille, 1792) [*Mutilla*] (*Mutilla formicaria* Latreille, 1792; *Methocha ichneumonides* Latreille, 1805; *Methocha mutillaria* Latreille, 1806; *Tengyra sanvitali* Latreille, 1809; *Gonatopus mutillarius* Nees, 1834; *Spinolia italicica* Costa, 1858; *Methocha ichneumonoides* auct.). Russia: EP (NW, C, E, S, NC), ES (IR, BR, ZB), FE (AM, PR, SA). – Europe (WE, NE, SE, EE), N Africa, Turkey, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Subfamily MYZININAE

Ectoparasitoids of subterranean beetle larvae of Scarabaeidae, Cerambycidae and Tenebrionidae. The subfamily is distributed worldwide. Number of taxa: Word – 24 genera; Palaearctic – 10; Russia – 3 genera, 12 species.

MERIA Illiger, 1807 (*Arapatka* Argaman, 1994; *Fuk-pokta* Argaman, 1994; *Tilkuya* Argaman, 1994; *Upanterka* Argaman, 1994). Type species *Tiphia tripunctata* Rossi, 1790. Parasitoids of tenebrionid larvae (Tenebrionidae). Widespread in the Palaearctic and Afrotropical regions. Number of species: World – 72, Palaearctic – 35, Russia – 6.

Meria askhabadensis Radoszkowski, 1886 (*Plesia baluchistanensis* Cameron, 1907). Russia: EP (S). – Armenia, Azerbaijan, Turkey, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NW).

Meria aurantiaca (Guérin-Méneville, 1837) [Myzine] (*Meria soikai* Guiglia, 1966; *M. thoracica*: Tobias, 1978). Russia: EP (S, NC). UR. – Europe (SE), Armenia, Azerbaijan, Turkey, Jordan, Iran, Turkmenistan, Kazakhstan.

Meria dorsalis (Fabricius, 1804) [Bethylus] (*Myzine polita* Tournier, 1889; *Meria cylindrica* auct.; *M. lineata* auct.). Russia: EP (E, S, NC, CR), UR, ES (TU, IR, BR, ZB). – Europe (WE, SE, EE), Armenia, Turkey, Kazakhstan, Mongolia.

Meria geniculata (Brullé, 1832) [Myzine] (*Pseudomeria graeca* S. Saunders, 1850; *P. swanetiae* Radoszkowski, 1861; *Myzine nigriceps* Mocsáry, 1883; *M. graeca* Mocsáry, 1883; *Tiphia picta* Schulthess, 1893; *Meria nigrigena* Guiglia, 1966). Russia: EP (S). – Europe (SE), Armenia, Azerbaijan, Turkey, Kazakhstan.

Meria nitidula Klug, 1810. Russia: EP (C, S, NC), UR, ES (BR). – Europe (WE, SE, EE), Turkey, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan.

Meria quadrimaculata Cameron, 1902 (*M. discussa* Guiglia, 1966; *M. investiganda* Guiglia, 1973; *M. perfecta* Guiglia, 1975; *M. similis* Guiglia, 1975). Russia: EP (S). – Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NW).

MYZINELLA Guiglia, 1959 (*Illoswia* Argaman, 1994). Type species *Myzinella patrizii* Guiglia, 1959. Hosts unknown. Palaearctic and Afrotropical regions. Number of species: World – 10; Palaearctic – 7; Russia – 1.

Myzinella flavidollis (F. Morawitz, 1896) [Meria]. Russia: EP (S). – Afghanistan, Turkmenistan, Uzbekistan, Kazakhstan, Mongolia.

POECILOTIPHIA Cameron, 1902 (*Dermasothes* Menozzi, 1940; *Fongiba* Argaman, 1994; *Hahiya* Argaman, 1994; *Ivazuga* Argaman, 1994; *Jurja* Argaman, 1994; *Nurmiya* Argaman, 1994). Type species

- Poecilotiphia albomaculata** Cameron, 1902. Palaearctic and Noth of Afrotropical region. Number of species: World – 48, Palaearctic – 39, Russia – 4.
- Poecilotiphia brevicauda brevicauda** (F. Morawitz, 1890) [Meria]. Russia: EP (S). – Kazakhstan.
- Poecilotiphia brevicauda kaszabi** (Guiglia, 1965) [Meria]. Russia: WS (AL), ES (TU, ZB). – Mongolia, China (NC).
- Poecilotiphia caucasia** Boni Bartalucci, 2016. Russia: EP (S, NC). – Azerbaijan, Turkey.
- Poecilotiphia ciliata** (F. Morawitz, 1894) [Meria]. Russia: EP (S). – Iran, Uzbekistan, Kazakhstan.
- Poecilotiphia rugosopunctata** (Tournier, 1889) [Myzine]. Russia: EP (S). – Kazakhstan.

Subfamily TIPHIINAE

Ectoparasitoids of subterranean scarab beetle larvae (Scarabaeidae). Distributed worldwide. Number of taxa: Word – 10 genera; Palaearctic – 5; Russia – 2 genera, 11 species.

PSEUDOTIPHIA Ashmead, 1903 (*Icronatha* Nagy, 1967; *Ludita* Nagy, 1967). Type species *Tiphia brevipennis* Lucas, 1846. Palaearctic and Afrotropical regions. Number of species: World – 20, Palaearctic – 16, Russia – 4.

Pseudotiphia beckeri (Tournier, 1889) [Tiphia]. Russia: EP (S).

Pseudotiphia caucasica (Mocsáry, 1883) [Tiphia]. Russia: EP (NC). – Europe (SE, EE), N Africa, Georgia, Armenia, Azerbaijan, Turkey, Syria, Jordan, Lebanon, Iran.

Pseudotiphia fulvipennis (F. Smith, 1879) [Tiphia]. Russia: EP (E, S, NC, CR). – Europe (SE, EE), Azerbaijan, Turkey.

Pseudotiphia villosa (Fabricius, 1793) [Tiphia] (*Tiphia major* Mocsáry, 1883; *T. olcesei* Tournier, 1889; *T. rufepunctata* Tournier, 1901; *T. morio* auct.). Russia: EP (C, E, S, NC, CR). – Europe (WE, SE, EE), N Africa, Turkey, Iran, Kazakhstan.

TIPHIA Fabricius, 1775 (*Serpapinta* Argaman, 1992; *Sasmarila* Argaman, 1992). Type species *Tiphia femorata* Fabricius, 1775. Distributed worldwide. The genus includes four subgenera: *Tiphia* Fabricius, 1775, *Sierocolpa* Nagy, 1967, *Jaynesia* Allen, 1969, *Punctotiphia* Tsuneki, 1986. Number of species: World – more than 700 species, Russia – 11.

Tiphia (Sierocolpa) minuta Vander Linden, 1827. Russia: EP (N, NW, C, E, S, NC), WS (AL), ES (IR), FE (PR). – Europe (SE, EE), Armenia, Azerbaijan, Turkey, Iran, Tajikistan, Uzbekistan, Kyrgyzstan, Mongolia.

Tiphia (Tiphia) agilis Smith, 1873 (*T. asericae* Allen et Jaynes, 1930). Russia: FE (AM, KH, PR). – China (NE, NC, CC, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu), N America (introduced).

Tiphia (Tiphia) femorata Fabricius, 1775 (*Sphex palipes* Schrank, 1781; *Bethylus pilipennis* Klug, 1808; *B. ater* Klug, 1810; *S. nigripes* Costa, 1858; *S. rufipes* Costa, 1858; *Tiphia infima* Tournier, 1889; *T. lativentris* Tournier, 1889; *T. rugosa* Tournier, 1889; *T. tournieri* Dalla Torre, 1891). Russia: EP, UR, ES (BR, ZB), FE (AM). – Europe (WE, NE, SE, EE), N Africa, Turkey, Iran, Tajikistan, Kyrgyzstan, Kazakhstan, Mongolia, N America (introduced).

Tiphia (Tiphia) ordinaria Smith, 1873 (*T. bicarinata* Cameron, 1902; *T. chosensis* Allen, 1969). Russia: FE (PR). – China (SE), Korean Peninsula, Japan (Hok, Hon, Kyu), India.

Tiphia (Tiphia) ovidorsalis Allen et Jaynes, 1930. Russia: FE (AM, PR). – China (NC), Korean Peninsula.

Tiphia (Tiphia) ovinigris Allen et Jaynes, 1930. Russia: FE (AM, KH, PR). – Korean Peninsula.

Tiphia (Tiphia) popilliavora Rohwer, 1924. Russia: FE (AM, PR). – China (NE, NC, CC, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu), N America (introduced).

Tiphia (Tiphia) sareptana Tournier, 1889. Russia: EP (C, E, S, NC, CR).

Tiphia (Tiphia) singularis Allen et Jaynes, 1930. Russia: FE (PR). – China (CC, SE).

Tiphia (Tiphia) unicolor Lepeletier de Saint Fargeau, 1845 (*Bethylus ruficornis* Klug, 1810; *Tiphia polita* Costa, 1858). Russia: EP (C, S, NC), WS (AL). – Europe (WE, SE, EE), Turkey.

Tiphia (Tiphia) vernalis Rohwer, 1924. Russia: FE (AM, PR). – China (CC, SE), Korean Peninsula, Japan (Hon, Kyu), N America (introduced).

Doubtful species

Meria sexpunctata Eversmann, 1849 “*Cepi in promotoriis Uralensibus*”.

64. FAMILY SIEROLOMORPHIDAE

A.S. LELEJ AND M.V. MOKROUsov

A small family which is known from three genera: *Sierolomorpha* Ashmead, 1903, *Proscleroderma* Kieffer, 1905 and fossil *Loreisomorpha* Rasnitsyn, 2000. *Sierolomorpha* comprises seven described species in North America, one in Hawaii and three in the Old World (Uzbekistan, Mongolia and Russian Far East). *Proscleroderma* comprises one subapterous species from Syria. The fossil *Loreisomorpha nascimbenei* Rasnitsyn, 2000 was described from Upper Cretaceous amber of New Jersey. Brothers considered Sierolomorphidae to be the sister-group of other vespoids (Brothers, Carpenter, 1993, Brothers, 1999), while Rasnitsyn (2002) considered Sierolomorphidae to be the sister-group of the fossil Falsiformicidae within the superfamily Scolioidea. The larvae of sierolomorphids are probably ectoparasitoids of other insects, but nothing is known about their biology (Brothers, Finnimore, 1993). Three species of *Sierolomorpha* were recorded from the Old

World (Nagy, 1971; Lelej, Mokrousov, 2015). Sexual dimorphism slight to marked: both sexes usually macropterous but females sometimes brachypterous or apterous; brachypterous and apterous forms with mesosoma similar to that of in macropterous form with pronotum freely articulating, and with mesonotum, metanotum, and propodeum distinct.

Number of taxa: World – 2 genera, 13 species, Palaearctic – 2/5, Russia – 1/2.

References. Eversmann, 1849; Nagy, 1971; Argaman, 1990; Brothers, Carpenter, 1993; Brothers, Finnimore, 1993; Gorbatovskij, Lelej, 1998; Brothers, 1999; Rasnitsyn, 2002; Lelej, Mokrousov, 2015.

SIEROLOMORPHA Ashmead, 1903. Type species *Sierola ambigua* Ashmead, 1893. Number of species: World – 12, Palaearctic – 4, Russia – 2.

Sierolomorpha abnormis (Eversmann, 1849) [Tiphia].
Russia: UR, WS (NS).

Sierolomorpha atropos Nagy, 1971. Russia: FE (PR). – Mongolia.

65. FAMILY MUTILLIDAE – VELVET ANTS

A.S. LELEJ

Mutillidae currently include 216 genera and about 4300 described species (Lelej, 2007; Lelej, Brothers, 2008; Aguiar et al., 2013, updated). In the Palaearctic region 525 species in 61 genera and in the Oriental region 640 species in 64 genera are reported (Lelej, 2002, updated; Lelej, 2005a), but the hosts information are known for only about 150 species and are mainly bees and wasps (Brothers, 1989). Because of extreme sexual dimorphism (female apterous, male macropterous, rare apterous), sex associations cannot be made on morphology alone; most species and even many genera are known from one sex only. This has resulted in many taxonomic challenges in trying to make associations, and resulted in many synonyms recognized through matching of males and females. The males is very much larger than the females in some species and may then carry the females before mating, or true phoretic copulation may occur. Most velvet ants are solitary ectoparasitoids of the enclosed immatures (usually larvae or pupae) of other insects: Hymenoptera, mainly aculeate (Brothers et al., 2000), Diptera (Amini et al., 2014), Lepidoptera, Coleoptera (Sergeev, Lelej, 2011), and Blattodea.

Number of taxa: World – 216 genera, about 4300 species, Palaearctic – 61/525, Russia – 20/74.

R e f e r e n c e s. Pallas, 1771, 1773; Nylander, 1847; Baer, 1848; Morawitz A., 1865; Radoszkowski, 1865, 1866; Sichel, Radoszkowski, 1869–1870; Morawitz F., 1891, 1893; André, 1902; Skorikov, 1927, 1935; Chen, 1957; Pulawski, 1963; Invrea, 1964; Tsuneki, 1972; Brorhers, 1975; Lelej, 1978, 1985, 2002, 2005, 2007, 2012b; Petersen, 1988; Suárez, 1988; Brothers, 1989; Lelej, Nemkov, 1997; Brothers et al., 2000; Lelej, Schmid-Egger, 2005; Bogusch, 2006; Pagliano, Strumia, 2007; Lelej, Brothers, 2008; Muskovitz, György, 2011; Sergeev, Lelej, 2011; Aguiar et al., 2013; Amini et al., 2014; Lelej et al., 2014, 2016, 2017; Lelej, Shlyakhtenok, 2015; Humala, Polevoi, 2015.

Subfamily MYRMOSINAE

Tribe KUDAKRUMIINI

Number of taxa: World – 5 genera, 16 species, Palaearctic – 1/4, Russia – 1/2.

PSEUDOMYRMOSA Suárez, 1980. Type species *Myrmosa minuta* Morawitz, 1894. Number of species: World – 5, Palaearctic – 5, Russia – 2.

Pseudomyrmosa gobicola Lelej, 1981. Collected in the nest aggregations of *Hylaeus* spp. (Colletidae) and small species of Crabronidae. Russia: **ES** (TU). – Mongolia.

Pseudomyrmosa minuta (Morawitz, 1894) [Myrmosa] (*Mutilla myrmosina* Dalla Torre, 1897). Russia: **EP** (S, NC). – Armenia, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan.

Tribe MYRMOSINI

Number of taxa: World – 7 genera, 40 species, Palaearctic – 6/32, Russia – 4/7.

KROMBEINELLA Pate, 1947 (*Paramyrmosa* Suárez, 1963, nom. praeocc., nec Saussure, 1880). Type species *Hylaeus thoracicus* Fabricius, 1793. Number of species: World – 17, Palaearctic – 17, Russia – 2.

Krombeinella longicollis (Tournier, 1889) [Myrmosa]. Russia: **EP** (S, NC, CR). – Europe (SE, EE), Turkey.

Krombeinella wolfi (Invrea, 1963) [Myrmosa]. Russia: **EP** (CR). – Europe (SE, EE), Georgia, Azerbaijan, Turkey, Kazakhstan.

MYRMOSA Latreille, 1797 (*Ischioceras* Provancher, 1882). Type species *Myrmosa atra* Panzer, 1801. Number of species: World – 6, Palaearctic – 2, Russia – 1.

Myrmosa atra atra Panzer, 1801 (*Mutilla melanoccephala* Fabricius, 1793, nom. praeocc., nec Schrank, 1781; *Myrmosa nigra* Latreille, 1809; *M. atra* Lepeletier, 1845, nom. preocc., nec Panzer, 1801; *M. nigra* Lepeletier, 1845, nom. praeocc., nec Latreille, 1809; *M. bicolor* Baer, 1848; *Mutilla melanaria* Blanchard, 1849; *Myrmosa melanoccephala* f. *rufomaculata* Hoffer, 1938). Parasitoid of *Crabro peltarius* (Schreber), *Crossocerus palmipes* (L.), *C. wesmaeli* (Vander Linden), *Diodontus tristis* (Vander Linden), *Oxybelus uniglumis* (L.), *Lindenius panzeri* (Vander Linden), and *L. albilabris* (F.) (Crabronidae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (KS, TU, KR, YA). – Europe (WE, NE, SE, EE), Turkey, ?Iran, Kyrgyzstan, Kazakhstan.

PARAMYRMOSA Saussure, 1880 (*Morysmula* Suárez, 1963). Type species *Myrmosa* (*Paramyrmosa*)

radoszkowskyi Saussure, 1880. Number of species: World – 8, Palaearctic – 8, Russia – 2.

Paramyrmosa brunnipes (Lepeletier, 1845) [Myrmosa] (*Myrmosa cognata* Costa, 1858). Parasitoid of *Lasioglossum majus* Nylander, *L. pauxillum* Schenck, *L. morio* (F.) (Halictidae), *Crabro peltarius* (Schreber), *Cerceris rybyensis* (L.) (Crabronidae). Russia: **EP** (C, S, NC, CR). – Europe (WE, SE, EE), Armenia, Turkey.

Paramyrmosa pulla (Nylander, 1847) [Myrmosa] (*Myrmosa czerskii* Skorikov, 1929). Russia: **ES** (TU, BR, ZB), **FE** (AM, KH, PR). – Mongolia, China (NE).

TAIMYRMOSA Lelej, 2005. Type species *Myrmosa eos* Lelej, 1981. Number of species: World – 4, Palaearctic – 3, Russia – 2.

Taimyrmosa eos (Lelej, 1981) [Myrmosa]. Russia: **FE** (PR). – Korean Peninsula.

Taimyrmosa mongolica (Suárez, 1974) [Myrmosa] (*M. tsunekii* Haneda, 1979). Russia: **ES** (IR, BR, YA, ZB), **FE** (AM, KH, PR, SA). – Mongolia, China (NE), Korean Peninsula, Japan (Hon, Shi).

Subfamily SPHAEROPHTHALMINAE

Number of taxa: World – 63 genera, 1417 species, Palaearctic – 2/4, Russia – 1/1.

Tribe SPHAEROPHTHALMINI

CYSTOMUTILLA André, 1896. Type species *Mutilla ruficeps* Smith, 1855. Number of species: World – 2, Palaearctic – 2, Russia – 1.

Cystomutilla ruficeps (Smith, 1855) [Mutilla] (*M. erythrocephala* Lucas, 1849, nom. praeocc., nec Latreille, 1792; *M. parvula* Kriechbaumer, 1896). Parasitoid of *Ectemnius rubicola* (Dufour et Perris) and *Pemphredon rugifer* (Dahlbom) (Crabronidae). Russia: **EP** (S, CR). – Europe (WE, SE, EE), Algeria, Georgia.

Subfamily DASYLABRINAE

Tribe DASYLABRINI

Number of taxa: World – 8 genera, 324 species, Palaearctic – 4/10, Russia – 1/13.

DASYLABRIS Radoszkowski, 1885 (*Allomutilla* Ashmead, 1899). Type species *Mutilla arenaria* Fabricius, 1787. The genus includes four

subgenera: *Baltilla* Lelej, 1976, *Craspedopyga* Lelej, 1976, *Dasylabris* Radoszkowski, 1885, and *Inbaltilla* Lelej, 1976. Number of species: World – 151, Palaearctic – 63, Russia – 13.

Dasylabris (Baltilla) adversa Skorikov, 1935 (*D. italica* var. *adversa* Skorikov, 1935; *D. kaszabi* Suárez, 1974). Collected in the nest aggregations of *Colletes* spp. and *Hylaeus* spp. (Colletidae). Russia: **EP** (C, E, S, NC), **ES** (TU, BR, ZB). – Europe (EE), Mongolia, China (NO).

Dasylabris (Baltilla) bicolor (Pallas, 1771) [Mutilla] (*M. albeola* Pallas, 1773; *M. caspica* Kolenati, 1846; *M. funeraria* Erichson, 1849; *D. bergi* Skorikov, 1935). Russia: **EP** (S, NC). – Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NW).

Dasylabris (Baltilla) gobicola Skorikov, 1935 (*D. kaszabi funerula* Suárez, 1974). Collected in the nest aggregations of *Colletes* spp. and *Hylaeus* spp. (Colletidae). Russia: **WS** (AL), **ES** (TU). – Mongolia, China (NW, NO).

Dasylabris (Baltilla) kozlovi Skorikov, 1935. Collected in the nest aggregations of *Anthophora* spp. (Apidae). Russia: **ES** (TU). – Mongolia, China (NW, NO, WP).

Dasylabris (Craspedopyga) manderstiernii manderstiernii (Radoszkowski, 1865) [Mutilla] (*D. mauropsis* Skorikov, 1935; *D. mauropsis* var. *canonota* Skorikov, 1935). Russia: **EP** (NC, CR). – Caucasus, Turkey, Syria, Lebanon, Iran, Turkmenistan.

Dasylabris (Dasylabris) intermedia Skorikov, 1935 (*D. rubrosignata ypsilon* Chen, 1957). Russia: **ES** (BR, ZB). – Afghanistan, Mongolia, China (NO, CE, SE), Korean Peninsula.

Dasylabris (Dasylabris) maura maura (Linnaeus, 1758) [Mutilla] (*M. austriaca* Panzer, 1799; *M. rubricans* Lepeletier, 1845; *M. argenteofasciata* Costa, 1859; *M. trisinuosa* Costa, 1886; *D. maura* f. *nigrociliata* Hoffer, 1938). Parasitoid of *Katamenes arbustorum* (Panzer) (Vespidae), *Sphex occitanicus* (Lepeletier et Serville), *Ammophila heydeni* Dahlbom (Sphecidae). Russia: **EP** (CR). – Europe (WE, SE, EE), Caucasus.

Dasylabris (Dasylabris) maura sungora (Pallas, 1773) [Mutilla] (*M. quadrimaculata* Radoszkowski, 1865, nom. praeocc., nec Lucas, 1849; *D. maura nostras* Skorikov, 1935; *D. maura nostras* var. *argentata* Skorikov, 1935; *D. maura nostras* var. *argyraxa* Skorikov, 1935; *D. maura nostras* var. *consuenta* Skorikov, 1935; *D. maura nostras* var. *iberica*

Skorikov, 1935; *D. maura nostras* var. *infuscata* Skorikov, 1935; *D. maura nostras* var. *quinquemaculata* Skorikov, 1935; *D. maura nostras* var. *tricincta* Skorikov, 1935). Russia: EP (C, E, S, NC), UR, WS (AL). – Kyrgyzstan, Kazakhstan, China (NW).

Dasylabris (Dasylabris) maura armeniaca (Kolenati, 1846) [Mutilla] (*D. mlokosewitzii* Radoszkowski, 1885; *D. dupicolor* Skorikov, 1935). Russia: EP (NC). – Armenia, Azerbaijan, Turkey, ?Palestine, Iran, Turkmenistan.

Dasylabris (Dasylabris) mongolica (Radoszkowski, 1861) [Mutilla] (*D. exarenaria* Skorikov, 1935; *D. turana* Skorikov, 1935). Russia: ES (TU). – Iran, Afghanistan, Central Asia, Kazakhstan, Mongolia, China (NW).

Dasylabris (Dasylabris) siberica (Christ, 1791) [Mutilla] (*M. rubrosignata* Radoszkowski, 1865; *M. maura* var. *quadrimaculata* Sichel et Radoszkowski, 1870, nom. praeocc., nec Lucas, 1848; *M. maura* var. *quadripunctata* Sichel et Radoszkowski, 1870, nom. praeocc., nec Olivier, 1811; *D. siberica* var. *insignata* Skorikov, 1935; *D. siberica* var. *incompleta* Skorikov, 1935). Collected in the nest aggregations of *Colletes* spp. (Colletidae) and *Anthophora* spp. (Apidae). Russia: WS (AL), ES (KS, TU, KR, BR, ZB), FE (AM, PR). – Mongolia, China (NO, CE), Korean Peninsula.

Dasylabris (Inbaltilla) petiolata (Baer, 1848) [Mutilla] (*D. italica* var. *miogramma* Skorikov, 1935). Russia: EP (NC, CR). – Europe (EE).

Dasylabris (Inbaltilla) regalis (Fabricius, 1793) [Mutilla] (*M. petiolaris* Radoszkowski, 1865; *M. concolora* Radoszkowski, 1865; *M. unipetiolaris* Radoszkowski, 1866; *D. italica* var. *triangulomaculata* Skorikov, 1935; *D. italica* f. *slovaca* Hoffer, 1938; *D. italica* f. *batai* Hoffer, 1938; *M. lugubris*: Sichel, Radoszkowski, 1870) Russia: EP (E, S, NC, CR), UR, WS (AL). – Europe (WE, SE, EE), Kazakhstan, China (NW).

Subfamily MYRMILLINAE

Number of taxa: World – 23 genera, 238 species, Palaearctic – 10/65, Russia – 1/8.

MYRMILLA Wesmael, 1851 (*Rudia* Costa, 1856). Type species *Mutilla distincta* Lepeletier, 1845 (= *M. calva* Villers, 1789). The genus includes three subgenera: *Eurygnathilla* Skorikov, 1927, *Myrmilla* Wesmael, 1851, and *Pseudomutilla* Costa, 1885.

Parasitoids of *Halictus* spp. (Halictidae). Number of species: World – 55, Palaearctic – 48, Russia – 8.

Myrmilla (Myrmilla) caucasica (Kolenati, 1846) [Mutilla] (*Mutilla corniculata* Sichel et Radoszkowski, 1869, part.; *Myrmilla jasonia* Skorikov, 1927; *M. kiritschenkoi* Skorikov, 1927; *Mutilla erythrocephala*: Radoszkowski, 1865; *M. cornuta*: Sichel, Radoszkowski, 1869). Russia: EP (S, NC, CR). – Europe (SE, EE), Caucasus, Turkey, Iran.

Myrmilla (Myrmilla) etzchmiadzini (Radoszkowski, 1885) [Mutilla] (*Myrmilla calva armeniaca* Skorikov, 1927). Russia: EP (NC). – Armenia, Azerbaijan, Iran.

Myrmilla (Myrmilla) lezginica (Radoszkowski, 1885) [Mutilla] (*Mutilla subcornuta* Morawitz, 1893; *Myrmilla kokujevi* Skorikov, 1927; *M. susiana* Skorikov, 1927). Russia: EP (NC, CR). – Europe (SE, EE), Georgia, Azerbaijan, Turkey, Syria, Palestine, Iran, Turkmenistan, Uzbekistan.

Myrmilla (Myrmilla) mutica (André, 1903) [Mutilla] (*Mutilla erythrocephala* var. *mutica* André, 1903). Russia: EP (CR). – Europe (SE, EE), Egypt.

Myrmilla (Pseudomutilla) glabrata (Fabricius, 1775) [Mutilla] (*Mutilla ciliata* Fabricius, 1793; *M. fraterna* Baer, 1848; *M. cephalica* Sichel et Radoszkowski, 1869; *Myrmilla sarmatica* Skorikov, 1927; *M. sarmatica* var. *belgovskii* Skorikov, 1927). Russia: EP (C, E, S, NC, CR), UR. – Europe (WE, SE, EE), Turkey, Syria, Uzbekistan, Kazakhstan.

Myrmilla (Pseudomutilla) meda Skorikov, 1927 (*M. appropinqua* Skorikov, 1927). Russia: EP (NC). – Azerbaijan, Iran, Turkmenistan.

Myrmilla (Pseudomutilla) skorikovi Lelej, 1985 (*M. meda caucasica* Skorikov, 1927, nom. praeocc., nec Kolenati, 1846). Russia: EP (S, NC). – Georgia, Azerbaijan, Iran.

Myrmilla (Pseudomutilla) vutshetitschi Skorikov, 1927 (*M. capitata kusdasi* Invrea, 1965). Russia: EP (S, NC, CR). – Europe (WE, SE, EE), Turkey, Syria.

Subfamily MUTILLINAE

Number of taxa: World – 98 genera, 2045 species, Palaearctic – 30/254, Russia – 12/42.

Tribe CTENOTILLINI

Number of taxa: World – 10 genera, 181 species, Palaearctic – 1/3, Russia – 1/1.

CTENOTILLA Bischoff, 1920 Type species *Mutilla pectinifera* André, 1893 (=*M. caeca* Radoszkowski, 1879). Number of species: World – 3, Palaearctic – 1, Russia – 1.

Ctenotilla caeca (Radoszkowski, 1879) [Mutilla] (*Mutilla pectinifera* André, 1893; *C. pectinifera* var. *obscura* Nonveiller, 1959. Russia: EP (CR). – Europe (SE, EE), Turkey, Armenia, Azerbaijan, Syria.

Tribe SMICROMYRMINI

Number of taxa: World – 27 genera, 449 species, Palaearctic – 13/190, Russia – 5/29.

EPHUTOMMA Ashmead, 1899 (*Eremomyrme* Suárez, 1965). Type species *Mutilla incerta* Radoszkowski, 1877 (=*M. turcestanica* Dalla Torre, 1897). Number of species: World – 16, Palaearctic – 15, Russia – 2.

Ephutomma angustata (Skorikov, 1935) [Smicromyrme] (*S. sogdiana* Lelej, 1971). Russia: EP (S). – Iran, Central Asia, Kazakhstan.

Ephutomma coronata (Lelej, 1976) [Eremomyrme]. Russia: UR. – Turkmenistan, Kazakhstan.

Ephutomma mongolensis (Suárez, 1974) [Smicromyrme]. Russia: ES (ZB). – Mongolia.

NEMKA Lelej, 1985 (*Horia Tsuneki*, 1993). Type species *Mutilla viduata* Pallas, 1773. Number of species: World – 22, Palaearctic – 7, Russia – 1.

Nemka viduata viduata (Pallas, 1773) [Mutilla] (*Tiphia stridula* Rossi, 1790; *M. coronata* Rossi, 1792; *M. coronata* Fabricius, 1793, nom. praeocc., nec Rossi, 1792; *M. paedemontana* Fabricius, 1794; *M. togata* Ahrens, 1812; *M. simplica* Radoszkowski, 1865; *M. pauperata* Sichel et Radoszkowski, 1869). Parasitoid of *Larra anathema* (Rossi), *Bembix* spp., *Bembecinus tridens* (Fabricius), *Gorytes* spp. (Crabronidae), *Dasypoda hirtipes* (Fabricius) (Melittidae). Russia: EP (E, S, NC, CR); UR, WS (AL). – Europe (WE, SE, EE), Caucasus, Turkey, Syria, Kazakhstan, Mongolia, China (NW).

PHYSETOPODA Schuster, 1949 (*Paramyrme* Lelej, 1977). Type species *Physetopoda insularis* Schuster, 1949. Number of species: World – 48, Palaearctic – 44, Russia – 8.

Physetopoda cingulata (Costa, 1858) [Mutilla] (*Smicromyrme rudolfae* Lelej, 1984). Russia: EP (S). – Europe (SE, EE), Kazakhstan.

Physetopoda daghestanica (Radoszkowski, 1885)

[Mutilla] (*Smicromyrme montata* f. *moravica* Hoffer, 1936; *S. variabilis* Hoffer, 1938; *S. variabilis* f. *maculata* Hoffer, 1938; *S. variabilis* f. *lineata* Hoffer, 1938; *S. variabilis* f. *obenbergeri* Hoffer, 1938; *S. variabilis* f. *conversa* Hoffer, 1938). Russia: EP (S, CR), WS (AL). – Europe (WE, SE, EE), Turkey, Kazakhstan.

Physetopoda halensis (Fabricius, 1787) [Mutilla]

(*M. ephippium* Fabricius, 1793, nom. praeocc., nec Rossi, 1792; *M. montana* Panzer, 1805; *M. schencki* Schmiedeknecht, 1907; *Smicromyrme punctata pseudomontana* Hoffer, 1936; *S. montana* f. *susterai* Hoffer, 1936; *S. compacta* f. *melaclena* Invrea, 1958; *S. subalpina* Ressl, 1960). Parasitoid of chrysomelid beetles *Labidostomis hummeralis* (Schneider), *Clytra quadripunctata* Linnaeus (Chrysomelidae). Russia: EP (C, E, S, NC, CR), UR, WS (AL), ES (KS). – Europe (WE, SE, EE), Kazakhstan.

Physetopoda oratoria (Chen, 1957) [Smicromyrme]

(*S. oratoria vaticinia* Chen, 1957; *S. oratoria miscisis* Chen, 1957; *Glossotilla rectangulata* Chen, 1957; *S. eoa* Lelej, 1977). Russia: ES (KS, TU, IR, YA, ZB), FE (AM, KH, PR). – Mongolia, China (NE, NO).

Physetopoda portschinskii (Radoszkowski, 1888)

[Mutilla]. Russia: EP (S, NC). – Armenia, Azerbaijan, Turkmenistan, Kazakhstan.

Physetopoda scutellaris (Latreille, 1792) [Mutilla]

(*M. bimaculata* Jurine, 1807; *M. subcomata* Wesmael, 1851). Russia: EP (CR), ES (KS, TU). – Europe (WE, SE, EE), ?N Africa, Armenia, Turkey.

Physetopoda similis (Lelej, 1984) [Paramyrme]. Russia: EP (CR). – Europe (EE), Armenia, Azerbaijan, Turkey.

Physetopoda turgajensis (Lelej, 1984) [Paramyrme]. Russia: EP (S), UR. – Kazakhstan, Mongolia.

SKORIKOVIA Ovtchinnikov, 2002. Type species

Mutilla elongata Radoszkowski, 1885. Number of species: World – 5, Palaearctic – 5, Russia – 3.

Skorikovia elongata (Radoszkowski, 1885) [Mutilla]

(*M. elongata* var. *pura* André, 1901; *Smicromyrme ambigua* Skorikov, 1935; *S. baigacuma* Skorikov, 1935; *S. gracilis* Skorikov, 1935; *S. kuznetzovi* Skorikov, 1935). Russia: EP (S, NC). – Azerbaijan, Turkey, Iran, Afghanistan, Central Asia, Kazakhstan.

Skorikovia pliginskiji (Lelej, 1984) [Smicromyrme].

Russia: EP (E, S, CR), UR. – Europe (SE, EE).

Skorikovia radoszkovskii (Skorikov, 1935) [Smicromyrme].

Russia: EP (NC). – Georgia, Azerbaijan, Turkey.

SMICROMYRME Thomson, 1870. Type species *Mutilla rufipes* Fabricius, 1787. The genus includes five subgenera: *Astomyrme* Schwartz, 1984, *Eremotilla* Lelej, 1985, *Erimyrme* Lelej, 1985, *Rhombotilla* Nagy, 1966, and *Smicromyrme* Thomson, 1870. Number of species: World – 232, Palaearctic – 110, Russia – 15.

Smicromyrme (Astomyrme) ausonius Invrea, 1950 (*S. sabatia* Invrea, 1953; *S. ausonia* var. *etrusca* Invrea, 1950). Parasitoid of flies *Zeuxia cinerea* Meigen (Tachinidae), *Eumerus strigatus* (Fallén) (Syrphidae). Russia: **EP** (S, NC, CR), **UR**. – Europe (SE, EE), Cyprus, Turkmenistan, Kazakhstan.

Smicromyrme (Eremotilla) coracinus Lelej, 1985. Russia: **UR**, **WS** (AL). – Kyrgyzstan, Kazakhstan, Mongolia.

Smicromyrme (Eremotilla) nigriceps Nonveiller, 1959 (*S. orientalis* Nonveiller, 1959, nom. praeocc., nec Mickel, 1933; *S. orientalis* var. *maculata* Nonveiller, 1959, nom. praeocc., nec Hoffer, 1938; *S. orientalis* f. *afasciata* Nonveiller, 1972; *S. pseudoorientalis* Nonveiller, 1979). Parasitoid of *Bembecinus tridens* (F.) (Crabronidae). Russia: **EP** (S, NC, CR). – Europe (SE), Caucasus, Iran, Turkmenistan.

Smicromyrme (Eremotilla) novaki Invrea, 1954 (*S. nonveillieri* Invrea, 1954; *S. curtiventris* var. *dalmatina* Invrea, 1954). Russia: **EP** (S, CR). – Europe (SE, EE). Azerbaijan.

Smicromyrme (Eremotilla) schwarzi Suárez, 1975. Russia: **EP** (NC, CR). – Europe (SE, EE), Georgia, Armenia, Aserbaijan, Turkey.

Smicromyrme (Eremotilla) triangularis (Radoszkowski, 1865) [Mutilla] (*S. pouzdranensis* Hoffer, 1936). Russia: **EP** (E, NC, CR), **UR**. – Europe (SE, EE).

Smicromyrme (Erimyrme) bidenticulatus Chen, 1957 (*S. hasanensis* Lelej, 1977; *S. hasanensis* f. *monopunctata* Lelej, 1977; *S. hasanensis* f. *apunctata* Lelej, 1977). Russia: **FE** (PR). – Mongolia, China (NO, CE), Korean Peninsula.

Smicromyrme (Erimyrme) sicanus (De Stefani, 1887) [Mutilla] (*S. pusilla septentrionalis* Hoffer, 1936; *S. pusilla septentrionalis* f. *gregory* Hoffer, 1936; *S. rufipes* var. *lutescens* Invrea, 1954). Parasitoid of *Cerceris arenaria* (L.), *Nitela spinolae* Latr., *Mischophus bicolor* Jutine, *Tachysphex pompiliformis* (Panzer), *Diodontus minutus* (F.) (Crabronidae). Russia: **EP** (S, CR), **UR**, **WS** (AL). – Europe (WE, SE, EE), Turkey, Syria, Kazakhstan.

Smicromyrme (Erimyrme) stepposus Lelej, 1984. Russia: **EP** (S, CR). Kazakhstan.

Smicromyrme (Smicromyrme) lewisi Mickel, 1935 (*Mutilla japonica* Cameron, 1900, nom. praeocc., nec Dalla Torre, 1897; *M. cameroni* Yano, 1932, nom. praeocc., nec Schulz, 1906; *S. lewisi yanoi* Mickel, 1935; *S. lewisi nigricula* Mickel, 1935; *S. momidzizyama* Tsuneki, 1972). Collected in the nest aggregations of *Oxybelus bipunctatus* Olivier (Crabronidae), *Hylaeus* spp., *Colletes* spp. (Colletidae). Russia: **ES** (KS, KR, TU, IR, BR, ZB), **FE** (AM, KH, PR, SA, KU). – Mongolia, China (NO, CE, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Tsushima, Tanegashima).

Smicromyrme (Smicromyrme) ruficollis ruficolis (Fabricius, 1793) [Mutilla] (*S. kiritshenkoi* Skorikov, 1935). Russia: **EP** (S, CR), **WS** (AL). – Europe (WE, SE, EE), Turkey, Kazakhstan.

Smicromyrme (Smicromyrme) rufipes (Fabricius, 1787) [Mutilla] (*M. nigra* Rossi, 1792; *M. ephippium* Rossi, 1792; *M. sellata* Panzer, 1797; *M. nigrita* Panzer, 1801; *M. petiolaris* Fabricius, 1804; *M. gibba* Baer, 1848; *M. nemoralis* Baer, 1848; *M. rufipes* var. *fulvogastra* André, 1901; *Smicromyrme rufipes* var. *nigricollaris* Skorikov, 1935; *S. rufipes* f. *zavadili* Hoffer, 1938; *S. rufipes* f. *lidmilae* Hoffer, 1938; *S. kuehlhorni* Invrea, 1963; *S. globosa* Franz, 1982). Parasitoid of *Tachysphex* spp., *Oxybelus uniglumis* (L.), *O. bipunctatus* Olivier, *Mischophus spurius* (Dahlbom) (Crabronidae), *Halictus* spp. (Halictidae), *Andrena* spp. (Andrenidae). Russia: **EP** (N, NW, C, E, S, CR), **UR**, **WS** (AL), **ES** (YA). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan.

Smicromyrme (Smicromyrme) sertus (Radoszkowski, 1885) [Mutilla]. Russia: **UR**.

Smicromyrme (Smicromyrme) tristis Lelej, 1984. Russia: **EP** (CR), **UR**. – Europe (WE, EE), Kazakhstan.

Smicromyrme (Smicromyrme) viktorovi Lelej, 1984. Russia: **EP** (S). – Georgia, Azerbaijan.

Tribe MUTILLINI

Number of taxa: World – 23 genera, 592 species, Palaearctic – 6/28, Russia – 3/9.

MUTILLA Linnaeus, 1758. Type species *Mutilla europaea* Linnaeus, 1758. Parasitoids of the *Bombyx* spp. (Apidae). Number of species: World – 157, Palaearctic – 11, Russia – 5.

Mutilla europaea Linnaeus, 1758 (*Apis aptera* Uddman, 1753; *M. similis* Harris, 1782; *M. schaefferi* Schrank, 1802; *M. coerulans* Lepeletier, 1845; *M. cyanea* Lepeletier, 1845; *M. panzeri* Lepeletier, 1845; *M. obscura* Nylander, 1847; *M. kaschiriensis* Baer, 1848; *M. notomelas* André, 1902; *M. europaea* f. *rufula* Hoffer, 1938, nom. praeocc., nec André, 1901; *M. europaea* f. *nigrescens* Hoffer, 1938, nom. praeocc., nec André, 1901; *M. europaea* f. *ruficeps* Hoffer, 1938, nom. praeocc., nec Smith, 1855). Parasitoid of *Bombus* spp. (Apidae). Russia: **EP** (N, NW, C, E, NC, CR), **UR**, **WS** (AL, NS). – Europe (WE, NE, SE, EE), N Africa, Turkey, Kazakhstan.

Mutilla marginata Baer, 1848 (*M. trifasciata* Radoszkowski, 1865; *M. marginata* f. *insueta* Invrea, 1964). Parasitoid of *Bombus* spp. (Apidae). Russia: **EP** (N, NW, C, E, CR), **UR**. – Europe (WE, SE, EE), Georgia, Syria.

Mutilla mikado Cameron, 1900 (*M. maesta* Chen, 1957). Parasitoid of *Bombus ardens* Smith, *B. diversus* Smith (Apidae). Russia: **ES** (KS, IR, ZB), **FE** (AM, KH, PR, SA). – China (NO, NE, CE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Mutilla quinquemaculata Cyrillo, 1787 (*M. quinquepunctata* Olivier, 1811; *M. quadrinotata* Klug, 1829). Parasitoid of *Anthidium florenthinum* (F.), *Osmia tricornis* Latreille (Megachilidae). Russia: **EP** (CR). – Europe (WE, SE), N Africa, Cyprus, Turkey, Palestine, Syria.

Mutilla saltensis Radoszkowski, 1885. Russia: **EP** (NC). – Caucasus, Iraq, Iran.

RONISIA Costa, 1858 (*Pycnotilla* Bischoff, 1920). Type species *Ronisia torosa* Costa, 1858 (= *Mutilla brutia* Petagna, 1787). Number of species: World – 29, Palaearctic – 8, Russia – 1.

Ronisia brutia brutia (Petagna, 1787) [Mutilla] (*M. sexmaculata* Cyrillo, 1787; *M. hungarica* Fabricius, 1793; *M. confusa* Lepeletier, 1845; *M. monspeliensis* Lepeletier, 1845; *M. sexmaculata* Baer, 1848, nom. praeocc., nec Cyrillo, 1787; *M. robusta* Smith, 1855; *Ronisia torosa* Costa, 1858; *Mutilla sericeiventris* Costa, 1864; *M. taurica* Radoszkowski, 1865, nom. praeocc., nec Gistel, 1857; *M. discoidalis* Radoszkowski, 1865; *M. morawitzii* Dalla Torre, 1897; *M. crimeae* Strand, 1917; *R. barbara minoensis* Nonveiller, 1972; *M. ghilianii*: Sichel, Radoszkowski, 1870). Parasitoid of *Polistes* Latreille (Vespidae), *Creightonella sericans* (Fonscolombe), *Chalicodoma parietinum* (Fourcroy), *Megachile albisepta*

(Klug) (Megachilidae). Russia: **EP** (E, S, NC, CR). – Europe (WE, SE, EE), Turkey, Kazakhstan.

TROPIDOTILLA Bischoff, 1920. Type species *Mutilla litoralis* Petagna, 1787. Number of species: World – 13, Palaearctic – 7, Russia – 3.

Tropidotilla grisescens (Lepeletier, 1845) [Mutilla] (*M. vulnericeps* Costa, 1860; *M. stribigata* Sichel et Radoszkowski, 1870). Russia: **EP** (CR). – Europe (WE, SE, EE), Algeria, Turkey.

Tropidotilla litoralis (Petagna, 1787) [Mutilla] (*M. italica* Fabricius, 1793; *M. indigena* Illiger, 1807; *M. graeca* Lepeletier, 1845; *M. iberica* Kolenati, 1846; *M. salentina* Costa, 1858; *M. quinquefasciata*: Radoszkowski, 1865). Parasitoid of *Polistes* spp., especially *P. gallicus* (Linnaeus) (Vespidae). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Palestine, Syria, Iran, Turkmenistan.

Tropidotilla sareptana (André, 1901) [Mutilla]. Russia: **EP** (S, NC). – Armenia, Azerbaijan, Turkey.

Tribe TROGASPIDIINI

Number of taxa: World – 38 genera, 823 species, Palaearctic – 10/33, Russia – 3/3.

ARTIOTILLA Invrea, 1950 (*Glossomyrme* Suárez, 1979). Type species *Mutilla biguttata* Costa, 1859. Number of species: World – 3, Palaearctic – 3, Russia – 1.

Artiotilla biguttata (Costa, 1859) [Mutilla] (*Mutilla dalmatica* André, 1893; *M. mervensis* Radoszkowski, 1893; *Smicromyrme fertoni* Invrea, 1953; *A. biguttata* f. *palestinenensis* Invrea, 1965). Russia: **EP** (NC). – Europe (SE), Azerbaijan, Cyprus, Turkey, Palestine, Iran, Turkmenistan.

NEOTROGASPIDIA Lelej, 1996. Type species *Mutilla pustulata* Smith, 1873. Number of species: World – 6, Palaearctic – 2, Russia – 1.

Neotrogaspidia hammeri (Suárez, 1959) [Trogaspidia]. Russia: **EP** (NC). – Europe (SE), Armenia, Azerbaijan, Turkey, Cyprus, Palestine, Iran, Turkmenistan.

TROGASPIDIA Ashmead, 1899. Type species *Mutilla medon* Smith, 1855. Number of species: World – 262, Palaearctic – 19, Russia – 1.

Trogaspidia catanensis (Rossi, 1794) [Mutilla] (*M. ballioni* Radoszkowski, 1866; *M. paripunctata*

Sichel et Radoszkowski, 1869; *M. interrupta*: Radoszkowski, 1865). Russia: **EP** (S, NC, CR). – Europe (SE, EE), Algeria, Armenia, Turkey, Kazakhstan.

Species *incertae sedis*

Mutilla taurica Gistel, 1857. Russia: **EP** (CR).

Species wrongly described from Russia

Mutilla aureomaculata Sichel et Radoszkowski, 1869.
Russia: **EP** (CR) [actually Africa].

66. FAMILY BRADYNOBENAIDAE**A.S. LELEJ AND M.V. MOKROUsov**

A small family which is widespread (absent from the Australasian region) and includes four subfamilies: Typhoctinae Schuster, 1949 (=Eotillinae Schuster, 1949) (four genera, 10 species), which is distributed in Nearctic and Neotropical regions; Chyphotinae Bradley et Bequaert, 1928 (one genus, 55 species) – Nearctic region; Apterogyninae André, 1899 (four genera, 116 species) – Afrotropical, Palaearctic and Oriental regions; Bradynobaninae de Saussure, 1892 (one genus, seven species) – Neotropical region (Lelej, 2003). Sexual dimorphism is extreme: female apterous with pronotum freely articulated, male macropterous with reduced wing venation. All species are apparently solitary. Information on a single species indicates that the larvae are ectoparasitoids of Solifuga (Arachnida), pupation occurring within the shelter of the host.

Number of taxa: World – 10 genera, 188 species, Palaearctic – 4/43, Russia – 1/2.

R e f e r e n c e s . Radoszkowski, 1879; Morawitz, 1890; Brothers, Finnimore, 1993; Argaman, 1994a; Pagliano, 2002; Lelej, 2003; Lelej, Mokrousov, 2015.

APTEROGYNA Latreille, 1809 (*Icalantica* Argaman, 1994; *Utapitoca* Argaman, 1994; *Zarabayca* Argaman, 1994). Type species *Apterogyna olivieri* Latreille, 1809. Number of species: World – 31, Palaearctic – 15, Russia – 2.

Apterogyna lateritia Morawitz, 1890 (*A. judaica* Invrea, 1950; *A. taurica* Panfilov, 1954; *A. nonveilleri* Invrea, 1957; *A. stancici* Invrea, 1959; *A. wahrmani* Invrea, 1965; *A. palestinensis* Invrea, 1965). Russia: EP (CR). – Europe (SE), Turkey, Israel, Turkmenistan, Tajikistan.

Apterogyna mlokosevitzii Radoszkowski, 1879 (*A. nigra* Dover, 1924; *A. volgensis* Panfilov, 1954; *A. mlokosevitzi* (!): Morawitz, 1894; *A. dorsostriata*: Skorikov, 1935; *Icalantica mlokosewitzii* (!): Argaman, 1994a). Russia: EP (S, NC). – Europe (SE), N Africa, Caucasus, Turkey, Syria, Jordan, Lebanon, Israel, Iran, Turkmenistan, Uzbekistan, Tajikistan, ?Pakistan.

SUPERFAMILY POMPILOIDEA

67. FAMILY POMPILIDAE – SPIDER WASPS

V.M. LOKTIONOV AND A.S. LELEJ

The family Pompilidae (spider wasps) is one of the largest families among the aculeate wasps. These wasps are distributed worldwide, but mostly in the tropical regions (Pitts et al., 2006). Five subfamilies (Ctenocerinae, Notocyphinae, Ceropalinae, Pepsinae, and Pompilinae) are now recognized for Pompilidae (Waichert et al., 2015), three latters are recorded from Russia. Spider wasps are parasitoids that use spiders as host provisioning each cell with a single paralyzed spider on which they lay an egg (Iwata, 1976). There are several genera (subfam. Ceropalinae entirely, genera *Poecilagenia* Haupt and *Nipponodipogon* Ishikawa from subfam. Pepsinae, and genus *Evagetes* Lepeletier de Saint Fargeau from subfam. Pompilinae), which have evolved a cleptoparasitic existence, with the modes of cleptoparasitism differing among them (Wasbauer, 1995; Shimizu, 2000; O'Neill, 2001; Shimizu et al., 2012). Number of the taxa: World – 254 genera, 4855 species, Palaearctic – about 65/ about 1000, Russia – 37/246.

References. Pallas, 1771; Assmuss, 1859; Eversmann, 1846, 1849; Radoszkowski, 1887a,b, 1888; Morawitz, 1890, 1891, 1892, 1893; Westerlund, 1892; Haupt, 1927, 1959; Nordström, Forsius, 1928; Gussakovskij, 1930, 1932; Pate, 1946; Wolf, 1967, 1972; Day, 1981, 1988; Wahis, 1986, 2013; Tobias, 1978; Krombein, 1979; Dathe, 1980b; Shimizu, 1994; Lelej, 1995, 2000; Engel, Grimaldi, 2006; Pitts et al., 2006; Elliot, 2007; Wiśniowski, 2009; Lelej, Loktionov, 2012; Aguiar et al., 2013; Baghirov, 2014; Loktionov, Lelej, 2014, 2015; Waichert et al., 2015; Kochetkov, 2017; Loktionov, Lelej, 2017; Lelej et al., 2017.

Subfamily CEROPALINAE

CEROPOALES Latreille, 1797 (*Agenioxenus* Ashmead, 1902; *Ceratopales* Schulz, 1906; *Hypsiceraeus* Morice, Durrant, 1914; *Aceropales* Priesner, 1969; *Priesnerius* Móczár, 1978). Type species *Evania maculata* Fabricius, 1775). Cleptoparasites of spider wasps (Pompilidae) and digger wasps (Crabronidae). The genus includes three subgenera: *Ceropales* Latreille, 1797, *Bifidoceropales* Wolf, 1965 and *Hemiceropales* Wolf, 1965. Distributed worldwide.

Number of species: World – 135, Palaearctic – about 35, Russia – 11.

Ceropales (Bifidoceropales) pygmaea appendiculata Yasumatsu, 1939 (*Bifidoceropales lehri* Lelej, 1985). Russia: FE (KH, PR). – China (NE).

Ceropales (Bifidoceropales) pygmaea tsunekii (Lelej, 1994) [Bifidoceropales]. Russia: FE (PR). – Korean Peninsula.

Ceropales (Ceropales) albicincta albicincta (Rossi, 1790) [Evania]. Russia: EP (C, S, NC, CR), UR, ES (ZB). – Europe (WE, SE, EE), Georgia, Armenia, Azerbaijan, Turkey, Cyprus, Israel, Lebanon, Jordan, Iran, Kyrgyzstan, Kazakhstan, Mongolia, ?Japan.

Ceropales (Ceropales) bicoloripes bicoloripes Móczár, 1967. Russia: FE (AM). – Europe (SE), Turkey, Israel, Turkmenistan, Mongolia.

Ceropales (Ceropales) bicoloripes dorni Móczár, 1987. Russia: ES (ZB). – Mongolia.

Ceropales (Ceropales) bogdanovii bogdanovii Radoszkowski, 1877. Russia: EP (S, NC). – Uzbekistan.

Ceropales (Ceropales) bogdanovii sabulosa F. Morawitz, 1891. Russia: EP (S). – Uzbekistan, Kazakhstan.

Ceropales (Ceropales) erythropoda Gussakovskij, 1926. Russia: WS (AL), ES (KR, IR, BR, ZB). – Tajikistan, Kazakhstan, Mongolia, China (NE).

Ceropales (Ceropales) helvetica helvetica Tournier, 1889. Russia: EP (S, NC, CR), WS (AL). – Europe (WE, SE, EE), Morocco, Algeria, Tunisia, Georgia, Azerbaijan, Turkey, Cyprus, Israel, Jordan, Syria, Iraq, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia.

Ceropales (Ceropales) maculata maculata (Fabricius, 1775) [Evania]. Russia: EP (N, NW, C, S, NC, CR), UR, WS (OM, AL), ES (KR, IR, BR, YA, ZB), FE (AM, KH, PR, SA, KU, MG, CH). – Europe (WE, SE, NE, EE), Northern Africa, Turkey, Cyprus, Georgia, Armenia, Azerbaijan, Lebanon, Syria, Israel, Jordan, Iran, Iraq, Turkmenistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NC, NE, CC, SW), Korean Peninsula, Japan (Hon).

Ceropales (Ceropales) magnifica Gussakovskij, 1926. Russia: FE (KH, PR). – China (NE).

Ceropales (Ceropales) sibirica Radoszkowski, 1888. Russia: WS (AL), ES (KR, IR, BR, ZB), FE (KH). – Mongolia.

Ceropales (Ceropales) variegata (Fabricius, 1798) [Evania] (*C. implicatus* Yasumatsu, 1939). Russia:

EP (C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (IR, BR, ZB), **FE** (KH). – Europe (WE, NE, SE, EE), Morocco, Algeria, Tunisia, Turkey, Georgia, Israel, Kyrgyzstan, Kazakhstan, China (NE).

Ceropales (Hemiceropales) cibrata cibrata Costa, 1881 (*C. versicolor* Gussakovskij, 1926). Russia: **EP** (S, CR), **ES** (ZB). – Europe (WE, SE, EE), North Africa, Turkey, Cyprus, Georgia, Azerbaijan, Israel, Saudi Arabia, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Africa.

Subfamily PEPSINAE

Tribe PEPSINI

CRYPTOCHEILUS Panzer, 1806 (*Cryptochilus* Rafflesque, 1815; *Calicurgus* Brullé, 1833; *Homonotus* Dahlbom, 1845, nom. praeocc., nec Dahlbom, 1843; *Stenoclavelia* Arnold, 1932; *Chilochares* Banks, 1941; *Adirostes* Banks, 1946; *Chlorocheilus* Wolf, 1965, nom. nudum; *Trachycheilus* Wolf, 1965, nom. nudum; *Chyphonocheilus* Wolf, 1965, nom. nudum; *Ichneumocheilus* Wolf, 1965, nom. nudum; *Priocnemiocheilus* Wolf, 1970). Type species *Sphex annulata* Fabricius, 1798. Parasitoids of the spiders from the families Araneidae, Lycosidae, Pisauridae, Agelenidae, Amaurobiidae, Clubionidae, Gnaphosidae, Sparassidae and Thomisidae. The genus includes four subgenera: *Adonta* Billberg, 1820 (= *Salius* Fabricius, 1804, nom. praeocc., nec Schrank, 1798); *Chyphonocheilus* Wolf, 1970; *Cryptochelus* Panzer, 1806, and *Ichneumocheilus* Wolf, 1970. Widely distributed in Palaearctic, Nearctic and Neotropical (Panama, Costa Rica) regions. Number of species: Palaearctic – more than 60, Russia – 23.

Cryptochelus (Adonta) costae (Radoszkowski, 1888) [*Salius*]. Russia: **EP** (E, CR), **UR**. – Europe (SE, EE), Iran, Kazakhstan.

Cryptochelus (Adonta) egregius (Lepeletier de Saint Fargeau, 1845) [*Calicurgus*]. Russia: **EP** (NC). – Europe (WE, SE, EE), Algeria, Caucasus, Turkey, Syria, Turkmenistan.

Cryptochelus (Adonta) elegans (Spinola, 1806) [*Pompilus*] (*Pompilus tripunctatus* Spinola, 1807). Russia: **EP** (?Caucasus), **UR**. – Europe (WE, SE, EE), Turkey.

Cryptochelus (Adonta) fabricii (Vander Linden, 1827) [*Pompilus*]. Russia: **EP** (NW, C, S, NC, CR), **UR**, **WS** (AL), **ES** (IR). – Europe (WE, NE, SE,

EE), Turkey, Caucasus, Jordan, Iran, Kyrgyzstan, Kazakhstan.

Cryptochelus (Adonta) fischeri (Spinola, 1839) [*Pompilus*] (*C. confinis* Haupt, 1927; *C. fallax* Priesner, 1965). Russia: **EP** (S, CR). – Europe (WE, SE, EE), Morocco, Algeria, Egypt, Turkey, Cyprus, Syria, Israel, Iran, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan.

Cryptochelus (Adonta) flavus (Eversmann, 1849) [*Priocnemis*] (*Salius desertorum* F. Morawitz, 1891). Russia: **EP** (S), **UR**. – Central Asia, Kazakhstan, Mongolia.

Cryptochelus (Adonta) freygessneri (Kohl, 1883) [*Priocnemis*]. Russia: **EP** (S, NC, CR), **WS** (AL). – Europe (SE, EE), Turkey, Turkmenistan.

Cryptochelus (Adonta) guttulatus (Costa, 1887) [*Priocnemis*]. Russia: **EP** (C), **UR**. – Europe (SE, EE), Turkey, Iran, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan.

Cryptochelus (Adonta) gyrrifrons (F. Morawitz, 1889) [*Salius*]. Russia: **ES** (KR, IR, BR), **FE** (AM, KH, PR). – Korean Peninsula, China (NC, CC).

Cryptochelus (Adonta) manchurianus Yasumatsu, 1935. Russia: **ES** (IR, YA, ZB), **FE** (AM, PR). – Mongolia, China (NE), Korean Peninsula.

Cryptochelus (Adonta) maruyamai Ishikawa, 1962. Russia: **FE** (AM, PR). – Japan (Hon, Shi, Kyu).

Cryptochelus (Adonta) notatus (Rossi, 1792) [*Sphex*] (*Pompilus affinis* Vander Linden, 1827). Russia: **EP** (NC). – Europe (WE, NE, SE, EE), Morocco, Algeria, Turkey, Cyprus, Iran, Turkmenistan, Kyrgyzstan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia.

Cryptochelus (Adonta) richardsi Móczár, 1953. Russia: **EP** (S, NC), **UR**. – Europe (WE, SE, EE), Turkey, Israel, Syria, Iran, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan.

Cryptochelus (Adonta) strigosus (Eversmann, 1849) [*Pompilus*]. Russia: **UR**. – Eastern Kazakhstan.

Cryptochelus (Adonta) transcaucasicus Šustera, 1924. Russia: **EP** (NC). – Azerbaijan.

Cryptochelus (Adonta) variabilis (Rossi, 1790) [*Sphex*] (*Pompilus decemguttatus* Jurine, 1807). Russia: **EP** (S, NC, CR), **UR**. – Europe (WE, SE, EE), Algeria, Cyprus, Turkey, Iran, Uzbekistan, Kyrgyzstan, Kazakhstan.

Cryptochelus (Adonta) variipennis Šustera, 1924. Russia: **EP** (?S, NC). – Europe (WE, SE), Turkey, Jordan, Syria.

Cryptocheilus (Adonta) versicolor (Scopoli, 1763)

[*Sphex*] (*Priocnemis variegatus*: Eversmann, 1849). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), Cyprus, Turkey, Kyrgyzstan.

Cryptocheilus (Chyphonocheilus) rubellus (Eversmann, 1846) [*Pompilius* (!)]. Russia: **EP** (S, NC, CR), **UR**. – Europe (WE, SE, EE), Morocco, Algeria, Tunisia, Turkey, Cyprus, Iran, Syria, Pakistan, Afghanistan, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan.**Cryptocheilus (Cryptocheilus) annulatus** (Fabricius, 1798) [*Sphex*] (*Pompilus alternatus* Lepeletier de Saint Fargeau, 1845). Russia: **EP** (CR). – Europe (WE, SE, EE), Morocco, Algeria, Tunisia, Turkey, Cyprus, Lebanon, Israel, Saudi Arabia, Iran, Tajikistan, Kyrgyzstan.**Cryptocheilus (Cryptocheilus) discolor** (Fabricius, 1793) [*Sphex*]. Russia: **EP** (NC). – Europe (SE, EE), North Africa, Turkey, Cyprus, Syria, Iran, Afghanistan, Uzbekistan, Turkmenistan, Tajikistan, Kyrgyzstan, Kazakhstan.**Cryptocheilus (Cryptocheilus) octomaculatus** (Rossi, 1790) [*Sphex*]. Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), Morocco, Algeria, Tunisia, Turkey, Cyprus, Syria, Israel, Iran, Kyrgyzstan, Turkmenistan, Tajikistan.**Cryptocheilus (Ichneumocheilus) ichneumonoides** (Costa, 1874) [*Priocnemis*]. Russia: **EP** (Wahis, 2013). – Europe (WE, SE, EE), Turkey, Syria.

Tribe PRIOCNEMINI

CALIADURGUS Pate, 1946 (*Calicurgus* Lepeletier de Saint Fargeau, 1845, nom. praeocc., nec Brullé, 1833). Type species *Pompilus fasciatellus* Spinola, 1807. Parasitoids of the spiders from the families Tetragnathidae and Araneidae. Worldwide distributed genus. Number of species: Palaearctic – 2, Russia – 2.**Caliadurgus fasciatellus** (Spinola, 1807) [*Pompilus*] (*Priocnemis fasciatellus*: Eversmann, 1849; *Calicurgus hyalinatus*: Tobias, 1978). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (AL), **ES** (IR, BR, YA), **FE** (AM, KH, PR, SA, KA). – Europe (WE, NE, SE, EE), Turkey, Cyprus, Kyrgyzstan, Kazakhstan, Mongolia, N America.**Caliadurgus ussuriensis** (Gussakovskij, 1932) [*Calicurgus*] (*Calicurgus shirakii* Yasumatsu, 1943). Russia: **FE** (AM, KH, PR, SA, KU). – China (NC, CC, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Tsushima).**CTENOPRIOCNEMIS** Ishikawa, 1962. Type species *Ctenopriocnemis filicornis* Ishikawa, 1962. Distributed in Eastern Palaearctic (south of the Russian Far East, Korean Peninsula and Japan). Number of species: World – 1.**Ctenopriocnemis filicornis** Ishikawa, 1962. Russia: **FE** (PR, KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).**EOPOMPILUS** Gussakovskij, 1932 (*Sialus* Matsumura, 1911). Type species *Eopompilus orientalis* Gussakovskij, 1932 (= *Sialus* (!) *internalis* Matsumura, 1911). Parasitoids of the spiders from the family Theridiidae. Distributed in Eastern Palaearctic (south of the Russian Far East, China, Korean Peninsula and Japan) and Oriental region (China: Taiwan and Yunnan). Number of species: World – 5, Palaearctic – 4, Russia – 3.**Eopompilus internalis** (Matsumura, 1911) [*Sialus* (!)] (*Sialus* (!) *bizonatus* Matsumura, 1912; *E. orientalis* Gussakovskij, 1932). Russia: **FE** (AM, KH, PR, SA, KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Tsushima, Tanegashima, Mishima).**Eopompilus luteus** Lelej, 1986. Russia: **FE** (PR). – China (NC, NE).**Eopompilus minor** Gussakovskij, 1932. Russia: **FE** (AM, PR, SA, KU). – China (NC, NE, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Tsushima, Yakushima).**PRIOCNEMIS** Schiødte, 1837 (*Priocnemus* Burmeister, 1872; *Priocnemis* Kirby, 1884; *Hemipogonius* Saussure, 1892; *Myrmecosalius* Ashmead, 1903; *Priocnemus* Banks, 1917; *Maculipennis* del Junco y Reyes, 1947; *Ceratocnemis* Wolf, 1963). Type species *Sphex exaltata* Fabricius, 1775. Parasitoids of the spiders from the families Atypidae, Segestriidae, Linyphiidae, Tetragnathidae, Araneidae, Lycosidae, Pisauridae, Oxyopidae, Zoridae, Agelenidae, Dictynidae, Amaurobiidae, Anaphaenidae, Liocranidae, Clubionidae, Gnaphosidae, Thomisidae and Salticidae. The genus includes four subgenera: *Leptopriocnemis* Ishikawa, 1962; *Moczarellus* Wolf, 1963; *Priocnemis* Schiødte, 1837; *Umbripennis* del Junco y Reyes, 1947 (= *Priocnemissus* Haupt, 1949). Worldwide distributed genus. Number of species: Palaearctic – about 100, Russia – 39.**Priocnemis (Leptopriocnemis) cyphonota** Pérez, 1905. Russia: **FE** (PR, KU). – Japan (Hok, Hon, Shi, Kyu).

- Priocnemis (Moczarellus) occipitalis** Gussakovskij, 1930 (*P. crassicapitis* Móczár, 1944). Russia: **EP** (S, NC). – Europe (EE).
- Priocnemis (Priocnemis) agenia** (Gussakovskij, 1926) [Salius]. Russia: **UR**.
- Priocnemis (Priocnemis) agilis** (Shuckard, 1837) [*Pompilus*] (*P. obtusiventris* Schiödte, 1837). Russia: **EP** (N, NW, C, S, NC, CR), **WS** (AL), **ES** (IR). – Europe (WE, NE, SE, EE), Turkey, Cyprus, Uzbekistan, Kyrgyzstan, Kazakhstan.
- Priocnemis (Priocnemis) amurensis** Lelej, 1988. Russia: **ES** (IR, BR), **FE** (AM, PR).
- Priocnemis (Priocnemis) belokobylskii** Lelej, 1988. Russia: **FE** (AM, KH, PR).
- Priocnemis (Priocnemis) cordivalvata** Haupt, 1927. Russia: **EP** (NW, C, NC), **ES** (KR, IR). – Europe (WE, NE, SE, EE), Mongolia.
- Priocnemis (Priocnemis) exaltata** (Fabricius, 1775) [*Sphex*] (*Priocnemis nudipes* Dahlbom, 1845; *P. gibbus*: Gussakovskij, 1930). Russia: **EP** (N, NW, C, E, CR), **UR**, **WS** (AL), **ES** (IR, ZB). – Europe (WE, NE, SE, EE), Turkey, Iran, Kyrgyzstan, Kazakhstan.
- Priocnemis (Priocnemis) fastigiata** Haupt, 1935. Russia: **EP** (S). – Europe (WE, SE, EE), Turkey.
- Priocnemis (Priocnemis) fenestrata** (Gussakovskij, 1926) [Salius]. Russia: **WS** (AL), **ES** (IR, YA), **FE** (AM, KH, PR). – Korean Peninsula.
- Priocnemis (Priocnemis) fennica** Haupt, 1927. Russia: **EP** (N, NW), **ES** (KS). – Europe (WE, NE, SE, EE), Turkey.
- Priocnemis (Priocnemis) ghilarovi** Lelej, 1988. Russia: **FE** (AM, PR).
- Priocnemis (Priocnemis) gracilis** (Gussakovskij, 1926) [Salius] (*P. festivus* Gussakovskij, 1930). Russia: **EP** (N, S).
- Priocnemis (Priocnemis) gussakovskiji** Wolf, 2004 (*P. confusor* Wahis, 2006). Russia: **EP** (C, ?S). – Europe (WE, NE, SE, EE), Turkey, Jordan, Kazakhstan.
- Priocnemis (Priocnemis) hyalinata** (Fabricius, 1793) [*Sphex*] (*Pompilus femoralis* Dahlbom, 1829). Russia: **EP** (NW, C, E, NC, CR), **UR**, **WS** (AL), **ES** (KR). – Europe (WE, NE, SE, EE), Turkey, Uzbekistan.
- Priocnemis (Priocnemis) kunashirensis** Lelej, 1988. Russia: **FE** (KU). – Japan (Hok, Hon).
- Priocnemis (Priocnemis) minuta** (Vander Linden, 1827) [*Pompilus*]. Russia: **EP** (C, E, NC, CR), **ES** (BR), **FE** (AM, PR). – Europe (WE, NE, SE, EE), Turkey, Cyprus, Turkmenistan, Kyrgyzstan, Kazakhstan.
- Priocnemis (Priocnemis) mitakensis** Ishikawa, 1954. Russia: **FE** (AM, PR). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Priocnemis (Priocnemis) nigricans** Gussakovskij, 1930 (*P. diversus*: Wolf, 1963; *P. parvulus parvulus*: Wolf, 1963). Russia: **EP** (NC, CR).
- Priocnemis (Priocnemis) parvula** Dahlbom, 1845 (*P. mocsaryi*: Gussakovskij, 1930; *P. minor*: Gussakovskij, 1930; Tobias, 1978). Russia: **EP** (N, NW, C, E, S, NC, CR), **WS** (OM, TK), **ES** (IR), **FE** (AM, KH, PR, SA, MG). – Europe (WE, NE, SE, EE), Turkey, Tunisia, Jordan, Uzbekistan, Kyrgyzstan.
- Priocnemis (Priocnemis) pseudopogonia** Gussakovskij, 1930. Russia: **ES** (BR), **FE** (KH, PR). – Korean Peninsula.
- Priocnemis (Priocnemis) punctatissimus** (F. Mörner, 1891) [Salius]. Russia: **EP** (S).
- Priocnemis (Priocnemis) pusilla** (Schiödte, 1837) [*Pompilus*]. Russia: **EP** (N, NW, C, NC, CR), **WS** (OM), **ES** (KR). – Europe (WE, NE, SE, EE), Georgia, Turkey, Iran, Kyrgyzstan, Turkmenistan, Mongolia.
- Priocnemis (Priocnemis) schiödtei** Haupt, 1927. Russia: **EP** (N, NW, C, S, NC, CR). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan, Mongolia.
- Priocnemis (Priocnemis) shidai** Ishikawa, 1962. Russia: **FE** (KU). – Japan (Hon).
- Priocnemis (Priocnemis) sugonjaevi** Lelej et Laktionov, 2015. Russia: **FE** (AM).
- Priocnemis (Priocnemis) unicolor** (Gussakovskij, 1926) [Salius]. Russia: **ES** (KS, KR, IR), **FE** (AM, PR, KU). – Japan (Hok, Hon).
- Priocnemis (Priocnemis) yasumatsui** Ishikawa, 1954 (*P. yezoensis* Ishikawa, 1959). Russia: **FE** (KH, PR, SA). – Japan (Hok, Hon).
- Priocnemis (Umbripennis) coriacea** Dahlbom, 1843. Russia: **EP** (C, S, NC, CR), **UR**. – Europe (WE, NE, SE, EE), Turkey.
- Priocnemis (Umbripennis) enslini** Haupt, 1927. Russia: **EP** (CR). – Europe (WE, NE, SE, EE).
- Priocnemis (Umbripennis) fallax** Verhoeff, 1892 (*P. vulgaris*: Gussakovskij, 1930; Tobias, 1978). Russia: **EP** (C, E, S). – Europe (WE, SE, EE), Tunisia, Turkey, Kyrgyzstan, Uzbekistan, Kazakhstan.
- Priocnemis (Umbripennis) gussakovskii** Lelej, 1988. Russia: **FE** (AM, KH, PR). – Korean Peninsula.
- Priocnemis (Umbripennis) hankoi** Móczár, 1944 (*P. mimulus*: Gussakovskij, 1930). Russia: **EP** (?S, ?South-East). – Europe (WE, NE, SE, EE), Turkey, Uzbekistan, Turkmenistan, Kyrgyzstan, Tajikistan, Kazakhstan.

- Priocnemis (Umbripennis) japonica** Gussakovskij, 1930 (*P. fuscus orientalis* Gussakovskij, 1930). Russia: **FE** (AM, KH, PR, SA, KU, MG). – Korean Peninsula, Japan (Hok, Hon).
- Priocnemis (Umbripennis) perturbator** (Harris, 1780) [Sphex] (*P. fuscus*: Eversmann, 1849; Assmuss, 1859; Gussakovskij, 1930). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (IR). – Europe (WE, NE, SE, EE), Cyprus, Turkey, Georgia, Iran, Uzbekistan, Kyrgyzstan.
- Priocnemis (Umbripennis) pillichi** Priesner, 1960. Russia: **EP** (NC). – Europe (WE, EE), Turkey.
- Priocnemis (Umbripennis) pseudojaponica** Lelej, 1988. Russia: **FE** (PR).
- Priocnemis (Umbripennis) susterai** Haupt, 1927 (*P. clémenti* Haupt, 1927). Russia: **EP** (S, NC, CR). – Europe (WE, NE, SE, EE), Azerbaijan, Kazakhstan, Kyrgyzstan.
- Priocnemis (Umbripennis) ussuriensis** Lelej, 1988. Russia: **FE** (AM, PR, SA). – China (CC), Korean Peninsula.
- Priocnemis (Umbripennis) vulgaris** (Dufour, 1841) [Pompilus] (*P. mimulus*: Gussakovskij, 1930; *P. mimula*: Tobias, 1978). Russia: **EP** (NW, C, S, NC). – Europe (WE, SE, EE), Turkey, Iran, Uzbekistan, Kyrgyzstan, Tajikistan, Kazakhstan.

Tribe DEUTERAGENIINI

DEUTERAGENIA Šusterá, 1912 (*Agenia* Schiødte, 1837, nom. praeocc., nec Descourtilz, 1825; *Pogonius* Dahlbom, 1845, nom. praeocc., nec Leach, 1815; *Anoplius* Lepeletier de Saint Fargeau, 1845, nom. praeocc., nec Dufour, 1834; *Adipogon* Banks, 1944; *Mesagenia* Haupt, 1959). Type species *Sphex variegata* Linnaeus, 1758. Parasitoids of the spiders from the families Segestriidae, Dysderidae, Oxyopidae, Agelenidae, Amaurobiidae, Clubionidae, Gnaphosidae, Thomisidae and Salticidae. Worldwide (except Australia) distributed genus. Number of species: World – 54, Palaearctic – about 25, Russia – 10.

Deuteragenia albiclypeata (Lelej, 1986) [Dipogon]. Russia: **FE** (PR, SA). – Korean Peninsula.

Deuteragenia bifasciata (Geoffroy, 1785) [Ichneumon] (*Pompilus hircanus* Fabricius, 1798; *Agenia intermedia* Dahlbom, 1843). Russia: **EP** (N, NW, C, E, NC, CR), **UR**, **WS** (AL), **ES** (IR, BR, ZB), **FE** (AM, KH, PR, SA, MG). – Europe (WE, SE, NE,

- EE), Georgia, Kyrgyzstan, Kazakhstan, Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Deuteragenia bokhaica** (Lelej, 1986) [Dipogon] (*Dipogon antropovi* Lelej, 1995). Russia: **FE** (PR).
- Deuteragenia immarginata** Gussakovskij, 1932. Russia: **FE** (PR, SA, KU). – Japan (Hok, Hon, Shi, Kyu).
- Deuteragenia lehri** Loktionov et Lelej, 2014. Russia: **FE** (PR).
- Deuteragenia nipponica** Yasumatsu, 1939. Russia: **FE** (AM, KH, PR). – Japan (Hok, Hon, Shi).
- Deuteragenia romankovae** (Lelej, 1986) [Dipogon]. Russia: **FE** (AM, PR). – Japan (Hok, Hon).
- Deuteragenia subintermedia** (Magretti, 1886) [Pogonius] (*D. nitida* Haupt, 1927). Russia: **EP** (C, E, NC). – Europe (WE, NE, SE, EE).
- Deuteragenia variegata** (Linnaeus, 1758) [Sphex] (*Agenia variegata*: Assmuss, 1859). Russia: **EP** (C, NC). – Europe (WE, NE, SE, EE), North Africa, Caucasus, Turkey, Cyprus, Iran, Turkmenistan.
- Deuteragenia vechti** (Day, 1979) [Dipogon] (*Dipogon bifasciatum*: Tobias, 1978). Russia: **EP** (N, NW, C, E, S), **UR**, **WS** (TK, KM, AL), **ES** (IR, BR, ZB), **FE** (AM, KH, PR, SA, KU, MG, CH). – Europe (WE, NE, EE), Kyrgyzstan, Kazakhstan, Japan (Hok, Hon).
- KURILOAGENIA** Loktionov et Lelej, 2014. Type species *Kuriloagenia ermolenkoi* Loktionov et Lelej, 2014. Distributed in Eastern Palaearctic (Russia: Kuril Islands). Number of species: World – 1.
- Kuriloagenia ermolenkoi** Loktionov et Lelej, 2014. Russia: **FE** (KU).
- NIPPONODIPOGON** Ishikawa, 1965. Type species *Dipogon iwatai* Ishikawa, 1965. The cleptoparasites of the spider wasps (Pompilidae) from the genera *Auplopus* Spinola and *Deuteragenia* Šusterá. Distributed in Eastern Palaearctic (south of the Russia Far East, Japan) and Oriental region (China: Guangdong, Yunnan, Hainan and Taiwan). Number of species: World – 9, Palaearctic – 7, Oriental – 2, Russia – 2.
- Nipponodipogon kurilensis** (Lelej, 1986) [Dipogon]. Russia: **FE** (KU).
- Nipponodipogon rossicus** (Lelej, 1986) [Dipogon]. Russia: **FE** (AM, PR).
- STIGMATODIPOGON** Ishikawa, 1965. Type species *Dipogon macrostigmatus* Ishikawa, 1959.

Distributed in Eastern Palaearctic (south of the Russian Far East and Japan). Number of species: World – 4, Palaearctic – 4, Russia – 2.

Stigmatodipogon budrisi Loktionov et Lelej, 2014.
Russia: **FE** (KU).

Stigmatodipogon petiolatus (Lelej, 1986) [Dipogon].
Russia: **ES** (IR), **FE** (AM, PR).

Tribe AGENIELLINI

Subtribe AGENIELLINA

POECILAGENIA Haupt, 1927 (*Meragenia* Banks, 1934; *Poecilageniella* Ishikawa, 1965; *Taiwagenia* Tsuneki, 1989; *Trachyglyptus*: Arnold, 1959). Type species *Calicurgus rubricans* Lepeletier de Saint Fargeau, 1845. The cleptoparasites of the spider wasps (Pompilidae) from the genus *Auplopus* Spinola. Distributed in Palaearctic, Afrotropical and Oriental regions. Number of species: World – 22, Palaearctic – 6, Russia – 4.

Poecilagenia hirashimai (Ishikawa, 1965) [*Poecilageniella*]. Russia: **FE** (AM, PR). – Japan (Hok, Hon).

Poecilagenia rubricans (Lepeletier de Saint Fargeau, 1845) [*Calicurgus*]. Russia: **EP** (NC), **FE** (AM, PR). – Europe (WE, SE, EE), Turkey.

Poecilagenia sculpturata (Kohl, 1898) [*Pseudagenia*]. Russia: **EP** (NC), **FE** (KH, PR). – Europe (WE, SE), Turkey, China (NE, CC), Japan (Hon, Shi, Kyu).

Poecilagenia shimidzui Lelej, 2000. Russia: **FE** (PR).

Subtribe AUPLOPODINA

AUPLOPUS Spinola, 1842 (*Aoplopus* Agassiz, 1846; *Pilpomus* Costa, 1859; *Pseudagenia* Kohl, 1884; *Stenagenia* Saussure, 1892; *Schizagenia* Cameron, 1908; *Tumagenia* Banks, 1934; *Calagenia* Banks, 1934; *Lophagenia* Banks, 1934). Type species *Pompilus femoratus* Fabricius, 1804. Parasitoids of the spiders from the families Segestriidae, Dysderidae, Tetragnathidae, Araneidae, Lycosidae, Pisauridae, Oxyopidae, Agelenidae, Anyphaenidae, Clubionidae, Sparassidae, Gnaphosidae, Thomisidae and Salticidae. The genus includes two subgenera: *Auplopus* Spinola, 1842 and *Conagenia* Banks, 1934 Worldwide distributed genus. Number of species: Palaearctic – about 30, Russia – 10.

Auplopus (Auplopus) albifrons (Dalman, 1823) [*Pompilus*] (*Agenia fallax* Eversmann, 1849). Russia: **EP** (S, NC), **UR**. – Europe (WE, NE, SE, EE), Turkey, Iran.

Auplopus (Auplopus) carbonarius carbonarius (Scopoli, 1763) [*Sphex*] (*Agenia carbonaria*: Assmuss, 1859). Russia: **EP** (N, NW, C, E, S, NC, CR), **WS** (AL), **ES** (IR, BR), **FE** (AM, KH, PR, MG). – Europe (WE, SE, NE, EE), Morocco, Algeria, Turkey, Cyprus, Jordan, Israel, Lebanon, Iran, Tajikistan, Kazakhstan, Mongolia, China (NC), Korean Peninsula, Japan (Hok, Non, Shi, Kyu, Tsushima).

Auplopus (Auplopus) mama Loktionov et Lelej, 2014.
Russia: **FE** (PR).

Auplopus (Auplopus) mandshuricus Lelej, 1990. Russia: **FE** (AM, PR). – Mongolia, China (NC).

Auplopus (Auplopus) pacificus Lelej, 1990. Russia: **FE** (AM, KH, PR). – Korean Peninsula.

Auplopus (Auplopus) pygialis (Pérez, 1905) [*Pseudagenia*]. Russia: **ES** (IR, BR), **FE** (AM, KH, PR, SA, KU). – Eastern Kazakhstan, Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Tsushima, Yakushima).

Auplopus (Auplopus) rectus (Haupt, 1927) [*Pseudagenia*]. Russia: **EP** (S, NC). – Europe (WE, SE, EE), Turkey, Cyprus, Lebanon, Jordan, Iran, Turkmenistan, Uzbekistan, Tajikistan, Kyrgyzstan, Kazakhstan, Mongolia.

Auplopus (Auplopus) takachihoi (Yasumatsu, 1943) [*Pseudagenia*]. Russia: **FE** (PR, SA). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Auplopus (Auplopus) yasumatsui Lelej, 1995. Russia: **FE** (PR).

Auplopus (Conagenia) constructor (Smith, 1873) [*Agenia*]. Russia: **FE** (AM, KH, PR). – China (CC, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Tsushima).

MACHAEROPTHRIX Haupt, 1938. Type species *Machaerothrix coactifrons* Haupt, 1938. Distributed in Eastern Palaearctic (south of the Russian Far East, Japan, China) and Oriental region (Sri Lanka; China: Guangdong; Philippines: Mindanao). Number of species: World – 6, Palaearctic – 3, Russia – 1.

Machaerothrix ussuriensis Lelej, 1986. Russia: **FE** (PR, KU).

Subfamily POMPILINAE

Tribe APORINI

Subtribe APORINA

APORUS Spinola, 1807 (*Actenopoda* Ashmead, 1902; *Melanaporus* Ashmead, 1902; *Odontaporus* Bradley, 1944). Type species *Aporus bicolor* Spinola, 1807. Parasitoids of the spiders from the family Atypidae. The genus includes two subgenera: *Aporus* Spinola, 1807 and *Cosmiaporus* Bradley, 1944. Worldwide (except Australia) distributed genus. Number of species: Palaearctic – 7, Russia – 3.

Aporus (Aporus) bicolor Spinola, 1807 (*Planiceps fulviventris* Costa, 1882). Russia: **EP** (NC). – Europe (WE, SE, EE), Cyprus, Turkey, Iran, Kazakhstan, Mongolia.

Aporus (Aporus) pollux (Kohl, 1888) [Pompilus]. Russia: **EP** (C). – Europe (WE, SE, EE), Tunisia, Turkey, Iran, Kyrgyzstan.

Aporus (Aporus) unicolor Spinola, 1807 (*A. femoralis* Vander Linden, 1827). Russia: **UR**, **ES** (?), **FE** (AM, PR, KU). – Europe (WE, SE, EE), Korean Peninsula, Japan (Hon).

Subtribe HOMONOTINA

HOMONOTUS Dahlbom, 1843 (*Isonotus* Dahlbom, 1842, nom. praeocc., nec Lepeletier de Saint Fargeau, Audinet-Serville, 1828; *Homotus* Smith, 1855; *Wesmaelinus* Costa, 1886; *Hemisalius* Sausure, 1892; *Wesmaelinus* de Dalla Torre, 1897; *Gilbertella* Turner, 1917, nom. praeocc., nec Waite, 1902; *Gilbertellana* Pate, 1946). Type species *Sphex sanguinolenta* Fabricius, 1793. Parasitoids of the spiders from the family Miturgidae. Widely distributed in the Palaearctic, Oriental and Afrotropical regions. Number of species: Palaearctic – 5, Russia – 2.

Homonotus iwatai Yasumatsu, 1932. Russia: **FE** (AM, PR, SA, KU). – Japan (Hon).

Homonotus sanguinolentus (Fabricius, 1793) [Sphex] (*Homonotus nasutus* F. Morawitz, 1888; *Salius sanguinolentus*: Assmuss, 1859). Russia: **EP** (NW, C, E, S, CR), **ES** (IR), **FE** (MG). – Europe (WE, NE, SE, EE), Cyprus, Turkey, Syria, Iran, Turkmenistan, Kyrgyzstan, Kazakhstan.

Subtribe FERREOLINA

FERREOLA Lepeletier de Saint Fargeau, 1845 (*Platyderes* Guérin-Méneville, 1844, nom. praeocc., nec Stephens, 1828; *Pseudoferreola* Radoszkowski, 1888; *Aporinus* Ashmead, 1902; *Prosalius* Aurivillius, 1907). Type species *Ferreola algira* Lepeletier de Saint Fargeau, 1845. Parasitoids of the spiders from the families Eresidae and Clubionidae. Distributed in the Palaearctic (North Africa, Europe, Russia, Central Asia), Afrotropical (Eritrea, Yemen) and Australian regions. Number of species: Palaearctic – about 10, Russia – 1.

Ferreola diffinis (Lepeletier de Saint Fargeau, 1845) [*Anoplius*] (*Homonotus rectangulatus* F. Morawitz, 1888). Russia: **EP** (N, NC), **ES** (IR). – Europe (WE, NE, SE, EE), Turkey, Afghanistan, Turkmenistan, Kyrgyzstan.

Tribe EPISYRONINI

BATOZONELLUS Arnold, 1937. Type species *Pompilus fuliginosus* Klug, 1834. Parasitoids of the spiders from the family Araneidae. Distributed in the Palaearctic (from European part of Russia in the west to Japan in the east), Afrotropical and Oriental regions. Number of species: Palaearctic – 2, Russia – 1.

Batozonellus truchmenus (F. Morawitz, 1888) [Pompilus]. Russia: **EP** (S). – Azerbaijan, Turkmenistan, Kyrgyzstan, Kazakhstan.

EPISYRON Schiødte, 1837 (*Epizuron* Ashmead, 1902; *Spilopompilus* Ashmead, 1902). Type species *Sphex rufipes* Linnaeus, 1758. Parasitoids of the spiders from the families Tetragnathidae, Lycosidae and Amaurobiidae. Worldwide distributed genus. Number of species: Palaearctic – about 40, Russia – 6.

Episyon albonotatum (Vander Linden, 1827) [Pompilus]. Russia: **EP** (N, NW, C, S, NC), **WS** (AL), **ES** (KR, IR, BR, ZB), **FE** (AM, KH, SA). – Europe (WE, NE, SE, EE), Morocco, Turkey, Syria, Iran, Turkmenistan, Uzbekistan, Kyrgyzstan, Tajikistan, Kazakhstan, Mongolia.

Episyon arrogans (Smith, 1873) [Pompilus] (*Pompilus funereipes* Costa, 1874; *P. sapporensis* Matsuura, 1912; *E. tripunctatus*: Gussakovskij, 1932). Russia: **EP** (C, NC), **ES** (IR), **FE** (AM, KH, PR,

KU). – Europe (WE, SE, EE), Turkey, China (NE, CC, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Tsushima, Izu, Ryukyu), Philippines.

Episyron candiotum Wahis, 1966. Russia: **EP** (S), **WS** (AL), **ES** (IR), **FE** (AM, PR). – Europe (SE, EE), Turkey, Syria, Transcaucasia, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE), Korean Peninsula.

Episyron gallicum (Tournier, 1889) [Pompilus]. Russia: **EP** (NC). – Europe (WE, NE, SE, EE), Turkey, Jordan, Syria, Iran, Uzbekistan, Kyrgyzstan, Kazakhstan.

Episyron kurilese Lelej, 1990. Russia: **FE** (KU). – Japan (Hok, Hon, Kyu, Tanegashima).

Episyron rufipes (Linnaeus, 1758) [Sphex] (*Pompilus karafutonis* Matsumura, 1911). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **ES** (IR, BR, ZB), **FE** (AM, KH, PR, SA, KA, MG). – Europe (WE, SE, NE, EE), Morocco, Turkey, Cyprus, Jordan, Syria, Saudi Arabia, Iran, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Yemen, Sudan, Ethiopia.

PARABATOZONUS Yasumatsu, 1936. Type species *Pompilus hakodadi* de Dalla Torre, 1897 (= *Pompilus jankowskii* Radoszkowski, 1887). Parasitoids of the spiders from the family Araneidae. Widely distributed in Palaearctic (from Europe and North Africa in the west to Japan in the east). Number of species: Palaearctic – 3, Russia – 2.

Parabatozonus jankowskii (Radoszkowski, 1887) [Pompilus] (*Pompilus bilunatus* Smith, 1873, nom. praeocc., nec Haliday, 1836; *P. hakodadi* de Dalla Torre, 1897; *P. fallaciosus* Matsumura, 1911). Russia: **FE** (PR). – China (NC), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Tsushima, Yakushima).

Parabatozonus lacerticida (Pallas, 1771) [Sphex]. Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (IR, BR, YA, ZB), **FE** (AM, KH, PR). – Europe (WE, SE, NE, EE), Morocco, Turkey, Cyprus, Georgia, Syria, Iran, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE), Korean Peninsula, Japan (Hok, Hon).

Tribe POMPILINI

AGENIOIDEUS Ashmead, 1902 (*Aporoideus* Ashmead, 1902; *Agenoideus* Banks, 1917; *Gymnochares* Banks, 1917; *Pseudevagete*s Wolf, 1970; *Rufopomilus* Wolf, 1970; *Alloridestus* Wolf, 1979; *Alloridestus* Wolf, 1985; *Sericopomilus*: Gussakovskij,

1932). Type species *Pompilus humilis* Cresson, 1867. Parasitoids of the spiders from the families Segestriidae, Dysderidae, Linyphiidae, Araneidae, Pisauridae, Agelenidae, Thomisidae and Salticidae. The genus includes two subgenera: *Agenioideus* Ashmead, 1902 and *Schizanoplus* Cameron, 1904 (= *Ridestus* Banks, 1912, = *Galactopterus* Arnold, 1937; = *Eggysomma* Haupt, 1962). Worldwide distributed genus. Number of species: Palaearctic – more than 100, Russia – 12.

Agenioideus (Agenioideus) amurensis (Gussakovskij, 1932) [Psammochares]. Russia: **FE** (AM, KH, PR).

Agenioideus (Agenioideus) cinctellus (Spinola, 1807) [Pompilus]. Russia: **EP** (N, NW, C, E, S, NC, CR). – Europe (WE, NE, SE, EE), Turkey, Cyprus, Kyrgyzstan, Mongolia, Japan (Hon).

Agenioideus (Agenioideus) haupti Loktionov et Lelej, 2014 (*Psammochares cinctellus* f. *rufa* Haupt, 1938, nom. praeocc., nec *Psammochares rufus* Haupt, 1927; *A. cinctellus*: Lelej, 1990, part.; Shimizu, Wahis, 2009, part.; Lelej, Loktionov, 2012, part.). Russia: **FE** (AM). – China (CC, SE), Korean Peninsula, Japan (Hon).

Agenioideus (Agenioideus) nubecula (Costa, 1874) [Pompilus]. Russia: **EP** (S, NC). – Europe (WE, NE, SE, EE), Tunisia, Turkey, Cyprus, Jordan, Iran, Kyrgyzstan, Tajikistan, Kazakhstan.

Agenioideus (Agenioideus) pacificus Lelej, 1994 (*A. cinctellus*: Lelej, 1990, part.; Shimizu, Wahis, 2009, part.). Russia: **FE** (AM, PR, SA).

Agenioideus (Agenioideus) sericeus (Vander Linden, 1827) [Pompilus]. Russia: **EP** (C, S, NC), **UR**. – Europe (WE, SE, EE), Turkey, Jordan, Iran, Uzbekistan, Tajikistan, Kyrgyzstan, Kazakhstan, Mongolia, China (?).

Agenioideus (Schizanoplus) ciliatus (Lepeletier de Saint Fargeau, 1845) [Pompilus] (*Psammochares rubigidorsum* Haupt, 1929). Russia: **EP** (S, CR). – Europe (WE, SE, EE), Turkey, Cyprus, Jordan, Syria, Uzbekistan, Turkmenistan, Kyrgyzstan, Kazakhstan.

Agenioideus (Schizanoplus) excisus (F. Morawitz, 1890) [Pompilus]. Russia: **EP** (S, NC). – Europe (SE, EE), Morocco, Algeria, Tunisia, Egypt, Turkey, Cyprus, Israel, Jordan, Iraq, Iran, Syria, Afghanistan, Turkmenistan, Kyrgyzstan, Kazakhstan.

Agenioideus (Schizanoplus) kohlii (Radoszkowski, 1888) [Pompilus] (*Pompilus ruficornis* F. Morawitz, 1890). Russia: **EP** (NC). – Iran, Turkmenistan, Kazakhstan.

Agenioideus (Schizanoplus) puniceus (Haupt, 1927) [Psammochares]. Russia: EP (S).

Agenioideus (Schizanoplus) ruficeps (Eversmann, 1849) [Pompilus]. Russia: EP (S). – Europe (WE, SE, EE), Morocco, Algeria, Azerbaijan, Turkey, Cyprus, Libya, Israel, Syria, Iran, Western Kazakhstan.

Agenioideus (Schizanoplus) udegeicus Lelej, 1990. Russia: FE (PR).

AMBLYELLUS Day, 1981. Type species *Pom-pilus hasdrubal* Kohl, 1894. Distributed in the Palaearctic (Western, Eastern and South Europe, North Africa, Russia, Near East, Central Asia) and Afrotropical (Yemen, South Africa) regions. Number of species: Palaearctic – 4, Russia – 2.

Amblyellus hasdrubal (Kohl, 1894) [Pompilus] (*Pom-piloides obtusus* Gussakovskij, 1935; *Aporinellus hamibal*: Tobias, 1978). Russia: EP (S, CR), UR. – Europe (WE, SE, EE), North Africa, Turkey, Armenia, Iran, Kyrgyzstan, Turkmenistan.

Amblyellus kirtschenkoi (Gussakovskij, 1930) [So-phropompilus]. Russia: EP (NC). – Turkmenistan.

ANOSPILUS Haupt, 1929. Type species *Pom-pilus orbitalis* Costa, 1863. Parasitoids of the spiders from the family Nemesiidae. Widely distributed in the Palaearctic region. Number of species: Palaearctic – about 20, Russia – 1.

Anospilus (Anospilus) carbonicolor (Gussakovskij, 1932) [Psammochares] (*Anoplius yamamotoi* Ishikawa, 1958; *Dicyrtomellus atratinus* Wolf et Móczár, 1972). Russia: FE (AM, PR). – Europe (SE), Mongolia, Korean Peninsula, Japan (Hok, Hon, Kyu, Tsushima).

APORINELLUS Banks, 1912 (*Ferreoloides* Haupt, 1929; *Eupom-piloides* Gussakovskij, 1935; *Cera-topompilus* Bradley, 1944). Type species *Aporus fasciatus* Smith, 1855. Parasitoids of the spiders from the families Oxyopidae, Philodromidae, Salticidae and Thomisidae. Worldwide (except Australia) distributed genus. Number of species: World – about 17, Palaearctic – about 5, Russia – 2.

Aporinellus (Aporinellus) moestus (Klug, 1834) [Pompilus] (*Pom-pilus sericeomaculata* Kohl, 1888). Russia: EP (NC, S, CR). – Europe (WE, SE, EE), Tunisia, Azerbaijan, Turkey, Cyprus, Georgia, Turkmenistan, Syria, Jordan, Uzbekistan, Kyrgyzstan, Tajikistan, Kazakhstan.

Aporinellus (Aporinellus) sexmaculatus (Spinola, 1805) [Pompilus] (*Pom-piloides sexmaculatus asiaticus* Gussakovskij, 1935; *Pom-pilus sexnotatus*: Radoszkowski, 1888). Russia: EP (E, S, NC, CR), WS (AL). – Europe (WE, SE, NE, EE), Egypt, Turkey, Armenia, Azerbaijan, Israel, Jordan, Lebanon, Syria, Iran, Iraq, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan.

ARACHNOSPILA Kincaid, 1900 (*Pycnopompilus* Ashmead, 1902). Type species *Arachnospila septentrionalis* Kincaid, 1900. Parasitoids of the spiders from the families Segestriidae, Lycosidae, Pisauridae, Agelenidae, Amaurobiidae, Anyphaenidae, Clubionidae, Gnaphosidae, Thomisidae, Salticidae and Philodromidae. The genus includes three subgenera: *Ammosphex* Wilcke, 1942 (=*Anopompilus* Dreisbach, 1949; =*Aridopompilus* Wolf, 1965; =*Boreopompilus* Wolf, 1965; =*Holarctopompilus* Wolf, 1965; =*Saxatilipompilus* Wolf, 1965); *Anoplochares* Banks, 1939; *Arachnospila* Kincaid, 1900. Worldwide (except Australia) distributed genus. Number of species: Palaearctic – about 90, Russia – 47.

Arachnospila (Ammosphex) abnormis (Dahlbom, 1842) [Pompilus]. Russia: EP (NW, C, S), WS (AL), ES (KS, TU, IR, BR), FE (AM, KH, PR, SA, KA, MG, CH). – Europe (WE, NE, SE, EE), Kyrgyzstan, Kazakhstan, Mongolia, China (NE).

Arachnospila (Ammosphex) alvarabnormis (Wolf, 1965) [Pompilus]. Russia: EP (?). – Europe (WE, EE), Cyprus.

Arachnospila (Ammosphex) anceps (Wesmael, 1851) [Pompilus] (*Sphex vagus* Harris, 1780, nom praeocc., non *Sphex vaga* Linnaeus, 1758; *Pom-pilus unguicularis* Thomson, 1870; *A. anceps serica* Wolf et Móczár, 1972). Russia: EP (N, NW, C, E, S, NC, CR), WS (AL), ES (KS, TU, KR, ZB), FE (AM, KA, MG, CH). – Europe (WE, NE, SE, EE), Turkey, Syria, Iran, Kyrgyzstan, Kazakhstan, Mongolia.

Arachnospila (Ammosphex) ausa (Tournier, 1890) [Pompilus]. Russia: EP (?). – Europe (WE, SE, EE), Tunisia, Turkey, Kyrgyzstan.

Arachnospila (Ammosphex) belokobylskii Loktionov et Lelej, 2011. Russia: FE (PR).

Arachnospila (Ammosphex) colpostoma (Kohl, 1886) [Pompilus]. Russia: EP (?). – Europe (WE, SE), Turkey, Kazakhstan.

Arachnospila (Ammosphex) consobrina (Dahlbom, 1843) [Pompilus]. Russia: EP (NW, C, E,

- NC). – Europe (WE, NE, SE, EE), Turkey, Israel, Syria, Iran, Turkmenistan, Uzbekistan, Tajikistan, Kyrgyzstan, Kazakhstan, Mongolia, China (?).
- Arachnospila (Ammosphex) dschingis** Wolf et Móczár, 1972. Russia: ES (KS, TU, IR, BR). – Kyrgyzstan, Mongolia, China (NC).
- Arachnospila (Ammosphex) eoabnormis** Lelej, 1995. Russia: ES (TU, KR), FE (AM, PR).
- Arachnospila (Ammosphex) gibbomima** (Haupt, 1929) [Psammochares]. Russia: EP (S). – Europe (EE), Turkey, Kyrgyzstan, Tajikistan, Kazakhstan, Mongolia.
- Arachnospila (Ammosphex) hedickei** (Haupt, 1929) [Psammochares] (*Pompilus pseudabnormis* Wolf, 1965; *Arachnospila pseudabnormis*: Dathe, 1980b). Russia: EP (N, NW, NC). – Europe (WE, NE, SE, EE), Turkey, Tajikistan, Kyrgyzstan.
- Arachnospila (Ammosphex) kasparyani** Loktionov et Lelej, 2015. Russia: ES (TU).
- Arachnospila (Ammosphex) kaszabi** Wolf, Móczár et 1972. Russia: ES (TU, KR, BR). – Mongolia.
- Arachnospila (Ammosphex) kurentzovi** Lelej, 1995. Russia: ES (BR), FE (AM, KH, PR, MG, CH).
- Arachnospila (Ammosphex) kurzenkoi** Lelej, 1995. Russia: WS (AL), ES (KS, TU, KR, IR, BR).
- Arachnospila (Ammosphex) kuwayamai** (Ishikawa, 1966) [Pompilus] (*Pompilus hirsutifrons* Ishikawa, 1966). Russia: FE (SA, KU). – Japan (Hon).
- Arachnospila (Ammosphex) maxim** Loktionov et Lelej, 2015. Russia: ES (KS, TU).
- Arachnospila (Ammosphex) moczari** Loktionov et Lelej, 2015. Russia: WS (AL), ES (TU).
- Arachnospila (Ammosphex) mongolica** Móczár, 1968. Russia: ES (KS, TU, KR, IR, BR, ZB), FE (AM, PR, SA). – Kyrgyzstan, Kazakhstan, Mongolia.
- Arachnospila (Ammosphex) mongolopinata** Wolf, 1981. Russia: WS (AL), ES (KS, TU), FE (AM, PR, SA). – Mongolia.
- Arachnospila (Ammosphex) nivalabnormis** (Wolf, 1965) [Pompilus]. Russia: EP (?). – Europe (WE, NE, SE, EE).
- Arachnospila (Ammosphex) opinata** (Tournier, 1890) [Pompilus] (*Pompilus carinatus* F. Morawitz, 1893). Russia: EP (N, NW). – Europe (WE, NE, SE, EE), Tunisia, Kyrgyzstan, Kazakhstan.
- Arachnospila (Ammosphex) orientausa** Loktionov et Lelej, 2011. Russia: ES (KR), FE (AM, PR).
- Arachnospila (Ammosphex) rasnitsyni** Loktionov et Lelej, 2011. Russia: ES (TU, BR).
- Arachnospila (Ammosphex) scythia** Loktionov et Lelej, 2015. Russia: ES (TU).
- Arachnospila (Ammosphex) sibirica** Loktionov et Lelej, 2015. Russia: WS (AL), ES (TU).
- Arachnospila (Ammosphex) subvittata** (F. Mörner, 1889) [Pompilus] (*Pompilus nagasei* Ishikawa, 1966). Russia: ES (KS, TU, KR, IR, ZB), FE (AM, KH, PR, KU). – China (NC), Korean Peninsula, Japan (Hon, Kyu).
- Arachnospila (Ammosphex) tobiasi** Loktionov et Lelej, 2011. Russia: ES (BR).
- Arachnospila (Ammosphex) trivialis** (Dahlbom, 1843) [Pompilus] (*Psammochares gibbus*: Gussakovskij, 1932). Russia: EP (N, NW, C, S, NC), WS (AL), FE (AM, KA, MG, CH). – Europe (WE, NE, SE, EE), Turkey, Near East, Kyrgyzstan, Tajikistan, Kazakhstan, Mongolia.
- Arachnospila (Ammosphex) wesmaeli** (Thomson, 1870) [Pompilus]. Russia: EP (N, NW, C, ?S), ES (KR). – Europe (WE, NE, SE, EE), Kyrgyzstan, Mongolia.
- Arachnospila (Ammosphex) westerlundi** (F. Mörner, 1893) [Pompilus] (*Pompilus guentheri* F. Mörner, 1893). Russia: EP (N, NW). – Europe (WE, NE, SE, EE), Cyprus.
- Arachnospila (Ammosphex) wolffi** Lelej, 1995. Russia: FE (PR).
- Arachnospila (Ammosphex) yasumatsui** Wolf et Móczár, 1972. Russia: ES (KS, TU, KR, IR, BR). – Mongolia, China (NC).
- Arachnospila (Ammosphex) zonsteini** Loktionov et Lelej, 2011. Russia: ES (BR).
- Arachnospila (Anoplochares) ainu** Lelej, 1995. Russia: FE (AM, PR, SA, KU). – Japan (Hok).
- Arachnospila (Anoplochares) fuscomarginata** (Thomson, 1870) [Pompilus] (*Pompilus signaticeps* F. Morawitz, 1893). Russia: EP (N, NW, C, S, NC, CR), UR. – Europe (WE, NE, SE, EE), Turkey, Iran, Uzbekistan, Tajikistan, Kyrgyzstan, Kazakhstan, Mongolia.
- Arachnospila (Anoplochares) minutula** (Dahlbom, 1842) [Pompilus] (*Pompilus minutus* Dahlbom, 1829, nom. praeocc., nec Vander Linden, 1827; *A. minutula* *mongolominutula* Wolf et Móczár, 1972). Russia: EP (N, NW, C, S, NC, CR), UR, WS (AL), ES (BR, ZB), FE (AM, KH, PR). – Europe (WE, NE, SE, EE), Turkey, Near East, Kyrgyzstan, Kazakhstan, Mongolia.
- Arachnospila (Anoplochares) mongolopissa** Wolf et Móczár, 1972. Russia: WS (AL), ES (IR, ZB), FE (AM, PR). – Mongolia, China (NC).

Arachnospila (Anoplochares) spissa (Schiødte, 1837)

[Pompilus]. Russia: **EP** (N, NW, C, S, NC), **WS** (AL), **ES** (IR, YA, ZB), **FE** (AM, SA, MG). – Europe (WE, NE, SE, EE), Turkey, Near East, Turkmenistan, Uzbekistan, Kyrgyzstan, Tajikistan, Kazakhstan, Mongolia, China (NC).

Arachnospila (Anoplochares) ussuriensis (Gussakovskij, 1932) [Psammochares] (*Pompilus hokkaidensis* Ishikawa, 1966). Russia: **FE** (AM, KH, PR, SA, KU). – Japan (Hok).**Arachnospila (Anoplochares) usurata** (Blüthgen, 1957) [Pompilus] (*A. asiatica*: Lelej, 1995; Wolf, Sorg, 2007; Wiśniowski, 2009; Wahis, 2013). Russia: **EP** (S), **ES** (IR), **FE** (AM, KH, PR); Europe (WE, SE, EE), Kazakhstan, Mongolia.**Arachnospila (Arachnospila) amurensis** (Motschulsky, 1860) [Pompilus]. Russia: **ES** (BR, ZB), **FE** (AM, KH, PR).**Arachnospila (Arachnospila) clericalis** (F. Morawitz, 1889) [Pompilus] (*A. mongolorufa* Wolf et Móczár, 1972). Russia: **WS** (AL), **ES** (TU, IR, BR, ZB). – Turkey, Kazakhstan, Mongolia, China (NC).**Arachnospila (Arachnospila) eisukei** (Ishikawa, 1969) [Pompilus] (*Psammochares herbignadus* (!): Kobayashi, 1931). Russia: **FE** (SA, KU). – Japan (Hok, Hon).**Arachnospila (Arachnospila) fumipennis** (Zetterstedt, 1838) [Pompilus] (*Psammochares malaisei* Nordström, Forsius, 1930). Russia: **EP** (N, C, NC, CR), **UR**, **WS** (AL), **ES** (KS, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KA, MG, CH). – Europe (WE, SE, NE, EE), North Africa, Turkey, Israel, Iran, Kyrgyzstan, Tajikistan, Kazakhstan, Mongolia, Japan (Hok, Hon), N America.**Arachnospila (Arachnospila) rufa** (Haupt, 1927) [Psammochares]. Russia: **EP** (N, NW, C, S), **UR**, **WS** (OM, AL), **ES** (TU). – Europe (WE, NE, SE, EE), Turkey, Near East, Kyrgyzstan, Kazakhstan, Mongolia.**Arachnospila (Arachnospila) sogdianoides** (Wolf, 1964) [Pompilus] (*A. malaisei*: Lelej, 2000, 2005b; Loktionov, Lelej, 2010; Loktionov, 2011; *A. sogdiana*: Tobias, 1978; Dathe, 1980b; Wahis, 1986). Russia: **EP** (N, C, NC), **ES** (KS, TU, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KA, MG). – Europe (WE, NE, SE, EE), Turkey, Near East, Uzbekistan, Tajikistan, Kyrgyzstan, Kazakhstan, Mongolia.**DICYRTOMELLUS** Gussakovskij, 1935. Type species *Sphex cingulata* Rossi, 1790 (nom. praecocc.,

nec Fabricius, 1775, replaced by *Pompilus luctuosus* Mocsáry, 1879, nom. praecocc., nec Cresson, 1865; replaced by *Dicyrtomellus argenteus* Wahis, 1983 =*Arachnospila tingitana* Wolf, 1966). Distributed in the Palaearctic region (Western, Eastern and South Europe, North Africa, Near East, European part of Russia, Central Asia, Kazakhstan, Mongolia). Number of species: Palaearctic – about 20, Russia – 3.

Dicyrtomellus kizilkumii (Radoszkowski, 1877) [Pompilus]. Russia: **EP** (NC). – Turkey, Iran, Afghanistan, Turkmenistan, Kazakhstan.**Dicyrtomellus litatus** (Tournier, 1889) [Pompilus] (*Pompilus brevigena* F. Morawitz, 1891; *D. wolfi* Móczár, 1968). Russia: **EP** (S). – Kazakhstan, Mongolia.**Dicyrtomellus tingitanus** (Wolf, 1966) [Arachnospila] (*Pompilus luctuosus* Móczáry, 1879, nom. praecocc., nec Cresson, 1865; *Dicyrtomellus argenteus* Wahis, 1983). Russia: **EP** (NC). – Europe (WE, SE, EE), Morocco, Turkey, Israel, Jordan, Syria, Iran, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan.**EVAGETES** Lepeletier de Saint Fargeau, 1845 (*Eyagates* Lucas, 1852; *Evagethes* Smith, 1858; *Sophropompilus* Howard, 1901; *Sophropompilus* Ashmead, 1902, nom. praecocc., nec Howard, 1901; *Nannopompilus* Ashmead, 1902; *Nanopompilus* Banks, 1914; *Trichosyron* Haupt, 1930; *Leuchimon* Haupt, 1930; *Asthenoctenus* Arnold, 1937, nom. praecocc., nec Simon, 1897; *Asthenoctenidia* Pate, 1946; *Psammocharoides* Móczár, 1946; *Streptosella* Dreisbach, 1950; *Streptosella* Evans, 1951; *Carinevagetes* Wolf, 1970; *Contemptevagetes* Wolf, 1970). Type species *Evagetes bicolor* Lepeletier de Saint Fargeau, 1845 (=*Aporus dubius* Vander Linden, 1827). The cleptoparasites of spider wasps (*Anoplus* Dufour, *Arachnospila* Kincaid, *Pompilus* Fabricius, *Aporinellus* Banks, *Parabatozonus* Yasumatsu, *Episyron* Schiødte, *Aporus* Spinola). Distributed mainly in Palaearctic (from Europe and North Africa in the west to Japan in the east), but also occur in Nearctic, Neotropical and Oriental regions. Number of species: World – 64, Palaearctic – 51, Russia – 20.**Evagetes alamannicus** (Blüthgen, 1944) [Pompilus]. Russia: **EP** (N, NC, CR). – Europe (WE, NE, SE, EE).**Evagetes crassicornis crassicornis** (Shuckard, 1837) [Pompilus]. Russia: **EP** (N, NW, C, E), **WS** (AL), **ES** (IR, BR, YA, ZB), **FE** (AM, SA, MG, CH). – Europe

- (WE, SE, NE, EE), Morocco, Turkey, Georgia, Kyrgyzstan, Kazakhstan, Mongolia, Canada, N America.
- Evagetes deirambo** Ishikawa, 1960 (*E. yezoensis* Ishikawa, 1960). Russia: **ES** (IR, BR), **FE** (AM, KH, PR, SA, MG). – Korean Peninsula, Japan (Hok, Hon).
- Evagetes dubius** (Vander Linden, 1827) [Aporus]. Russia: **EP** (NW, C, E, S, NC, CR), **ES** (IR, BR), **FE** (AM, PR). – Europe (WE, NE, SE, EE), Turkey, Cyprus, Turkmenistan, Kyrgyzstan, Tajikistan, Kazakhstan, Mongolia.
- Evagetes elongatus** (Lepeletier de Saint Fargeau, 1845) [Anoplus] (*Pompilus magrettii* Kohl, 1886). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), Morocco, Algeria, Turkey, Georgia, Syria, Kyrgyzstan, Kazakhstan.
- Evagetes gibbulus** (Lepeletier de Saint Fargeau, 1845) [Pompilus] (*Pompilus pectinipes* var. *pilosellus* Wesmael, 1851; *Evagetes pilosellus*: Dathe, 1980b). Russia: **EP** (NW, C, S, NC), **WS** (AL). – Europe (WE, NE, SE, EE), Turkey, Kyrgyzstan, Kazakhstan, Mongolia.
- Evagetes iconionus** Wolf, 1970. Russia: **EP** (C, CR). – Europe (WE, SE, EE), Morocco, Turkey, Georgia, Kyrgyzstan, Kazakhstan, China (?).
- Evagetes ishikawai** Lelej, 1995. Russia: **ES** (IR, BR, ZB), **FE** (AM, PR).
- Evagetes littoralis** (Wesmael, 1851) [Pompilus]. Russia: **EP** (NW, C, E, S, NC), **ES** (IR, BR, ZB), **FE** (AM, KH, PR). – Europe (WE, SE, EE), Kyrgyzstan.
- Evagetes orientalis** Lelej et Loktionov, 2009 (*Evagetes sahlbergi*: Lelej, 1992, part.; 1995: 248, part.; 2005b, part.). Russia: **FE** (AM, KH, PR, SA, KU).
- Evagetes pectinipes** (Linnaeus, 1758) [Sphecius]. Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (TK), **ES** (IR, BR, ZB), **FE** (AM, PR, SA). – Europe (WE, NE, SE, EE), North Africa, Turkey, Uzbekistan, Kyrgyzstan, Kazakhstan.
- Evagetes pontomoravicus** (Šusterá, 1938) [Psammochares] (*Evagetes atratulus* Tobias, 1978, part.). Russia: **EP** (S, NC, CR), **ES** (BR, ZB). – Europe (SE, EE), North Africa, Turkey, Cyprus, Turkmenistan, Kyrgyzstan, Kazakhstan, Mongolia.
- Evagetes proximus** (Dahlbom, 1845) [Pompilus] (*E. daisetsuzanus* Ishikawa, 1960; *E. proximus mongoloproximus* Wolf et Móczár, 1972). Russia: **EP** (NW, C, NC, CR), **WS** (AL), **ES** (IR, BR, ZB), **FE** (AM, KH, PR, MG). – Europe (WE, NE, SE, EE), Turkey, Iran, Tajikistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NC), Japan (Hok).

- Evagetes sahlbergi** (F. Morawitz, 1893) [Pompilus]. Russia: **EP** (N, NW, C, NC), **FE** (PR, KA, MG, CH). – Europe (WE, NE, SE, EE), Kyrgyzstan, Tajikistan, Mongolia.
- Evagetes siculus** (Lepeletier de Saint Fargeau, 1845) [Pompilus] (*Pompilus contemptus* Tournier, 1889). Russia: **EP** (S, NC, CR). – Europe (WE, NE, SE, EE), North Africa, Turkey, Cyprus, Syria, Kyrgyzstan, Turkmenistan, Uzbekistan.
- Evagetes sikhotealinensis** (Lelej, 1990) [Anospilus] (*Evagetes proximus* part.). Russia: **ES** (IR, ZB), **FE** (AM, KH, PR, SA, KA). – Kazakhstan, Mongolia, China (NC).
- Evagetes subglaber** (Haupt, 1941) [Psammochares] (*Psammochares subnudus* Haupt, 1941). Russia: **EP** (C, NC, CR), **WS** (AL). – Europe (WE, NE, SE, EE), Turkmenistan, Kyrgyzstan, Western Mongolia.
- Evagetes transbaicalicus** Lelej, 1995. Russia: **ES** (BR, ZB).
- Evagetes trispinosus** (Kohl, 1886) [Pompilus]. Russia: **WS** (AL). – Europe (WE, SE, EE), Tunisia, Armenia, Azerbaijan, Georgia, Turkey, Cyprus, Israel, Jordan, Lebanon, Syria, Iraq, Iran, Kyrgyzstan, Kazakhstan, Mongolia.
- Evagetes tumidosus** (Tournier, 1890) [Pompilus] (*Pompilus pedicellaris* F. Morawitz, 1891). Russia: **EP** (S, CR), **ES** (IR, ZB), **FE** (PR). – Europe (WE, SE, EE), Turkey, Kyrgyzstan, Turkmenistan, Tajikistan, Kazakhstan.
- POMPILUS** Fabricius, 1798 (*Chionopompilus* Priesner, 1955). Type species *Pompilus pulcher* Fabricius, 1798. Parasitoids of the spiders from the families Atypidae, Araneidae, Lycosidae, Pisauridae, Zoropsidae, Zoridae, Ctenidae, Clubionidae, Gnaphosidae, Thomisidae and Salticidae. Widely distributed in Palaearctic, Afrotropical, Oriental and Australian regions. Number of species: World – 7, Palaearctic – 1.
- Pompilus cinereus** (Fabricius, 1775) [Sphecius] (*Sphex plumbea* Fabricius, 1787; *Pompilus pulcher* Fabricius, 1798). Russia: **EP** (N, NW, C, E, S, NC, CR), **FE** (AM, PR). – Europe (WE, SE, NE, EE), Cyprus, Algeria, Tunisia, Egypt, Turkey, Israel, Pakistan, China (NC, SE), Korean Peninsula, Japan (Hon, Kyu, Ryukyu). Widely distributed in Afrotropical, Oriental and Australian regions.
- SCHISTONYX** Saussure, 1887. Type species *Cyphononyx velutinus* Saussure, 1887. Distributed in Palaearctic (South Europe, North Africa, Central

Asia) and Afrotropical regions. Number of species: Palaearctic – about 10, Russia – 1.

Schistonyx umbrosus (Klug, 1834) [Pompilus]. Russia: EP (NC). – Europe (WE), Morocco, Tunisia, Egypt, Syria, Israel, Jordan, Iran, Central Asia, Mauritania.

Tribe ANOPLIINI

ANOPLIUS Dufour, 1834 (*Pompioloides* Radoszkowski, 1887; *Pompiolodes* Ashmead, 1902; *Aphilocetus* Ashmead, 1902; *Paranoplius* Haupt, 1929). Type species *Sphex nigerrima* Scopoli, 1763. Parasitoids of the spiders from the families Lycosidae, Pisauridae, Oxyopidae, Ctenidae, Agelenidae, Amaurobiidae, Liocranidae, Gnaphosidae, Thomisidae, Salticidae and Philodromidae. The genus includes two subgenera: *Anoplius* Dufour, 1834 and *Arachnophroctonus* Howard, 1901 (= *Psamnochares* Latreille, 1797; = *Arachnophila* Ashmead, 1902, nom. praeocc., nec Salvadori, 1874; = *Pomphilus* Ashmead, 1902; = *Anoplindellus* Banks, 1934; = *Arachnodaicter* Pate, 1946). Worldwide distributed genus. Number of species: Palaearctic – about 30, Russia – 15.

Anoplius (Anoplius) aberrans Gussakovskij, 1932 (*Paranoplius separatus* Haupt, 1929, nom. praeocc. nec *Pomphilus separatus* Taschenberg, 1869; *A. lusonicus* Tsuneki, 1988). Russia: WS (AL), ES (IR, BR), FE (AM, KH, PR, SA, KU, KA, MG). – Uzbekistan, Korean Peninsula, China (SE), Japan (Hok, Hon, Shi, Kyu), Philippines (Luzon), Indonesia (Celebes).

Anoplius (Anoplius) alpinobalticus Wolf, 1965. Russia: EP (C, E), FE (PR). – Europe (WE, NE, SE, EE), ?Mongolia.

Anoplius (Anoplius) caviventris (Aurivillius, 1907) [Pompilus]. Russia: EP (C, E). – Europe (WE, NE, SE, EE), Turkey.

Anoplius (Anoplius) concinnus (Dahlbom, 1843) [Pompilus] (*Pomphilus bifidus* F. Morawitz, 1891; *P. distinguendus* F. Morawitz, 1891). Russia: EP (N, NW, C, E, S, NC), WS (AL), ES (KS, IR, BR), FE (AM, KH, PR). – Europe (WE, SE, NE, EE), Turkey, Cyprus, Israel, Iran, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia.

Anoplius (Anoplius) fratellus (Pérez, 1905) [Pompilus] (*A. aberrans* Gussakovskij, 1932, part.; *Anoplius (Anoplius) saigusai* Lelej in Lelej, Yamane, 1994). Russia: FE (AM, KH, PR). – Korean Peninsula, Japan (Hon, Kyu).

Anoplius (Anoplius) iwatai Yasumatsu, 1939. Russia: FE (AM, KH, PR, SA, KU). – Mongolia, Japan (Hon, Kyu).

Anoplius (Anoplius) nigerrimus (Scopoli, 1763) [Sphecius] (*Pomphilus niger*: Eversmann, 1849, Assmuss, 1859). Russia: EP (N, NW, C, E, NC, CR), UR, WS (AL), ES (IR, BR, YA), FE (AM, KH, PR, SA, KU, KA). – Europe (WE, SE, NE, EE), Turkey, Iran, Uzbekistan, Tajikistan, N America.

Anoplius (Anoplius) petiolaris Gussakovskij, 1932. Russia: FE (KH, PR, SA, KU). – Mongolia, Japan (Hok, Hon, Kyu).

Anoplius (Anoplius) ryukyuensis Tsuneki, 1990. Russia: FE (AM, KH, PR, KU). – China (SE), Japan (Hok, Kyu, Tsushima, Ryukyu).

Anoplius (Anoplius) sachalinensis Lelej, 1994. Russia: FE (AM, KH, PR, SA, KU). – Japan (Hok).

Anoplius (Anoplius) sundukovi Loktionov et Lelej, 2014 (*A. eous*: Loktionov, 2010, part.; Loktionov, 2011, part.; Lelej, Loktionov, 2012, part.; Shlyakhtenok et al., 2012, part.). Russia: FE (PR).

Anoplius (Anoplius) tenuicornis (Tournier, 1889) [Pompilus]. Russia: EP (N, NW, NC), WS (AL), ES (KR, IR), FE (AM, KH, PR, SA, KA, MG). – Europe (WE, NE, SE, EE), Central Asia, Mongolia, North America.

Anoplius (Arachnophroctonus) aeruginosus (Tournier, 1890) [Pompilus] (*Pomphilus dispar* Dahlbom, 1843, nom. praeocc., nec Latreille, 1809). Russia: EP (NW, C, NC), ES (KR). – Europe (WE, NE, SE, EE), Morocco, Turkey, Central Asia.

Anoplius (Arachnophroctonus) infuscatus (Vander Linden, 1827) [Pompilus] (*Pomphilus reflexus* Smith, 1873; *A. infuscatus mongolinfuscatus* Wolf, 1981; *A. esakii* Ishikawa, 1958). Russia: EP (N, NW, C, E, S, NC, CR), UR, ES (IR), FE (AM, KH, PR, SA, KU). – Europe (WE, SE, NE, EE), Algeria, Egypt, Cyprus, Jordan, Iran, Turkmenistan, Uzbekistan, Kyrgyzstan, Mongolia, China (NC), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryukyu).

Anoplius (Arachnophroctonus) viaticus (Linnaeus, 1758) [Sphecius] (*Pomphilus ogumae* Matsumura, 1911). Russia: EP (N, NW, C, E, S, NC, CR), UR, WS (TM, OM, TK, AL), ES (KR, IR, BR, YA, ZB), FE (AM, KH, PR, SA). – Europe (WE, SE, NE, EE), Egypt, Georgia, Turkey, Cyprus, Jordan, Iran, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NC), ?Korean Peninsula, Japan (Hok, Hon, Kyu).

LOPHOPOMPILUS Radoszkowski, 1887 (*Pompi-logaster* Howard, 1901; *Pompiologastra* Ashmead, 1902). Type species *Pompilus grandis* Eversmann, 1846. Parasitoids of the spiders from the families Araneidae, Lycosidae, Pisauridae, Gnaphosidae and Sparassidae. Distributed in Palaearctic and Nearctic regions. Number of species: World – 9, Palaearctic – 4, Russia – 2.

Lophopompilus grandis (Eversmann, 1846) [*Pompilus* (!)]. Russia: **EP** (S, CR), **UR**, **WS** (AL), **ES** (TU), **FE** (AM, PR). – Kazakhstan, Mongolia, China (NW).

Lophopompilus samariensis (Pallas, 1771) [Sphex] (*Pompilus zelleri* Dahlbom, 1843). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (OM, AL), **ES** (TU, IR, BR, ZB), **FE** (AM, KH, PR, SA). – Europe (WE, SE, EE), Turkey, Iran, Kyrgyzstan, Kazakhstan, Japan (Hon, Shi, Kyu, Tsushima, Yakushima).

Tribe PSAMMODERINI

ARACHNOTHEUTES Haupt, 1927 (*Microdrapetes* Arnold, 1937; *Phauloclavelia* Haupt, 1962; *Chionopterus* Priesner, 1965). Type species *Pompilus rufithorax* Costa, 1881. Distributed in Palaearctic (from Europe in the west to Mongolia in the east). Number of species: Palaearctic – 12, Russia – 1.

Arachnotheutes rufithorax (Costa, 1881) [*Pompilus*] (*Pompilus affinis* Eversmann, 1849, nom. praeocc., nec Vander Linden, 1827; *P. galactopterus* Kohl, 1888; *P. breviscapus* F. Morawitz, 1891; *P. cardinalis* F. Morawitz, 1891; *P. leucurus* Morawitz, 1891; *Agenioideus europaeus* Priesner, 1965; *Phauloclavelia rufipes* Priesner, 1966). Russia: **EP** (S). – Europe (WE, SE, EE), Turkey, Kyrgyzstan, Kazakhstan, Mongolia.

EOFERREOLA Arnold, 1935 (*Paraferreola* auct., nec Šuster, 1912; *Tea* Pate, 1946). Type species *Anoplus soleanus* Cameron, 1905. Parasitoids of the spiders from the family Eresidae. Distributed in Palaearctic (from Europe and North Africa in the west to the Russian Far East in the east), Nearctic and Afrotropical regions. Number of species: Palaearctic – 8, Russia – 4.

Eoferreola erythraea (Pallas, 1773) [Sphex]. Russia: **EP** (E, S, NC), **UR**, **ES** (KR). – Turkey, Kyrgyzstan, Turkmenistan, Kazakhstan.

Eoferreola manticata (Pallas, 1771) [Sphex] (*Salius hellmanni* Eversmann, 1849; *Ferreola caucasica* Radoszkowski, 1888; *F. rossica* Radoszkowski, 1888). Russia: **EP** (E, S, NC, CR), **UR**, **WS** (AL), **FE** (PR). – Europe (WE, SE, EE), Morocco, Algeria, Tunisia, Egypt, Azerbaijan, Turkey, Israel, Jordan, Lebanon, Syria, Iran, Uzbekistan, Turkmenistan, Tajikistan, Kyrgyzstan, Kazakhstan.

Eoferreola rhombica (Christ, 1791) [Sphex] (*Pompilus fasciatus* Eversmann, 1849). Russia: **EP** (S, NC, CR), **UR**, **WS** (AL), **ES** (KR, IR, BR), **FE** (AM). – Europe (WE, SE, EE), Azerbaijan, Turkey, Uzbekistan, Kyrgyzstan, Tajikistan, Kazakhstan, Mongolia.

Eoferreola variabilis (Eversmann, 1849) [*Pompilus*] (*Ferreola distincta* Smith, 1855; *Salius grandis* Radoszkowski, 1887). Russia: **EP** (S, NC, CR), **UR**. – Europe (SE), Algeria, Egypt, Turkey, Syria, Iran, Uzbekistan, Kyrgyzstan, Kazakhstan.

Tribe EIDOPOMPILINI

MICROPHADNUS Cameron, 1905 (*Plagioceps* Haupt, 1930). Type species *Microphadnus bicolor* Cameron, 1905. Distributed in Palaearctic (from Europe and North Africa in the west to Mongolia in the east), Afrotropical (South Africa, Madagascar) and Australian regions. Number of species: Palaearctic – 4, Russia – 1.

Microphadnus pumilus (Costa, 1886) [Aporus]. Russia: **EP** (NC). – Europe (WE, SE), Morocco, Tunisia, Egypt, Turkey, Lebanon, Israel, Syria, Afghanistan, Uzbekistan, Kyrgyzstan, Tajikistan, Kazakhstan, Mongolia.

STIGMAPORUS Zonstein, 2001. Type species *Gonaporus wolfi* Zonstein, 2001. Distributed in Palaearctic (European part of Russia, Central Asia, Kazakhstan, Mongolia and China: Xinjiang). Number of species: World – 4, Palaearctic – 4, Russia – 1.

Stigmaporus volgadon Loktionov et Lelej, 2016. Russia: **EP** (S).

TACHYAGETES Haupt, 1930 (*Pareixenetus* Haupt, 1962; *Dasyagetes* Priesner, 1955; *Exagetes* Priesner, 1955; *Epagetes* Priesner, 1955). Type species *Evagetes filicornis* Tournier, 1889. Parasitoids of the spiders from the families Lycosidae, Clubionidae, Gnaphosidae and Salticidae. Distributed in Palaearctic region (from Europe and North Africa

in the west to Mongolia in the east). Number of species: Palaearctic – about 90, Russia – 6.

Tachyagetes (Tachyagetes) argentatus Haupt, 1930.
Russia: UR. – Turkmenistan, Uzbekistan.

Tachyagetes (Tachyagetes) consimilis Tobias, 1978.
Russia: EP (S).

Tachyagetes (Tachyagetes) dudichi Móczár, 1944.
Russia: EP (NC). – Europe (EE).

Tachyagetes (Tachyagetes) filicornis (Tournier, 1889)
[Evagetes]. Russia: EP (S, NC). – Europe (WE, SE,
EE), Morocco, Turkey, Cyprus, Iran, Uzbekistan,
Tajikistan, Kyrgyzstan, Turkmenistan, Kazakhstan,
Mongolia.

Tachyagetes (Tachyagetes) grandis (Tournier, 1889)
[Evagetes]. Russia: EP (S, NC, CR). – Kyrgyzstan,
Turkmenistan, Kazakhstan.

Tachyagetes (Tachyagetes) gratiosus (Radoszkowski, 1893) [Aporus]. Russia: EP (NC). – Turkey,
Iran, Turkmenistan, Uzbekistan, Tajikistan, Kyrgyzstan,
Kazakhstan.

TELOSTEGUS Costa, 1887 (*Elaphrosyron* Haupt,
1929; *Protelostegus* Priesner, 1955). Type species
Aporus major Costa, 1886 (=*Aporus inermis*
Brullé, 1833). Distributed in Palaearctic (from Europe
and North Africa in the west to Mongolia in the
east), Afrotropical and Australian regions. Number
of species: Palaearctic – about 20, Russia – 1.

Telostegus inermis (Brullé, 1833) [Aporus] (*Aporus
major* Costa, 1886). Russia: EP (S, NC). – Europe
(WE, SE, EE), Morocco, Algeria, Turkey, Cyprus,
Iran, Kyrgyzstan, Tajikistan, Kazakhstan.

XENAPORUS Ashmead, 1902 (*Taeniaporus* Haupt,
1930). Type species *Pompilus amoenus* Klug,
1834. The genus includes two subgenera: *Baguenaia*
Giner Mari, 1942 (=*Nannaporus* Priesner, 1966;
=*Melanaporus* Priesner, 1967; =*Ciliaporus* Wahis,
1972) and *Xenaporus* Ashmead, 1902. Distributed
in Palaearctic (from Europe and North Africa in
the west to Mongolia in the east) and Nearctic
regions. Number of species: Palaearctic – about
35, Russia – 1.

Xenaporus (Baguenaia) eremocanus Wolf, 1990.
Russia: ES (TU). – Turkmenistan, Uzbekistan,
Kyrgyzstan, Kazakhstan, Mongolia.

Doubtful or wrongly recorded taxa

Ceropales pygmaea pygmaea Kohl, 1880. Russia:
“Eastern Siberia”.

Ceropales ruficornis Gussakovskij, 1931. Russia: “Russian
(Asia)”, “Russian SSR”.

Eoferreola anatolica Priesner, 1973. Russia: “Russia:
Kulp”; “Russia”.

Hemipepsis sinensis (Smith, 1855). Russia: “Russia”.
Pompilus trivialis var. **maris** Eversmann, 1849. Doubtful
taxon, probably belongs to the *Arachnospila*
(*Ammosphex*) spp.

Telostegus nigritulus (Klug, 1834). Russia: “in provin-
ciis Orenburg., Saratov. et Astrachanensi” [actually
North Africa].

SUPERFAMILY VESPOIDEA**68. FAMILY VESPIDAE**

A.V. ANTROPOV AND A.V. FATERYGA

Vespidae are the wasps of predominantly medium size, though rarely relatively small (2.5 mm) to very large (up to 50 mm). Their most distinct differences are the inner eye orbits with more or less pronounced notches, the pronotum laterally extending back to the tegulae and with deep oval or triangular dorsal emargination, and the forewings folding longitudinally at rest (except Stenogastrinae, Euparagiinae, and part of Masarinae), with discoidal cell 1 longer than half the wing length. The body of vespids has usually typical “wasp” coloration, consisting of more or less regular yellow or white spots and bands on black background, but it may be black with reddish markings, or completely black-brown or yellow, while some species of *Ropalidia* Guérin-Méneville, 1831 may be even green.

The family includes six extant subfamilies: Euparagiinae, Masarinae, Eumeninae, Stenogastrinae, Polistinae, and Vespinae. The first three groups include only solitary insects, while the last three demonstrate the gradual development of sociality. Russian fauna is represented only by the subfamilies Eumeninae (potter wasps), Masarinae (pollen wasps), Polistinae (paper wasps), and Vespinae (yellowjackets and hornets). There are also two subfamilies known only by fossil records: Priorvespinae and Protovespinae.

The subfamily Euparagiinae is a small group of relic vespids containing the single extant genus *Euparagia* Cresson, 1879 with 10 species, which are known only from the southwest of North America. These wasps have habits similar to the eumenine genus *Odynerus*. Their females excavate vertical burrows in ground and surmount them by mud turrets (or “chimneys”); the nest contains several cells provisioned with larvae of Curculionidae.

The members of the subfamily Masarinae are most diverse and abundant in arid regions of the World. They are present in all zoogeographical regions, however only one species of the genus *Quartinia* André, 1884 is known from the Oriental region. This subfamily includes two tribes: Gayellini, with two Neotropical genera, and Masarini, which is additionally divided on three subtribes: Priscomasarina, with one monotypic Afrotropical genus, Paragiina, with two Australian genera, and Masarina, with nine genera distributed

in the Nearctic (1), the Neotropical (2), the Palaearctic (1), the Afrotropical (1), and both Palaearctic and Afrotropical (4) region. Masarines are unique among solitary vespids in feeding their larvae exclusively with pollen and nectar. They nest in ground and excavate burrows surmounted by mud entrance turrets or build aerial nests consisted of several ovoid (*Gayella* Spinola, 1851) or cylindrical (*Celonites* and *Pseudomasaris* Ashmead, 1902) cells usually covered with additional mud layer. Nectar is used by females as bonding agent in nests of *Celonites* and *Pseudomasaris*, while ground nesting masarines usually use water during the nest construction. Species of the genus *Quartinia*, nesting in friable soil, use silk to confirm the inner walls of the nest burrow and the cells as well as to construct the turret. All masarines lay an egg into empty cell prior to its provisioning; an egg is usually not suspended by a filament (except *Gayella*). **Number of taxa** of Masarinae: World – 14 genera, 371 species, Palaearctic – 5/115, Russia – 1/2.

The members of the subfamily Eumeninae are also solitary wasps, except just several species that demonstrate a primitively social behavior (nest-sharing, reuse of brood cells, progressive provisioning, and strong aggressive competition among resident females for empty cells). This is the most divers subfamily distributed worldwide. It contains three tribes: Zethini (18 genera, 354 species), Eumenini (20/365), and Odynerini (167/3039). The nesting habits of the eumenines are extremely diverse. Species of the tribe Zethini usually build nests in preexisting cavities and use pressed pith or leaf cuts as building material for partitions between nest cells. However some species in the genera *Zethus* Fabricius, 1804 and *Calligaster* de Saussure, 1852 build aerial nests of leaf cuts. Species of the tribe Eumenini build nest cells in the shape of juglike pots with narrow collar, these cells are made usually of mud (mix of dry ground and water) or rarely (*Katamenes*) of pebbles confirmed with mud. Each nest can contain one or several cells and is usually covered with additional layer of mud (or pebbles with mud in *Katamenes*). Tribe Odynerini has the most diverse nesting habits. Some genera nest in ground and excavate vertical or horizontal burrows surmounted by nest turrets, others nest in preexisting cavities (hollow stems, galls, beetle burrows in wood, snail shells and abandoned nests of other wasps and bees), and the rest build aerial nests consisting of several cells, usually with additional cover. Building material in Odynerini is generally mud made of dry ground mixing with water. However, species

in the genera *Pterocheilus* and *Onychopterocheilus* nest in friable soil and do not use water to construct their nests; species of the genus *Leptochilus* and some species of *Odynerus* nesting in preexisting cavities use dry pressed pith, ground particles or gravel. One species of Odynerini, *Pseudabispa paragioides* Meade-Waldo, 1910 is known as the cleptoparasite of *Abispa ephippium* Fabricius, 1775. In most cases the females of Eumeninae lay an egg into empty cell prior to its provisioning; an egg is usually suspended to cell wall or sealing by a filament. However, some species of *Synagris* Latreille, 1802, *Calligaster*, *Paraleptomenes* Girdani Soika, 1970, *Paravespa*, and *Tropidodynerus* do not attach their eggs. The cells of eumenine wasps are provisioned with paralyzed caterpillars or larvae of beetles (usually Curculionidae or Chrysomelidae) or sawflies. Most species are mass provisioning, while about two tens species are progressively provisioning. **Number of taxa** of Eumeninae: World – 205 genera, 3758 species, Palaearctic – 78/963, Russia – 33/158.

Social vespids form more or less numerous families, including one to several fertile “queens” laying eggs, and “workers”, which are their offspring, performing the auxiliary duties of guarding and building the nest and feeding the larvae. The “workers” are sterile females, as a rule not laying eggs.

The subfamily Stenogastrinae (hover wasps), containing 7 genera and 63 medium-sized (11–21 mm long) slender species with a very long and narrow metasomal segment 1, is known only from the Oriental region and the island of New Guinea. Unlike other social vespids their nests has no primary petiole, and the ventrally open cells are attached directly to the substratum – thin branches of trees, vine tendrils, loose strands of thatch under the roofs of houses, or rarely under stones. The cells are made of chewed plants in composition with the material of the previous cells and sometimes with more or less amount of wet clay. The wasps increase the length of each cell as far as the larvae grow. Individual families of stenogastrine species are very small, but sometimes they form nesting clusters, where the “workers” may move between multiple nests.

The subfamilies Polistinae and Vespinae are distributed worldwide, though the first is most variable in the New World (21 genus among 25 of the World fauna), while the second is mostly represented in the Old World, mainly in the Oriental region. The combs and walls of the cells are made of chewed wooden fibers, sometimes mixed with wet soil and sand.

Subfamily Polistinae includes four tribes. The members of Epiponini (19 genera, 242 species) and Mischoctytarini (1/249) exclusively inhabit subtropical and tropical regions of the New World, those of Ropalidiini (4/288, including 3/34 in Palaearctic) inhabit the same regions of the Old World, and only the species of Polistini, including the single genus *Polistes*, are distributed worldwide (World – 224 species, Palaearctic – 33, Russia – 14). Typically Polistinae form eusocial families consisting of a “queen” and more or less numerous “workers”, which are not distinctly differ in size. The vast majority of polistines build the nests consisting of the single comb fasten to substratum with a firm petiole and lacking in outer envelope. They usually feed larvae with masticated caterpillars. At the same time, some representatives of Epiponini demonstrate high diversity in behavior and construction of nests. The species of *Apoica* Lepeletier de Saint Fargeau, 1836 are truly nocturnal wasps. Some species of *Agelaia* Lepeletier de Saint Fargeau, 1836 build giant nests, containing several combs without envelope, and their colonies may number more than a million stinging wasps. Finally, some species of Epiponini cover their combs with an outer envelope, which may be soft in *Polybia* Lepeletier de Saint Fargeau, 1836 or *Synoeca* de Saussure, 1852, but hard cartoon-like in *Chartergus* Lepeletier de Saint Fargeau, 1836 and *Brachygastera* Perty, 1833. The last genus is also known as a honey producer. The Old World members of Polistinae build nests with the cells disposed in one flat comb or sometimes in vertical rows along the pendent shoots of plants, but always without envelopes. **Number of taxa** of Polistinae: World – 25 genera, 1003 species, Palaearctic – 4/67, Russia – 1/14.

The nests of Vespinae built of masticated wooden fibers, usually consist of several combs, connected by columns and covered with outer envelope with the entrance in the lower part. Such nests may be free and attached to the tree branches or placed in various shelters from the hollows of the tree-trunks and underground cavities to human buildings. Vespinae feed their larvae with various masticated insects and spiders, though some species use all accessible sources of proteins, including dead invertebrates and vertebrates. Vespinae are usually active during day-light hours, and only three species of the genus *Provespa* Ashmead, 1903 and the East Asian hornet *Vespa binghami* are truly nocturnal wasps. **Number of taxa** of Vespinae: World – 4 genera, 69 species, Palaearctic – 3/36, Russia – 3/23.

The life cycle of the families of social wasps in the moderate climate (though often also in tropics) is annual. After wintering the females inseminated in the end of the previous season build new nests and feed the first generation of "workers". After that, all the care to bring the materials for the construction of new cells, and food for the larvae becomes the responsibility of "workers". The "queens" of Vespinae, with the rare exceptions (for example, in *Vespa ducalis*), differ by distinctly larger size, and they completely dominate "workers". New generation of females and males appears in the end of active season. After copulation young females look for the places for wintering, while the rest members of the family die.

Among social vespids there are several obligate nesting parasites (in Russia these are *Vespula austriaca*, *Dolichovespula adulterina*, *D. omissa*, *Polistes atrimandibularis*, and *P. sulcifer*), who do not build their own nests, but usurp the nests of the allied species. Obligate parasites have not "workers", and their first generation consists of capable to copulate males and females. Other species (*Vespa dybowskii* and *Polistes nimpha*) may be facultative nesting parasites, which can usurp the nests of the allied species, though are able to form their own families and build the nests.

Many species of Vespidae are important regulators of quantity of various harmful insects, such as Lepidoptera, Orthoptera, and Diptera, and may be effective pollinators. *Brachygastra mellifica* (Say, 1837) is a traditional source of honey in Central America, while some species of *Belonogaster* de Saussure, 1854 are used in traditional medicine in Africa, and the larvae and pupae of *Vespa* are the parts of nourishing diet of the local inhabitants in the South Eastern Asia. At the same time, some species of *Vespa* which hunt honey bees may cause serious damage to apiculture, and species of *Vespa* and *Vespula* sometimes cause severe damage to the harvests of sweet fruit, particularly to grapes and pears, in the period of their ripening. Finally, some species of Polistinae and Vespinae represent certain medical problem, because their attacks, which are usually caused by the carelessness of people, annually lead to hundreds of lethal outcomes in different countries.

Number of taxa of Vespidae: Word – 256 genera, 5274 species; Palaearctic – 90/1181; Russia – 38/197.

R e f e r e n c e s. Morawitz, 1867, 1895; Horne, Smith, 1872; André, 1881–1886; Mocsáry, Szépligeti, 1901; Alfken, 1915; Birula, 1925, 1927, 1930a,b; Bialynicki-Birula, 1926; Kostylev, 1928, 1929, 1940a,b; Bequaert,

1930; Zimmermann, 1930; Gussakovskij, 1932; de Beaumont, 1944; Guiglia, 1948; Cavro, 1950; Richards O., Richards M., 1951; Giordani Soika, 1958, 1976; Blüthgen, 1961; Richards O., 1962, 1973; Løken, 1964; Gusenleitner, 1966a,b, 1972a,b, 1977, 1981, 1991, 2004, 2007a,b, 2013a,b; Kemper, Döhring, 1967; van der Vecht, Fischer, 1972; Spradbery, 1973; Yamane Sk., 1975; Kurzenko, 1977a,b, 1978a,b, 1981, 1995, 2004, 2012; Tobias, Kurzenko, 1978; Day, 1979; Archer, 1980, 1981a,b, 1982, 1989a,b, 1992, 1995, 1997, 1998a,b, 1999a,b, 2006, 2008; Dathe, 1980b; Yamane Sk. et al., 1980; Akre, 1982; Carpenter, 1986, 1996, 2001; Lee, 1986; Cumming, 1989; Scobiola-Palade, 1989; van der Vecht, Carpenter, 1990; Turillazzi, 1991; Kim et al., 1994; Gess, 1996; Carpenter, Kojima, 1996, 1997, 2002; Ono, 1997; Dubatolov, 1998, 1999, 2011; Dubatolov et al., 2002; Castro, 2003; Carpenter, Nguyen, 2003; Dubatolov, Novomodnyi, 2005; Cervo, 2006; Dvořák, Roberts, 2006; Dvořák, Castro, 2007; Castro et al., 2008; Ebrahimi, Carpenter, 2008; Gereys, 2008; Amolin, 2009; Castro, Dvořák, 2009, 2010; Dubatolov, Dolgikh, 2009; Aliyeva, 2010; Carpenter, Glare, 2010; Carpenter et al., 2011; Rudoiskatel et al., 2011; Abasheev, 2012; Archer, Penney, 2012; Hoa et al., 2012; Kim, 2012, 2013; Mahmood et al., 2012; Malikova, 2012; Shibaev, Polumordvinov, 2012; Yildirim, Gusenleitner, 2012; Zhou et al., 2012; Abasheev, Dabaev, 2014; Amolin, Artokhin, 2014; Antropov, Proshchalykin, 2014; Hermes et al., 2014; Li, Chen, 2014; Abasheev, Buyanjargal, 2015; Knyazev et al., 2015; Neumeyer, Praz, 2015; Yamane, Terayama, 2016; Abasheev, 2016; Buyanjargal et al., 2016; Ma et al., 2016; Mauss et al., 2016; Pham, Kumar, 2016; Ruchin, Antropov, 2016; Fateryga, 2017, 2018; Fateryga et al., 2017; Fateryga, Popov, 2017; Kumar et al., 2017; Perrard et al., 2017.

Subfamily MASARINAE – POLLEN WASPS

A.V. FATERYGA

CELONITES Latreille, 1802 (*Chelonites* (!): Leach, 1815; *Coelonites* (!): Blanchard, 1840; *Eucelonites* Richards, 1962). Type species *Masaris apiformis* Fabricius, 1793. Females build aerial mud nests consisted of cylindrical cells usually covered with additional mud layer; nests are usually attached to stones or twigs (some Afrotropical species place their nest cells within excavated or preexisting holes in ground). The genus is distributed in the Palaearctic and Afrotropical regions. Number of species: World – 61; Palaearctic – 39; Russia – 2.

Celonites sibiricus Gусенлеитнер, 2007. Russia: **WS** (AL).

Celonites tauricus Костyleв, 1935 (*C. abbreviatus*: Костyleв, 1928, 1929; *C. abbreviatus tauricus* Костyleв, 1935; *C. spinosus* Гусенлеитнер, 1966; *C. abbreviatus invitus* Гусенлеитнер, 1973; *C. abbreviatus invitis* (!): Schmidt, 1984). On flowers of *Teucrium*, *Thymus*, and *Ziziphora* (Lamiaceae). Russia: **EP** (CR). – Europe (SE: Greece), Armenia, Azerbaijan, Turkey, Cyprus, Syria, Iran.

Subfamily EUMENINAE – POTTER WASPS

A.V. FATERGYA

ALASTOR Lepeletier de Saint Fargeau, 1841 (*Antalastor* de Saussure, 1856; *Eualastor* Dalla Torre, 1904; *Belalastor* Атанасов, 1967). Type species *Alastor atropos* Lepeletier de Saint Fargeau, 1841. Females nest in preexisting cavities such as hollow stems or empty snail shells; partitions between cells are made of gravel and mud; provision is larvae of Curculionidae. The genus is distributed in the Palaearctic, Afrotropical, and Oriental regions and includes four subgenera: *Alastor* s.str.; *Alastorellus* Giordani Soika, 1991; *Megalastor* Blüthgen, 1951; *Parastalor* Blüthgen, 1939. Number of species: World – 79; Palaearctic – 28; Russia – 1.

Alastor (Alastor) mocsaryi (André, 1884) [Odynerus] (*A. atropos*: Morawitz, 1895, Kostylev, 1928; *A. biegelebeni* Giordani Soika, 1942; *A. biegelebeni pannonicus* Blüthgen, 1956; *A. kuehlhorni* Giordani Soika, 1958; *A. biegelebeni* (!): Tobias, Kurzenko, 1978). Russia: **EP** (E, S, NC, CR). – Europe (WE, SE, EE), Caucasus, Turkey, Syria, Lebanon.

ALLODYNERUS Blüthgen, 1938 (*Delphinaloides* Bohart, 1951; *Acarepipona* Giordani Soika, 1985). Type species *Odynerus floricola* de Saussure, 1853. Females nest in preexisting cavities such as beetle burrows in wood or hollow stems; partitions between cells are made of mud; provision is caterpillars, mainly of Gelechiidae. The genus is distributed in the Palaearctic, Afrotropical and Oriental regions. Number of species: World – 11; Palaearctic – 9; Russia – 4.

Allodynerus delphinalis (Giraud, 1866) [Odynerus] (*O. tristis* Thomson, 1870; *O. novellus* Morawitz, 1895; *Allodynerus delphinalis fallax* Blüthgen, 1953;

A. delphinalis sardous Blüthgen, 1953). Russia: **EP** (E, S, NC, CR), **UR**, **WS** (AL), **FE** (AM, PR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Lebanon, Iran, Central Asia, Kazakhstan, China (NE), Korean Peninsula, Japan (Hon).

Allodynerus floricola (de Saussure, 1853) [Odynerus] (*Odynerus floricola* ab. *immaculata* Mader, 1936; *Allodynerus floricola inaequalis* Giordani Soika, 1970). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), N Africa, Armenia, Turkey, Cyprus, Jordan.

Allodynerus mandschuricus Blüthgen, 1953. Russia: **ES** (BR), **FE** (PR). – Mongolia, China (NE), Korean Peninsula, Japan (Hon).

Allodynerus rossii (Lepeletier de Saint Fargeau, 1841) [Odynerus] (?*Odynerus hospes* Dufour et Perris, 1840; *O. notabilis* Morawitz, 1895, nom. praeocc., nec Walker, 1871; *O. umbilicalis* Dalla Torre, 1904). Russia: **EP** (C, ?E, S, NC, CR), **ES** (?BR, ?ZB). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Iran, Turkmenistan.

ANCISTROKERUS Wesmael, 1836 (*Aucistrocerus* (!): Rudow, 1876; *Ancystrocerus* (!): Dalla Torre, 1894; *Euancistrocerus* Dalla Torre, 1904). Type species *Vespa parietum* Linnaeus, 1758. Females nest in various preexisting cavities: hollow stems, beetle burrows in wood, abandoned nests of other wasps and bees; several species (*A. densipilosellus*, *A. japonicus*, *A. oviventris*) build aerial nests consisted of mud cells attached to stones and covered with additional layer of mud. Provision is caterpillars of various Lepidoptera or rarely larvae of beetles (Chrysomelidae or Curculionidae) or sawflies (Tenthredinidae). The genus is distributed worldwide except Australia. Number of species: World – 116; Palaearctic – 57; Russia – 19.

Ancistrocerus antilope (Panzer, 1798) [Vespa] (?*Vespa insolens* Harris, 1780; *Odynerus pictus* Curtis, 1826; *O. capra* de Saussure, 1857; *Ancistrocerus nearcticus* Cameron, 1906; *A. lecontei* Cameron, 1908; *A. navajo* Bequaert, 1925; *A. antilope allegrus* Bequaert, 1943; *A. antilope spenceri* Bequaert, 1943). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TK, NS, AL), **ES** (KR, IR, BR, ZB), **FE** (AM, KH, PR, SA). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Afghanistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE), Japan (Hon), N America, India.

Ancistrocerus auctus (Fabricius, 1793) [Vespa]. Russia: **EP** (C, S, NC, CR). – Europe (WE, SE, EE), Caucasus, Turkey, Syria, Jordan, Israel, Iraq, Iran, Central Asia, Kazakhstan.

Ancistrocerus claripennis Thomson, 1874 (?*Vespa quadrata* Panzer, 1799; *Ancistrocerus claripennis* var. *subrenimacula* Blüthgen, 1938; *A. claripennis ponticus* Giordani Soika, 1970). Russia: **EP** (N, NW, C, NC). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Tajikistan, China (NC).

Ancistrocerus densepilosellus Cameron, 1911. Russia: **WS** (NS, AL), **ES** (IR, BR), **FE** (AM, KH, PR, KU). – Mongolia, ?China (NE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu).

Ancistrocerus dusmetiolas (Strand, 1914) [*Odynerus*] (*Odynerus sociabilis* Dusmet, 1903, nom. praeocc., nec Perkins, 1899; *O. alius* Kostylev, 1935). Russia: **WS** (AL), **ES** (KR). – Europe (WE, SE, EE), Turkey, Afghanistan, Central Asia, Kazakhstan.

Ancistrocerus gazella (Panzer, 1798) [*Vespa*] (?*Vespa emarginata* Fabricius, 1793; *Ancistrocerus pictipes* Thomson, 1874; *Odynerus parietum*: Kostylev, 1928; *A. verhaereni* Bondroit, 1943; *A. claripennis*: Tobias, Kurzenko, 1978). Russia: **EP** (C, E, S, NC, CR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Cyprus, Israel, Iran, Afghanistan, ?Pakistan, N America (introduced), New Zealand (introduced).

Ancistrocerus hangaicus Kurzenko, 1977. Russia: **WS** (AL), **ES** (IR, BR, ZB), **FE** (MG). – Mongolia.

Ancistrocerus ichneumonideus (Ratzeburg, 1844) [*Vespa*] (*A. thomsoni* Blüthgen, 1938; *A. ichneumonideus kaszabi* Giordani Soika, 1970). Russia: **EP** (NW, C, NC), **UR**, **WS** (AL), **ES** (KR, IR, BR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Kazakhstan, Mongolia.

Ancistrocerus japonicus (von Schulthess, 1908) [*Odynerus*]. Russia: **FE** (PR). – Korean Peninsula, Japan (Hon, Shi, Kyu, Ryu).

Ancistrocerus kazbekianus (Kostylev, 1940) [*Odynerus*]. Russia: **EP** (NC). – Georgia.

Ancistrocerus leleji Kurzenko, 1995. Russia: **FE** (PR).

Ancistrocerus mongolicus (Kostylev, 1940) [*Odynerus*]. Russia: **WS** (AL), **ES** (?KR, IR), **FE** (AM, KH, PR). – Mongolia, Korean Peninsula.

Ancistrocerus nigricornis (Curtis, 1826) [*Odynerus*] (?*Vespa sexpunctata* Christ, 1791; *Odynerus callosus* Thomson, 1870; *O. excisus* Thomson, 1870; *Ancistrocerus nigricornis* var. *polonica* Blüthgen, 1952). Russia: **EP** (NW, C, ?E, S, NC, CR), **UR**, **WS** (NS, KM, AL), **ES** (KR, IR, BR, YA, ZB), **FE** (KH, PR, SA). – Europe (WE, NE, SE, EE), N Africa, Turkey, Iran, Central Asia, Kazakhstan, Mongolia, China (NE), Japan (Hok, Hon), India.

Ancistrocerus oviventris (Wesmael, 1836) [*Odynerus*] (*Odynerus constans* Herrich-Schäffer, 1839; *O. viduus* Herrich-Schäffer, 1839; *O. pictus* var. *hibernica* Blüthgen, 1937; *O. oviventris* var. *rufipes* Blüthgen, 1938; *A. oviventris siculosus* Blüthgen, 1955; *A. yamanei* Giordani Soika, 1986; *A. oviventris caucasicus* Guseinleitner, 1994). Russia: **EP** (N, NW, C, S, NC, CR), **UR**, **WS** (TK, AL), **ES** (IR, BR, YA, ZB), **FE** (KH, PR, SA, MG). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Central Asia, Kazakhstan, Mongolia, China (NE), Korean Peninsula, Japan (Hok).

Ancistrocerus parietinus (Linnaeus, 1761) [*Vespa*] (*Sphex domestica* Christ, 1791; *Odynerus affinis* Herrich-Schäffer, 1839; *O. thuringiacus* Blüthgen, 1937). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (TK, NS, AL), **ES** (KR, IR, BR, ZB), **FE** (AM, KH, PR, SA). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Iran, Kazakhstan, Mongolia, Korean Peninsula, Japan (Hok).

Ancistrocerus parietum (Linnaeus, 1758) [*Vespa*] (?*Vespa parietum* β *nigra* Poda, 1761; ?*V. aeneipennis* Christ, 1791; ?*Odynerus geoffroyanus* Spinola, 1808; ?*V. angulata* Donovan, 1810; *O. flavipes* Curtis, 1826; *O. similis* Curtis, 1826; *O. ochlerus* de Saussure, 1852; *O. parietum* var. *incisoides* Verhoeff, 1890; *O. parietum* var. *incisus* Verhoeff, 1890; *O. parietum* var. *parietoides* Verhoeff, 1890). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (OM, TK, NS, AL), **ES** (KR, IR, BR, ZB), **FE** (AM, KH, PR, SA). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, ?Iran, Central Asia, Kazakhstan, Mongolia, China (NE, NC), Korean Peninsula, N America (introduced).

Ancistrocerus raddei (Kostylev, 1940) [*Odynerus*]. Russia: **WS** (AL), **ES** (ZB). – Mongolia, China (NC, NW).

Ancistrocerus scoticus (Curtis, 1826) [*Odynerus*] (*Odynerus albotruncatus* Zetterstedt, 1838; *O. albotruncatus* (!): Dalla Torre, 1894; *Ancistrocerus scoticus iberogallicus* Blüthgen, 1956; *A. scoticus meridianus* Guseinleitner, 1977). Russia: **EP** (N, NW, NC), **UR**, **WS** (TK, NS, AL), **ES** (KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KA, MG). – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, Iran, Kazakhstan, Mongolia, China (NE, NC), Korean Peninsula.

Ancistrocerus trifasciatus (Müller, 1776) [*Vespa*] (*Vespa trifasciata* Fabricius, 1787; ?*V. iuncea* Christ, 1791; *Odynerus trimarginatus* Zetterstedt, 1838;

O. tricinctus Herrich-Schäffer, 1839; *O. trifasciatus orientalis* Kostylev, 1938, nom. praeocc., nec *O. orientalis* Dalla Torre, 1889; *O. shibuyai* Yasumatsu, 1938; *Ancistrocerus trifasciatus* var. *moe-schleri* Blüthgen, 1938; *A. trifasciatus kostylevi* van der Vecht, 1972). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KA). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Kyrgyzstan, Kazakhstan, Mongolia, China (NE), Korean Peninsula, Japan (Hok, Hon).

ANTEPIPONA de Saussure, 1855 (*Antepiponus* (!): de Saussure, 1875; *Antepipone* (!): Dalla Torre, 1894; *Odontodynerus* Blüthgen, 1938; *Dichodynerus* Blüthgen, 1938; *Metastenancistrocerus* Blüthgen, 1938; *Mehelyella* Bohart, 1951). Type species *Odynerus silaos* de Saussure, 1853. Females nest in ground, often on vertical cliffs, and excavate burrows surmounted by turrets; provision is caterpillars of Psychidae, Tortricidae and Gelechiidae. The genus is distributed in the Palaearctic, Afrotropical, and Oriental regions. Number of species: World – 165; Palaearctic – 54; Russia – 4.

Antepipona cibrata (Morawitz, 1885) [*Lionotus*] (*Odynerus biarcuatus* Morawitz, 1895). Russia: **EP** (NC). – Armenia, Azerbaijan, Turkey, Afghanistan.

Antepipona deflenda (S. Saunders, 1853) [*Ancistrocerus*] (*Odynerus parvulus* Lepeletier de Saint Fargeau, 1841; *O. parvulus* var. *ruthenicus* Morawitz, 1867; *O. bidenticulatus* Maidl, 1922; *Odontodynerus deflendus lepeletieri* Blüthgen, 1961). Russia: **EP** (?E, S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Cyprus, Jordan, Lebanon, Israel, Iraq, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, China (?NC).

Antepipona nigricornis (Morawitz, 1885) [*Lionotus*] (*Nannodynerus binominatus* Blüthgen, 1939; *Alodynus hispanicus* Gusenleitner, 1993; *Antepipona gusenleitneri* Tussac, 1996). Russia: **EP** (NC, CR). – Europe (WE, SE), Turkey, Turkmenistan.

Antepipona orbitalis (Herrich-Schäffer, 1839) [*Odynerus*] (*Odynerus ballioni* Morawitz, 1867; *O. rubripes* André, 1884; *O. parvulus* var. *alpicola* Berland, 1928). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (NS), **ES** (IR, BR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Kazakhstan, Mongolia.

BRACHYODYNERUS Blüthgen, 1938. Type species *Odynerus magnificus* Morawitz, 1867. Females

excavate nest burrows in lateral walls of ground cracks; provision is caterpillars of Coleophoridae. The genus is distributed in the Palaearctic. Number of species: Palaearctic – 9; Russia – 2.

Brachyodynerus magnificus (Morawitz, 1867) [Odynerus] (*Odynerus morawitzi* André, 1884; *O. magnificus luxorius* Kostylev, 1929). Russia: **EP** (S, NC, CR). – Europe (EE), Armenia, Azerbaijan, Turkey, Turkmenistan, Kazakhstan.

Brachyodynerus quadrimaculatus (André, 1884) [*Odynerus*]. Russia: **EP** (S, NC, CR). – Europe (SE: Greece), Azerbaijan, Turkey.

BRACHYPIPONA Gusenleitner, 1967 (*Desertodynerus* Kurzenko, 1977). Type species *Pseudepipona schmidti* Gusenleitner, 1967. Bionomics of the genus is unknown. The genus is distributed in the Palaearctic. Number of species: Palaearctic – 7; Russia – 1.

Brachypipona laticeps (Morawitz, 1895) [Odynerus]. Russia: **EP** (S). – Kazakhstan.

CHLORODYNERUS Blüthgen, 1951. Type species *Odynerus chloroticus* Spinola, 1838. Bionomics of the genus is unknown. The genus is distributed in the Palaearctic, Afrotropical, and Oriental regions. Number of species: World – 26; Palaearctic – 19; Russia – 1.

Chlorodynerus ypsilon (Kostylev, 1929) [Odynerus] (*Euodynerus ypsilon rhodius* Blüthgen, 1954). Russia: **EP** (NC). – Europe (SE), Caucasus, Turkey, Cyprus, Syria, Israel.

DELTA de Saussure, 1855 (*Phi* de Saussure, 1855, nec 1854; *Alfieria* Giordani Soika, 1934; *Alfelia* (!): Neave, 1939; *Erinys* Zirngiebl, 1953, nec Rye, 1876). Type species *Vespa maxillosa* De Geer, 1773. Females build aerial nests consisted of several juglike mud cells covered with additional mud layer; provision is caterpillars of Geometridae, Noctuidae, Lyceanidae and Pieridae. The genus is distributed worldwide. Number of species: World – 48; Palaearctic – 13; Russia – 1.

Delta unguiculatum (Villers, 1789) [*Vespa*] (*V. coangustata* Rossi, 1790; *Sphex cursor* Christ, 1791; *S. lapicida* Christ, 1791; *V. infundibuliformis* Olivier, 1791, nom. praeocc., nec Geoffroy, 1785; *V. dumetorum* Panzer, 1799; *Eumenes olivieri* Lepeletier de Saint Fargeau, 1841; *E. huberti* de Saussure,

1852; *E. aschabadensis* Radoszkowski, 1893; *E. unguiculus* (!) var. *sporadensis* von Schulthess, 1929; ?*E. tadzhicum* Gussakovskij, 1935; *E. unguiculatus* var. *libanicus* Giordani Soika, 1941; *E. unguiculatus* var. *mauritanicus* Giordani Soika, 1941). Russia: EP (NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Cyprus, Syria, Lebanon, Israel, Saudi Arabia, Iran, Central Asia.

DISCOELIUS Latreille, 1809 (*Discaelias* (!): Leach, 1815; *Discaelius* (!): Leach, 1815; *Dicoelius* (!): Haldy, 1836; *Discollius* (!): Froggatt, 1892; *Tritodiscoelius* Dalla Torre, 1904). Type species *Vespa zonalis* Panzer, 1801. Females nest in preexisting cavities such as beetle burrows in wood or hollow stems; partitions between cells are made of partially masticated leaf cuts; provision is caterpillars of various Microlepidoptera or rarely larvae of Pamphiliidae. The genus is distributed in the Palaearctic and Oriental regions. Number of species: World – 11; Palaearctic – 3, Russia – 2.

Discoelius dufouri Lepeletier de Saint Fargeau, 1841 (*D. manchurianus* Yasumatsu, 1934; *D. priesneri* Mader, 1936; *D. dufouri atripes* Giordani Soika, 1970; *D. planiventris* Giordani Soika, 1971). Russia: EP (NW, C, E, S, NC, CR), UR, WS (TM, TK, NS, AL), ES (BR, YA, ZB), FE (AM, KH, PR, SA). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Kazakhstan, Mongolia, China (NE, SE), Korean Peninsula.

Discoelius zonalis (Panzer, 1801) [Vespa] (*D. naryschkini* Morawitz, 1894; *D. japonicus* Pérez, 1905). Russia: EP (C, E, S, NC, CR), UR, WS (TM, NS, AL), ES (BR, ZB), FE (PR, KU). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Mongolia, China (NE, CC, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu).

EUMENES Latreille, 1802 (*Alpha* de Saussure, 1855, nec 1854; *Eumenis* (!): Kriechbaumer, 1879; *Eumenes* (!): Fox, 1899; *Eumes* (!): Bertoni, 1910; *Eumenidion* von Schulthess, 1913; *Eumenidium* (!): Sharp, 1915). Type species *Vespa coarctata* Linnaeus, 1758. Females build aerial nests consisted of one or several juglike mud cells sometimes covered with additional mud layer or rarely placed within pre-existing cavities; provision is usually caterpillars of Geometridae or various Microlepidoptera. The genus is distributed worldwide and includes two subgenera: *Eumenes* s.str.; *Zeteumenoides* Giordani

Soika, 1972. Number of species: World – 107; Palaearctic – 43; Russia – 23.

Eumenes (Eumenes) affinissimus de Saussure, 1852 (*E. coelestis* *montana* Kostylev, 1940; ?*E. solidus* Gusenleitner, 1972). Russia: WS (AL). – Pakistan, Tajikistan, Kyrgyzstan, Kazakhstan, Mongolia, China, SE Asia, India.

Eumenes (Eumenes) aquilonius Sk. Yamane, 1977 (*E. rubronotatus* *aquilonius* Sk. Yamane, 1977). Russia: FE (PR). – Korean Peninsula, Japan (Hok).

Eumenes (Eumenes) coarctatus (Linnaeus, 1758) [Vespa] (*E. lunulata* Fabricius, 1804; *E. pomiformis* *dernaensis* Blüthgen, 1938; *E. pomiformis* *limissicus* Blüthgen, 1938; *E. pomiformis* *ordubadensis* Blüthgen, 1938; *E. pomiformis* *barbatulus* Blüthgen, 1943; *E. pedunculata* var. *punctata* Hellén, 1944; *E. pedunculatus* var. *turaniformis* Blüthgen, 1959; *E. coarctatus* *corsicus* Gusenleitner, 1972; *E. coarctatus* *maroccanus* Gusenleitner, 1972; *E. lunulatus* var. *tenebricosus* Gusenleitner, 1972; *E. lunulatus* var. *balcanicus* Gusenleitner, 1972; *E. coarctatus* *nigaricus* Giordani Soika, 1986; *E. coarctatus* *dernensis* (!) Gusenleitner, 1999). Russia: EP (NW, C, E, S, NC, CR), UR, WS (TM, OM, NS, AL), ES (KR, IR, BR, YA, ZB), FE (AM, KH, PR, MG). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Cyprus, Syria, Jordan, Israel, Iran, Central Asia, Kazakhstan, Mongolia, China (NE, NC, NW, CC, SW), Japan (Hon, Shi, Kyu).

Eumenes (Eumenes) coronatus (Panzer, 1799) [Vespa] (*E. atricornis* Fabricius, 1804; *E. costata* (!): Luciani, 1845; *Eumenis* (!) *mediterranea* var. *neesi* Kriechbaumer, 1879; *Eumenes coarctatus* var. *opulenta* Blüthgen, 1938; *E. coarctatus* *detonsus* Blüthgen, 1943; *E. coarctatus* ab. *nigrotibia* Hellén, 1944; *E. coarctatus* var. *niger* Szulczewski, 1950; *E. coarctatus* *ibericus* Blüthgen, 1956; *E. coronatus* *corrugatus* Gusenleitner, 1972). Russia: EP (NW, C, E, NC, CR), UR, WS (AL), ES (IR, BR, ZB), FE (AM, KH, PR, KU). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Israel, Iran, Central Asia, Kazakhstan, Mongolia, China (NE, SW), Korean Peninsula, Japan (Kyu).

Eumenes (Eumenes) crimensis Blüthgen, 1938 (*E. dubius* *crimensis* Blüthgen, 1938; *E. sareptanus* *scabrosus* Gusenleitner, 1972). Russia: EP (CR). – Iran, Turkmenistan, Uzbekistan, Kazakhstan.

Eumenes (Eumenes) dubius de Saussure, 1852 (*E. dubius* *dubius* var. *palaestinensis* Blüthgen, 1938;

- E. dubius dubius* var. *macedonica* Blüthgen, 1952). Russia: EP (NC, CR). – Europe (WE, EE, SE), N Africa, Caucasus, Turkey, Cyprus, Syria, Jordan, Lebanon, Israel, Iraq, Iran, Turkmenistan, Tajikistan, Kazakhstan, S America (introduced).
- Eumenes (Eumenes) fraterculus** Dalla Torre, 1894 (*E. fraterna* Smith, 1873, nom. praeocc., nec Say, 1824). Russia: FE (AM, KH, PR). – China (NE, CC), Japan (Hon, Shi, Kyu, Ryu).
- Eumenes (Eumenes) glacialis** Giordani Soika, 1940. Russia: ES (IR). – Mongolia.
- Eumenes (Eumenes) kiangsuensis** Giordani Soika, 1941. Russia: FE (PR). – China (NE, CC, SE), Korean Peninsula.
- Eumenes (Eumenes) mediterraneus** Kriechbaumer, 1879 [*Eumenis*] (*E. affinissima* race *quettaensis* Cameron, 1907; *Labus superbus* Meade-Waldo, 1910; *E. mediterraneus bengasinus* Blüthgen, 1938; *E. mediterraneus cyprius* Blüthgen, 1938; *E. houskai* Giordani Soika, 1952; *E. mediterraneus anatolicus* Giordani Soika, 1952; *E. mediterraneus manchurianus* Giordani Soika, 1971; *E. mediterraneus* var. *opacus* Guseinleitner, 1972). Russia: EP (S, NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Cyprus, Syria, Jordan, Lebanon, Israel, Saudi Arabia, Yemen, UAE, Oman, Iraq, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NE, NC, NW, CC), Korean Peninsula, Tahiti (introduced).
- Eumenes (Eumenes) micado** Cameron, 1904 (*E. samuray* von Schulthess, 1908; *E. samurayi* (!) *rufescens* Giordani Soika, 1973). Russia: FE (KU). – ?China (NC), Japan (Hok, Hon, Shi, Kyu, Ryu).
- Eumenes (Eumenes) mongolicus** Morawitz, 1889 (*E. coarctatus* var. *mongolicus* Morawitz, 1889). Russia: WS (AL), ES (IR, BR, ZB). – Kyrgyzstan, Mongolia, China (NC, WP).
- Eumenes (Eumenes) papillarius** (Christ, 1791) [Sphex] (*E. bipunctis* de Saussure, 1852; *E. bimaculatus* André, 1884; *E. papillarius* var. *baltica* Blüthgen, 1938; *E. papillarius monticola* Blüthgen, 1956; *E. papillarius rubricornis* Guseinleitner, 1972). Russia: EP (C, E, S, NC, CR), UR. – Europe (WE, NE, SE, EE), Caucasus, Turkey, Syria, Lebanon, Israel, Iran, Pakistan, Central Asia, Kazakhstan, Mongolia.
- Eumenes (Eumenes) pedunculatus** (Panzer, 1799) [Vespa] (*E. eburneopictus* Giordani Soika, 1940; *E. pedunculatus turanus* Blüthgen, 1943; *E. pedunculatus* var. *lapponica* Hellén, 1944; *E. karafutonis* Sk. Yamane, 1977). Russia: EP (NW, C, E), UR, WS (OM, TK, NS, AL), ES (KR, IR, BR, YA, ZB), FE (PR, SA). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan, Mongolia, China (SW).
- Eumenes (Eumenes) pomiformis** (Fabricius, 1781) [Vespa] (*Eumenis* (!) *mediterranea* var. *heri* Kriechbaumer, 1879; *Eumenes fastidiosissimus* Giordani Soika, 1943; *E. pomiformis turcicus* Giordani Soika, 1952). Russia: EP (S, NC, CR), UR. – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Lebanon, UAE, Pakistan, Central Asia, Kazakhstan, China (SW), Korean Peninsula, India.
- Eumenes (Eumenes) punctaticeps** Giordani Soika, 1943 (*E. robustus* Kostylev, 1940, nom. praeocc., nec Isely, 1917; *E. calabricus* Giordani Soika, 1943; *E. kostylevi* Kurzenko, 1976). Russia: EP (CR), WS (AL). – Europe (WE, SE, EE), Turkey, Iran, Central Asia, Kazakhstan.
- Eumenes (Eumenes) punctatus** de Saussure, 1852 (*E. formosensis nigrior* Giordani Soika, 1973; *E. asioboreus* Kim et Sk. Yamane, 2001). Russia: FE (KH, PR). – Pakistan, Mongolia, China (NE, NC, CC, SW, SE), Korean Peninsula, Japan (Tsushima), India, Sri Lanka.
- Eumenes (Eumenes) rubronotatus** Pérez, 1905 (*E. dimidiaticeps* Giordani Soika, 1973). Russia: FE (PR). – China (NE, CC, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu), SE Asia.
- Eumenes (Eumenes) sareptanus** André, 1884 (*E. pomiformis* f. *insolata* Müller, 1923; *E. dubius sareptanus* var. *germanica* Blüthgen, 1938). Russia: EP (E, S, NC, CR), UR, WS (OM, NS, AL). – Europe (WE, SE, EE), Caucasus, Turkey, Central Asia, Kazakhstan.
- Eumenes (Eumenes) septentrionalis** Giordani Soika, 1940 (*E. septentrionalis khangmarenensis* Giordani Soika, 1966; *E. glacialis*: Kurzenko, 1995). Russia: ES (IR, BR, ZB), FE (AM, KH, PR). – Mongolia, China (NE, WP).
- Eumenes (Eumenes) subpomiformis** Blüthgen, 1938 (*E. subpomiformis crassipunctatus* Blüthgen, 1956). Russia: EP (NC). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Lebanon, Israel, Central Asia, Kazakhstan.
- Eumenes (Eumenes) transbaicalicus** Kurzenko, 1984. Russia: ES (BR). – Mongolia, China (NE).
- Eumenes (Eumenes) tripunctatus** (Christ, 1791) [Sphex] (*E. venusta* Fischer-Waldheim, 1843). Russia: EP (E, S, NC, CR), UR. – Europe (EE), Iran, Turkmenistan, Kazakhstan, Mongolia, China (NC, NW, WP).

EUODYNERUS Dalla Torre, 1904. Type species

Vespa dantici Rossi, 1790. Females nest mainly in various preexisting cavities: hollow stems, beetle burrows in wood and abandoned nests of other wasps and bees, partitions between cells are made of mud; several Nearctic species excavate nest burrows in ground and surmount them by turrets; provision is various caterpillars (mainly of Microlepidoptera) or rarely larvae of Chrysomelidae. The genus is distributed worldwide and includes three subgenera: *Euodynerus* s.str.; *Incolelipona* Giordani Soika, 1995; *Pareuodynerus* Blüthgen, 1938. Number of species: World – 104; Palaearctic – 44; Russia – 9.

Euodynerus (Euodynerus) caspicus (Morawitz, 1873) [Leionotus] (*Lionotus cardinalis* Morawitz, 1885; *E. caspicus astrachanensis* Blüthgen, 1942). Russia: **EP** (S). – Armenia, Azerbaijan, Iran, Turkmenistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NC, NW).

Euodynerus (Euodynerus) dantici (Rossi, 1790) [*Vespa*] (*Odynerus postscutellatus* Lepeletier de Saint Fargeau, 1814; *O. tinctus* Walker, 1871; *O. dantici brachytomus* Kostylev, 1940; *E. dantici iberogalicus* Blüthgen, 1942; *E. dantici pamiricus* Blüthgen, 1942; *O. dantici* var. *lagostae* Giordani Soika, 1942; *E. espagnoli* Vergés, 1965; *O. dantici violaceipennis* Giordani Soika, 1973; *E. dantici nigrescens* Gusenleitner, 1979; *O. dantici poggii* Giordani Soika, 1986). Russia: **EP** (S, NC, CR), **UR**, **ES** (BR, ZB), **FE** (PR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Syria, Jordan, Lebanon, ?Israel, Iran, Afghanistan, Central Asia, Kazakhstan, Mongolia, China (NE, NC, NW, CC, SE), Korean Peninsula, Japan (Hon, Shi, Kyu, Ryu), SE Asia.

Euodynerus (Euodynerus) disconotatus (Lichtenstein, 1884) [*Odynerus*] (*Lionotus sulfuripes* Morawitz, 1885; *Odynerus kokpeticus* Radoszkowski, 1886; *O. collariventris* Giordani Soika, 1942; *E. disconotatus albidus* Blüthgen, 1951; *E. disconotatus laniensis* Giordani Soika, 1979). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), Caucasus, Turkey, Cyprus, Jordan, Saudi Arabia, Iraq, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan.

Euodynerus (Euodynerus) fastidiosus (de Saussure, 1853) [*Odynerus*] (*Odynerus germabicus* Radoszkowski, 1893; *O. garmabicus* (!): Dalla Torre, 1904). Russia: **EP** (S, NC, CR). – Europe (WE, SE), N Africa, Caucasus, Turkey, Syria, Lebanon, Oman, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, China (NC).

Euodynerus (Euodynerus) velutinus Blüthgen, 1951

(*Odynerus curictensis*: Kostylev, 1940). Russia: **EP** (CR). – Europe (SE: Greece), Caucasus, Turkey.

Euodynerus (Pareuodynerus) nippicus (von Schultess, 1908) [Lionotus] (*Lionotus tomentosus* var. *nippicus* von Schultess, 1908; *?Rhynchium seuli* Radoszkowski, 1890; *Odynerus micado* (!): Matsu-mura, 1911; *O. flavidypeatus* Sonan, 1930; *Rh. satumanus* Sonan, 1930; *Euodynerus notatus tonkinensis* Giordani Soika, 1973; *E. nippicus* (!): Giordani Soika, 1982; *E. notatus flavigornis* Sk. Yamane, 1987; *E. notatus ryukyuensis* Tano, 1987; *E. notatus*: Kurzenko, 1995, 2012). Russia: **ES** (ZB), **FE** (AM, KH, PR, SA, MG). – ?Mongolia, China (NE, NC, CC, SW, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu), SE Asia.

Euodynerus (Pareuodynerus) notatus (Jurine, 1807) [*Vespa*] (*?Odynerus bissstrigatus* Spinola, 1808; *O. nigripes* Herrich-Schäffer, 1839; *?O. maculatus* Lepeletier de Saint Fargeau, 1841; *O. clypearis* Thomson, 1870; *O. pubescens* Thomson, 1870; *O. unguaris* Thomson, 1870; *?O. pubescens* var. *cupreus* von Schultess, 1897; *Euodynerus notatus* var. *pernotata* Blüthgen, 1938; *E. notatus cyrenicus* Giordani Soika, 1986). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (IR, BR, YA, ZB), **?FE** (AM, KH, PR, SA, MG). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Iran, Kazakhstan, Mongolia, ?China (NC).

Euodynerus (Pareuodynerus) posticus (Herrich-Schäffer, 1841) [*Odynerus*] (*Odynerus innumeralis* de Saussure, 1853; *O. graphicus* de Saussure, 1853; *O. differens* Morawitz, 1895; *Euodynerus cephalicus* Blüthgen, 1944; *Pseudepipona postica punctatissima* Giordani Soika, 1952; *E. posticus posticus* var. *notatiformis* Blüthgen, 1955; *E. posticus punctatissimus* var. *nahariensis* Blüthgen, 1955). Russia: **EP** (S, NC, CR). – Europe (WE, EE, SE), N Africa, Caucasus, Turkey, Israel, Iran.

Euodynerus (Pareuodynerus) quadrifasciatus (Fabricius, 1793) [*Vespa*] (*Vespa quadricincta* Fabricius, 1793; *V. simplex* Fabricius, 1793; *Odynerus lindenii* Lepeletier de Saint Fargeau, 1841; *O. tomentosus* Thomson, 1870; *Pseudepipona sachalinensis* Yasumatsu, 1938; *Euodynerus quadrifasciatus* var. *pseudonotata* Blüthgen, 1939; *E. quadrifasciatus atripes* Giordani Soika, 1976; *E. quadrifasciatus rubrosignatus* Gusenleitner, 1984; *E. quadrifasciatus rufipes* Gusenleitner, 1984; *E. quadrifasciatus eburnus* Sk. Yamane, 1987). Russia: **EP** (N, NW, C,

E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, AL), **ES** (KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, MG). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Kazakhstan, Mongolia, China (NE), Korean Peninsula, Japan (Hok, Hon).

EUSTENANCISTROKERUS Blüthgen, 1938. Type species *Odynerus blanchardianus* de Saussure, 1855. Females nest in ground and excavate burrows surmounted by turrets; provision is caterpillars of Coleophoridae. The genus is distributed in the Palaearctic and Afrotropical regions and includes three subgenera: *Eustenancistrocerus* s.str.; *Hemistenancistrocerus* Blüthgen, 1938; *Parastenancistrocerus* Blüthgen, 1938. Number of species: World – 20; Palaearctic – 17; Russia – 2.

Eustenancistrocerus (Eustenancistrocerus) tegularis (Morawitz, 1885) [*Lionotus*]. Russia: **EP** (NC). – ?Europe (SE: Greece), Caucasus, ?Iran, Central Asia, Kazakhstan.

Eustenancistrocerus (Parastenancistrocerus) amadanensis (de Saussure, 1855) [*Odynerus*] (*Odynerus turca* de Saussure, 1863; *O. transitorius* Morawitz, 1867; *O. lobatus* André, 1883; *Ancistrocerus iranensis* Giordani Soika, 1943; *Eustenancistrocerus transitorius hoberlandti* Giordani Soika, 1952; *E. transitorius mauritanensis* Giordani Soika, 1953; *E. lobatus flavellus* Giordani Soika, 1952). Russia: **EP** (S, NC, CR). – Europe (EE, SE), N Africa, Caucasus, Turkey, Cyprus, Syria, Israel, Saudi Arabia, Iraq, Iran, Pakistan, Central Asia, Kazakhstan.

GYMNOMERUS Blüthgen, 1938. Type species *Odynerus laevipes* Shuckard, 1837. Females nest in hollow stems where they build a compact row of cylindrical mud cells without plugs, the nest is stopped by an empty final cell filled with particles of soil and various plant material; provision is larvae of *Hypera* Germar (Curculionidae). The monotypic Palaearctic genus.

Gymnomerus laevipes (Shuckard, 1837) [*Odynerus*] (*Odynerus cognatus* Dufour, 1839; *O. rubicola* Dufour, 1839; *Pterocheilus simplicipes* Herrich-Schäffer, 1839; *O. scandinavus* de Saussure, 1856; *O. sibiricus* Mocsáry, 1883; *O. levipes* (!): Dalla Torre, 1894; *O. algeriensis* Dusmet, 1928). Russia: **EP** (NW, C, E, ?S, NC, CR), **UR**, **WS** (TM, NS, AL), **ES** (BR, YA), **FE** (AM, PR, SA). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Iran, Kazakhstan, Mongolia.

HEMIPTEROCHILUS Ferton, 1909 (*Pseudopterochilus* van der Vecht, 1972). Type species *Odynerus terricola* Mocsáry, 1883. Females nest in ground and excavate vertical burrows surmounted by a small collar; provision is caterpillars of Noctuidae and Pyraustidae. The genus is distributed in the Palaearctic. Number of species: Palaearctic – 12; Russia – 1.

Hemipterochilus bembeciformis (Morawitz, 1867) [*Odynerus*] (*Odynerus terricola* Mocsáry, 1883; *O. sinuatofasciatus* Costa, 1888; *Hoplomerus tinniens*: Bialynicki-Birula, 1926; *H. tinniens* var. *rubromaculatus* Bialynicki-Birula, 1926; *Hoplomerus tinniens turcestanicus* Bialynicki-Birula, 1926; *Pterocheilus bembeciformis* var. *circensis* Giordani Soika, 1942). Russia: **EP** (E, S, NC, CR), **UR**. – Europe (WE, SE, EE), Caucasus, Turkey, Central Asia, Kazakhstan.

JUCANCISTROKERUS Blüthgen, 1938 (*Iucancistrocerus* (!): Blüthgen, 1951). Type species *Odynerus jucundus* Mocsáry, 1883. Females nest in vertical clay cliffs and excavate burrows surmounted by turrets; provision is larvae of Curculionidae. The genus is distributed in the Palaearctic and includes two subgenera: *Eremodynerus* Blüthgen, 1939; *Jucancistrocerus* s.str. Number of species: Palaearctic – 13; Russia – 2.

Jucancistrocerus (Jucancistrocerus) caspicus Giordani Soika, 1970. Russia: **EP** (S, NC). – Europe (EE: Ukraine), Caucasus, Turkey, Iran, Kazakhstan.

Jucancistrocerus (Jucancistrocerus) minutepunctatus Giordani Soika, 1970. Russia: **EP** (S). – Armenia, Azerbaijan.

KATAMENES Meade-Waldo, 1910. Type species *Katamenes watsoni* Meade-Waldo, 1910. Females build aerial nests attached to stones and consisted of one or several juglike cells build of pebbles confirmed and covered with mud, the nest is usually covered by additional layer of pebbles confirmed with mud; provision is caterpillars of Noctuidae, Pieridae, Lyceanidae and Pterophoridae. The genus is distributed in the Palaearctic, Afrotropical, and Oriental regions. Number of species: World – 14; Palaearctic – 13; Russia – 5.

Katamenes arbustorum (Panzer, 1799) [*Vespa*] (*Eumenes amedei* Lepeletier de Saint Fargeau, 1841; *E. frivaldszkyi* Herrich-Schäffer, 1841; *E. tabida* Eversmann, 1854; *E. arbustorum* var. *burlinii* Giordani

Soika, 1949; *K. arbustorum soikai* Borsato, 1993). Russia: EP (E), UR. – Europe (WE, SE, EE), ?Kazakhstan.

Katamenes dimidiatus (Brullé, 1832) [Eumenes] (?*E. fulva* var. β Eversmann, 1854; *E. montana* Nurse, 1904; *K. watsoni* Meade-Waldo, 1910; *E. arbustorum*: Kostylev, 1928, 1929; ?*K. sesquicinctus radoszkovskii* Blüthgen, 1962; *K. dimidiatus arrestus* Giordani Soika, 1970). Russia: EP (S, CR), UR. – Europe (SE, EE), Caucasus, Turkey, Cyprus, Syria, Israel, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NC).

Katamenes flavigularis (Blüthgen, 1951) [Eumenes] (*E. arbustorum flavigularis* Blüthgen, 1951; *E. arbustorum* var. *tabidus*: Kostylev, 1928; *K. arbustorum*: Tobias, Kurzenko, 1978). Russia: EP (NC, CR). – Europe (SE, EE), ?Caucasus, Turkey, Syria, Lebanon, Iran, Afghanistan, Tajikistan.

Katamenes sichelii (de Saussure, 1852) [Eumenes] (?*Eumenes fluctuans* de Saussure, 1852; *E. fulva* Eversmann, 1854; *E. baerii* Radoszkowski, 1865; *E. sichelii* var. *biblicus* Giordani Soika, 1935; *E. sichelii* var. *fulvooides* Giordani Soika, 1949; *E. sichelii* var. *luristanensis* Giordani Soika, 1949; *E. sichelii* *tauriae* Giordani Soika, 1960; *K. sichelii hispanicus* Giordani Soika, 1966; *K. sichelii pseudofulvus* Giordani Soika, 1966; *K. sichelii coranicus* Giordani Soika, 1970; *K. sichelii timanensis* Giordani Soika, 1986). Russia: EP (S), UR. – Europe (SE), N Africa, Caucasus, Turkey, Jordan, Israel, Saudi Arabia, UAE, Iraq, Iran, Afghanistan, Central Asia, Kazakhstan, Mongolia, China (WP), India.

Katamenes tauricus (de Saussure, 1855) [Eumenes] (*E. latipes* Sickmann, 1894; *E. arbustorum* var. *baicalensis* Kokujev, 1927; *E. latipes* var. *viratus* Giordani Soika, 1949; *E. sesquicinctus viratus* var. *caspica* Blüthgen, 1952). Russia: EP (CR), WS (AL), ES (TU, IR, BR). – Iran, Afghanistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NC), India.

LEPTOCHILUS de Saussure, 1853. Type species *Pterochilus mauritanicus* Lepeletier de Saint Fargeau, 1841. Females nest in various preexisting cavities: hollow stems, beetle burrows in wood or empty snail shells; partitions between cells are made of sand or gravel, several species use also macerated pith in addition to mineral material or build partitions of shell bits and sand confirmed with mud or resin; provision is caterpillars of various Microlepidoptera or beetle larvae (usually Curculionidae

or Chrysomelidae). The genus is distributed worldwide except Australia and includes six subgenera: *Euleptochilus* Blüthgen, 1943; *Leptochilus* s.str.; *Lionotulus* Blüthgen, 1938; *Neoleptochilus* Blüthgen, 1961; *Sarochilus* Guseinleitner, 1970; *Zendalia* Robertson, 1928. Number of species: World – 197; Palaearctic – 137; Russia – 6.

Leptochilus (Euleptochilus) limbiferus (Morawitz, 1867) [Odynerus] (*L. limbiferus anatolicus* Blüthgen, 1955; *L. limbiferus achaeus* Guseinleitner, 1970). Russia: EP (NC). – Europe (SE, EE), Armenia, Turkey.

Leptochilus (Lionotulus) alpestris (de Saussure, 1855) [Odynerus] (*O. trinacriae* André, 1883; *Leptochilus alpestris* var. *grandii* Giordani Soika, 1947; *L. alpestris alpestris* var. *bulsanensis* Blüthgen, 1953; *L. alpestris iberobarbarus* Blüthgen, 1953; *L. alpestris iberobarbarus* var. *rufina* Blüthgen, 1953; *L. guichardi* Giordani Soika, 1973; *L. astrictus* Giordani Soika, 1986). Russia: EP (S, NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Jordan, Turkmenistan.

Leptochilus (Lionotulus) ergenicus (Kostylev, 1940) [Odynerus]. Russia: EP (S). – Kazakhstan.

Leptochilus (Lionotulus) membranaceus (Morawitz, 1867) [Odynerus] (*L. membranaceus luxuriosus* Blüthgen, 1955). Russia: EP (E, S, NC, CR). – Europe (SE, EE), Caucasus, Turkey, Israel, Iran, Turkmenistan, Uzbekistan, Kazakhstan.

Leptochilus (Lionotulus) signatus Guseinleitner, 1995 (*L. signatus gueruensis* Guseinleitner, 1995). Russia: EP (NC). – Turkey.

Leptochilus (Neoleptochilus) regulus (de Saussure, 1855) [Odynerus] (*O. gallicus* de Saussure, 1855; ?*O. calabricus* Gribodo, 1881; *O. siculus* De Stefani, 1883; *O. costae* Costa, 1884; *O. buyssoni* Medina, 1891; ?*O. pauli* Medina, 1892; *O. perforatus* Dusmet, 1903; *O. obtusidentatus* Maidl, 1922; *Microdynerus wuczeticzi* Kostylev, 1928). Russia: EP (NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Cyprus, Syria, Jordan, Iran.

MICRODYNERUS Thomson, 1874. Type species *Odynerus exilis* Herrich-Schäffer, 1839. Females of the nominotypical subgenus nest in preexisting cavities (hollow stems or beetle burrows in wood), partitions between cells are made of mud; species of the subgenus *Pseudomicrodynerus* excavate burrows in pith of plant stems and branches; nesting of the subgenus *Alastorynerus* is unknown;

provision is larvae of Curculionidae or caterpillars of Microlepidoptera. The genus is distributed in the Holarctic and includes three subgenera: *Alastorynerus* Blüthgen, 1938; *Microdynerus* s.str.; *Pseudomicrodynerus* Blüthgen, 1938. Number of species: World – 53; Palaearctic – 37; Russia – 7.

Microdynerus (Alastorynerus) microdynerus (Dalla Torre, 1889) [Odynerus] (*M. alastoroides* Morawitz, 1885, nom. praeocc., nec de Saussure, 1852; *M. tridentatus* Kostylev, 1935). Russia: EP (S, CR). – Europe (SE, EE), Caucasus, Turkey, Jordan, Iran, Tajikistan, Kazakhstan.

Microdynerus (Microdynerus) exilis (Herrich-Schäffer, 1839) [Odynerus] (*O. bivittatus* Lepetier de Saint Fargeau, 1841). Russia: EP (E, ?S, NC). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey.

Microdynerus (Microdynerus) longicollis Morawitz, 1895 (*Odynerus patagiatus* von Schulthess, 1897; *Microdynerus patagiatus sicanius* Blüthgen, 1964). Russia: EP (S, NC, CR). – Europe (WE, SE, EE), Turkey, Turkmenistan.

Microdynerus (Microdynerus) mirandus (Giordani Soika, 1947) [Leptochilus]. Russia: EP (CR). – Europe (SE, EE), Turkey.

Microdynerus (Microdynerus) nugdunensis (de Saussure, 1855) [Odynerus] (*M. nugdunensis sicelis* Blüthgen, 1964; *Leptochilus micans* Giordani Soika, 1986). Russia: EP (E, NC, CR). – Europe (WE, SE, EE), Azerbaijan, Turkey.

Microdynerus (Microdynerus) timidus (de Saussure, 1856) [Odynerus] (*Vespa minuta* Fahricius, 1798, nom. praeocc., nec Linnaeus, 1767; ?*Odynerus rus-sky* Dusmet, 1917; *Microdynerus timidus adustus* Guseinleitner, 1977). Russia: EP (CR). – Europe (WE, NE, SE, EE), N Africa.

Microdynerus (Pseudomicrodynerus) parvulus (Herrich-Schäffer, 1838) [Odynerus] (*O. helvetius* de Saussure, 1855; *Microdynerus bifidus* Morawitz, 1885). Russia: EP (C, E, S, NC, CR), UR, WS (AL). – Europe (WE, SE, EE), Caucasus, Turkey, Iran, Uzbekistan, Kazakhstan.

ODYNERUS Latreille, 1802 (*Odynera* (!): Illiger, 1807; *Epipone* Kirby et Spence, 1815, nec Latreille, 1804; *Oplopus* Wesmael, 1836, nec Laporte, 1832; *Oplo-merus* Westwood, 1840, nec Dejean, 1833; *Hoplo-merus* (!): Agassiz, 1846; *Hoplopus* (!): Agassiz, 1846; *Epiponus* (!): de Saussure, 1875; *Hoplonus* (!): Dalla Torre, 1889; *Euepipona* Dalla Torre, 1904;

Odyneurus (!): Wolcott, 1924). Type species *Vespa spinipes* Linnaeus, 1758. Females of most species nest in ground (on both vertical and horizontal surfaces) and excavate burrows (containing several cells) surmounted by turrets; provision is larvae of *Hypera* Germar (Curculionidae) or rarely larvae of Tenthredinidae (in subgenus *Monoplomerus*). Some Nearctic species nest in preexisting holes in ground and use pressed soil to make partitions between cells (they do not construct turrets and do not use water during the building works); their provision is caterpillars of Gelechiidae. The genus is distributed in the Holarctic and Afrotropical regions and includes three subgenera: *Monoplomerus* Blüthgen, 1941; *Odynerus* s.str.; *Spinicoxa* Blüthgen, 1938. Number of species: World – 45; Palaearctic – 39; Russia – 12.

Odynerus (Odynerus) alpinus von Schulthess, 1897 (*O. spinipes* var. *alpinus* von Schulthess, 1897). Russia: EP (NC), WS (AL), ES (KR, IR, BR, YA, ZB), FE (KH, MG). – Europe (WE, SE, EE), Turkey, Tajikistan, Mongolia.

Odynerus (Odynerus) femoratus de Saussure, 1856 (*O. duskei* Morawitz, 1895). Russia: EP (S, NC, CR), ES (IR, BR). – Europe (WE, SE, EE), Caucasus, Turkey, Syria, Israel, Iran.

Odynerus (Odynerus) melanocephalus (Gmelin, 1790) [Vespa] (*Pterocheilus dentipes* Herrich-Schäffer, 1839; *P. serripes* Taschenberg, 1866; *O. sareptanus* André, 1884; *Hoplomerus armeniacus* Morawitz, 1885; *H. melanocephalus* var. *combinata* Blüthgen, 1940; *H. armeniacus* var. *plurinotata* Blüthgen, 1941; *H. melanocephalus tadschicus* Blüthgen, 1941; *H. melanocephalus melanocephalus* var. *luxuriosa* Blüthgen, 1955; *O. melanocephalus robustus* Eck, 1971; *O. melanocephalus cypriacus* Eck, 1971; *O. melanocephalus latus* Eck, 1971). Russia: EP (C, E, S, NC, CR), UR, WS (NS, KM, AL). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Cyprus, Syria, Jordan, Israel, Iraq, Iran, Afghanistan, Central Asia, Kazakhstan.

Odynerus (Odynerus) poecilus de Saussure, 1856 (*O. ruficornis* Rudow, 1876). Russia: EP (NC). – Europe (WE, SE, EE), Azerbaijan, Turkey.

Odynerus (Odynerus) serricus (Blüthgen, 1961) [Oplomerus] (*Odynerus serripes* Morawitz, 1867, nom. praeocc., nec *Pterocheilus serripes* Taschenberg, 1866). Russia: EP (C, E, S), ?WS (OM, TK, AL). – ?Europe (EE), Turkey, Kyrgyzstan, Kazakhstan.

Odynerus (Odynerus) spinipes (Linnaeus, 1758)
[*Vespa*] (*V. quinquefasciata* Fabricius, 1793; *Odynerus muticus* Zetterstedt, 1838; *O. spinipes* ab. *flaviscapus* Mader, 1936; *Hoplomerus spinipes* var. *scutellaris* Blüthgen, 1940; *H. spinipes amurensis* Blüthgen, 1941). Russia: **EP** (NW, C, E, NC), **UR**, **WS** (TK, AL), **ES** (IR, BR), **FE** (KH, PR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Tajikistan, Kazakhstan, Mongolia.

Odynerus (Odynerus) tristis (Blüthgen, 1939) [*Hoplomerus*] (*O. tristis tianschanicus* Kurzenko, 1977). Russia: **EP** (NC). – Tajikistan, Kyrgyzstan, Kazakhstan, China (NW).

Odynerus (Spinicoxa) albopictus de Saussure, 1856
(*Hoplomerus calcaratus* Morawitz, 1885; *H. rufospinosus* Morawitz, 1895; *Odynerus rufospinosus* var. *argivus* von Schultess, 1924; *O. albopictus kazakhstanicus* Kurzenko, 1977). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), ?N Africa, Caucasus, Turkey, Syria, Jordan, Israel, Iraq, Iran, Turkmenistan, Uzbekistan, Kazakhstan.

Odynerus (Spinicoxa) cuneiformis Kostylev, 1940.
Russia: **ES** (TU, BR), **FE** (AM, KH). – Mongolia.

Odynerus (Spinicoxa) fulvitarsis (Morawitz, 1895)
[*Hoplomerus*]. Russia: **EP** (S, NC). – Caucasus, Turkey, Turkmenistan, Uzbekistan, Kazakhstan.

Odynerus (Spinicoxa) reniformis (Gmelin, 1790)
[*Vespa*] (?*Vespa sexfasciata* Fabricius, 1781; *V. melanochra* Gmelin, 1790; *Pterocheilus coxalis* Herrich-Schäffer, 1839; *Odynerus reaumurii* Dufour, 1839; *O. dufouri* Lepeletier de Saint Fargeau, 1841; *O. velox* de Saussure, 1853; ?*O. depressus* André, 1883; *Oplomerus reniformis balcanicus* Blüthgen, 1960). Russia: **EP** (NW, C, E), **UR**, **WS** (TK, AL), **ES** (TU, BR, YA, ZB). – Europe (WE, NE, SE, EE), N Africa, Turkey, Israel, Kazakhstan.

Odynerus (Spinicoxa) simillimus Morawitz, 1867
(*O. dentifer* Drogoszewski, 1934). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (NS, AL), **ES** (KR, IR, YA). – Europe (WE, EE), Caucasus, Turkey, Israel, Turkmenistan, Kazakhstan, Mongolia.

ONYCHOPTEROCHEILUS Blüthgen, 1955. Type species *Odynerus daw* Dusmet, 1903. Females excavate nest burrows in friable soil without using water; provision is caterpillars. The genus is distributed in the Palaearctic and includes five subgenera: *Acutopterocheilus* Kurzenko, 1988; *Asiapterocheilus* Kurzenko, 1988; *Ghilardocheilus* Kurzenko, 1988; *Neopterocheilus* Kurzenko, 1988;

Onychopterocheilus s.str. Number of species: Palaearctic – 60; Russia – 4.

Onychopterocheilus (Acutopterocheilus) sareptanus (Kurzenko, 1976) [*Pterocheilus*]. Russia: **EP** (S).

Onychopterocheilus (Onychopterocheilus) crabroniformis (Morawitz, 1867) [*Odynerus*] (*Pterocheilus albopictus*: Morawitz, 1895, Kostylev, 1940). Russia: **EP** (S), **UR**, **WS** (AL), **ES** (KR). – Kazakhstan.

Onychopterocheilus (Onychopterocheilus) eckloni (Morawitz, 1885) [*Pterocheilus*]. Russia: **ES** (TU, IR, BR, ZB). – Mongolia, China (NC).

Onychopterocheilus (Onychopterocheilus) pallasii (Klug, 1805) [*Pterocheilus*] (*P. latipalpis* Lepeletier de Saint Fargeau, 1841). Russia: **EP** (E, S, NC, CR), **UR**, **WS** (OM, NS, AL). – Europe (EE), Turkey, Kazakhstan, Mongolia.

PARAGYMNOMERUS Blüthgen, 1938. Type species *Odynerus spiricornis* Spinola, 1808. Females nest in ground (on both vertical and horizontal surfaces) and excavate burrows contained several cells and surmounted by turrets; provision is larvae of Pamphiloidea. The genus is distributed in the Palaearctic. Number of species: Palaearctic – 7; Russia – 1.

Paragymnomerus signaticollis (Morawitz, 1888)
[*Hoplomerus*] (*H. semenowi* Morawitz, 1895; *H. signaticollis semenovi* (!): Bialynicki-Birula, 1926; *H. signaticollis* var. *lutescens* Bialynicki-Birula, 1926; *Odynerus signaticollis tauricus* Kostylev, 1940). Russia: **EP** (CR). – Europe (EE), Azerbaijan, Turkey, Turkmenistan, Tajikistan, Uzbekistan.

PARARRHYNCHIUM de Saussure, 1855 (*Prorrhynchium* de Saussure, 1855; *Prorrhynchium* (!): de Saussure, 1856; *Pararrhynchium* (!): de Saussure, 1862; *Parrhynchium* (!): Dalla Torre, 1894). Type species *Rhynchium ornatum* Smith, 1852. Females nest in preexisting cavities such are hollow stems, partitions between cells are made of mud; provision is caterpillars. The genus is distributed in the Palaearctic and Oriental regions. Number of species: World – 11; Palaearctic – 4; Russia – 1.

Pararrhynchium paradoxum (Gussakovskij, 1932)
[*Odynerus*] (?*Pararrhynchium* (!) *paradoxum laetum* Giordani Soika, 1986). Russia: **FE** (PR). – China (CC, SE), Korean Peninsula, ?India.

PARAVESPA Radoszkowski, 1886 (*Theletor* Kokujev, 1913). Type species *Paravespa komarowii*

Radoszkowski, 1886. Females nest in ground and excavate vertical burrows surmounted by turrets; provision is caterpillars of Noctuidae. The genus is distributed in the Palaearctic and Afrotropical regions and includes two subgenera: *Gestrodynerus* Giordani Soika, 1961; *Paravespa* s.str. Number of species: World – 17; Palaearctic – 4; Russia – 1.

Paravespa (Paravespa) rex (von Schulthess, 1924) [Odynerus] (*O. rex* var. *clarior* von Schulthess, 1924; *O. rex* var. *obscurior* von Schulthess, 1924; *Hoplomerus jakobsoni* Bialynicki-Birula, 1926; *P. regina* (!): Blüthgen, 1939; *P. rex* var. *rufina* Blüthgen, 1955; *P. rex* var. *submimetica* Blüthgen, 1955). Russia: EP (CR). – Turkey, Iran, Uzbekistan, Kyrgyzstan, Kazakhstan.

PARODONTODYNERUS Blüthgen, 1938 (*Parodontodynerus* (!): Blüthgen, 1961). Type species *Eumenes ephippium* Klug, 1817. Females build nests consisted of several barrel-shaped mud cells often placed within preexisting cavities such as hollow stems of caverns in stones; provision is beetle larvae (Coleoptera). The genus is distributed in the Palaearctic. Number of species: Palaearctic – 5; Russia – 1.

Parodontodynerus ephippium (Klug, 1817) [Eumenes] (?*Pterocheilus laetus* Herrich-Schäffer, 1839; *Odynerus dubius* de Saussure, 1852; *O. opacus* Morawitz, 1867; *O. schulthessi* Dusmet, 1903; *Odon-todynerus ephippium anatoliae* Giordani Soika, 1951; *Parodontodynerus ephippium dalanicus* Kurzenko, 1976). Russia: EP (S, NC, CR). – Europe (WE, SE, EE), Caucasus, Turkey, Lebanon, Iran, Turkmenistan, Uzbekistan, Kazakhstan.

PSEUDEPIPONA de Saussure, 1856 (*Leptepipona* Blüthgen, 1951; *Metepipona* Blüthgen, 1951; *Trichepipona* Blüthgen, 1951; *Pseudopipona* (!): Giordani Soika, 1952; *Pseudepipone* (!): Bytinski-Salz, Guseinleitner, 1971). Type species *Odynerus herrichii* de Saussure, 1856. Females excavate nests in ground, on horizontal surfaces; provision is caterpillars of Microlepidoptera (mainly Gelechiidae). The genus is distributed in the Holarctic and Afrotropical regions and includes two subgenera: *Deuterepipona* Blüthgen, 1951; *Pseudepipona* s.str. Number of species: World – 34; Palaearctic – 32; Russia – 10.

Pseudepipona (Deuterepipona) niveopicta Giordani Soika, 1970. Russia: EP (NC). – Turkey.

Pseudepipona (Deuterepipona) superba (Morawitz, 1867) [Odynerus] (*O. hyalinipennis* André, 1884). Russia: EP (S, CR). – ?Azerbaijan.

Pseudepipona (Deuterepipona) tricolor Guseinleitner, 1976. Russia: EP (NC). – Iran, Kazakhstan.

Pseudepipona (Pseudepipona) augusta (Morawitz, 1867) [Odynerus] (*O. aurantiacus* Mocsáry, 1877). Russia: EP (?E, S, NC, CR), UR, WS (AL), ES (KR). – Europe (EE), Caucasus, Turkey, Kazakhstan, Mongolia, China (NC).

Pseudepipona (Pseudepipona) beckeri (Morawitz, 1867) [Odynerus] (*O. falsus*: Kostylev, 1928; *Pseudepipona lativentris*: Tobias, Kurzenko, 1978). Russia: EP (?E, S, CR). – Kazakhstan.

Pseudepipona (Pseudepipona) herrichii (de Saussure, 1856) [Odynerus] (*O. variegatus* Herrich-Schäffer, 1839, nom. praeocc., nec Fabricius, 1793; *O. basalis* Smith, 1857; *O. aldrichi* Fox, 1892; *Pseudepipona herrichii* var. *derufata* Blüthgen, 1951; *P. variegata mongolica* Giordani Soika, 1970; *P. herrichii afromontana* Guseinleitner, 1977). Russia: EP (E, S, NC, CR), UR, WS (OM, NS, AL), ES (KR, IR, BR, YA, ZB), FE (AM, PR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Syria, Central Asia, Kazakhstan, Mongolia, China (NE, NC, SW), N America.

Pseudepipona (Pseudepipona) peculiaris (Morawitz, 1895) [Odynerus]. Russia: EP (S). – Iran, Kazakhstan.

Pseudepipona (Pseudepipona) rubiginosa (André, 1884) [Odynerus]. Russia: EP (S). – ?Georgia.

Pseudepipona (Pseudepipona) sellata (Morawitz, 1885) [*Lionotus*]. Russia: EP (S, NC). – Azerbaijan, Turkey, Iran, Turkmenistan, Kazakhstan.

Pseudepipona (Pseudepipona) tricarinata (Kokujev, 1912) [Odynerus] (*O. herrichii* var. *tricarinatus* Kokujev, 1912). Russia: EP (S, NC). – Azerbaijan, Turkey, Kazakhstan, China (NC).

PSILIGLOSSA S. Saunders, 1872 (*Stenoglossa* de Saussure, 1852, nec Chaudoir, 1848; *Psiloglossa* (!): Dalla Torre, 1894). Type species *Raphiglossa odyneroides* S. Saunders, 1850. Females nest in hollow stems, partitions between cells are made of plant pith, soil particles or other pressed material; provision is beetle larvae (Coleoptera). The genus is distributed in the Palaearctic. Number of species: Palaearctic – 4; Russia – 1.

Psiliglossa odyneroides (S. Saunders, 1850) [Raphiglossa] (*P. odyneroides kozhantshikovi* Kostylev,

1940). Russia: EP (NC). – Europe (EE, SE), Caucasus, Turkey, Turkmenistan.

PTEROCHEILUS Klug, 1805 (*Pterochilus* (!): Illiger, 1807; *Pterochile* (!): Blanchard, 1840; *Pterochylus* (!): de Saussure, 1853; *Odontopterochilus* van der Vecht, 1971; *Nannopterochilus* Blüthgen, 1961). Type species *Vespa phalerata* Panzer, 1797. Females excavate nest burrows in friable soil without using water; provision is caterpillars (Psychidae and Noctuidae). The genus is distributed in the Holarctic and Afrotropical regions and includes four subgenera: *Megapterocheilus* Bohart, 1940; *Micropterocheilus* Bohart, 1940; *Onchopterocheilus* Bohart, 1940; *Pterocheilus* s.str. Number of species: World – 85; Palaearctic – 32; Russia – 2.

Pterocheilus (Pterocheilus) phaleratus (Panzer, 1797) [Vespa] (*P. interruptus* Klug, 1805; *P. klugii* Panzer, 1809; *Pterochilus* (!) *chevrieranus* de Saussure, 1856; *Pterochilus* (!) *formosus* Frivaldszky, 1876; *Pterochilus* (!) *bezzii* Cobelli, 1892; *Pterochilus* (!) *phaleratus yeguasicus* Blüthgen, 1951; *Pterocheilus phaleratus tuberculatus* Blüthgen, 1955; *P. phaleratus kaszabi* Giordani Soika, 1970; *P. phaleratus luteiscapus* Guseinleitner, 1979; *P. phaleratus tibialis* Guseinleitner, 1991; *P. phaleratus suarezi* Guseinleitner, 1994; *P. phaleratus boreoibericus* Castro, 2003). Russia: EP (C, ?E, S, CR), UR, WS (NS, AL). – Europe (WE, NE, SE, EE), Georgia, Azerbaijan, Turkey, Kazakhstan, Mongolia.

Pterocheilus (Pterocheilus) sibiricus (Morawitz, 1867) [Odynerus] (*P. sibiricus* var. *fulvuleja* Blüthgen, 1955). Russia: ES (TU, BR, ZB). – Mongolia, China (NC).

RHYNCHIUM Spinola, 1806 (*Ryghium* (!) Spinola, 1806; *Rhynchium*: Billberg, 1820, emend. supported by IZN, 1965, Opinion 747; *Rychium* (!): Billberg, 1820; *Rynchium* (!): Sturm, 1829; *Rhygchium* (!): de Saussure, 1853; *Rhynchium* (!): de Saussure, 1863; *Eurrhynchium* Dalla Torre, 1904; *Rygochium* (!): Willink, 1982). Type species *Ryghium europaeum* Spinola, 1806. Females nest in various pre-existing cavities: hollow stems, beetle burrows in wood and abandoned nests of other wasps and bees, partitions between cells are made of mud; provision is various caterpillars (mainly of Microlepidoptera). The genus is distributed in the Palaearctic, Afrotropical, Oriental, and Australian regions. Number of species: World – 21; Palaearctic – 5; Russia – 1.

Rhynchium quinquecinctum (Fabricius, 1787) [Vespa (*Rh. tahitense* de Saussure, 1867; *Rh. fukaii* Cameron, 1911; *Rh. haemorrhoidale samurayi* Giordani Soika, 1973; *Rh. quinquecinctum murotai* Tano, 1983; *Rh. quinquecinctum nambui* Sk. Yamane, 1983). Russia: FE (KH, PR). – China (NE, NC, SE), Korean Peninsula, Japan (Hon, Shi, Kyu, Ryu), New Caledonia, Tahiti.

STENANCISTROKERUS de Saussure, 1863 (*Stenancystrocerus* (!): Dalla Torre, 1894; *Atropancistrocerus* Blüthgen, 1938). Type species *Odynerus atropos* Lepeletier de Saint Fargeau, 1841. Females build nests in vertical loam cliffs; provision is larvae of Curculionidae. The genus is distributed in the Palaearctic and Oriental regions and includes two subgenera: *Paratropancistrocerus* Blüthgen, 1938; *Stenancistrocerus* s.str. Number of species: World – 9; Palaearctic – 8; Russia – 1.

Stenancistrocerus (Paratropancistrocerus) obstrictus (Morawitz, 1895) [Odynerus]. Russia: EP (S, NC). – Caucasus.

STENODYNERUS de Saussure, 1863 (*Stemodynerus* (!): Rohwer, 1913; *Nannodynerus* Blüthgen, 1938; *Parhypodynerus* Giordani Soika, 1974). Type species *Odynerus chinensis* de Saussure, 1863. Females nest mainly in various preexisting cavities such as hollow stems, galls and abandoned nests of other wasps and bees, partitions between cells are made of mud; several Nearctic species excavate nest burrows in ground and surmount them by turrets; provision is various caterpillars (mainly of Microlepidoptera) or rarely larvae of Chrysomelidae. The genus is distributed in the Holarctic, Oriental, and Neotropical regions. Number of species: World – 166; Palaearctic – 44; Russia – 16.

Stenodynerus aequisculptus (Kostylev, 1940) [Odynerus] (*Nannodynerus meyeri* Blüthgen, 1952; *Pseudepipona constrictiventris* Giordani Soika, 1952; *N. aequisculptus taurus* Guseinleitner, 1966; *S. aequisculptus inaequisculptus* Giordani Soika, 1979; *S. aequisculptus cretensis* Guseinleitner, 1985). Russia: EP (NC). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Cyprus, Syria, Lebanon, Israel, Pakistan, Central Asia, Kazakhstan.

Stenodynerus bluethgeni van der Vecht, 1971 (*Odynerus pictus* Herrich-Schäffer, 1839, nom. praecocc., nec Curtis, 1826; *O. dentisquama*: Morawitz, 1895, Kostylev, 1928). Russia: EP (NW, S, NC, CR), UR,

WS (NS), ES (KR, IR). – Europe (WE, SE, EE), Caucasus, Turkey, Iran, Turkmenistan, Kazakhstan, China (NE, NC).

Stenodynerus chevrieranus (de Saussure, 1855) [Odynerus] (*O. dufourianus* de Saussure, 1855; *O. germanicus*: Kostylev, 1928; *O. germanicus ponticus* Kostylev, 1928; *Nannodynerus chevrieranus* var. *eximia* Blüthgen, 1953). Russia: **EP** (C, ?E, S, NC, CR), **UR**. – Europe (WE, SE, EE), Caucasus, Turkey, Iran, Central Asia, Kazakhstan.

Stenodynerus clypeopictus (Kostylev, 1940) [Odynerus] (*O. japonicus* Matsumura, 1926, nom. praeocc., nec von Schulthess, 1908; *Nannodynerus hungaricus* Blüthgen, 1961). Russia: **EP** (NC), **FE** (KH, PR). – Europe (WE, SE, EE), Caucasus, Turkey, Kyrgyzstan, Kazakhstan, Mongolia, China (NC), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu).

Stenodynerus fastidiosissimus (de Saussure, 1855) [Odynerus] (*O. difficilis* Morawitz, 1867; *O. proximus* Morawitz, 1867, nom. praeocc., nec de Saussure, 1857; *O. laborans* Costa, 1882; *O. insularis* André, 1883, nom. praeocc., nec Smith, 1857; *O. maroccanus* Dalla Torre, 1889; *O. ferrugineitarsis* De Stefani, 1889; *O. insulicola* Dalla Torre, 1894; *O. punctuosus* Ferton, 1901; *Nannodynerus vergesi* Giordani Soika, 1961; *Stenodynerus difficilis judaicus* Gusenleitner, 1970; *S. fastidiosissimus rufescens* Giordani Soika, 1977). Russia: **EP** (S, NC, CR), **UR**, **WS** (AL). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iraq, Iran, Central Asia, Kazakhstan.

Stenodynerus frauenfeldi (de Saussure, 1867) [Odynerus] (*O. nigricepsatus* Sonan, 1930). Russia: **FE** (PR, SA, KU). – China (NE, NC, CC, SW), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu).

Stenodynerus funebris (André, 1884) [Odynerus] (*O. limbonotatus* Kostylev, 1940). Russia: **ES** (BR, YA), **FE** (AM, KH, PR). – China (NE, NC), Korean Peninsula.

Stenodynerus orenburgensis (André, 1884) [Odynerus] (*Lionotus orbitalis* Thomson, 1874, nom. praeocc., nec *Odynerus orbitalis* Herrich-Schäffer, 1839). Russia: **EP** (C, S, NC, CR), **UR**, **WS** (OM, AL), **ES** (IR, BR, YA). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Kyrgyzstan, Kazakhstan, Mongolia, China (NE).

Stenodynerus pappi Giordani Soika, 1976 (*Parancistrocerus ussuriensis* Kurzenko, 1982; *S. pappi luteifasciatus* Kim et Sk. Yamane, 2004). Russia: **FE** (PR). – China (NE, NC, CC, SW, SE), Korean Peninsula.

Stenodynerus picticus (Thomson, 1874) [*Lionotus*]. Russia: **EP** (N, NW, C), **UR**, **WS** (TK, AL), **ES** (IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KA, MG). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan, Mongolia.

Stenodynerus pullus Gusenleitner, 1981. Russia: **ES** (BR, ZB), **FE** (AM, PR). – Turkey, Mongolia, China (NC), Korean Peninsula.

Stenodynerus punctifrons (Thomson, 1874) [*Lionotus*]. Russia: **EP** (C), **UR**, **WS** (OM, NS, AL), **ES** (TU, IR, BR, YA, ZB), **FE** (AM, PR). – Europe (WE, SE, EE), Caucasus, Turkey, ?Iran, Kyrgyzstan, Kazakhstan, Mongolia.

Stenodynerus sapidus Giordani Soika, 1970. Russia: **EP** (S, NC). – Iran.

Stenodynerus steckianus (von Schulthess, 1897) [Odynerus] (*O. dentisquama* var. *steckianus* von Schulthess, 1897; *O. teutonicus* Blüthgen, 1937). Russia: **EP** (S, NC, CR), **UR**. – Europe (WE, SE, EE), Azerbaijan, Turkey, Kazakhstan.

Stenodynerus tokyanus (Kostylev, 1940) [Odynerus] (*S. tokyanus flavoscutellatus* Sk. Yamane et Gusenleitner, 1982). Russia: **FE** (PR). – Korean Peninsula, Japan (Hon, Shi, Kyu, Ryu).

Stenodynerus xanthomelas (Herrich-Schäffer, 1839) [Odynerus] (*O. germanicus* de Saussure, 1855). Russia: **EP** (C, E, S, NC, CR), **WS** (AL). – Europe (WE, SE, EE), Caucasus, Turkey, Iran, China (NE).

SYMMORPHUS Wesmael, 1836 (*Protodynerus* de Saussure, 1855; *Synomorphus* (!): Rohwer, 1917; *Odynerus*: Blüthgen, 1938; *Koptodynerus* Blüthgen, 1943). Type species *Odynerus elegans* Wesmael, 1833. Females nest in various preexisting cavities (hollow stems, galls, beetle burrows in wood, abandoned nests of other wasps and bees), partitions between cells are made of mud; provision is generally larvae of Chrysomelidae, some species hunt for caterpillars of Microlepidoptera (*S. debilitatus*) or larvae of *Cionus* Clairville (Curculionidae) (*S. gracilis*). The genus is distributed in the Holarctic, Oriental, and Neotropical regions and includes two subgenera: *Parasymmorphus* Cumming et van der Vecht, 1986; *Symmorphus* s.str. Number of species: World – 48; Palaearctic – 27; Russia – 14.

Symmorphus (Symmorphus) allobrogus (de Saussure, 1855) [Odynerus] (*S. bifasciatus*: Tobias, Kurzenko, 1978). Russia: **EP** (N, NW, C, NC), **UR**, **WS** (TK, NS, KM, AL), **ES** (KR, BR, YA, ZB), **FE** (PR, MG). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Kyrgyzstan, Kazakhstan, China (NC).

Symmorphus (Symmorphus) angustatus (Zetterstedt, 1838) [Odynerus] (*Odynerus alternanus* Zetterstedt, 1838; *O. suecicus* de Saussure, 1855; *O. laeviventris* Thomson, 1874; *Symmorphus huktozanus* Tsuneki, 1986; *S. nansetsurei* Tsuneki, 1986; *S. iwatai* Sk. Yamane, 1990). Russia: **EP** (N, NW, C), **UR**, **WS** (TK, AL), **ES** (IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KA, MG). – Europe (WE, NE, SE, EE), Caucasus, Kazakhstan, Mongolia, China (NE), Korean Peninsula, Japan (Hon).

Symmorphus (Symmorphus) apiciornatus (Cameron, 1911) [Ancistrocerus] (*S. seoulensis* Tsuneki, 1986). Russia: **FE** (PR). – China (NE, NC, CC, SW, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Symmorphus (Symmorphus) bifasciatus (Linnaeus, 1761) [Vespa] (*Vespa sinuata* Fabricius, 1793, nom. praeocc., nec Geoffroy, 1785; *Odynerus sinuatus* var. *mutinensis* Baldini, 1894; *Symmorphus sparsus* Morawitz, 1895; *S. sinuatissimus* Richards, 1935; *S. mutinensis austriacus* Giordani Soika, 1975; *S. mutinensis yezoanus* Tsuneki, 1977). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (NS, AL), **ES** (KR, IR, ZB), **FE** (AM, KH, PR, SA). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Central Asia, Kazakhstan, Mongolia, China (NE, NC, CC, SW), Korean Peninsula, Japan (Hok, Hon).

Symmorphus (Symmorphus) connexus (Curtis, 1826) [Odynerus]. Russia: **EP** (C, NC), **WS** (AL). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Kazakhstan.

Symmorphus (Symmorphus) crassicornis (Panzer, 1798) [Vespa] (*Odynerus arcticus* de Saussure, 1855; *O. herrichianus* de Saussure, 1855; *O. nursei* Bingham, 1898; *O. hospes* Nurse, 1902, nom. praeocc., nec Dufour et Perris, 1840; *O. hospitans* Dalla Torre, 1904; *O. crassicornis* var. *baltica* Blüthgen, 1937). Russia: **EP** (N, NW, C, S, NC, CR), **UR**, **WS** (NS, AL), **ES** (KR, IR, ZB). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Central Asia, Kazakhstan, Mongolia, ?China (NE), India.

Symmorphus (Symmorphus) debilitatus (de Saussure, 1855) [Odynerus]. Russia: **EP** (C, E, NC, CR). – Europe (WE, NE, SE, EE), Turkey.

Symmorphus (Symmorphus) decens (Kostylev, 1940) [Odynerus]. Russia: **FE** (PR). – Korean Peninsula, Japan (Hon, Shi, Kyu).

Symmorphus (Symmorphus) foveolatus Gussakovskij, 1932. Russia: **FE** (KH, PR, SA). – China (SW), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu).

Symmorphus (Symmorphus) fuscipes (Herrich-Schäffer, 1838) [Odynerus] (*S. karelicus* Morawitz,

1895). Russia: **EP** (N, NW, C), **UR**, **WS** (TK, NS, AL), **ES** (KR, IR, BR, YA, ZB), **FE** (AM, PR, SA, MG). – Europe (WE, NE, EE), Azerbaijan, Kazakhstan, Mongolia, China (NE).

Symmorphus (Symmorphus) gracilis (Brullé, 1832) [Odynerus] (*O. elegans* Wesmael, 1833; *Symmorphus elegans libanicus* Giordani Soika, 1963). Russia: **EP** (C, E, S, NC, CR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Syria, Lebanon, Israel, Iran, Turkmenistan, Tajikistan, Kazakhstan.

Symmorphus (Symmorphus) lucens (Kostylev, 1938) [Odynerus] (*S. ishikawai* Giordani Soika, 1975). Russia: **UR**, **WS** (AL), **ES** (IR, BR, ZB), **FE** (AM, KH, PR, SA). – Mongolia, China (NC), Korean Peninsula, Japan (Hon).

Symmorphus (Symmorphus) mizuhonis Tsuneki, 1977 (*S. kurentzovi* Kurzenko, 1982; *S. iiyamai* Tsuneki, 1986; *S. shiroyamai* Tsuneki, 1986; *S. piceanus* Tsuneki, 1986; *S. sassai* Tsuneki, 1986). Russia: **WS** (AL), **ES** (IR, BR, ZB), **FE** (KH, PR, SA). – Kazakhstan, China (NE, NC, CC, SW, SE), Korean Peninsula, Japan (Hon).

Symmorphus (Symmorphus) murarius (Linnaeus, 1758) [Vespa] (*Odynerus nidulator* de Saussure, 1855). Russia: **EP** (NW, C, NC, CR), **UR**, **WS** (TM), **ES** (KR, IR, BR, ZB), **FE** (PR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Iran, Central Asia, Kazakhstan, Korean Peninsula.

SYNEUODYNERUS Blüthgen, 1951 (*Syneodynerus* (!): Kurzenko, 1981). Type species *Odynerus egregius* Herrich-Schäffer, 1839. Females nest in preexisting cavities such as hollow stems or beetle burrows in wood, partitions between cells are double and made of mud; provision is caterpillars of Pyraustidae. The genus is distributed in the Palaearctic, Afrotropical, and Australian region. Number of species: World – 11; Palaearctic – 5; Russia – 1.

Syneodynerus egregius (Herrich-Schäffer, 1839) [Odynerus] (*O. bohemani* de Saussure, 1855; *O. unicolor* Maidl, 1922; *O. egregius* var. *discosignatus* Mader, 1936; *Euodynerus egredius barbarus* Blüthgen, 1942; *E. egredius arzhakanensis* Giordani Soika, 1970; *E. egredius egredior* Gusenleitner, 1970). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Cyprus, Syria, Jordan, Israel, Iran, Turkmenistan.

TROPIDODYNERUS Blüthgen, 1939. Type species *Polistes interrupta* Brullé, 1832. Females

excavate vertical nest burrows on horizontal ground surface, each nest contains one cell; nest entrance is obstructed by the female with mud pellets before each her departure from the nest; provision is larvae of Curculionidae. The genus is distributed in the Palaearctic and Oriental regions and includes two subgenera: *Tropidepipona* Giordani Soika, 1995; *Tropidodynerus* s.str. Number of species: World – 7; Palaearctic – 5; Russia – 1.

Tropidodynerus (Tropidodynerus) interruptus (Brullé, 1832) [Polistes] (*Pterocheilus interruptus* Herrich-Schäffer, 1839, nom. praeocc., nec Klug, 1805; *Odynerus bulgaricus* Mocsáry, 1883; *Hoplomerus mandibularis* Morawitz, 1885; *Tropidodynerus interruptus tricolor* Blüthgen, 1956; *T. interruptus hispanicus* Giordani Soika, 1966). Russia: **EP** (E, S, NC, CR). – Europe (WE, SE, EE), Armenia, Azerbaijan, Turkey, Kazakhstan.

Subfamily POLISTINAE – PAPER WASPS

A.V. ANTROPOV

POLISTES Latreille, 1802 (*Polistes*: Palisot de Beauvois, 1818, lapsus calami; *Polistes* (*Eupolistes*) Dalla Torre, 1904; *Pseudopolistes* Weyrauch, 1937, unavailable name, Art. 13.3 (ICZN, 1999); *Polistes* (*Sulcopolistes*) Blüthgen, 1938; *Polistula* Weyrauch, 1938, unavailable name, Art. 13.3 (ICZN, 1999); *Polistula* Weyrauch, 1939, *Pseudopolistes* Weyrauch, 1939, *Polistes* (*Leptopolistes*) Blüthgen, 1943. Type species “*Polistes gallica*, Fab.” [= *Vespa gallica* Linnaeus, 1767]. The genus includes four subgenera: *Gyrostoma* Kirby, 1828 (*Cyclostoma* Kirby, 1826, nom. praeocc., nec Lamarck, 1779; *Megapolistes* van der Vecht, 1968; *Nygapolistes* Richards, 1973; *Megapostes* [!]: Lee, 1982), *Polistella* Ashmead, 1904 (*Stenopolistes* van der Vecht, 1972), *Polistes* s.str., *Aphanilopterus* Meunier, 1888 (*Polistoides* Dalla Torre, 1904, nom. praeocc., nec Loew, 1873; *Onerarius* Richards, 1973; *Fuscopolistes* Richards, 1973; *Polistarchus* Richards, 1973; *Palisotius* Richards, 1973; *Epicnemius* Richards, 1973). Number of species: World – 224; Palaearctic – 33; Russia – 14.

Polistes (Gyrostoma) rothneyi Cameron, 1900 (*P. rufolineatus* Cameron, 1900; *P. rothneyi yayeyamae* Matsumura, 1908; *P. rothneyi grahami* van der Vecht, 1968; *P. rothneyi hainanensis* van der

Vecht, 1968; *P. rothneyi iwatai* van der Vecht, 1968; *P. rothneyi gressitti* van der Vecht, 1968; *P. rothneyi tibetanus* van der Vecht, 1968; *P. rothneyi koreanus* van der Vecht, 1968; *P. rothneyi sikkimensis* van der Vecht, 1968). Russia: **FE** (PR). – Pakistan, China (WP, SE), Korean Peninsula, Japan (Ryu), Nepal, Bhutan, SE Asia, India.

Polistes (Polistella) diakonovi Kostylev, 1940. Russia: **FE** (PR). – Korean Peninsula.

Polistes (Polistella) snelleni de Saussure, 1862 (*P. puncticollis* F. Morawitz, 1892; *P. snelleni* f. *nigrihumerus* Uchida, 1936). Russia: **ES** (BR, ZB), **FE** (AM, KH, PR). – Mongolia, China (NE), Korean Peninsula, Japan (Hok, Hon, Kyu).

Polistes (Polistes) albellus Giordani Soika, 1976 (*P. foederatus albellus* Giordani Soika, 1976; *P. bischoffi* Weyrauch, 1937, part.; *P. helveticus* Neumeyer, 2014). Russia: **EP** (C, E, S), **UR**, **WS** (NS, AL), **ES** (KS, TU, IR, BR, ZB), **?FE** (PR). – Europe (WE, SE, EE), Kazakhstan, Mongolia.

Polistes (Polistes) associus Kohl, 1898 (*P. associa* Kohl, 1898). Russia: **EP** (S, NC). – Europe (WE, SE, EE), Azerbaijan, Turkey, Israel, Iran, Pakistan, Tajikistan, India, China (NW).

Polistes (Polistes) atrimandibularis Zimmermann, 1930 (*P. atrimandibularis albidus* Blüthgen, 1957). Russia: **EP** (S, NC). – Europe (SE, EE), Egypt, Armenia, Turkey, Iran.

Polistes (Polistes) biglumis (Linnaeus, 1758) [Vespa] (*V. rupestris* Linnaeus, 1758; *V. bimaculata* Geoffroy in Fourcroy, 1785; *P. Geoffroyi* Lepeletier de Saint Fargeau et Audinet-Serville, 1825; *P. dubia* Kohl, 1898, nom. praeocc., nec de Saussure, 1867; *P. Kohli* Dalla Torre, 1904, replacement name for *P. dubia* Kohl, 1898; *V. bimacula* [!] “Fourcroy”: Dalla Torre, 1904; *P. bimaculatus* var. *Arduinoi* Guiglia, 1948; *P. pamirensis* Soika: Zirngiebl, 1955; *P. pamirensis* var. *soikai* Zirngiebl, 1955; *P. pamirensis* var. *interruptus* Zirngiebl, 1955; *P. bimaculatus* var. *nigrinotum* Zirngiebl, 1955; *P. biglumis alpium* Blüthgen, 1957; *P. geoffroyi* [!]: Das, Gupta, 1984). Russia: **EP** (C, S, NC). – Europe (WE, NE, SE, EE), N Africa, Abkhasia, Azerbaijan, Turkey, Syria, Israel, Iraq, Iran, Pakistan, Kyrgyzstan, Kazakhstan, Mongolia, Korean Peninsula, India.

Polistes (Polistes) bischoffi Weyrauch, 1937 (*P. Bischoffi* Weyrauch, 1937). Russia: **EP** (C, S, NC), **FE** (AM, KU). – Europe (WE, NE, SE, EE), N Africa, Turkey, Kazakhstan, Iran, Mongolia.

Polistes (Polistes) chinensis (Fabricius, 1793) [Vespa] (*P. biglumis* var. *antennalis* Pérez, 1905; *P. insepultus* Giordani Soika, 1976; *P. caspicus* Yoshikawa, 1962, nom. nud.). Russia: **WS** (AL), **ES** (TU, BR, ZB), **FE** (AM, KH, PR, ?KA). – Turkey, Kazakhstan, Iran, Mongolia, China (SE), Korean Peninsula, Japan (Hok, Hoh, Kyu, Ryu). Introduced into New Zealand.

Polistes (Polistes) dominula (Christ, 1791) [Vespa] (*P. italicica* Herrich-Schäffer, 1840, nom. nud.; *P. pectoralis* "Meg." Herrich-Schäffer, 1841; *P. gallica* var. *Lefebvrei* Guérin-Méneville, 1844; *P. Bucharensis* Erichson, 1849; *P. Lefebvrii* [!]: de Saussure, 1853; *P. maculatus* Rudow, 1889; *P. Merceti* Dusmet, 1903; *P. gallicus* var. *rufescens* du Buysson, 1912; *P. gallica* var. *ornata* Weyrauch, 1938, nom. praeocc., nec *Polistes ornata* Lepeletier de Saint Fargeau, 1836; *P. gallica* var. *pacifica* Weyrauch, 1939, nom. praeocc., nec *P. pacifica* Fabricius, 1804; *P. gallicus pseudopacificus* Giordani Soika, 1970, replacement name for *P. gallica* var. *pacifica* Weyrauch, 1939; *P. gallicus muchei* Gusenleitner, 1976; *Vespa dominuta* [!]: Das, Gupta, 1984). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), N Africa, Armenia, Georgia, Turkey, Syria, Jordan, Israel, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NW), India. Introduced into the Canary Islands, Australia, N and S America.

Polistes (Polistes) gallicus (Linnaeus, 1767) [Vespa] (*P. pictior* Radoszkowski, 1872, nom. nud.; *P. foederata* Kohl, 1898; *P. gallicus* var. *mongolicus* du Buysson, 1911; *P. foederata* var. *obscuricornis* Mader, 1936; *Polistula omissa* Weyrauch, 1938; *Polistes omissus* var. *ordubadensis* Zirngiebl, 1955; *P. gallicus* var. *monogolicus* [!]: Yoshikawa, 1962; *P. omissus kaszabi* Giordani Soika, 1970). Russia: **EP** (C, S, NC, CR), **WS** (AL). – Europe (WE, SE, EE), N Africa, Ethiopia, Israel, Jordan, Turkey, Iran, Afghanistan, Pakistan, Armenia, Azerbaijan, Turkmenistan, Kyrgyzstan, Kazakhstan, Mongolia.

Polistes (Polistes) nimpha (Christ, 1791) [Vespa] (*V. diadema* Latreille, 1802, nom. praeocc., nec Christ, 1791; *Polistes opinabilis* Kohl, 1898; *P. nimpha* [!]: Weyrauch, 1939; *P. nimpha* var. *Moltonii* Guiglia, 1944; *P. nimpha* var. *moltonii* [!]: Yoshikawa, 1962; *P. nimpha irakensis* Gusenleitner, 1976). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (TM, KM, AL), **ES** (KS, TU, KR, BR, YA, ZB), **FE** (AM, KH). – Europe (WE, NE, SE, EE), N Africa, Israel, Jordan, Turkey, Georgia, Armenia, Azerbaijan, Iraq,

Iran, Pakistan, Kazakhstan, Kyrgyzstan, Mongolia, China (NE), India.

Polistes (Polistes) riparius Sk. Yamane et S. Yamane, 1987. Russia: **EP** (C), **UR**, **WS** (TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU). – Mongolia, Kazakhstan, Mongolia, China (NE), Korean Peninsula, Japan (Hok, Hon).

Polistes (Polistes) sulcifer Zimmermann, 1930 (*P. sulcifera* [!]: Guiglia, Capra, 1933; *Pseudopolistes sulcifer* var. *similator* Zirngiebl, 1955). Russia: **EP** (NC). – Europe (WE, SE, EE), N Africa, Armenia, Azerbaijan, Egypt, Israel, Turkey, Iran.

Subfamily VESPINAE – HORNETS AND JELLOW JACKETS

A.V. ANTROPOV

VESPA Linnaeus, 1758 – Hornets (*Macrovespa* Dalla Torre, 1904; *Nyctovespa* van der Vecht, 1959). Type species "Vespa crabro Fab." [= *Vespa crabro* Linnaeus, 1758]. Number of species: World – 22; Palaearctic – 15; Russia – 8.

Vespa analis Fabricius, 1775 – Yellow-tailed Hornet (*V. Crabro sphinx* Christ, 1791; *V. tyrranica* F. Smith, 1857; *V. japonica* F. Smith, 1868, nom. praeocc., nec Radoszkowski, 1857, nec de Saussure, 1858; *V. parallela* André, 1884; *V. insularis* Dalla Torre, 1894, replacement name for *V. japonica* F. Smith, 1868; *V. tridentata* Cameron, 1903; *V. nigrons* du Buysson, 1903; *V. parallela* var. *Biroi* du Buysson, 1905; *V. analis* var. *tenebrosa* du Buysson, 1905; *V. analis* var. (subsp.) *barbouri* Bequaert, 1939; *V. analis* var. (subsp.) *kuangiana* Bequaert, 1939; *V. analis eisa* Sk. Yamane, 1987; *V. analis nagatomii* Sk. Yamane, 1987; *V. maguanensis* Dong, 2001; *V. hekouensis* Dong et Wang, 2003). Russia: **FE** (AM, KH, PR). – China (NE, CC, SW, SE), Korean Peninsula, Japan (Hok, Hon, Ryu), Nepal, SE Asia, India.

Vespa binghami du Buysson, 1905 – Hornet of Bingham (*V. Binghami* du Buysson, 1905; *V. suprunenkoi* Birula, 1925). Russia: **FE** (KH, PR, SA). – China (CC, SW, WP,), Korean Peninsula, SE Asia, India.

Vespa crabro Linnaeus, 1758 – Common Hornet (*V. Crabro* Linnaeus, 1758; *V. vexator* Harris, 1776; *V. Crabro major* Retzius, 1783; ?*V. pratensis* Geoffroy in Fourcroy, 1785; *V. Crabro germana* Christ, 1791; *V. Crabroniformis* F. Smith, 1852; *V. crabro* var. *borealis* Radoszkowski, 1863, nom. praeocc.,

nec *V. borealis* Kirby, 1837, nec *V. borealis* Zetterstedt, 1838, nec *V. borealis* Zetterstedt, 1840, nec *V. borealis* F. Smith, 1843; *V. crabro* var. *anglica* Gribodo, 1892, nom. praeocc., nec *V. anglica* F. Smith, 1843; *V. oberthuri* du Buysson, 1902; *V. flavofasciata* Cameron, 1903; *V. crabro* var. *tartarea* du Buysson, 1905; *V. Crabro* var. *Altaica* Pérez, 1910; *V. Crabro* var. *Caspica* Pérez, 1910; *V. crabro nigra* Birula, 1925, nom. praeocc., nec *V. nigra* Geoffroy, 1785; *V. crabro vulgata* Birula, 1925; *V. crabro meridionalis* Birula, 1925; *V. crabro chinensis* Birula, 1925, nom. praeocc., nec *V. chinensis* Fabricius, 1793; *V. crabro* var. *birulai* Bequaert, 1931, replacement name for *V. crabro chinensis* Birula, 1925; *V. crabro* var. *gribodoi* Bequaert, 1931, replacement name for *V. crabro* var. *anglica* Gribodo, 1892). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, ZB), **FE** (AM, KH, PR, SA). – Europe (WE, NE, SE, EE), Georgia, Azerbaijan, Turkey, Turkmenistan, Kazakhstan, Iran, Mongolia, China (NE, CC, SW, SE), Korean Peninsula, Japan (Hok, Hon). Introduced into N America and Guatemala.

Vespa ducalis F. Smith, 1852 – Black-tailed Hornet (*V. ducalis* var. *pulchra* du Buysson, 1905; *V. matsu-murai* Sonan, 1935; *V. esakii* Sonan, 1935; *V. tropica loochooensis* Bequaert, 1936; *V. tropica pseudosoror* van der Vecht, 1959). Russia: **ES** (ZB), **FE** (AM, KH, PR). – China (NE, NC, CC, SW, SE), Korean Peninsula, Japan (Hok, Hon, Ryu), Nepal, SE Asia, India.

Vespa dybowskii André, 1884 – Hornet of Dybowski (*Vespa Dybowskii* "Rad. in litt." André, 1884; *V. Dubowskii*: Dalla Torre, 1904; *V. Walkeri* du Buysson, 1905; *V. dybowskii mutata* Ma, 1937). Russia: **ES** (ZB), **FE** (AM, KH, PR). – China (NW, CC), Korean Peninsula, Japan (Hok, Hon, Ryu), Myanmar, Thailand.

Vespa mandarinia F. Smith, 1852 – Asian Giant Hornet (*V. Mandarinia* F. Smith, 1852; *V. magnifica* F. Smith, 1852; *V. japonica* Radoszkowski in Motschulsky, 1857; *V. bellona* F. Smith, 1871; *V. magnifica* var. *latilineata* Cameron, 1903; *V. mandarina*: Dalla Torre, 1894; *V. magnifica* var. *nobilis* Sonan, 1929; *V. magnifica sonani* Matsumura, 1930). Russia: **FE** (AM, KH, PR). – China (NW, CC, SW, SE), Korean Peninsula, Japan (Hok, Hon, Kyu, Ryu), Nepal, Bhutan, SE Asia, India, Sri Lanka.

Vespa orientalis Linnaeus, 1771 – Oriental Hornet (*V. turcica* Drury, 1773; *V. quadripunctata* Forskål, 1775; *V. Crabro fusca* Christ, 1791; *V. Ägyptiaca*

Vallot, 1802; *V. Nilotica* Vallot, 1802; *V. Jurinei* de Saussure, 1854; *V. orientalis* var. *Ägyptiaca* André, 1884, nom. praeocc., nec *V. Ägyptiaca* Vallot, 1802; *V. indica*: Wroughton, 1889; *V. orientalis* var. *Zavattarii* Guiglia et Capra, 1933; *V. orientalis* var. *somalica* Giordani Soika, 1934; *V. orientalis arabica* Giordani Soika, 1957). Russia: **EP** (S, NC, CR). – Europe (SE, EE), N Africa, Georgia, Armenia, Turkey, Syria, Jordan, Lebanon, Israel, Saudi Arabia, Bahrain, Yemen, UAE, Oman, Iraq, Iran, Afghanistan, Pakistan, Turkmenistan, Tajikistan, Uzbekistan, Nepal, India. Introduced into China (SE), Madagascar, and Mexico.

Vespa simillima F. Smith, 1868 – Japanese Hornet (*V. mongolica* André, 1884; *V. xanthoptera* Cameron, 1903; *V. micado* Cameron, 1903; *V. Mongolica* var. *Sexpunctata* Pérez, 1905; *V. Mongolica* var. *flavata* Pérez, 1910). Russia: **FE** (AM, KH, PR, SA, KU, KA). – China (NE), Korean Peninsula, Japan (Hok, Hon, Kyu, Ryu). Introduced into Canada (BC) but has apparently not become established.

DOLICHOVESPULA Rohwer, 1916 – "Long-cheeked" Jellow Jackets [*Pseudovespa* (*Pseudovespula*) Bischoff, 1931; *Dolichovespula* (*Boreovespula*) Blüthgen, 1943; *Dolichovespula* (*Metavespula*) Blüthgen, 1943]. Type species *Vespa maculata* Linnaeus, 1763. Number of species: World – 19; Palaearctic – 10; Russia – 8.

Dolichovespula adulterina (du Buysson, 1905) [Vespa] – False wasp (*V. norwegica* var. *adulterina* du Buysson, 1905; *V. norwegica saxonica* natio *colchica* Birula, 1930, unavailable name, Art. 45.5, 45.5.1 (ICZN, 1999); *D. adulterina* *montivaga* Sk. Yamane, 1982). Russia: **EP** (N, NW, C, E, NC), **WS** (TK, NS, AL), **ES** (KM, TM, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KA, MG, CH). – Europe (WE, NE, SE, EE), Georgia, Kyrgyzstan, Kazakhstan, Turkey, Mongolia, China (SW, ?SE), Korean Peninsula, Japan (Hok, Hon). All records of this species from North America should be referred to *D. arctica* (Rohwer, 1916) (*Vespa borealis* Lewis, 1897).

Dolichovespula intermedia (Birula, 1930) [Vespula] (*Vespula sylvestris* var. *intermedia* Birula, 1930; *D. asiatica* Archer, 1981, unnecessary replacement name for *Vespula sylvestris* var. *intermedia* Birula, 1930, nec *Vespa rufa* var. *intermedia* du Buysson, 1905). Russia: **WS** (AL). – Central Asia, Kazakhstan, Afghanistan, Pakistan, Mongolia, China (NC, NW, SE), Japan (Hok, Hon), India.

Dolichovespula media (Retzius, 1783) [Vespa] – Medium wasp (*V. Crabro medius* Retzius, 1783; *V. Geerii* Lepeletier de Saint Fargeau, 1836; *V. crassa* Herrich-Schäffer, 1841; *V. similis* Schenck, 1853; *V. rufoscutellata* Schenck, 1853; *V. flavicincta* Schenck, 1853, nom. praeocc., nec Gravenhorst, 1807; *V. Geeri*: Dalla Torre, 1904; *V. media* var. *lineolata* Pérez, 1910; *Vespa media media* var. *lugubris* Birula, 1927, unavailable name, Art. 45.5, 45.5.1 (ICZN, 1999); *D. media* var. *conjugens* Paul, 1943; *D. media sugare* Ishikawa, 1969; *D. borealis* Lee, 1986, nom. praeocc., nec *V. borealis* Kirby, 1837, nec *V. borealis* Zetterstedt, 1838, nec *V. borealis* Lewis, 1897). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, ?KA). – Europe (WE, NE, SE, EE), Morocco, Georgia, Turkey, Syria, Kazakhstan, Mongolia, China (NE), Korean Peninsula, Japan (Hok, Hon). Occasionally introduced into New Zealand.

Dolichovespula norwegica (Fabricius, 1781) [Vespa] – Norwegian wasp (*V. britannica* Leach, 1814; *V. marginata* Kirby, 1837, nom. praeocc., nec Gmelin, 1790; *V. borealis* Zetterstedt, 1838, nom. praeocc., nec Kirby, 1837; *V. norvegica*: Olivier, 1792; *V. peruana* de Saussure, 1867; *V. saxonica* var. *arctica* Friese, 1919, nec *Vespa arctica* Rohwer, 1916; *D. norwegica* var. *Zetterstedti* Blüthgen, 1937, replacement name for *V. saxonica* var. *arctica* Friese, 1919). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (TM, OM, NS, TK, AL), **ES** (KS, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KA, MG, CH). – Europe (WE, NE, EE), Georgia, Turkey, Kyrgyzstan, Kazakhstan, Mongolia, China (NE, NW). All records of this species from North America should be referred to *D. albida* (Sladen, 1918).

Dolichovespula omissa (Bischoff, 1931) [Vespa] – Carefree wasp [*V. tripunctata* Schenck, 1861, nom. praeocc., nec Fabricius, 1787; *Vespa norvegica saxonica* morpha *ingrica* Birula, 1930, unavailable name, Art. 45.5, 45.5.1 (ICZN, 1999); *Vespa ingrica* Bequaert, 1932, available name, Art. 45.5.1 (ICZN, 1999)]. Russia: **EP** (NW, C, E), **UR**, **WS** (AL), **ES** (KS). – Europe (WE, NE, EE), Georgia, Turkey; Iran, Kazakhstan.

Dolichovespula pacifica (Birula, 1930) [Vespa] – Pacific wasp (*V. norvegica pacifica* Birula, 1930; *D. loekenae* Eck, 1980). Russia: **EP** (N, NW, C), **UR**, **WS** (TM, OM, NS, TK, AL), **ES** (TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, MG, KA). – Europe

(NE), Kazakhstan, Korean Peninsula, Japan (Hok, Hon, Shi).

Dolichovespula saxonica (Fabricius, 1793) [Vespa] – Saxonian wasp (*V. bavarica* von Schrank, 1802; *V. tridens* Schenck, 1853; *Vespa norvegica saxonica* morpha *monticola* Birula, 1930, unavailable name, Art. 45.5, 45.5.1 (ICZN, 1999); *D. saxonica nipponica* Sk. Yamane, 1975; *D. saxonica kamtschatkensis* Eck, 1983; *D. saxonica nigrescens* Eck, 1983). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KH, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KA, MG, CH). – Europe (WE, NE, SE, EE), Turkey, Iran, Kazakhstan, Mongolia, China (NE, NC, NW, SW), Korean Peninsula, Japan (Hok, Hon).

Dolichovespula sylvestris (Scopoli, 1763) [Vespa] – Forest wasp (*V. parietum* Harris, 1776, nom. praeocc., nec Linnaeus, 1758; *V. holsatica* Fabricius, 1793; *V. frontalis* Latreille, 1802; *V. campanaria* Fowler, 1833; *V. anglica* "Leach" F. Smith, 1843, available name, Art. 11.6.1 (ICZN, 1999); *V. pilosella* A. Costa, 1858, nom. praeocc., nec Gravenhorst, 1807; *V. silvestris*: Dalla Torre, 1894; *V. silvestris* var. *sumptuosa* du Buysson, 1905; *D. xinjiangensis* Lee, 1986). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, AL), **ES** (KS, TU, KR, IR, BR, ZB), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), Morocco, Georgia, Armenia, Turkey, Syria, Iran, Afghanistan, Pakistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE, NC, NW, CC, SE), India.

VESPULA Thomson, 1869 – "Short-cheeked" Jellow Jackets [*Vespa (Pseudovespa)* Schmiedeknecht, 1881; *Dolichovespula (Paravespula)* Blüthgen, 1938; *Paravespula (Allovespula)* Blüthgen, 1943; *Vespa (Rugovespula)* Archer, 1982]. Type species *Vespa austriaca* Panzer, 1799. Number of species: World – 25; Palaearctic – 11; Russia – 7.

Vespa austriaca (Panzer, 1799) [Vespa] – Austrian wasp (*V. borealis* F. Smith, 1843, nom. praeocc., nec Kirby, 1837, nec Zetterstedt, 1840; *V. arborea* F. Smith, 1849, replacement name for *V. borealis* F. Smith, 1843; *V. biloba* Schilling, 1850; *V. arborea* "Ross" Dalla Torre, 1894, nom. praeocc., nec F. Smith, 1849; *Vespa shinanoensis* Takamizawa, 2005, nom. nud.). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (TM, NS, AL), **ES** (KS, KR, IR, BR, YA), **FE** (AM, KH, PR, SA, KU, KA). – Europe (WE, NE, EE), Georgia, Azerbaijan, Turkey, Pakistan,

Kyrgyzstan, Kazakhstan, Mongolia, China (NE, NW), Korean Peninsula, Japan (Hok, Hon), India. All records of this species from North America should be referred to *V. infernalis* (de Saussure, 1854).

Vespa flaviceps (F. Smith, 1870) [Vespa] – Yellow-fronted wasp (*V. japonica* de Saussure, 1858, nom. praeocc., nec Radoszkowski, 1857; *V. Lewisi*, Sauss. Guepes Soc. (MS.) F. Smith, 1873, nom. nud.; *V. lewisi*, Saus. MS.) Cameron, 1903, available name, Art. 11.6.1.(ICZN, 1999); *V. Saussurei* Schulz, 1906, replacement name for *V. japonica* de Saussure, 1858; *V. karenkona* Sonan, 1929; *V. 4-maculata* Sonan, 1929; *V. vulgaris* var. *flavitor* Stolfa, 1934; *Vespa japonica pionganensis* Giordani Soika, 1976; *Vespa vulgaris* var. *flavitor*: Das, Gupta, 1984; *Vespa gracilia* Lee, 1986). Russia: **FE** (AM, KH, PR). – China (NE, NW, CC, SW, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu), Nepal, SE Asia, India.

Vespa germanica (Fabricius, 1793) [Vespa] – German wasp (?*V. maculata* Scopoli, 1763, nom. praeocc., nec Linnaeus, 1763; ?*V. macularis* Olivier, 1792). Russia: **EP** (C, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, AL), **ES** (KS, TU, KR, BR, ZB), **FE** (AM, KH, PR, SA). – Europe (WE, NE, SE, EE), N Africa, Israel, Jordan, Lebanon, Georgia, Armenia, Turkey, Syria, Iran, Iraq, Afghanistan, Pakistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NE, NC, NW, SW, WP, SE), Korean Peninsula, India. Introduced into Iceland, Madeira, Azores, Canaries, South Africa, Australia, Tasmania, New Zealand, Ascension Is., Norfolk Is., Hawaiian Is., N America, S America (Argentina, Chile).

Vespa koreensis (Radoszkowski, 1887) [Vespa] – Korean wasp (*V. Koreensis* Radoszkowski, 1887; *V. germanica* var. *stizoides* du Buysson, 1905; ?*V. orbata* var. *aurulenta* du Buysson, 1905; *Pseudovespa birulai* Gussakovskii, 1932; *Vespa koreensis salebrosa* Archer, 1982; *V. hainanensis* Lee,

1986). Russia: **FE** (AM, KH, PR). – China (NE, CC, SW, SE), Korean Peninsula, Thailand, Laos, India.

Vespa rufa (Linnaeus, 1758) [Vespa] – Red wasp (*V. Schrenckii* Radoszkowski, 1861; *V. Sibirica* André, 1884; *V. schrenki*: Dalla Torre, 1904; *Vespa rufa grahami* Archer, 1981; *V. obscura* Lee, 1986; *V. yichunensis* Lee, 1986; *V. yulongensis* Dong et Wang, 2002; *V. gongshanensis* Dong, 2005). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KA, MG, CH). – Europe (WE, NE, SE, EE), Turkey, Georgia, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Iran, Mongolia, China (NE, NW, SW, SE), Korean Peninsula, Japan (Hok, Hon, Shi), Nepal. All records of this species from North America should be referred to *V. intermedia* (du Buysson, 1905).

Vespa shidai Ishikawa, Sk. Yamane et Wagner, 1980 – Wasp of Shida (*V. shidai amamiana* Sk. Yamane, 1987). Russia: **FE** (AM, KH, PR, KU). – China (NE), Korean Peninsula, Japan (Hok, Hon, Shik, Kyu, Ryu).

Vespa vulgaris (Linnaeus, 1758) [Vespa] – Common wasp (*V. sexcincta* Panzer, 1799; *V. communis* de Saussure, 1857, nom. praeocc., nec von Schrank, 1785; *V. vulgaris* var. *pseudogermanica* Stolfa, 1932; *V. mixta* "Schenck" Edwards, 1980, nom. nud.; *Vespa rufosignata* Eck, 1998; *V. rufosignata vetus* Eck, 1999; *V. yunlongensis* Dong et Wang, 2003). Russia: **EP** (NW, N, C, E, S, NC, CR), **UR**, **WS** (TM, TK, NS, KM, AL), **ES** (TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KA, MG, CH). – Europe (WE, NE, SE, EE), Turkey, Georgia, Kyrgyzstan, Kazakhstan, Iran, Mongolia, China (NE, SW), Korean Peninsula, Japan (Hok, Hon), India. Introduced into Iceland, New Zealand, Australia, and occasionally into Maui and Hawaii. All records of this species from North America should be referred to *V. alascensis* (Packard, 1870).

SUPERFAMILY FORMICOIDEA

69. FAMILY FORMICIDAE – ANTS

D.A. DUBOVIKOFF AND Z.M. YUSUPOV

Ants are eusocial aculeate hymenopterous insects and currently include 13240 extant species belongs to 334 genera of 17 subfamilies and 738 fossil species from 151 genera of 14 subfamilies (3 subfamilies are known only by fossils) (Bolton, 2016). There are 1777 species in 97 genera of 11 subfamilies in the Palaearctic region (AntWeb, 2017). Ants, as eusocial insects, have a clear cast differentiation, e.g. sexual (females (queens) and males) and workers (or also soldiers). Colonies include one (or more in polygynous colonies) queen and from dozens to millions of workers. Ants are one of the most evolutionarily successful groups of invertebrates. They are of great importance in various ecosystems, primarily as effective predators, but also as consumers and distributors of plant seeds, play a significant role in soil formation and much more. Ants nesting is much different: in soil, leaflitter, dead wood, in epiphytes, different plants and with different special constructions like mounds. Synonymic and nomenclature data are given according to «An online catalog of the ants of the world» (Bolton, 2016). The total number of taxa is given excluding fossil and infraspecific taxa, unless explicitly indicated.

Sections for the genera *Leptothorax*, *Myrmica*, *Temnothorax* and most part of *Tetramorium* species were prepared by Z.M. Yusupov, the rest of the text was prepared by D.A. Dubovikoff.

Number of taxa: World – 485 genera, 13978 species (including fossils), Palaearctic – 94/1350, Russia – 44/264.

References. Ruzsky, 1905, 1907; Karawajew, 1926; Kuznetsov-Ugamsky, 1929; Dlussky, Soyunov, Zabelin, 1990; Kupianskaya, 1990, 1995, 2012; Radchenko, 1991, 1992, 1996a, 1996b, 1997a,b,c,d, 2004, 2005a,b, 2016; Seifert, 1992, 2000, 2003, 2007, 2012; Arakelian, 1994; Czechowski et al., 2002; Dubovikoff, 2002, 2005; Csösz et al., 2007, 2010, 2015; Pfeiffer et al., 2007; Seifert, Schultz, 2009; Yusupov, 2009, 2014; Radchenko, Elmes, 2010; Czechowski et al., 2012; Douwes et al., 2012; Guénard, Dunn, 2012; Radchenko, Yusupov, 2012, 2017; Borowiec, 2014; Radchenko, Scupola, 2015; Radchenko et al., 2015; Bolton, 2016; Radchenko et al., 2016; Ward et al., 2016; AntWeb, 2017; Grebennikov, Zryanyi, 2017; Dubovikoff, Yusupov, 2017.

Subfamily DOLICHODERINAE

Number of taxa: World – 28 genera, 707 species, Palaearctic – 8/53, Russia – 4/16.

BOTHRIOMYRMEX Emery, 1869. Type species

Bothriomyrmex costae Emery, 1869. Temporary social parasites of *Tapinoma* species. Number of species: World – 24, Palaearctic – 22, Russia – 5.

Bothriomyrmex anastasiae Dubovikov, 2002. Russia: EP (NC).

Bothriomyrmex communista Santschi, 1919 (*B. meridionalis* var. *communista* Santschi, 1919). Russia: EP (NC, CR). – Europe (WE, SE, EE).

Bothriomyrmex corsicus Santschi, 1923 (*B. meridionalis* var. *corsica* Santschi, 1923; *B. gallicus* Emery, 1925; *B. gibbus* Soudek, 1925; *B. menozzii* Emery, 1925). Russia: EP (NC, CR). – Europe (WE, SE, EE).

Bothriomyrmex modestus Radchenko, 1985. Russia: EP (S). – Europe (EE).

Bothriomyrmex turcomanicus Emery, 1925 (*B. syrius turcomanicus* Emery, 1925). Russia: EP (S, NC). – Turkmenistan, Kazakhstan.

DOLICHODERUS Lund, 1831 (*Hypoclinea* Mayr, 1855). Type species *Formica attelaboides* Fabricius, 1775. Number of species: World – 131, Palaearctic – 2, Russia – 2.

Dolichoderus quadripunctatus (Linnaeus, 1771) [Formica] (*Hypoclinea quadripunctata* (Linnaeus, 1771), auct.). Russia: EP (C, E, S, NC, CR), UR. – Europe (WE, NE, SE, EE), Armenia, Iran.

Dolichoderus sibiricus Emery, 1889 (*D. quadripunctatus sibiricus* Emery, 1889). Russia: ES (KS, KR, BR), FE (AM, KH, PR, KU). – Mongolia, China (NE, NC), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

LIOMETOPUM Mayr, 1861. Type species *Formica microcephala* Panzer, 1798. Number of species: World – 7, Palaearctic – 3, Russia – 2.

Liometopum microcephalum (Panzer, 1798) [Formica]. Russia: EP (S, NC). – Europe (WE, SE, EE).

Liometopum orientale Karawajew, 1927 (*L. microcephalum* var. *orientalis* Karawajew, 1927). Russia: FE (PR). – Korean Peninsula.

TAPINOMA Förster, 1850. Type species *Tapinoma collina* Förster, 1850 =*T. erraticum* (Latrelle, 1798). Number of species: World – 69, Palaearctic – 18, Russia – 7.

Tapinoma christophi Emery, 1925. Russia: **EP** (S).

Tapinoma emeryanum Kuznetsov-Ugamsky, 1927
(*T. emeryanum dagestanicum* Kuznetsov-Ugamsky, 1927). Russia: **EP** (NC). – Turkmenistan, Kazakhstan.

Tapinoma erraticum (Latreille, 1798) [Formica]
(*T. breve* Emery, 1925; *T. tauridis* Emery, 1925). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), Caucasus, Turkmenistan, Kazakhstan.

Tapinoma kinburni Karawajew, 1937. Russia: **EP** (C, S). – Europe (EE).

Tapinoma melanocephalum (Fabricius, 1793)
[Formica]. Tramp species. Russia: **EP** (Moscow, introduced). – Cosmopolitan.

Tapinoma sinense Emery, 1925. Russia: **ES** (IR), **FE** (KH, PR). – Mongolia, China (NE, NC, NW, CC, SW, WP, SE), Korean Peninsula.

Tapinoma subboreale Seifert, 2012 (*T. erraticum ambiguum* Emery, 1925; *T. ambiguum* Emery, 1925, auct.). Russia: **EP** (C, S, NC). – Europe (WE, NE, SE, EE).

Subfamily FORMICINAE

Number of taxa: World – 51 genera, 3050 species, Palaearctic – 23/382, Russia – 12/98.

CAMPONOTUS Mayr, 1861. Type species *Formica ligniperda* Latreille, 1802. Number of species: World – 1021, Palaearctic – 113, Russia – 16.

Camponotus (Camponotus) herculeanus (Linnaeus, 1758) [Formica] (*C. herculeanus* var. *shitkovi* Ruzsky, 1904; *C. herculeanus* var. *altaica* Ruzsky, 1915; *C. herculeanus caucasicus* Arnoldi, 1967). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (TM, OM, NS). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Kyrgyzstan, Kazakhstan.

Camponotus (Camponotus) japonicus Mayr, 1866
(*C. herculeanus* var. *aterrima* Emery, 1894; *C. herculeanus japonicus* var. *manczhuricus* Ruzsky, 1915; *C. japonicus* var. *sanguinea* Karawajew, 1929; *C. japonica* var. *miltotus* Wheeler, 1929; *C. japonica wui* Wheeler, 1929; *C. aterrimus* Emery, 1895: Kupianskaya, 2012). Russia: **ES** (KS, KR, IR, BR, ZB), **FE** (AM, KH, PR, SA, KU). – Mongolia, China (NE, NC, NW), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Camponotus (Camponotus) ligniperda (Latreille, 1802) [Formica]. Russia: **EP** (N, NW, C, NC), **UR**. – Europe (WE, NE, SE, EE), Caucasus, Turkey.

Camponotus (Camponotus) obscuripes Mayr, 1879

(*C. herculeanus ligniperdus* var. *obscuripes* Mayr, 1879). Russia: **FE** (SA, KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Camponotus (Camponotus) sachalinensis Forel, 1904 (*C. herculeanus* var. *sachalinensis* Forel, 1904; *C. herculeanus* var. *wymperi* Forel, 1904; *C. herculeanus* var. *montana* Ruzsky, 1905 (part); *C. herculeanus* var. *altaica* Ruzsky, 1915; *C. herculeanus* var. *eudokiae* Ruzsky, 1926; *C. herculeanus* var. *jacutica* Karawajew, 1929). Russia: **WS** (AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (KH, PR, SA, KU, KA, MG). – Mongolia, China (NE), Korean Peninsula, Japan (Hok, Hon, Shi).

Camponotus (Camponotus) saxatilis Ruzsky, 1895
(*C. herculeanus saxatilis* Ruzsky, 1895; *C. pennsylvanicus saxatilis* Ruzsky, 1905). Russia: **EP** (E, S), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, MG). – Mongolia, China (NE), Korean Peninsula.

Camponotus (Camponotus) vagus (Scopoli, 1763)
[Formica] (*C. vagus* var. *kodorica* Forel, 1913). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL). – Europe (WE, NE, SE, EE), N Africa, Turkey, Kazakhstan.

Camponotus (Myrmentoma) atricolor (Nylander, 1849) [Formica] (*C. merula* Losana, 1834; *C. piceus* (Leach, 1825), part, auct.). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Iran, Kazakhstan.

Camponotus (Myrmentoma) fallax (Nylander, 1856)
[Formica] (*C. marginatus* Latreille, 1798 (part, auct.); *C. marginatus* var. *ruzskyi* Emery, 1898; *C. marginatus* var. *kamensis* Ruzsky, 1903). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Kazakhstan.

Camponotus (Myrmentoma) quadrinotatus Forel, 1886 (*C. marginatus* var. *quadrinotatus* Forel, 1886; *C. marginatus* var. *rubicolor* Ruzsky, 1925; *C. marginatus quadrinotatus* var. *nigricolor* Ruzsky, 1925; *C. marginatus brunni* Ruzsky, 1926). Russia: **FE** (KH, PR). – China (NC, NW, CC, SW, WP, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Camponotus (Myrmentoma) interjectus Mayr, 1877. Russia: **EP** (NC). – Israel, Iraq, Iran, Afghanistan, Central Asia.

Camponotus (Myrmentoma) lateralis (Olivier, 1792) [Formica]. Russia: **EP** (NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Iran, Turkmenistan.

Camponotus (Tanaemyrmex) aethiops (Latrelle, 1798) [Formica] (*F. migrata* Nylander, 1849; *Camponotus aethiops* var. *glaber* Emery, 1925). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran, Afghanistan, Central Asia, Kazakhstan.

Camponotus (Tanaemyrmex) fedtschenkoi Mayr, 1877 (*C. tichomirovi* Ruzsky, 1905; *C. fedtschenkoi* var. *adi* Stitz, 1930). Russia: **EP** (S, NC). – Armenia, Azerbaijan, Syria, Iran, Afghanistan, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan.

Camponotus (Tanaemyrmex) turkestanicus Emery, 1887 (*C. sylvaticus* st. *turkestanicus* Emery, 1887; *C. attila* Stärcke, 1935). Russia: **EP** (S, NC). – Iran, Afghanistan, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan.

Camponotus (Tanaemyrmex) turkestanus André, 1881 (*C. sylvaticus* var. *turkestanus* André, 1882; *C. ferganensis* Pisarski, 1967). Russia: **EP** (S, NC). – Israel, Iran, Afghanistan, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan.

CATAGLYPHIS Förster, 1850. Type species *Cataglyphis fairmairei* Förster, 1850. Number of species: World – 91, Palaearctic – 65, Russia – 3.

Cataglyphis aenescens (Nylander, 1849) [Formica] (*Myrmecocystus cursor* r. *tancrei* Forel, 1901; *M. cursor* var. *caspius* Ruzsky, 1902; *M. altisquamis* var. *jacobsoni* Ruzsky, 1905; *Cataglyphis aenescens aterrima* Pisarski, 1967; *C. aenescens georgica* Arnoldi, 1968; *C. aenescens chatkalenis* Tarbinsky, 1976). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (OM, NS, AL), **ES** (KS, TU, BR). – Europe (SE, EE), Caucasus, Turkey, Iran, Afghanistan, Central Asia, Kazakhstan, Mongolia, China (NE, NC, NW).

Cataglyphis nodus (Brullé, 1833) [Formica] (*Cataglyphis nodus caucasicola* Arnoldi, 1964). Russia: **EP** (NC). – Europe (SE, EE), Caucasus, Turkey, Israel, Saudi Arabia, Iran, United Arab Emirates, Afghanistan, Central Asia.

Cataglyphis pallida Mayr, 1877. Russia: **EP** (S, NC). – Afghanistan, Central Asia, Kazakhstan.

COLOBOPSIS Mayr, 1861. Type species *Formica truncata* Spinola, 1808. Number of species: World – 94, Palaearctic – 1, Russia – 1.

Colobopsis truncatus (Spinola, 1808) [Formica] (*Camponotus (Colobopsis) truncatus* (Mayr, 1861), auct.). Russia: **EP** (C, E, S, NC, CR). – Europe (WE, SE), N Africa, Caucasus, Turkey, Turkmenistan.

FORMICA Linnaeus, 1758. Type species *Formica rufa* Linnaeus, 1761. Number of species: World – 176, Palaearctic – 53, Russia – 32.

Formica (Coptoformica) exsecta Nylander, 1846 (*F. exsecta* var. *rubens* Forel, 1874; *F. nemoralis* Dlussky, 1964). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (IR, BR, ZB), **FE** (AM, KH, SA, KA, MG). – Europe (WE, NE, SE, EE), Mongolia, China (NE).

Formica (Coptoformica) fennica Seifert, 2000. Russia: **EP** (N, NC). – Europe (NE).

Formica (Coptoformica) foreli Bondroit, 1918 (*F. exsecta pressilabris* var. *foreli* Bondroit, 1918; *F. tamarae* Dlussky, 1964). Russia: **EP** (C, NC). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan.

Formica (Coptoformica) forsslundi Lohmander, 1949 (*F. brunneonitida* Dlussky, 1964; *F. fossilabris* Dlussky, 1965). Russia: **EP** (N, C, NC), **ES** (TU). – Europe (WE, NE, EE).

Formica (Coptoformica) manchu Wheeler, 1929 (*F. exsecta* var. *manchu* Wheeler, 1929; *F. longiceps* Dlussky, 1964, nom. praeocc., nec Smith, 1863; *F. dlusskyi* Bolton, 1995). Russia: **ES** (BR, YA, ZB), **FE** (AM, PR). – Mongolia, China (NE), Korean Peninsula.

Formica (Coptoformica) pisarskii Dlussky, 1964. Russia: **ES** (BR, YA, ZB), **FE** (AM). – Mongolia.

Formica (Coptoformica) pressilabris Nylander, 1846 (*F. exsecta pressilabris* var. *rufomaculata* Ruzsky, 1895; *F. exsecta pressilabris* Kuznetsov-Ugamsky, 1928; *F. rufomaculata*: Kupianskaya, 1995). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (AL), **ES** (IR, BR, ZB), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), Turkey, Mongolia, China (NE), Korean Peninsula.

Formica (Coptoformica) suecica Adlerz, 1902. Russia: **EP** (N), **WS** (OM). – Europe (NE).

Formica (Formica) aquilonia Yarrow, 1955. Russia: **EP** (N, NW, C), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, KR, IR, BR, YA, ZB), **FE** (KH, PR, SA, MG). – Europe (WE, NE, SE, EE), Mongolia, China (NE), Korean Peninsula.

Formica (Formica) kupyanskayae Bolton, 1995 (*F. opaca* Kupianskaya, 1980, nom. praeocc., nec Nylander, 1856). Russia: **FE** (AM, KH, PR, SA). – Korean Peninsula.

Formica (Formica) lugubris Zetterstedt, 1838 (*F. congerens* Nylander, 1846; *F. rufa* var. *rufo-pratensis* Forel, 1874; *F. pratensis unicolor* Ruzsky, 1926; *F. rufa* var. *nylanderi* Bondroit, 1919). Russia: **EP** (N, NW, C), **UR**, **WS** (TM, OM, TK, NS, KM, AL),

- ES** (KR, IR, BR, YA, ZB), **FE** (KH, PR, KA, MG). – Europe (WE, NE, SE, EE), Mongolia, China (NE), Korean Peninsula.
- Formica (Formica) polycetena** Förster, 1850. Russia: **EP** (N, NW, C, E, S), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, KR, IR, BR, YA, ZB). – Europe (WE, NE, SE, EE), Kazakhstan.
- Formica (Formica) pratensis** Retzius, 1783 (*F. pratensis* var. *ciliata* Ruzsky, 1926). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, AL), **ES** (KR, IR, BR, ZB), **FE** (AM). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Central Asia, Kazakhstan, Mongolia, China (NE).
- Formica (Formica) rufa** Linnaeus, 1761 (*F. rufa* var. *meridionalis* Nasonov, 1889). Russia: **EP** (N, NW, C, E, S), **UR**, **WS** (TM, OM, NS, AL), **ES** (KS, KR, IR). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan.
- Formica (Formica) truncorum** Fabricius, 1804 (*F. truncicola* Nylander, 1846; *F. rufa* var. *rufotruncicola* Ruzsky, 1896; *F. truncorum* var. *menozzii* Krausse, 1926). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TM, NS, AL), **ES** (KS, TU, KR, IR, BR, ZB), **FE** (AM, KH, PR, SA, KU). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Afghanistan, Central Asia, Kazakhstan, Mongolia, China (NE), Korean Peninsula, Japan (Hon).
- Formica (Formica) yessensis** Forel, 1901 (*F. truncicola* var. *yessensis* Forel, 1901). Russia: **FE** (PR). – China (NW, SE), Korean Peninsula, Japan (Hon).
- Formica (Raptiformica) sanguinea** Latreille, 1798 (*F. dominula* Nylander, 1846; *F. sanguinea* var. *fuscipes* Emery, 1895; *F. sanguinea* var. *mollesonae* Ruzsky, 1903; *F. sanguinea* var. *clarior* Ruzsky, 1905; *F. sanguinea arenicola* Kuznetsov-Ugamskij, 1928). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (OM, TK, NS, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, MG). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Afghanistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE, NW, CC), Korean Peninsula, Japan (Hon).
- Formica (Serviformica) candida** F. Smith, 1878 (*F. fusca picea* var. *implana* Ruzsky, 1914; *F. fusca picea* var. *orientalis* Ruzsky, 1914; *F. picea* auct.). Russia: **WS** (AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, MG). – Kyrgyzstan, Kazakhstan, Mongolia, China (NE, NW, CC), Korean Peninsula, Japan (Hon, Kyu).
- Formica (Serviformica) cinerea** Mayr, 1853 (*F. cinerea* var. *armenica* Ruzsky, 1905; *F. imitans* Ruzsky, 1902, auct.). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, AL). – Europe (WE, NE, SE, EE), Caucasus, Kazakhstan.
- Formica (Serviformica) cinereofusca** Karawajew, 1929 (*F. cinerea* var. *cinereofusca* Karawajew, 1929). Russia: **EP** (NC). – Europe (SE, EE), Caucasus.
- Formica (Serviformica) clarissima** Emery, 1925 (*F. rufibarbis subpilosa* var. *clarior* Ruzsky, 1914, nom. praeocc., nec Ruzsky, 1905; *F. rufibarbis* var. *clarissima* Emery, 1925; *F. subpilosa ruzskyi* Dlussky, 1965: Kupianskaya, 2012). Russia: **ES** (BR, ZB), **FE** (AM, KH). – Mongolia, China (NE).
- Formica (Serviformica) cunicularia** Latreille, 1798 (*F. cunicularia fuscoides* Dlussky, 1967). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (OM, TK, NS, AL). – Europe (SE, EE), Caucasus, Turkey, Iran, Central Asia, Kazakhstan.
- Formica (Serviformica) fusca** Linnaeus, 1758. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, ZB). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Central Asia, Kazakhstan, Mongolia, China (NE), N America (introduced).
- Formica (Serviformica) gagates** Latreille, 1798. Russia: **EP** (C, S, NC, CR). – Europe (WE, SE, EE), Georgia, Turkey.
- Formica (Serviformica) gagatoides** Ruzsky, 1904 (*F. fusca* var. *gagatoides* Ruzsky, 1904). Russia: **EP** (N, NW), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (KH, KA, MG). – Europe (NE), Japan (Hon).
- Formica (Serviformica) glauca** Ruzsky, 1896 (*F. rufibarbis* var. *glauca* Ruzsky, 1896; *F. rufibarbis* var. *katuniensis* Ruzsky, 1915). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (OM, TK, NS, KM, AL), **ES** (KS, KR, IR). – Europe (SE, EE), Caucasus, Turkey, Central Asia, Kazakhstan, Mongolia.
- Formica (Serviformica) japonica** Motschulsky, 1866 (*F. fusca* var. *nipponensis* Forel, 1901; *F. fusca* var. *japonica* Wheeler, 1913). Russia: **FE** (AM, PR, SA, KU). – Mongolia, China (NE, NC, NW, CC, SW, WP, SE), Korean Peninsula, Japan (Hon, Shi, Kyu).
- Formica (Serviformica) lemani** Bondroit, 1917 (*F. fusca* var. *borealis* Waschkewitsch, 1924). Russia: **EP** (N, NW, NC), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KA, MG, CH). – Europe (WE, NE, SE, EE), Mongolia, China (NE), Korean Peninsula, Japan (Hon, Shi).

Formica (Serviformica) picea Nylander, 1846
(*F. transcaucasica* Nasonov, 1889). Russia: **EP** (N, NW, C, E, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL). – Europe (WE, NE, SE, EE), Georgia, Armenia, Turkey, Kazakhstan.

Formica (Serviformica) rufibarbis Fabricius, 1793.
Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (OM, TK, NS, KM, AL). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran.

Formica (Serviformica) subpilosa Ruzsky, 1902
(*F. rufibarbis subpilosa* Ruzsky, 1902; *F. cinerea* var. *bipilosa* Karawajew, 1926). Russia: **EP** (S, NC). – Caucasus, Central Asia.

Formica (Serviformica) uralensis Ruzsky, 1895. Russia: **EP** (C, NC), **UR**, **WS** (OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, ZB), **FE** (AM, PR, SA, KU). – Europe (WE, NE, EE), Mongolia, China (NE), Korean Peninsula.

LASIUS Fabricius, 1804. Type species *Formica nigra* Linnaeus, 1758. Number of species: World – 112, Palaearctic – 55, Russia – 31.

Lasius (Austrolasius) carniolicus Mayr, 1861. Russia: **EP** (C, S), **UR**, **WS** (Radchenko, 2016), **ES** (Radchenko, 2016), **FE** (PR, SA). – Europe (WE, NE, SE, EE), Kazakhstan.

Lasius (Cautolasius) flavus (Fabricius, 1782) [Formica] (*L. flavus* var. *fuscooides* Ruzsky, 1902; *L. flavus odoratus* Ruzsky, 1905; *L. flavus* var. *olivacea* Karawajew, 1926). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU). – Europe (WE, NE, SE, EE), Caucasus, Iran, Kazakhstan, Korean Peninsula, Japan (Hon, Shi, Kyu), N America.

Lasius (Chthonolasius) bicornis (Förster, 1850) [Formica]. Russia: **EP** (C, E, S, NC). – Europe (WE, NE, SE, EE), Caucasus.

Lasius (Chthonolasius) citrinus Emery, 1922 (*L. affinis* Schenck, 1852, nom praeocc., nec Leach, 1825; *L. bicornis* var. *citrine* Emery, 1922; *L. affinis* auct.). Russia: **EP** (C, E, S, NC, CR), **ES** (BR), **FE** (AM, KH, PR, SA, KU). – Europe (WE, SE, EE), Caucasus, Kazakhstan, Korean Peninsula.

Lasius (Chthonolasius) distinguendus (Emery, 1916) [Formicina] (*F. umbrata distinguenda* Emery, 1916; *Lasius hybrida* (Bondroit, 1918)). Russia: **EP** (C, E, S, NC), **ES** (ZB), **FE** (AM, PR). – Europe (WE, SE, EE), Georgia, Armenia, Kazakhstan, Mongolia, Korean Peninsula.

Lasius (Chthonolasius) jensi Seifert, 1982 (*L. umbratus*: Kupianskaya, 1995). Russia: **FE** (AM, PR). – Europe (WE, EE), Armenia, Korean Peninsula.

Lasius (Chthonolasius) longiceps Seifert, 1988. Russia: **EP** (C, S), **UR**. – Europe (EE), Kazakhstan.

Lasius (Chthonolasius) meridionalis (Bondroit, 1920) [Formicina]. Russia: **EP** (C, S, NC), **FE** (PR). – Europe (WE, NE, SE, EE), Georgia, Armenia, Kazakhstan, Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Lasius (Chthonolasius) mixtus (Nylander, 1846) [Formica]. Russia: **EP** (NW, C, E, S, NC, CR), **WS** (TM, OM, TK, NS, KM, AL), **ES** (ZB), **FE** (AM, KH, PR). – Europe (WE, SE, EE), Georgia, Armenia, Kazakhstan, Korean Peninsula.

Lasius (Chthonolasius) sabularum (Bondroit, 1918) [Formicina]. Russia: **EP** (C, S, NC), **ES** (IR). – Europe (WE, NE, SE, EE).

Lasius (Chthonolasius) umbratus (Nylander, 1846) [Formica] (*L. umbratus* var. *exacutus* Ruzsky, 1902). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (ZB), **FE** (AM, PR, SA, KU). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Kazakhstan, Korean Peninsula, Japan (Hok, Hon, Shi, Kyu), N America.

Lasius (Dendrolasius) capitatus (Kuznetsov-Ugamsky, 1928) [Acanthomyops] (*A. fuliginosus capitatus* Kuznetsov-Ugamsky, 1928). Russia: **FE** (PR). – Japan (Hok, Hon, Shi).

Lasius (Dendrolasius) fuji Radchenko, 2005 (*L. fuliginosus*: Kupianskaya, 1995). Russia: **FE** (AM, KH, PR, SA, KU). – China (NE), Korean Peninsula, Japan (Hok, Hon, Shi).

Lasius (Dendrolasius) fuliginosus (Latreille, 1798) [Formica]. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (OM, TK, NS, KM, AL). – Europe (WE, NE, SE, EE), Caucasus, Kazakhstan.

Lasius (Dendrolasius) morisitai Yamauchi, 1978. Russia: **FE** (PR). – Korean Peninsula, Japan (Hon).

Lasius (Dendrolasius) nipponensis Forel, 1912 (*L. fuliginosus* var. *nipponensis* Forel, 1912; *L. crispus* Wilson, 1955). Russia: **FE** (PR). – Korean Peninsula, Japan (Hok, Hon, Shi).

Lasius (Dendrolasius) orientalis Karawajew, 1912 (*L. fuliginosus* var. *orientalis* Karawajew, 1912; *L. teranishii* Wheeler, 1928; *L. ouchii* Teranishi, 1940). Russia: **FE** (AM, PR, SA, KU). – Korean Peninsula, Japan (Hok, Hon).

Lasius (Dendrolasius) spathepus Wheeler, 1910. Russia: **FE** (PR). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Lasius (Lasius) alienus (Förster, 1850) [Formica].

Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (OM, TK, NS, KM, AL), **ES** (BR, ZB), **FE** (AM, KH, PR, SA). – Europe, Caucasus, Turkey, Israel, Iran, Kazakhstan, Mongolia, Japan (Hok, Hon, Shi, Kyu), N America.

Lasius (Lasius) brunneus (Latreille, 1798) [Formica].

Russia: **EP** (C, E, S, NC, CR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Israel, Iran.

Lasius (Lasius) emarginatus (Olivier, 1792) [Formica]

(*L. alienus* var. *pontica* Stärcke, 1944). Russia: **EP** (C, S, NC, CR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Israel, Iran.

Lasius (Lasius) hayashi Yamauchi et Hayashida, 1970. Russia: **FE** (KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).**Lasius (Lasius) japonicus** Santschi, 1941 (*L. emarginatus* var. *japonicus* Santschi, 1941; *L. niger*: Kupianskaya, 1995). Russia: **FE** (AM, KH, PR, SA, KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).**Lasius (Lasius) neglectus** Van Loon, Boomsma et Andrasfalvy, 1990. Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), Turkey, Israel, Iran, Uzbekistan, Kyrgyzstan.**Lasius (Lasius) niger** (Linnaeus, 1758) [Formica] (*L. niger* var. *alienonigra* Forel, 1874). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (BR, ZB). – Europe (WE, NE, SE, EE), Kazakhstan, Mongolia, N America.**Lasius (Lasius) koreanus** Seifert, 1982. Russia: **FE** (KU). – Korean Peninsula.**Lasius (Lasius) obscuratus** Stitz, 1930 (*L. brunneus* var. *obscurata* Stitz, 1930). Russia: **EP** (NC). – Tajikistan.**Lasius (Lasius) paralienus** Seifert, 1992. Russia: **EP** (C, S, NC, CR). – Europe (WE, NE, SE, EE), Turkey.**Lasius (Lasius) platythorax** Seifert, 1991. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL). – Europe (WE, NE, SE, EE), Iran.**Lasius (Lasius) psammophilus** Seifert, 1992. Russia: **EP** (C, S, NC, CR). – Europe (WE, NE, SE, EE).**Lasius (Lasius) turcicus** Santschi, 1921. Russia: **EP** (NC). – Europe (WE, SE, EE), Georgia, Turkey, Iran.**LEPISIOTA** Santschi, 1926 (*Acantholepis* Mayr, 1861). Type species *Plagiolepis rothneyi* Forel, 1894. Number of species: World – 83, Palaearctic – 25, Russia – 1.**Lepisiota semenovi** Ruzsky, 1905 [Acantholepis]

(*Acantholepis frauenfeldi* var. *semenovi* Ruzsky, 1905). Russia: **EP** (NC). – Europe (SE, EE), Caucasus, Turkey, Iran, Central Asia, Kazakhstan.

NYLANDERIA Emery, 1906. Type species *Formica vividula* Nylander, 1846. Number of species: World – 110, Palaearctic – 9, Russia – 3.**Nylanderia colchica** (Pisarski, 1960) [Paratrechina]. Russia: **EP** (NC). – Abkhazia, Georgia.**Nylanderia flavipes** (Smith, 1874) [Tapinoma]. Russia: **EP** (NC, introduced), **FE** (KU). – Abkhazia (introduced), China (NE, NC, CC, SW, WP, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu).**Nylanderia vividula** (Nylander, 1846) [Formica]. Russia: **EP** (C, S, NC, CR) (introduced). – Tramp species. Cosmopolitan.**PLAGIOLEPIS** Mayr, 1861. Type species *Formica pygmaea* Latreille, 1798. Number of species: World – 62, Palaearctic – 14, Russia – 3.**Plagiolepis karawajewi** Radchenko, 1989. Russia: **EP** (CR).**Plagiolepis pallescens** Forel, 1889. Russia: **EP** (NC). – Europe (SE, EE), N Africa, Caucasus, Turkey, Central Asia.**Plagiolepis taurica** Santschi, 1920 (*P. maura* var. *taurica* Santschi, 1920; *P. barbara* var. *pyrenaica* Emery, 1921; *P. satunini* Karawajew, 1931). **EP** (C, E, S, NC, CR). – Europe (WE, SE, EE), Caucasus, Iran, Central Asia, Kazakhstan.**POLYERGUS** Latreille, 1804. Type species *Formica rufescens* Latreille, 1798. Number of species: World – 14, Palaearctic – 3, Russia – 3.**Polyergus nigerrimus** Marikovsky, 1963. Russia: **ES** (TU, BR), **FE** (KH). – Mongolia.**Polyergus rufescens** (Latreille, 1798) [Formica]. Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (OM, AL). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Kyrgyzstan, Kazakhstan.**Polyergus samurai** Yano, 1911. Russia: **FE** (AM, PR). – China (NE, NC, CC, SW, WP, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).**PRENOLEPIS** Mayr, 1861. Type species *Tapinoma nitens* Mayr, 1853. Number of species: World – 13, Palaearctic – 1, Russia – 1.**Prenolepis nitens** (Mayr, 1853) [Tapinoma]. Russia: **EP** (NC, CR). – Europe (WE, SE, EE), Georgia, Turkey.

PROFORMICA Ruzsky, 1902. Type species *Formica nasuta* Nylander, 1856. Number of species: World – 25, Palaearctic – 25, Russia – 3.

Proformica epinotalis Kuznetsov-Ugamsky, 1927. Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (EE), Iran, Turkmenistan, Kyrgyzstan, Kazakhstan.

Proformica mongolica (Emery, 1901) [Formica]. Russia: **ES** (KS, TU, KR, IR, BR, ZB). – Kyrgyzstan, Kazakhstan, Mongolia, China (NE, NC, NW), Korean Peninsula.

Proformica ossetica Dubovikoff, 2005. Russia: **EP** (NC).

ROSSOMYRMEX Arnoldi, 1928. Type species *Rossomyrmex proformicarum* Arnoldi, 1928. Number of species: World – 4, Palaearctic – 4, Russia – 1.

Rossomyrmex proformicarum Arnoldi, 1928. Russia: **EP** (S, NC). – Kazakhstan.

Subfamily MYRMICINAE

Number of taxa: World – 142 genera, 6605 species, Palaearctic – 34/1000, Russia – 23/142.

ANERGATES Forel, 1874. Type species *Myrmica atratula* Schenck, 1852. The social parasites of *Tetramorium* species. Number of species: World – 1, Palaearctic – 1, Russia – 1.

Anergates atratulus (Schenck, 1852) [Myrmica]. The social parasite of *Tetramorium caespitum*. Russia: **EP** (C, S, NC, CR). – Europe (WE, NE, SE, EE), Armenia, Turkey, Kazakhstan, N America (introduced).

APHAENOGASTER Mayr, 1853. Type species *Aphaenogaster sardoa* Mayr, 1853. Number of species: World – 187, Palaearctic – 78, Russia – 8.

Aphaenogaster famelica (Smith, 1874) [Ischnomyrmex] (*A. famelica frontosa* Wheeler, 1928; *A. famelica ruida* Wheeler, 1928; *A. vapidula* Wheeler, 1928). Russia: **FE** (KU). – China (CC, SW, WP, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Aphaenogaster gibbosa (Latreille, 1798) [Formica] (*A. striola* (Roger, 1859); *A. gibbosa muschtaidica* Emery, 1908). Russia: **EP** (NC). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iraq, Iran, Turkmenistan.

Aphaenogaster japonica Forel, 1911 (*A. schmitzi japonica* Forel, 1911; *A. smythiesii sinensis* Wheeler,

1928; *A. sinensis*: Kupianskaya, 1995). Russia: **FE** (KU). – China (CC, SW, WP, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Aphaenogaster kurdica Ruzsky, 1905 (*A. subterranea* var. *kurdica* Ruzsky, 1905). Russia: **EP** (NC). – Georgia, Turkey.

Aphaenogaster obsidiana (Mayr, 1861) [Atta]. Russia: **EP** (NC). – Europe (SE), Georgia, Turkey.

Aphaenogaster splendida (Roger, 1859) [Atta] (*Aphaenogaster splendida transcaucasica* Karawajew, 1926). Russia: **EP** (NC, CR). – Europe (SE), N Africa, Armenia, Azerbaijan, Turkey, Israel, Iran.

Aphaenogaster subterranea (Latreille, 1798) [Formica]. Russia: **EP** (NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey.

Aphaenogaster subterraneoides Emery, 1881 (*A. pallida* var. *subterraneoides* Emery, 1881). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey.

CARDIOCONDYLA Emery, 1869. Type species *Cardiocondyla elegans* Emery, 1869. Number of species: World – 70, Palaearctic – 31, Russia – 5.

Cardiocondyla koshewnikovi Ruzsky, 1902. Russia: **EP** (S, NC). – Kyrgyzstan, Kazakhstan, Mongolia, China (NW).

Cardiocondyla mauritanica Forel, 1890 (*C. nuda* var. *mauritanica* Forel, 1890). Russia: **EP** (CR). – Europe (SE), N Africa, Afghanistan, Turkey, introduced into many countries.

Cardiocondyla sahlbergi Forel, 1913 (*C. elegans* var. *sahlbergi* Forel, 1913). Russia: **EP** (NC). – Europe (EE), N Africa, Caucasus, Turkey, Israel, Iran, Uzbekistan.

Cardiocondyla stambuloffii Forel, 1892 (*C. bogdanovi* Ruzsky, 1905; *C. stambuloffii taurica* Karawajew, 1927). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (SE, EE), Armenia, Turkey, Iran.

Cardiocondyla ulianini Emery, 1889 (*C. elegans* Emery, 1869, auct.; *C. elegans* var. *ulianini* Emery, 1889; *C. elegans schkaffi* Arnoldi, 1928). Russia: **EP** (E, S, NC, CR). – Europe (EE), Caucasus, Saudi Arabia, Iran, Afghanistan, Kyrgyzstan, Kazakhstan, Mongolia.

CHALEPOXENUS Menozzi, 1923 (*Leonomyrma* Arnoldi, 1968). Type species *Chalepoxenus gribodoi* Menozzi, 1923. The social parasites of *Temnothorax* species. Number of species: World – 9, Palaearctic – 9, Russia – 1.

Chalepoxenus tauricus Radchenko, 1989. The social parasite of *Temnothorax tauricus*. Russia: **EP** (CR).

CREMATOGASTER Lund, 1831. Type species *Formica scutellaris* Olivier, 1792. Number of species: World – 506, Palaearctic – 36, Russia – 3.

Crematogaster schmidti (Mayr, 1853) [Acrocoelia]. Russia: **EP** (CR, NC). – Europe (WE, SE, EE), Caucasus, Turkey, Iran, Turkmenistan.

Crematogaster sordidula (Nylander, 1849) [Myrmica] (*C. sordidula* var. *caspica* Emery, 1912). Russia: **EP** (NC). – Europe (WE, SE, EE), Caucasus, Turkey, Israel, Iran, Turkmenistan.

Crematogaster subdentata Mayr, 1877. Russia: **EP** (S introduced, NC, CR introduced). – Caucasus, Turkey, Iran, Afghanistan, Central Asia, Kazakhstan, Mongolia.

FORMICOXENUS Mayr, 1855. Type species *Myrmica nitidula* Nylander, 1846. The social parasite species, in nest of *Formica* s. str. (xenobiotic). Number of species: World – 7, Palaearctic – 2, Russia – 2.

Formicoxenus nitidulus (Nylander, 1846) [Myrmica] (*F. laeviuscula* Förster, 1850). Russia: **EP** (NW, C, E, NC), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (IR, YA, ZB), **FE** (KH, PR). – Europe (WE, NE, SE, EE), Turkey.

Formicoxenus sibiricus (Forel, 1899) [Leptothorax] (*F. orientalis* Dlussky, 1963). Russia: **ES** (BR, ZB), **FE** (PR).

HARPAGOXENUS Forel, 1893 (*Tomognathus* Mayr, 1861). Type species *Myrmica sublaevis* Nylander, 1849. The social parasites of *Leptothorax* species (dulosis). Number of species: World – 3, Palaearctic – 1, Russia – 1.

Harpagoxenus sublaevis (Nylander, 1852) [Myrmica] (*M. hirtula* Nylander, 1849). The social parasite of *Leptothorax acervorum*. Russia: **EP** (N, NW, C, E, NC), **UR**, **ES** (BR, ZB), **FE** (PR). – Europe (WE, NE, SE, EE).

LEPTOTHORAX Mayr, 1855. Type species *Formica acervorum* Fabricius, 1793. Number of species: World – 19, Palaearctic – 9, Russia – 6.

Leptothorax acervorum (Fabricius, 1793) [Formica] (*L. acervorum* var. *superus* Ruzsky, 1905; *L. acervorum* var. *nigrescens* Ruzsky, 1905; *L. acervorum* *kamtschaticus* Ruzsky, 1920; *Mycothorax acervorum* *orientalis* Kuznetsov-Ugamsky, 1928). Russia:

EP (N, NW, C, E, NC), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KA, MG, CH). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Central Asia, Kazakhstan, Mongolia, China (NE, NC), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Leptothorax gredleri Mayr, 1855. Russia: **EP** (NC). – Europe (WE, NE, SE, EE).

Leptothorax muscorum (Nylander, 1846) [Myrmica] (*L. muscorum* var. *fagi* Ruzsky, 1905; *L. muscorum* var. *betulae* Ruzsky, 1916). Russia: **EP** (N, NW, C, E, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Mongolia, China (NE, NC).

Leptothorax oceanicus (Kuznetsov-Ugamsky, 1928) [Mycothorax] (*L. muscorum oceanicum*: Kupianskaya, 1990). Russia: **FE** (AM, KH, PR). – China (NE), Korean Peninsula.

Leptothorax pacis (Kutter, 1945) [Doronomyrmex]. Russia: **EP** (N), **WS** (KM). – Europe (WE, NE, SE).

Leptothorax scamni Ruzsky, 1905 (*L. muscorum* var. *scamni* Ruzsky, 1905). Russia: EP (NC). – Caucasus, Turkey.

MANICA Jurine, 1807. Type species *Formica rubida* Latreille, 1802. Number of species: World – 6, Palaearctic – 2, Russia – 1.

Manica rubida (Latreille, 1802) [Formica]. Russia: **EP** (NC, CR). – Europe (WE, SE, EE), Georgia, Armenia, Turkey.

MESSOR Forel, 1890. Type species *Formica barbara* Linnaeus, 1767. Number of species: World – 106, Palaearctic – 69, Russia – 4.

Messor aciculatus (Smith, 1874) [Aphaenogaster]. Russia: **FE** (PR). – Mongolia, China (NE, NC, NW), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Messor denticulatus Kuznetsov-Ugamsky, 1927 (*M. barbarus meridionalis* var. *denticulatus* Kuznetsov-Ugamsky, 1927). Russia: **EP** (S, NC). – Caucasus, Iran, Central Asia, Kazakhstan.

Messor kasakorum Arnoldi, 1970 (*M. kasakorum nadezhdae* Arnoldi, 1970). Russia: **EP** (S). – Kazakhstan.

Messor structor (Latreille, 1798) [Formica] (*Myrmica mutica* Nylander, 1849; *Messor structor* var. *clivorum* Ruzsky, 1905; *M. tataricus* Ruzsky, 1905; *M. structor turanicus* Kuznetsov-Ugamsky, 1927;

M. rufitarsis jakowlevi Arnoldi, 1977; *M. clivorum* auct., *M. rufitarsis* auct.). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), Caucasus, Israel, Iran, Afghanistan, Central Asia, Kazakhstan.

MONOMORIUM Mayr, 1855. Type species *Monomorium monomorium* Bolton, 1987, replacement name for *Monomorium minutum* Mayr, 1855. Number of species: World – 358, Palaearctic – 76, Russia – 2.

Monomorium barbatulum Mayr, 1877. Russia: **EP** (S). – Israel, Saudi Arabia, Iran, Oman, United Arab Emirates, Afghanistan, Turkmenistan, Kyrgyzstan, Kazakhstan.

Monomorium pharaonis (Linnaeus, 1758) [Formica]. Synanthropic species with cosmopolitan distribution, in Russia only in heated rooms. Russia, introduced. – Cosmopolitan.

MYRMECINA Curtis, 1829. Type species *Myrmecina latrellii* Curtis, 1829. Number of species: World – 53, Palaearctic – 7, Russia – 2.

Myrmecina graminicola (Latreille, 1802) [Formica] (*Myrmica striatula* Nylander, 1849). Russia: **EP** (C, E, S, NC, CR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel.

Myrmecina nipponica Wheeler, 1906 (*M. graminicola nipponica* Wheeler, 1906; *M. graminicola* Kupianskaya, 1995). Russia: **FE** (AM, KH, PR, KU). – China (SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

MYRMICA Latreille, 1804. Type species *Formica rubra* Linnaeus, 1758. Number of species: World – 188, Palaearctic – 75, Russia – 44.

Myrmica ademonia Bolton, 1995 (replacement name for *M. aspersa* Kupianskaya, 1990). Russia: **FE** (KH, PR, KU). – Korean Peninsula.

Myrmica angulinodis Ruzsky, 1905 (*M. scabrinodis angulinodis* Ruzsky, 1905; *M. incurvata* Collingwood, 1976; *M. baikalensis* Dmitrienko, 1979; *M. kasczenkoi*: Kupianskaya, 1979; *M. forcipata*: Kupianskaya, 1990). Russia: **WS** (NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, MG). – Mongolia, China (NE), Korean Peninsula.

Myrmica arnoldii Dlussky, 1963. Russia: **ES** (TU, KR, IR, BR, ZB). – Mongolia.

Myrmica bakurianica Arnoldi, 1970 (*M. pilosiscapus bakurianica* Arnoldi, 1970). Russia: **EP** (NC). – Georgia.

Myrmica bergi Ruzsky, 1902 (*M. rugosa* var. *kirgistica* Ruzsky, 1903; *M. bergi* var. *barchanica* Ruzsky, 1905; *M. bergi* var. *fortior* Crawley, 1920; *M. kamyschiensis* Arnoldi, 1934; *M. bergi persiana* Weber, 1947). Russia: **EP** (S, NC, CR), **WS** (AL). – Europe (EE), Azerbaijan, Iran, Central Asia.

Myrmica commarginata Ruzsky, 1905. Russia: **ES** (KR, BR, ZB). – Mongolia.

Myrmica deplanata Emery, 1921 (*M. lobicornis* var. *deplanata* Emery, 1921; *M. moravica* Soudek, 1922; *M. lobicornis* var. *plana* Karawajew, 1927; *M. moravica* var. *dagmarae* Sadil, 1939). Russia: **EP** (C, S, NC, CR), **WS** (AL). – Europe (WE, SE, EE), Caucasus, Iran, Central Asia.

Myrmica displicentia Bolton, 1995 (*M. bicolor* Kupianskaya, 1986). Russia: **FE** (MG, KA).

Myrmica divergens Karawajew, 1931 (*M. bergi* var. *divergens* Karawajew, 1931). Russia: **WS** (AL), **ES** (KR, IR, BR, YA, ZB). – Mongolia.

Myrmica eidmanni Menozzi, 1930 (*M. lobicornis littoralis* Weber, 1948). Russia: **WS** (AL), **ES** (KS, TU, KR, IR, BR, ZB), **FE** (AM, KH, PR). – Mongolia, China (NE), Korean Peninsula.

Myrmica elbrusi Radchenko et Yusupov, 2012. Russia: **EP** (NC).

Myrmica excelsa Kupianskaya, 1990 (*M. sinica* Wu et Wang, 1995; *M. cadusa* B. Kim, Park et J. Kim, 1997). Russia: **FE** (PR). – China (NE), Korean Peninsula, Japan (Hok, Hon).

Myrmica forcipata Karawajew, 1931. Russia: **ES** (KR, IR, BR, YA, ZB). – Mongolia.

Myrmica gallienii Bondroit, 1920 (*M. rugulosa limanica* Arnoldi, 1934; *M. jacobsoni* Kutter, 1963; *M. limanica obensis* Arnoldi, 1970). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (Radchenko, Elmes, 2010). – Europe (WE, NE, EE).

Myrmica hellenica Finzi, 1926 (*M. rugulosa* var. *hellenica* Finzi, 1926; *M. rugulosa* var. *rugulososcabrinodis* Karawajew, 1929; *M. rugulosa caucasica* Arnoldi, 1934). Russia: **EP** (NC, CR). – Europe (SE, EE), Caucasus, Turkey.

Myrmica hirsuta Elmes, 1978. Russia: **EP** (C). – Europe (WE, NE, SE, EE).

Myrmica jennyae Elmes, Radchenko et Aktac, 2002. Russia: **EP** (NC). – Turkey.

Myrmica kamtschatica Kupianskaya, 1986 (*M. aborigenica* Zhigulskaya, 1991). Russia: **WS** (KM, AL), **ES** (Radchenko, Elmes, 2010), **FE** (AM, KH, PR, MG, KA). – Mongolia, Korean Peninsula.

- Myrmica karavajevi** (Arnoldi, 1930) [Symbiomyrma] (*Sifolinia karavajevi* Arnoldi, 1930). Russia: **EP** (NW, C). – Europe (WE, NE, SE, EE).
- Myrmica kasczenkoi** Ruzsky, 1905 (*M. scabrinodis kasczenkoi* Ruzsky, 1905). Russia: **WS** (KM, AL), **ES** (KS, TU, KR, IR, BR, ZB). – Mongolia.
- Myrmica koreana** Elmes, Radchenko et Kim, 2001. Russia: **WS** (KM, AL), **ES** (KS, TU, KR, IR, BR, ZB), **FE** (AM, KH, PR). – Kazakhstan, Mongolia, China (NE), Korean Peninsula.
- Myrmica kotokui** Forel, 1911 (*M. rubra kotokui* Forel, 1911; *M. ruginodis orientalis* Karawajew, 1926; *M. orientalis*: Kupianskaya, 1990). Russia: **FE** (PR). – China (NE), Korean Peninsula, Japan (Hok, Hon).
- Myrmica kozakorum** Radchenko et Elmes, 2010. Russia: **EP** (S, NC, CR). – Europe (EE).
- Myrmica kurokii** Forel, 1907 (*M. rubra kurokii* Forel, 1907). Russia: **FE** (PR). – China (NE, NC), Korean Peninsula, Japan (Hok, Hon).
- Myrmica lobicornis** Nylander, 1846 (*M. lobicornis alpestris* Arnoldi, 1934). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (Radchenko, 2016), **ES** (KS, TU, KR, IR, BR, YA, ZB). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Kazakhstan, Mongolia.
- Myrmica lonae** Finzi, 1926 (*M. scabrinodis lonae* Finzi, 1926). Russia: **EP** (N, NW, C, E), **UR**, **WS** (Radchenko, 2016), **ES** (BR). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan.
- Myrmica luteola** Kupianskaya, 1990. Russia: **FE** (PR). – China (NE), Korean Peninsula, Japan (Hok, Hon).
- Myrmica pisarskii** Radchenko, 1994. Russia: **WS** (KM, AL), **ES** (KS, TU, KR, IR, BR, ZB). – Mongolia.
- Myrmica ravasinii** Finzi, 1923. Russia: **EP** (NC). – Europe (SE), Georgia, Armenia, Turkey.
- Myrmica rubra** (Linnaeus, 1758) [Formica] (*M. laevinodis* Nylander, 1846; *M. longiscapus* Curtis, 1854; *M. laevinodis* var. *europea* Finzi, 1926; *M. laevinodis* var. *bruesi* Weber, 1947; *M. microrubra* Seifert, 1993). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Central Asia.
- Myrmica ruginodis** Nylander, 1846 (*M. diluta* Nylander, 1849). Russia: **EP** (N, NW, C, E, S, NC, CR) **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, MG), **SA**, **KU**. – Europe (WE, NE, SE, EE), Caucasus, Turkey, Kazakhstan, Mongolia, China (NE, NC), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Myrmica rugulosa** Nylander, 1849. Russia: **EP** (NW, C, E, NC, CR), **UR**. – Europe (WE, NE, SE, EE), Caucasus.
- Myrmica sabuleti** Meinert, 1861. Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (Radchenko, 2016). – Europe (WE, NE, SE, EE), Caucasus, Turkey.
- Myrmica salina** Ruzsky, 1905 (*M. scabrinodis* var. *salina* Ruzsky, 1905). Russia: **UR**, **WS** (Radchenko, Elmes, 2010). – Kazakhstan.
- Myrmica scabrinodis** Nylander, 1846 (*M. scabrinodis* var. *rugulosoides* Forel, 1915; *M. scabrinodis* var. *ahngeri* Karawajew, 1926). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (Radchenko, 2016), **ES** (KR, BR, ZB). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Kazakhstan.
- Myrmica schencki** Viereck, 1903 (*M. rubra* var. *schencki* Viereck, 1903; *M. schenkioides* Boer, Noordijk, 2005). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (Radchenko, 2016), **ES** (KR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Central Asia.
- Myrmica slovaca** Sadil, 1952 (*M. curvithorax* Bondroit, 1920). Russia: **EP** (C, S, NC, CR), **UR**, **WS** (Radchenko, 2016). – Europe (WE, SE, EE), Caucasus, Turkey, Kazakhstan.
- Myrmica specioides** Bondroit, 1918 (*M. scabrinodis* var. *sancta* Karawajew, 1926; *M. sancta tshuliensis* Arnoldi, 1976). Russia: **EP** (C, S, NC, CR), **UR**, **WS** (Radchenko, Elmes, 2010). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Turkmenistan, Kazakhstan.
- Myrmica stangeana** Ruzsky, 1902 (*M. bergi stangeana* Ruzsky, 1902; *M. stangeana medvedevi* Arnoldi, 1970). Russia: **EP** (E, S), **UR**, **WS** (AL). – Europe (EE), Kazakhstan.
- Myrmica sulcinodis** Nylander, 1846 (*M. sulcinodis* var. *nigripes* Ruzsky, 1895; *M. myrmecophila* Wasmann, 1910; *M. sulcinodis* var. *sulcinodoscabrinodis* Forel, 1915). Russia: **EP** (N, NW, C, NC, CR), **UR**, **WS** (Radchenko, 2016), **ES** (Radchenko, 2016), **FE** (AM, KH, PR, SA, MG). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Kazakhstan, Mongolia, Korean Peninsula.
- Myrmica transsibirica** Radchenko, 1994 (*M. taediosa* Bolton, 1995). Russia: **WS** (KM, AL), **ES** (Radchenko, Elmes, 2010), **FE** (AM, KH, PR, SA). – China (NE), Korean Peninsula, Japan (Hok).
- Myrmica tschekanovskii** Radchenko, 1994. Russia: **ES** (KR, YA).
- Myrmica vandeli** Bondroit, 1920 (*M. scabrinodis* var. *vandeli* Bondroit, 1920). Russia: **EP** (NC). – Europe (WE, NE, SE, EE).

Myrmica zojae Radchenko, 1994. Russia: **UR, WS (AL)**.

MYRMOXENUS Ruzsky, 1902 (*Epimyrma* Emery, 1915). Type species *Myrmoxenus gordiagini* Ruzsky, 1902. The social parasites of *Temnothorax* species. Number of species: World – 12, Palaearctic – 12, Russia – 2.

Myrmoxenus gordiagini Ruzsky, 1902 (*Myrmetaerus microcellatus* Soudek, 1925). Russia: **WS (TK, NS, AL)**. – Europe (SE, EE), Kazakhstan.

Myrmoxenus ravouxi (André, 1896) [Formicoxenus] (*Epimyrma goesswaldi* Menozzi, 1930). Russia: **EP (NC, CR)**. – Europe (WE, SE, EE).

PHEIDOLE Westwood, 1839. Type species *Atta providens* Sykes, 1835. Number of species: World – 1004, Palaearctic – 19, Russia – 2.

Pheidole fervida Smith, 1874. Russia: **FE (PR, KU)**. – China (CC, SW, WP, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Pheidole pallidula (Nylander, 1849) [Myrmica] (*Ph. pallidula* var. *arenarum* Ruzsky, 1905; *Ph. pallidula koshevnikovi* Ruzsky, 1905). Russia: **EP (S, NC)**. – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran, Afghanistan, Central Asia, Kazakhstan.

SOLENOPSIS Westwood, 1840 (*Diplorhoptrum* Mayr, 1855). Type species *Solenopsis mandibularis* Westwood, 1840. Number of species: World – 195, Palaearctic – 33, Russia – 5.

Solenopsis deserticola Ruzsky, 1905 [*Diplorhoptrum*] (*D. deserticola* auct.). Russia: **EP (S, NC)**. – Caucasus, Central Asia, Kazakhstan.

Solenopsis fugax (Latreille, 1798) [Formica] (*Myrmica flavidula* Nylander, 1849; *Solenopsis fugax orientalis* Ruzsky, 1905; *S. orbula* var. *latroides* Ruzsky, 1905; *S. (Diplorhoptrum) fugax* var. *pontica* Santschi, 1934; *Diplorhoptrum fugax* auct.). Russia: **EP (C, E, S, NC, CR)**. – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran, Afghanistan, Central Asia, Kazakhstan.

Solenopsis ilinei Santschi, 1936 (*S. orbula* var. *oculata* Karawajew, 1926; *S. (Diplorhoptrum) orbula* var. *ilinei* Santschi, 1936). Russia: **EP (CR)**. – Azerbaijan.

Solenopsis japonica Wheeler, 1928 (*S. fugax* var. *japonica* Wheeler, 1928; *Diplorhoptrum japonica*: Kupianskaya, 1995). Russia: **FE (KU)**. – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Solenopsis juliae (Arakelian, 1991) [*Diplorhoptrum*]. Russia: **EP (NC)**. – Armenia.

STENAMMA Westwood, 1839. Type species

Stenamma westwoodi Westwood, 1839. Number of species: World – 84, Palaearctic – 14, Russia – 6.

Stenamma debile (Förster, 1850) [Myrmica] (*S. westwoodii* Westwood, 1839, auct.; *S. golosejevi* Karawajew, 1926; *S. ucrainicum* Arnoldi, 1928). Russia: **EP (C, E, S, NC, CR)**. – Europe (WE, NE, SE, EE), Armenia, Turkey.

Stenamma georgii Arnoldi, 1975. Russia: **EP (NC)**. – Abkhazia, Georgia.

Stenamma kuriense Arnoldi, 1975. Russia: **FE (KU)**.

Stenamma lippulum (Nylander, 1849) [Myrmica] (*S. westwoodi* var. *hirtula* Emery, 1898; *S. caucasicum* Arnoldi, 1975). Russia: **EP (NC)**. – Abkhazia, Georgia, Azerbaijan.

Stenamma striatulum Emery, 1895 (*S. westwoodi* var. *striatulum* Emery, 1895; *S. westwoodi* var. *tscherkesicum* Arnoldi, 1928). Russia: **EP (NC)**. – Europe (SE, EE), Turkey.

Stenamma ussuricense Arnoldi, 1975. Russia: **FE (PR)**. – Korean Peninsula.

STRONGYLOGNATHUS Mayr, 1853. Type species *Eciton testaceum* Schenck, 1852. The social parasites of *Tetramorium* species (dulosis). Number of species: World – 24, Palaearctic – 24, Russia – 5.

Strongylognathus arnoldii Radchenko, 1985. Russia: **EP (CR)**.

Strongylognathus christophi Emery, 1889 (*S. huberi ruzskyi* Emery, 1909). Russia: **EP (E, S, NC, CR, UR)**. – Caucasus, Kazakhstan.

Strongylognathus karawajewi Pisarski, 1966. Russia: **EP (S, NC, CR)**. – Caucasus, Turkmenistan.

Strongylognathus rehbinderi Forel, 1904. Russia: **EP (S, NC)**. – Abkhazia.

Strongylognathus testaceus (Schenck, 1852) [Eciton] (*Myrmus emarginatus* Schenck, 1853; *Strongylognathus diveri* Donisthorpe, 1936). Russia: **EP (C, E, S, NC, CR, UR, WS (NS, AL))**. – Europe (WE, NE, SE, EE), Caucasus, Turkey, Kazakhstan.

STRUMIGENYS Smith, 1860 (*Pyramica* Roger, 1862; *Epitritus* Emery, 1869). Type species *Strumigenys mandibularis* Smith, 1860. Number of species: World – 837, Palaearctic – 19, Russia – 1.

Strumigenys argiola (Emery, 1869) [Epitritus] (*Pyramica argiola* auct.). Russia: **EP (NC)**. – Europe (WE, SE), N Africa, Caucasus, Turkey, Israel.

TELEUTOMYRMEX Kutter, 1950. Type species *Teleutomyrmex schneideri* Kutter, 1950. Number of species: World – 2, Palaearctic – 2, Russia – 1.

Teleutomyrmex schneideri Kutter, 1950. Russia: **EP** (S). – Europe (WE), Turkmenistan.

TEMNOTHORAX Mayr, 1861. Type species *Myrmica recedens* Nylander, 1856. Number of species: World – 390, Palaearctic – 200, Russia – 29.

Temnothorax affinis (Mayr, 1855) [Leptocephalothorax]. Russia: **EP** (C, E, S, NC, CR). – Europe (WE, NE, SE, EE), Caucasus, Turkey.

Temnothorax alinae (Radchenko, 1994) [Leptocephalothorax]. Russia: **FE** (PR).

Temnothorax alpinus (Ruzsky, 1902) [Leptocephalothorax] (*L. wernerii* (Radchenko, 1994)). Russia: **EP** (NC). – Georgia.

Temnothorax arnoldii Radchenko et Fedoseeva, 2015. Russia: **EP** (NC).

Temnothorax brauneri (Ruzsky, 1905) [Leptocephalothorax] (*L. serviculus* var. *brauneri* Ruzsky, 1905). Russia: **EP** (NC). – Georgia.

Temnothorax clypeatus (Mayr, 1853) [Myrmica]. Russia: **EP** (CR). – Europe (WE, SE, EE).

Temnothorax congruus (Smith, 1874) [Leptocephalothorax]. Russia: **FE** (PR, KU). – China (NE), Korean Peninsula, Japan (Hon, Shi, Kyu).

Temnothorax corticalis (Schenck, 1852) [Myrmica]. Russia: **EP** (S, NC, CR). – Europe (WE, NE, SE, EE), N Africa, Caucasus.

Temnothorax crasecundus Seifert et Csósz, 2015. Russia: **EP** (NC, CR). – Europe (SE, EE), Caucasus, Turkey.

Temnothorax crassispinus (Karawajew, 1926) [Leptocephalothorax] (*L. nylanderii* var. *crassispina* Karawajew, 1926; *L. nylanderii slavonicus* Seifert, 1995). Russia: **EP** (C, E, S). – Europe (WE, SE, EE).

Temnothorax discoloratus (Arnoldi, 1977) [Leptocephalothorax]. Russia: **EP** (NC).

Temnothorax dlusskyi Radchenko, Yusupov et Fedoseeva, 2015. Russia: **EP** (NC).

Temnothorax interruptus (Schenck, 1852) [Myrmica]. Russia: **EP** (NC). – Europe (WE, NE, SE, EE).

Temnothorax jailensis (Arnoldi, 1977) [Leptocephalothorax]. Russia: **EP** (CR).

Temnothorax kaszabi (Pisarski, 1969) [Leptocephalothorax] (*L. tuberum sachalinensis* Kupianskaya, 1990). Russia: **WS** (AL), **ES** (TU, YA, ZB), **FE** (SA). – Mongolia, Korean Peninsula.

Temnothorax knipovitschi (Karawajew, 1916) [Leptocephalothorax] (*L. tuberum knipovitschi* Karawajew, 1916; *L. nikitae* Arnoldi, 1977; *L. junipereti* Arnoldi, 1977). Russia: **EP** (E, S, NC, CR), **UR**. – Europe (EE), Caucasus, Kazakhstan.

Temnothorax kuriensis (Radchenko, 1994) [Leptocephalothorax]. Russia: **FE** (KU).

Temnothorax mongolicus (Pisarski, 1969) [Leptocephalothorax] (*L. serviculus mongolicus* Pisarski, 1969). Russia: **ES** (ZB), **FE** (AM). – Mongolia, Korean Peninsula.

Temnothorax nassonovi (Ruzsky, 1895) [Leptocephalothorax] (*L. nassonovi* var. *subnudus* Ruzsky, 1905; *L. galeatus* Wheeler, 1927; *L. nassonovi firsovi* Kuznetsov-Ugamsky, 1928). Russia: **EP** (E, S), **UR**, **WS** (OM, TK, NS, AL), **ES** (BR, ZB), **FE** (AM, KH, PR, SA). – Europe (EE), Kazakhstan, Mongolia, China (NE, NC), Korean Peninsula.

Temnothorax parvulus (Schenck, 1852) [Myrmica]. Russia: **EP** (NC). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Iran, Turkmenistan.

Temnothorax recedens (Nylander, 1856) [Myrmica]. Russia: **EP** (S, NC, CR). – Europe (SE), N Africa, Caucasus, Turkey, Iran, Turkmenistan.

Temnothorax satunini (Ruzsky, 1902) [Leptocephalothorax]. Russia: **EP** (NC). – Caucasus, Turkey.

Temnothorax semenovi (Ruzsky, 1903) [Leptocephalothorax]. Russia: **EP** (S, NC, CR). – Central Asia.

Temnothorax serviculus (Ruzsky, 1902) [Leptocephalothorax]. Russia: **EP** (E, S), **UR**, **WS** (Radchenko, 1996b), **ES** (Radchenko, 1996b). – Kazakhstan, Mongolia.

Temnothorax tauricus (Ruzsky, 1902) [Leptocephalothorax] (*L. unifasciatus* var. *tauricus* Ruzsky, 1902; *L. pithyusae* Arnoldi, 1977; *L. pithyusae babuganicus* Arnoldi, 1977). Russia: **EP** (NC, CR). – Europe (EE).

Temnothorax tembotovi Radchenko et Yusupov, 2015. Russia: **EP** (NC).

Temnothorax tuberum (Fabricius, 1775) [Formica] (*F. tuberosa* Latreille, 1802; *Leptocephalothorax tuberum stipaceus* Ruzsky, 1905; *L. tuberum* var. *nigriceps* Karawajew, 1930; *L. tuberum alaicus* Tarbinsky, 1976; *L. pratostepposus* Arnoldi, 1977; *L. tuberum acutinodis* Arnoldi, 1977; *L. tuberum ciscaucasicus* Arnoldi, 1977). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (NS, AL). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Central Asia.

Temnothorax unifasciatus (Latreille, 1798) [Formica] (*Leptocephalothorax tuberum kirillovi* Ruzsky, 1905; *L. tuberum* var. *salina* Karawajew, 1937; *L. unifasciatus*

ucrainicus Arnoldi, 1977). Russia: EP (C, E, S, NC, CR), UR. – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Iran, Turkmenistan.

Temnothorax volgensis (Ruzsky, 1905) [Leptocephalus (*L. nassonovi volgensis* Ruzsky, 1905)]. Russia: EP (C, E, S), UR. – Europe (EE), Kazakhstan.

TETRAMORIUM Mayr, 1855. Type species *Formica caespitum* Linnaeus, 1758. Number of species: World – 564, Palaearctic – 120, Russia – 10.

Tetramorium caespitum (Linnaeus, 1758) [Formica]. Russia: EP (N, NW, C, E, S, NC, CR), UR, WS (OM, TK, NS, AL), ES (KS, TU, KR, IR, BR, ZB). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Syria, Lebanon, Central Asia.

Tetramorium chefketi Forel, 1911 (*T. caespitum* var. *chefketi* Forel, 1911; *T. caespitum* var. *sarkissianii* Forel, 1911; *T. caespitum* st. *turcomanicum* Santschi, 1921; *T. taurocaucasicum* Arnoldi, 1968). Russia: EP (S, NC), UR, WS (AL). – Europe (SE, EE), Caucasus, Turkey, Iran, Central Asia, Kazakhstan.

Tetramorium densopilosum Radchenko et Arakelian, 1990. Russia: EP (S, NC). – Georgia, Armenia, Turkey.

Tetramorium ferox Ruzsky, 1903 (*T. caespitum* var. *ferox* Ruzsky, 1903; *T. caespitum* st. *ferox* var. *perspicax* Santschi, 1921; *T. confinis* Radchenko et Arakelian, 1990; *T. perspicax* Radchenko, 1992). Russia: EP (S, NC, CR). – Europe (WE, SE, EE), Caucasus, Turkey.

Tetramorium feroxoides Dlussky et Zabelin, 1985 (*T. bursakovi* Radchenko, 1992). Russia: EP (S, NC). – Caucasus, Turkey, Iran, Central Asia, Kazakhstan.

Tetramorium hungaricum Röszler, 1935 (*T. caespitum* *hungarica* Röszler, 1935). Russia: EP (S, CR). – Europe (WE, EE).

Tetramorium impurum (Förster, 1850) [Myrmica] (*T. caespitum* var. *penninum* Santschi, 1927; *T. staerckeii* Kratochvíl, 1944). Russia: EP (S, NC). – Europe (WE, NE, SE, EE).

Tetramorium inerme Mayr, 1877 (*T. caespitum* var. *inermis* Mayr, 1877). Russia: EP (S, NC). – Iran, Central Asia, Kazakhstan.

Tetramorium moravicum Novák et Sadil, 1941 (*T. rhenanum* Schulz, 1996). Russia: EP (S, NC, CR). – Europe (WE, SE, EE), Caucasus, Turkey.

Tetramorium tsushimaiae Emery, 1925 (*T. caespitum* *jacoti* Wheeler, 1927). Russia: FE (AM, KH, PR, KU). – China (NE), Korean Peninsula, Japan (Hon, Shi, Tsushima, Kyu, Yakushima).

TRICHOMYRMEX Mayr, 1865. Type species *Trichomyrmex rogeri* Mayr, 1865. Number of species: World – 24, Palaearctic – 10, Russia – 1.

Trichomyrmex perplexus (Radchenko, 1997) [Monomorium]. Russia: EP (NC). – Europe (SE), Caucasus, Turkey, United Arab Emirates, Iran.

Subfamily PONERINAE

Number of taxa: World – 142 genera, 6605 species, Palaearctic – 12/33, Russia – 4/7.

BRACHYPONERA Emery, 1900. Type species *Euponera (Brachyponera) croceicornis* Emery, 1900. Number of species: World – 19, Palaearctic – 2, Russia – 1.

Brachyponera chinensis (Emery, 1895) [Ponera] (*P. nigrita chinensis* Emery, 1895). Russia: EP (NC introduced), ?FE (PR). – Abkhazia (introduced), Georgia (introduced), China (CC, SW, WP, SE), Korean Peninsula, Japan (Hon, Shi, Kyu, Ryu).

CRYPTOPONE Emery, 1893. Type species *Cryptopone testacea* Emery, 1893. Number of species: World – 24, Palaearctic – 1, Russia – 1.

Cryptopone ochracea (Mayr, 1855) [Ponera]. Russia: EP (NC, CR). – Europe (WE, SE, EE), Saudi Arabia.

HYPOPONERA Santschi, 1938. Type species *Ponera abeillei* André, 1881. Number of species: World – 152, Palaearctic – 9, Russia – 2.

Hypoponera eduardi (Forel, 1894) [Ponera]. Russia: EP (NC). – Europe (WE, SE, EE), N Africa, Turkey, Israel, Saudi Arabia, Iran, United Arab Emirates.

Hypoponera punctatissima (Roger, 1859) [Ponera]. Tramp species. Russia, in heated buildings. – Cosmopolitan.

PONERA Latreille, 1804. Type species *Formica coarctata* Latreille, 1802. Number of species: World – 57, Palaearctic – 5, Russia – 3.

Ponera coarctata (Latreille, 1802) [Formica] (*P. coarctata* var. *lucida* Emery, 1898; *P. coarctata coarctata colchica* Arnoldi, 1932; *P. taurica* Arnoldi, 1932). Russia: EP (C, E, S, NC, CR), UR. – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran, Kyrgyzstan, Kazakhstan.

Ponera japonica Wheeler, 1906 (*P. japonica* var. *croccea* Santschi, 1941). Russia: FE (PR, KU). – China

(SW, WP, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Ponera testacea Emery, 1895 (*P. coarctata* var. *testacea* Emery, 1895). Russia: EP (S, NC, CR). – Europe (WE, SE, EE).

Subfamily PROCERATIINAE

Number of taxa: World – 3 genera, 144 species, Palaearctic – 1/7, Russia – 1/1.

PROCERATIUM Roger, 1863. Type species *Proceratium silaceum* Roger, 1863. Number of species: World – 83, Palaearctic – 7, Russia – 1.

Proceratium melinum (Roger, 1860) [Ponera] (*Sysphingta europaea* Forel, 1886; *S. fialai* Kratochvíl, Novák et Snoflák, 1944, *S. rossica* Arnoldi, 1930). Russia: EP (S, NC, CR). – Europe (WE, SE, EE), Turkey.

SUPERFAMILY APOIDEA**SECTION SPHECIFORMES – DIGGER WASPS**

Number of taxa: Word – 4 families, 276 genera, 9889 species, Palaearctic – 4/101/2302, Russia – 3/87/684.

**70. FAMILY AMPULICIDAE – AMPULICID
DIGGER WASPS**

YU.N. DANILOV

The cockroach wasps are a small, mainly tropical group. These wasps hunt cockroaches. The wasps may enter houses in search of domestic roaches. After a cockroach is captured and weakly paralyzed, the wasp grasps it by the antennal bases with her mandibles and walking back-wards drags the prey to any suitable niche. The nest is stoppered with bits of leaves, grass, and other debris.

The family is distributed mainly in tropical and subtropical regions. Number of taxa: Word – 6 genera and 202 species, Palaearctic – 3/38, Russia – 2/4.

References. Bohart, Menke, 1976; Mokrousov, 2009; Kolesnikov, 1977; Mokrousov, 2010; Mokrousov, Popov, 2016; Shorenko, 2007.

Subfamily AMPULICINAE

AMPULEX Jurine, 1807 (*Pronaeus* Latreille, 1809; *Lorrhoeum* Shuckard, 1837; *Rhinopsis* Westwood, 1844; *Waagenia* Kriechbaumer, 1874; *Chlorampulix* de Saussure, 1892). Type species *Chlorion compressum* of Latreille and of Fabricius (= *Chlorion compressum* (Fabricius, 1804) = *Sphex compressus* Fabricius, 1781). Number of species: World – 132, Palaearctic – 21, Russia – 1.

Ampulex fasciata Jurine, 1807 (*A. europaea* Giraud, 1858). Russia: EP (CR). – Europe (WE, SE, EE).

DOLICHURUS Latreille, 1809 (*Thyreosphex* Ashmead, 1904). Type species *Pompilus corniculus* Spinola, 1808. Number of species: World – 50, Palaearctic – 16, Russia – 3.

Dolichurus bicolor Lepeletier de Saint Fargeau, 1845. Russia: EP (CR). – Europe (WE, NE, SE, EE), Turkey.

Dolichurus corniculus (Spinola, 1808) [Pompilus] (*Pompilus clavipes* Dahlbom, 1829; *D. dahlbomii* Tischbein, 1852). Russia: EP (NW, C, E, NC). – Europe (WE, NE, SE, EE), Turkey.

Dolichurus haemorrhous A. Costa, 1886. Russia: EP (NC, CR). – Europe (WE, SE), N Africa.

71. FAMILY SPHECIDAE – SPHECID DIGGER WASPS

Yu.N. DANILOV

Sphecidae - mainly large, actively flying, clearly visible insects. Sphecid wasps characterized by caring of posterity. The females make the nests and provide it by provision – paralyzed or dead insects from the orders of Orthoptera, Dycloptera, Lepidoptera, Hymenoptera, and spiders. The adult insects feed on nectar and pollinating the flowering plants. The elongated mouthparts of most Sphecidae allows them to visit flowers with deeply located nectaries. Sphecidae make their nests in the ground, often in sandy soils; the females of *Sceliphron* make the mud-nests and *Chalybion* arrange nests in the ready cavities. Most of Sphecidae prefer the open landscapes; they are rare in forests.

The family is distributed worldwide, mainly in arid, semiarid and tropical regions. Number of taxa: Word – 19 genera and 779 species, Palaearctic – 13/260, Russia – 11/68.

Reference. Bohart, Menke, 1976; Danilov, 2012; Danilov, 2014; Dollfuss, 2010, 2013; Menke, Pulawski, 2000; Nemkov, 2009; Nemkov, 2012.

Subfamily AMMOPHILINAE

AMMOPHILA W. Kirby, 1798 (*Miscus Jurine, 1807; Coloptera Latreille, 1845; Argyrammophila Gussakovskij, 1928; Apycnemia Leclercq, 1961*). Type species *Sphex sabulosus* Linnaeus, 1758. Number of species: World – 234, Palaearctic – 103, Russia – 23.

Ammophila adelpha Kohl, 1884. Russia: EP (S). – Iran, Uzbekistan, Kyrgyzstan, Kazakhstan, China (NW).

Ammophila asiatica Tsuneki, 1971. Russia: WS (AL), ES (TU, BR, ZB). – Mongolia.

Ammophila campestris Latreille, 1809. Russia: EP (C, E, S, NC, CR), UR, WS (OM, TK, NS, AL), ES (TU, KR, IR), FE (AM, PR). – Europe (WE, NE, SE, EE), N Africa, Turkey, Central Asia, Kazakhstan, Mongolia, China (NE, NC, NW), Korean Peninsula.

Ammophila deserticola Tsuneki, 1971. Russia: UR, WS (OM, NS, KM, AL), ES (KS, TU, KR, IR, BR, YA, ZB), FE (AM, KH, PR). – Kyrgyzstan, Mongolia, China (NC).

Ammophila elongata Fischer de Waldheim, 1843. Russia: EP (S), UR, FE (AM). – Turkey, Iran, Turkmenistan, Kazakhstan, Mongolia.

Ammophila gracillima Taschenberg, 1869 (*A. longicollis* Kohl, 1884; *A. debilis* F. Morawitz, 1889). Russia: EP (S). – N Africa, Turkey, Syria, Jordan, Israel, Iran, Afghanistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NE, NC, WP), India, Africa.

Ammophila haladai Dollfuss, 2013. Russia: WS (OM, NS), ES (KS). – Turkey.

Ammophila heydeni Dahlbom, 1845. Russia: EP (C, E, S, NC, CR), UR, WS (OM, NS, AL). – Europe (WE, SE, EE), N Africa, Armenia, Turkey, Syria, Jordan, Israel, Iran, Afghanistan, Central Asia, Kazakhstan, Mongolia, China (NW).

Ammophila hungarica Mocsáry, 1883 (*A. fallax* Kohl, 1884). Russia: EP (NC, CR), WS (OM, NS, AL). – Europe (WE, SE, EE), N Africa, Armenia, Turkey, Syria, Jordan, Iran, Afghanistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Ammophila infesta F. Smith, 1873. Russia: FE (PR). – China (NE, NC, CC), Korean Peninsula, Japan (Hon).

Ammophila mongolensis Tsuneki, 1971. Russia: WS (AL), ES (KS, BR, YA, ZB). – Turkey, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NC).

Ammophila occipitalis F. Morawitz, 1890 (*A. ruficollis* F. Morawitz, 1890). Russia: EP (S, NC). – Jordan, Iraq, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Ammophila pubescens Curtis, 1836 (*Miscus arvensis* Dahlbom, 1843; *A. arnaudi* Tsuneki, 1976). Russia: EP (N, NW, C, E, NC, CR), UR, WS, ES, FE (AM, KH, PR). – Europe (WE, NE, SE, EE), Turkey, Iran, Kazakhstan, Mongolia, China (NE, NC, NW, SW), N America.

Ammophila sabulosa (Linnaeus, 1758) [Sphex] (*A. sabulosa kamtschatica* Gussakovskij, 1932). Russia: EP, UR, WS, ES, FE (KA, MG). – Europe (WE, NE, SE, EE), N Africa, Turkey, Syria, Iran, Central Asia, Kazakhstan, Mongolia, China (NC).

Ammophila sareptana Kohl, 1884. Russia: EP (S, NC, CR). – Europe (EE), Turkey, Kyrgyzstan, Kazakhstan.

Ammophila separanda F. Morawitz, 1891. Russia: EP (S, NC). – Azerbaijan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Ammophila sickmanni Kohl, 1901. Russia: ES (ZB), FE (AM, PR). – Mongolia, China (NE, NC, NW, CC, SW, SE), Korean Peninsula.

Ammophila sinensis Sickmann, 1894. Russia: **EP** (NC), **WS** (NS, AL), **ES** (TU, KR, BR, ZB). – Turkey, Iran, Kazakhstan, Mongolia, China (NC, SE).

Ammophila striata Mocsáry, 1879. Russia: **ES**, **FE** (PR). – Turkey, Kazakhstan. Records in Russia doubtful.

Ammophila tekkensis Gussakovskij, 1930. Russia: **EP** (S). – Iran, Turkmenistan, Uzbekistan, Kazakhstan.

Ammophila terminata F. Smith, 1856 (*A. apicalis* Brullé, 1839; *A. mocsaryi* Frivaldszky, 1876; *A. kirgisica* F. Morawitz, 1891; *A. gegen* Tsuneki, 1971). Russia: **EP** (C, E, S, CR), **UR**, **WS** (OM, NS, AL), **ES** (TU, KR, IR, BR, ZB). – Europe (WE, NE, SE, EE), N Africa, Turkey, Central Asia, Kazakhstan, Mongolia, China (NC, NW).

Ammophila vagabunda F. Smith, 1856 (*A. solowiyofkae* Matsumura, 1911; *A. sabulosa nipponica* Tsuneki, 1967). Russia: **FE** (AM, KH, PR, SA, KU). – Mongolia, China (NE, NC, NW, CC, SW, WP, SE), Korean Peninsula, Japan (Hon, Kyu), SE Asia, India.

Ammophila vetuberosa Q. Li et J. Yang, 1994. Russia: **ES** (IR). – Mongolia, China (NC, NW).

EREMOCHARES Gribodo, 1882. Type species *Eremochares doriae* Gribodo, 1882 (= *Ammophila dives* Brullé, 1833). Number of species: World – 5, Palaearctic – 5, Russia – 1.

Eremochares dives (Brullé, 1833) [Ammophila] (*Parapsammophila retowskii* Konow, 1887). Russia: **EP** (S, NC, CR). – Europe (SE), N Africa, Azerbaijan, Turkey, Syria, Lebanon, Israel, Iraq, Iran, Afghanistan, Pakistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, China (NW), India.

HOPLAMMOPHILA de Beaumont, 1960 (*Micadophila* Tsuneki, 1962). Type species *Ammophila armata* (Illiger, 1807) (= *Sphex armatus* Illiger, 1807). Number of species: World – 4, Palaearctic – 4, Russia – 1.

Hoplammophila aemulans (Kohl, 1901) [Ammophila] (*Ammophila rhinoceros* Strand, 1913). Russia: **FE** (PR). – China (NE, NC, CC, SW, SE), Korean Peninsula, Japan (Hon, Kyu), SE Asia.

PODALONIA Fernald, 1927 (*Psammophila* Dahlbom, 1842). Type species *Ammophila violaceipennis* Lepeletier de Saint Fargeau, 1845. Number of species: World – 66, Palaearctic – 42, Russia – 15.

Podalonia affinis (W. Kirby, 1798) [Ammophila] (*Ammophila nigriventris* Gussakovskij, 1934). Russia:

EP (C, E, S, NC, CR), **UR**, **WS** (OM, NS, AL), **ES** (KS, TU, KR, IR, BR, ZB), **FE** (AM, KH, PR). – Europe (NE, EE), N Africa, Caucasus, Turkey, Jordan, Israel, Iran, Central Asia, Kazakhstan, Mongolia, China (NE, NC, SW), Korean Peninsula.

Podalonia alpina (Kohl, 1888) [Ammophila]. Russia: **EP** (NC), **WS** (AL), **ES** (TU). – Europe (WE, SE, EE), N Africa, Turkey, Iran, Afghanistan, Tajikistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NC).

Podalonia altaicensis (Tsuneki, 1971) [Ammophila]. Russia: **ES** (TU). – Mongolia.

Podalonia asiatica Danilov, 2017. Russia: **ES** (TU, BR). – Mongolia, China (NC, NW, WP).

Podalonia atrocyanea (Eversmann, 1849) [Psammophila]. Russia: **EP** (CR), **UR**, **ES** (KS, TU, ZB). – N Africa, Jordan, Israel, Turkmenistan, Uzbekistan, Kazakhstan, Mongolia, China (NC).

Podalonia caucasica (Mocsáry, 1883) [Psammophila]. Russia: **WS** (OM, NS, AL), **ES** (TU, KR, IR, BR, ZB). – Europe (EE), Georgia, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NC, NW), India.

Podalonia chalybea (Kohl, 1906) [Ammophila]. Russia: **ES** (IR). – Mongolia, China (NC).

Podalonia ebenina (Spinola, 1839) [Ammophila] (*Psammophila micipsa* Morice, 1900). Russia: **EP** (NC). – Europe (WE, SE), N Africa, Caucasus, Turkey, Syria, Jordan, Lebanon, Israel, Iraq, Iran, Afghanistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Podalonia fera (Lepeletier de Saint Fargeau, 1845) [Ammophila] (*Psammophila polita* Mocsáry, 1883; *Ammophila morawitzi* Ed. André, 1886). Russia: **EP** (C, S, NC, CR), **UR**. – Europe (WE, SE, EE), Armenia, Turkey, Syria, Iran, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan.

Podalonia flava (Kohl, 1901) [Ammophila] (*Ammophila rothi* de Beaumont, 1951; *A. obo* Tsuneki, 1971). Russia: **EP** (E, S, NC), **WS** (OM, NS, AL), **ES** (KS, TU, IR, BR, ZB), **FE** (PR, SA). – Europe (SE), N Africa, Turkey, Uzbekistan, Kazakhstan, Mongolia, China (NC).

Podalonia gobiensis (Tsuneki, 1971) [Ammophila]. Russia: **ES** (TU, ZB). – Mongolia, China (NC, NW).

Podalonia hirsuta (Scopoli, 1763) [Sphex] (*Ammophila mervensis* Radoszkowski, 1887). Russia: **EP**, **UR**, **WS**, **ES** (TU, IR, BR, ZB). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Lebanon, Israel, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NC, NW, WP), India.

Podalonia hirsutaffinis (Tsuneki, 1971) [Ammophila]. Russia: **ES** (TU). – Kyrgyzstan, Kazakhstan, Mongolia, China (WP).

Podalonia luffii (E. Saunders, 1903) [Ammophila] (*Psammophila arenaria* Lüderwaldt, 1897). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (NS, AL), **ES** (TU, KR, BR). – Europe (WE, NE, SE, EE), Turkey, Syria, Central Asia, Kazakhstan, Mongolia.

Podalonia tydei (Le Guillou, 1841) [Ammophila]. Russia: **EP** (C, E, S, NC, CR). – Europe (WE, SE, EE), N Africa, Turkey, Syria, Jordan, Israel, Iraq, Iran, Afghanistan, Central Asia, Kazakhstan, Mongolia, China (NW), India, Africa, Australia.

Subfamily SCELIPHRINAE

CHALYBION Dahlbom, 1843 (*Hemicalybion* Kohl, 1918). Type species “*Chalybion caeruleum*” (= *Sphex caeruleus* Linnaeus, 1767 = *S. caeruleus* Linnaeus, 1763 (nec *S. caeruleus* Linnaeus, 1758) = *S. cyaneus* Fabricius, 1775 (nec *S. cyaneus* Linnaeus, 1758) = *Pelopaeus* (sic) *californicus* de Saussure, 1867). Number of species: World – 46, Palaearctic – 11, Russia – 3.

Chalybion japonicum (Gribodo, 1883) [*Pelopaeus* (*Sceliphron inflexum* Sickmann, 1894)]. Russia: **FE** (PR). – China (NE, NC, CC, SE), Korean Peninsula, Japan (Hon, Kyu, Ryu), SE Asia, India.

Chalybion turanicum (Gussakovskij, 1935) [*Sceliphron*]. Russia: **WS** (AL). – Iran, Central Asia, Kazakhstan.

Chalybion walteri (Kohl, 1889) [*Pelopaeus*]. Russia: **EP** (NC). – Europe (SE), N Africa, Turkey, Syria, Jordan, Iran, Central Asia, Kazakhstan.

SCELIOPHRON Klug, 1801 (*Pelopaeus* Latreille, 1802; *Prosceliphron* van der Vecht in van der Vecht et van Breugel, 1968; *Hensenia* Pagliano et Scaramozzino, 1990). Type species *Sphex spirifex* Linnaeus, 1758. Number of species: World – 36, Palaearctic – 13, Russia – 6.

Sceliphron caementarium (Drury, 1773) [*Sphex*]. Russia: **EP** (CR). – Europe (WE, SE, EE), Iran, China (SW), Korean Peninsula, Japan (Hon), N America, India, S America, Australia.

Sceliphron curvatum (F. Smith, 1870) [*Pelopaeus*]. Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), Georgia, Turkey, Iraq, Afghanistan, Tajikistan, Uzbekistan, Kyrgyzstan, Japan (Ryu), India, S America.

Sceliphron deforme (F. Smith, 1856) [*Pelopaeus*] (*Pelopaeus atripes* F. Morawitz, 1888). Russia: **EP** (C, E, NC), **WS** (NS), **ES** (KR), **FE** (AM, KH, PR). – Europe (SE), Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE, NC, CC, SW, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu), SE Asia, India.

Sceliphron destillatorium (Illiger, 1807) [*Pepsis*]. Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (ZB). – Europe (WE, SE, EE), N Africa, Turkey, Syria, Israel, Iran, Afghanistan, Central Asia, Kazakhstan, Mongolia, China (NC, NW, CC, WP).

Sceliphron madraspatanum (Fabricius, 1781) [*Sphex*] (*Pelopaeus tubifex* Latreille, 1809). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), N Africa, Turkey, Syria, Israel, Iraq, Iran, Afghanistan, Central Asia, Kazakhstan, China (NE, NC, CC, SW, WP, SE), Korean Peninsula, Japan (Hon, Kyu, Ryu), SE Asia, India, Africa.

Sceliphron spirifex (Linnaeus, 1758) [*Sphex*]. Russia: **EP** (CR). – Europe (WE, SE, EE), N Africa, Georgia, Turkey, Lebanon, Israel, Iran, Turkmenistan, India, Africa.

Subfamily SPHECINAE

Tribe PRIONYCHINI

CHILOSPHEX Menke in R. Bohart et Menke, 1976. Type species *Sphex argyrius* Brullé, 1833. Number of species: World – 2, Palaearctic – 2, Russia – 1.

Chilosphex argyrius (Brullé, 1833) [*Sphex*] (*Sphex emarginatus* Brullé, 1833). Russia: **EP** (NC). – Europe (WE, SE, EE), N Africa, Turkey, Israel, Iraq, Iran, Turkmenistan.

PALMODES Kohl, 1890. Type species *Chlorion occitanicum* (Lepeletier de Saint Fargeau et Audinet-Serville, 1828) (= *Sphex occitanicus* Lepeletier de Saint Fargeau et Audinet-Serville, 1928). Number of species: World – 20, Palaearctic – 9, Russia – 4.

Palmodes melanarius (Mocsáry, 1883) [*Sphex*] (*Sphex anatolicus* Kohl, 1888; *S. picicornis* F. Morawitz, 1890). Russia: **EP** (E, S, NC, CR). – Europe (SE), N Africa, Georgia, Azerbaijan, Turkey, Syria, Iraq, Iran, Central Asia, Kazakhstan, China (NW), India.

Palmodes occitanicus (Lepeletier de Saint Fargeau et Audinet-Serville, 1828) [*Sphex*]. Russia: **EP** (C,

S, NC, CR), **WS** (AL), **ES** (KS, TU, IR, BR, YA, ZB), **FE** (AM). – Europe (WE, SE, EE), N Africa, Turkey, Syria, Israel, Iran, Central Asia, Kazakhstan, Mongolia, China (NE, NC, CC), Korean Peninsula.

Palmodes orientalis (Mocsáry, 1883) [Sphex]. Russia: **EP** (E, S, NC, CR), **UR**, **WS** (AL), **ES** (TU, KR, BR, ZB). – Turkey, Iran, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NW, SW).

Palmodes strigulosus (A. Costa, 1861) [Sphex]. Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), Georgia, Armenia, Turkey, Israel, Iran, Central Asia, Kazakhstan.

PRIONYX Vander Linden, 1827 (*Priononyx* Dahlbom, 1843; *Enodia* Dahlbom, 1843; *Harpactopus* F. Smith, 1856; *Parasphecia* F. Smith, 1856; *Gastrosphaeria* A. Costa, 1858; *Calosphecia* Kohl, 1890). Type species *Ammophila kirbii* Vander Linden, 1827. Number of species: World – 56, Palaearctic – 28, Russia – 6.

Prionyx kirbii (Vander Linden, 1827) [*Ammophila*] (*Sphex albiseptus* Lepeletier de Saint Fargeau et Audinet-Serville, 1828). Russia: **EP** (C, S, NC, CR), **WS** (AL). – Europe (WE, SE, EE), N Africa, Georgia, Turkey, Syria, Israel, Iran, Central Asia, Kazakhstan, China (NC, SE), Africa.

Prionyx lividocinctus (A. Costa, 1861) [*Enodia*]. Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), N Africa, Turkey, Central Asia, Kazakhstan, Mongolia, China (NC).

Prionyx niveatus (Dufour, 1854) [Sphex] (*Podium maracandicum* Radoszkowski, 1877; *Sphex suavis* F. Morawitz, 1893; *S. afghaniensis* de Beaumont, 1970). Russia: **EP** (S). – Europe (SE), N Africa, Turkey, Jordan, Israel, Iran, Afghanistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, Africa.

Prionyx nudatus (Kohl, 1885) [Sphex]. Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (NS, AL), **ES** (TU). – Europe (SE, EE), N Africa, Turkey, Iran, Afghanistan, Central Asia, Kazakhstan, China (NC).

Prionyx subfuscatus (Dahlbom, 1845) [Sphex] (*Sphex desertorum* Eversmann, 1849). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (TU, IR, BR), **FE** (PR). – Europe (WE, SE, EE), N Africa, Georgia, Turkey, Israel, Iran, Central Asia, Kazakhstan, Mongolia, China (NE, NC, NW, CC, SW, WP, SE), Korean Peninsula, India, Africa.

Prionyx viduatus viduatus (Christ, 1791) [Sphex] (*Sphex micans* Eversmann, 1849; *S. leuconotus* F. Morawitz, 1890). Russia: **EP** (C, S, NC, CR), **UR**, **WS** (OM). – Europe (WE, SE, EE), N Africa, Turkey, Israel, Iran, Afghanistan, Central Asia, Kazakhstan, China (NE, NC, NW, SE), Japan (Ryu), SE Asia, India, Africa.

Prionyx viduatus mocsaryi (Kohl, 1883) [Sphex] (*Enodia argentata* Mocsáry, 1883; *Sphex gobiensis* Tsuneki, 1971). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (TU). – Europe (SE, EE), N Africa, Turkey, Turkmenistan, Tajikistan, Kazakhstan, Mongolia, China (NC).

Tribe SPHECINI

ISODONTIA Patton, 1880. Type species *Isodontia philadelphica* (Lepeletier de Saint Fargeau, 1845) (= *Sphex philadelphicus* Lepeletier de Saint Fargeau, 1845). Number of species: World – 62, Palaearctic – 6, Russia – 3.

Isodontia mexicana (de Saussure, 1867) [Sphex] (*Sphex apicalis* de Saussure, 1867). Russia: **EP** (CR). – Europe (WE, SE, EE), Iran, N America.

Isodontia nigella (F. Smith, 1856) [Sphex]. Russia: **FE** (PR). – China (NE, NC, CC, SW, SE), Korean Peninsula, Japan (Hon, Shi, Kyu, Ryu), India, Australia.

Isodontia paludosa (Rossi, 1790) [Sphex]. Russia: **EP** (NC). – Europe (WE, SE, EE), N Africa, Turkey, Jordan, Iran, Turkmenistan.

SPHEX Linnaeus, 1758. Type species *Sphex flavipennis* Fabricius, 1793. Number of species: World – 130, Palaearctic – 20, Russia – 5.

Sphex atropilosus Kohl, 1885 (*Sphex atrohirtus* Kohl, 1890). Russia: **EP** (C, E, S, NC), **WS** (NS). – Europe (WE, SE, EE), N Africa, Turkey, Turkmenistan, Tajikistan, Kazakhstan.

Sphex flavipennis Fabricius, 1793 (*Sphex rufocinctus* Brullé, 1833). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), N Africa, Georgia, Turkey, Syria, Israel, Iran, Afghanistan, Central Asia, Kazakhstan, China (NW).

Sphex funerarius Gussakovskij, 1934 (*Sphex maxillosus* Fabricius, 1793). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (OM, TK, NS, AL), **ES** (TU, KR, IR, BR, ZB), **FE** (KH). – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, Syria, Israel, Iran,

Afghanistan, Central Asia, Kazakhstan, Mongolia, China (NC, NW).

Sphex leuconotus Brullé, 1833 (*Sphex afer* Lepetier de Saint Fargeau, 1845; *S. sordidus* Dahlbom, 1845; *S. tristis* Kohl, 1885; *S. plumipes* Radoszkowski, 1886; *S. pachysoma* Kohl, 1890). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), N Africa, Azerbaijan,

Turkey, Syria, Jordan, Israel, Iraq, Iran, Afghanistan, Central Asia, Kazakhstan.

Sphex pruinosus Germar, 1817. Russia: **EP** (S, NC). – Europe (WE, SE, EE), N Africa, Turkey, Syria, Lebanon, Israel, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, India, Africa.

72. FAMILY CRABRONIDAE – CRABRONID DIGGER WASPS

A.V. ANTROPOV

The family Crabronidae contains a large variety of species, which are very diverse externally – from robust Oxybelini to slender forms with strongly elongated abdomen in Trypoxylini. The body size is also variable from 2 mm in Miscophini, Ammoplanini, or Eremiaspheiinae to 50 mm in Bembicini. Their color varies from completely black or black with red elements to completely light yellow. Such diverse, from the first view, composition of the family is not a sign of artificiality of the taxon, but might speak of its relatively long evolution preserving some basal representatives. Proceeding from the latest data, the family Crabronidae is a monophyletic taxon, a sister-group of “bees” (Apidae s.lat.), with which they, obviously, have a common ancestor. An analysis of the structure of mouthparts and internal skeleton, and the discovery of representatives of the group in the fossil deposits and resins of different ages, starting at least from the Cretaceous, confirm this viewpoint.

The characteristic feature of Crabronidae consists in taking care of the offspring, usually expressed in building the nests before provisioning and oviposition. However, the commonly used name of the family – “sand wasps”, does not reflect the whole diversity of their nesting places. Only a part of species dig nests in the soil, while the majority uses hollow stems of reeds and grasses, thin branches with a soft core, abandoned nests of other Hymenoptera in ground and wood, galls, and abandoned burrows of xylophages in dead wood. Some Crabronidae are capable to gnaw burrows in rotten or even hard wood, and others build separate or combined in groups cells or galleries using wet clay and even the plant fluff. Moreover, among both terricolous and xylicolous Crabronidae there are species with presocial behavior, uniting in nesting clusters, and even using a common entrance to the complex of independent nests. Perhaps the most interesting features of crabronid biology is the nesting behavior including the guarding nesting sites by the males of *Oxybelus* and *Palarus* and directly the nests by the youngest generation in *Cerceris* and by males in some *Pison* and *Trypoxyylon*. The latter can even participate in nest building. Finally, the species of the genus *Microstigmus* demonstrate eusocial behavior, forming the families with the division of functions between their

members. At the same time, the species of *Larra* does not build nests, temporarily paralyzing their victims (mole crickets), and their larvae develop as ectoparasites on a living host.

The vast majority of crabronids is active hunters, which ensure the development of their larvae with paralyzed, usually juvenile, insects of almost all free-living orders, as well as spiders. The degree of specialization in the choice of prey in various groups of Crabronidae varies from orders and families to genera, and only in rare cases to species or certain sex. Usually each cell in the nest contains a number of prey, sufficient for the full development of the larva, but the progressive feeding of each larva to its pupation is characteristic for some species. Finally, an unusual and unique among apoid wasps *Krombeinictus nordenae* Leclercq, 1996 from Sri Lanka is coadapted with internodes of acacia *Humboldtia laurifolia* Vahl (Fabaceae), and its female rears one larva at a time and feeds it progressively with clumped masses of fine-grained pollen rather than paralyzed arthropods.

The main enemies of crabronids are their nest parasites – flies (Anthomyiidae, Asilidae, Bombyliidae, Milichiidae, Phoridae, Sarcophagidae, Tachinidae), hymenopterans (Braconidae, Chrysidae, Eulophidae, Eurytomidae, Mutilidae, Pteromalidae, Torymidae), beetles (Cleridae, Dermestidae, Meloidae, Rhipiphoridae), and ticks (Winterschmidtidae). Sometimes the larvae may be destroyed by various Ichneumonidae, who are usually the parasites of the provisioned prey. At the same time, several nyssonins and bembicins are the cleptoparasites of the related or remote crabronid genera: *Epinysson* of *Hoplisoides* (Bembicini), *Nysson* of *Gorytes*, *Argogorytes*, *Oryttus*, *Dienoplus*, *Hoplisoides*, and *Lestiphorus* (Bembicini), *Brachystegus* and *Zanysson* of *Tachytes* (Larrini), *Acanthostethus* of *Sericophorus* (Miscophini), *Metanysson* of *Cerceris* (Cercerini), and *Stizoides* of *Palmodes*, *Prionyx* (Prionychini), *Sphecius* (Sphecini), and *Bembix* (Bembicini). Adults of many crabronids may be the victims of the crabronid wasps *Palarus* and *Philanthus*. *Palarus variegatus* (Fabricius, 1781) can even successfully attack the females of bee-wolves as it is the only wasp known to be immune to the venom of *Philanthus triangulum* (Fabricius, 1775).

Many species of Crabronidae are effective regulators of quantity of various harmful insects, such as Homoptera, Orthoptera, and Diptera. At the same time, some species of *Palarus* and *Philanthus* which hunt honey bees may cause serious damage to apiculture.

Palarus species may also be harmful when hunting various pollinators and entomophagous wasps.

Crabronidae is distributed worldwide, mainly in tropical and subtropical regions. Crabronids, including 90 percentages of apoid wasps, is the largest group of insects among all families of wasps. Number of taxa: World – 250 genera, about 8900 species; Palaearctic – 84/2000; Russia – 74/613.

References. Brothers, 1975; Bohart, Menke, 1976; Lomholdt, 1982; Gauld, Bolton, 1988; Alexander, 1990, 1992; Melo, 1999; Pulawski, 2016.

Subfamily ASTATINAE

P.G. NEMKOV

ASTATA Latreille, 1796 (*Astatus* Latreille, 1796; *Dimorpha* Panzer, 1806). Type species *Tiphia abdominalis* Panzer, 1798 (= *Sphex boops* Schrank, 1781). Number of species: World – 82, Palaearctic – 35, Russia – 11.

Astata apostata Mercet, 1910. Russia: EP (C, S). – Europe (WE, SE, EE), N Africa, Turkey, Iran, Kazakhstan.

Astata boops boops (Schrank, 1781) [Sphex] (*Tiphia abdominalis* Panzer, 1798; *Dimorpha oculata* Jurine, 1807; *Astata victor* Curtis, 1829; *A. vanderlindeni* Robert, 1833; *A. fuscipes* Robert, 1833, unavailable (ICZN, 1999, Art. 11.5); *A. agilis* F. Smith, 1875). Russia: EP (N, NW, C, E, S, NC, CR), UR, ES (TK, NS, AL), WS (KR, IR, YA, ZB), FE (AM, KH, PR). – Europe (WE, NE, SE, EE), N Africa, Abkhazia, Turkey, Israel, Yemen, Oman, Iraq, Iran, Afghanistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NE, NC), Korean Peninsula, India.

Astata brevitarsis Pulawski, 1958. Russia: EP (S, CR). – Europe (SE, EE), Turkey.

Astata costae costae A. Costa, 1867 (*Astatus gracilis* Gussakovskij, 1927). Russia: EP (S, CR), ES (ZB). – Europe (WE, SE, EE), N Africa, Turkey, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Astata jucunda Pulawski, 1959. Russia: EP (S, NC, CR), WS (AL). – Europe (SE, EE), Turkey, Iran.

Astata kashmirensis kashmirensis Nurse, 1909 (*A. stecki* de Beaumont, 1942). Russia: EP (C, S, NC, CR), WS (AL), FE (PR). – Europe (WE, SE, EE), N Africa, Turkey, Iran, Turkmenistan, Uzbekistan, Kazakhstan, China (NE), India.

Astata miegii scapularis (Kohl, 1889) [Astatus] (*Astatus pelops* Morice, 1902). Russia: EP (S, CR). – Europe (SE, EE), Armenia, Turkey.

Astata minor Kohl, 1885. Russia: EP (N, NW, C, E, S, CR), WS (NS, AL), ES (IR), FE (AM, PR). – Europe (WE, NE, SE, EE), N Africa, Turkey, Israel, Iran, Turkmenistan, Uzbekistan, Kazakhstan.

Astata quettae Nurse, 1903 (*Dimorpha fletcheri* R. Turner, 1917; *Astatus hirsutulus* Gussakovskij, 1927; *A. tibialis* Gussakovskij, 1927; *Astata hungarica* Pulawski, 1958). Russia: EP (S), WS (AL). – Europe (EE), Turkey, Iran, Pakistan, Kazakhstan, Mongolia, China (NW), India.

Astata pontica Pulawski, 1958. Russia: EP (S). – Europe (SE, EE), Turkey.

Astata rufipes rufipes Mocsáry, 1883 (*Astatus sareptanus* Gussakovskij, 1927). Russia: EP (S, CR). – Europe (WE, SE, EE), Turkey, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

DIPLOPLECTRON W. Fox, 1893. Type species *Liris brunneipes* Cresson, 1881. Number of species: World – 20, Palaearctic – 4, Russia – 2.

Diploplectron asiaticum Pulawski, 1965. Russia: EP (S). – Israel, Turkmenistan, Kazakhstan, Mongolia.

Diploplectron pulawskii Kazenas, 1975. Russia: EP (S). – Kazakhstan.

DRYUDELLA Spinola, 1843. Type species *Dryudella ghilianii* Spinola, 1843 (= *Astata tricolor* Vander Linden, 1829). Number of species: World – 57, Palaearctic – 45, Russia – 7.

Dryudella lineata Mocsáry, 1879. Russia: EP (C, E, S). – Europe (EE), Kazakhstan.

Dryudella picticornis (Gussakovskij, 1927) [*Astata*]. Russia: EP (S), UR. – Europe (EE), Turkey, Kazakhstan.

Dryudella pinguis (Dahlbom, 1832) [Larra] (*Larra pinguis* Zetterstedt, 1838, nom. praeocc., nec *L. pinguis* Dahlbom, 1832; *Astata jaculator* F. Smith, 1845). Russia: EP (C, E), WS (AL), ES (IR, YA), FE (SA, KA). – Europe (WE, NE, EE), Kazakhstan, N America.

Dryudella pseudofemoralis Nemkov, 1990. Russia: ES (ZB).

Dryudella similis (Gussakovskij, 1927) [Astatus]. Russia: EP (E, S). – Kazakhstan, China (NW).

Dryudella stigma stigma (Panzer, 1809) [Dimorpha] (*Astata intermedia* Dahlbom, 1843; *A. stigma major* Grönblom, 1946). Russia: EP (N, NW, C, E), WS

(AL), ES (KR, IR, BR, YA, ZB), FE (PR). – Europe (WE, NE, SE, EE), Kazakhstan.

Dryudella tricolor tricolor (Vander Linden, 1829) [Astata] (*D. ghilianii* Spinola, 1843; *Dimorpha cincta* Spinola, 1834; *Dryudella spinolae* A. Costa, 1867; *Astata calopteryx* Gussakovskij, 1952). Russia: EP (C, E, S, NC), WS (AL), ES (KR, ZB). – Europe (WE, SE, EE), N Africa.

Subfamily BEMBICINAE

P.G. NEMKOV

Tribe ALYSSONTINI

ALYSSON Panzer, 1806 (*Alysson Jurine* in Panzer, 1801, suppressed by ICBN, 1939 (Opinion 135); *Alyson Jurine*, 1807). Type species *Pompilus spinosus* Panzer, 1801. Number of species: World – 39, Palaearctic – 14, Russia – 7.

Alysson cameroni Yasumatsu et Masuda, 1932. Russia: FE (KU). – Japan (Hon).

Alysson monticola ezoensis Tsuneki, 1977. Russia: FE (KU). – Japan (Hon).

Alysson pertheesi pertheesi Gorski, 1852 (*A. festivus* Mocsáry, 1879; *A. fuscatus jaroslavensis* Kokujev, 1906; *A. harbinensis* Tsuneki, 1967; *A. pertheesi typica* Tsuneki et Itami, 1973; *A. pertheesi nigra* Tsuneki et Itami, 1973). Russia: EP (NW, E, S), WS (AL), ES (KR, IR, BR), FE (AM, KH, PR, SA, KU). – Europe (WE, NE, EE), Mongolia, China (NE), Korean Peninsula, Japan (Hon).

Alysson ratzeburgi Dahlbom, 1843. Russia: EP (N, NW, E), WS (TM), ES (KS, KR, IR, ZB), FE (PR). – Europe (WE, NE, SE, EE), Korean Peninsula, Japan (Hon).

Alysson spinosus (Panzer, 1801) [*Pompilus*] (*Sphex bimaculatus* Panzer, 1798; *A. katkovi* Kokujev, 1906). Russia: EP (NW, C, E, S, NC). – Europe (WE, NE, SE, EE), Georgia, Turkey, Kazakhstan.

Alysson tricolor Lepeletier de Saint Fargeau et Audinet-Serville, 1825. Russia: EP (C). – Europe (WE, SE, EE), Turkey.

Alysson verhoeffi Tsuneki, 1967. Russia: FE (AM, KH, PR). – China (NE), Korean Peninsula.

DIDINEIS Wesmael, 1852. Type species *Alyson lunicornis* of Vander Linden, 1829 (= *Pompilus lunicornis* Fabricius, 1798). Number of species: World – 24, Palaearctic – 14, Russia – 7.

Didineis bactriana Gussakovskij, 1937 (*D. ogloblini* Gussakovskij, 1937). Russia: EP (S). – Turkmenistan, Uzbekistan, Mongolia.

Didineis bucharica Gussakovskij, 1937 (*D. zimini* Gussakovskij, 1937). Russia: EP (S, NC), UR. – United Arab Emirates, Uzbekistan.

Didineis clavimana Gussakovskij, 1937 (*D. turanica* Gussakovskij, 1937). Russia: EP (S, CR). – Europe (SE, EE), Israel, Uzbekistan.

Didineis crassicornis Handlirsch, 1888. Russia: EP (C, NC). – Europe (WE, SE, EE), N Africa, Turkey.

Didineis lunicornis (Fabricius, 1798) [*Pompilus*] (*Alyson kennedii* Curtis, 1836; *Didineis kozhevnikovi* Kokujev, 1906; *D. ruthenica* Gussakovskij, 1937). Russia: EP (C, E, S), UR. – Europe (WE, NE, SE, EE), N Africa, Abkhazia.

Didineis sibirica Gussakovskij, 1937 (*Alysson sibiricus nipponicus* Tsuneki, 1968). Russia: ES (IR, BR, ZB), FE (PR). – Japan (Hon).

Didineis wuestneii Handlirsch, 1888. Russia: EP (S). – Europe (SE, EE), Azerbaijan, Turkey.

Tribe NYSSONINI

BRACHYSTEGUS A. Costa, 1859. Type species *Nysson dufouri* Lepeletier, 1845 (= *Nysson scalaris* Illiger, 1807). Number of species: World – 21, Palaearctic – 4, Russia – 1.

Brachystegus scalaris (Illiger, 1807) [*Nysson*] (*Nysson interruptus* Latreille, 1803, nom. praeocc., nec *N. interruptus* (Fabricius, 1798); *N. rufipes* Olivier, 1812; *N. auratus* Schummel, 1834; *N. dufourii* Dufour, 1841; *N. dufourii* Lepeletier de Saint Fargeau, 1845, nom. praeocc., nec *N. dufourii* Dufour, 1841; *N. dufouri* Dahlbom, 1845, nom. praeocc., nec *N. dufourii* Dufour, 1841; *N. dufourianum* Blanchard, 1849; *N. transcaspicus* Radoszkowski, 1893). Russia: EP (C, E, S, NC, CR), WS (AL). – Europe (WE, SE, EE), N Africa, Georgia, Turkey, Yemen, Iran, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan.

NIPPONONYSSON Yasumatsu et Maidl, 1936. Type species *Nippononysson rufopictus* Yasumatsu et Maidl, 1936. Number of species: World – 3, Palaearctic – 2, Russia – 1.

Nippononysson rufopictus Yasumatsu et Maidl, 1936. Russia: FE (AM, PR, KU). – Japan (Hon, Kyu, Ryu).

NYSSON Latreille, 1802, validated by ICBN, 1979 (Opinion 1115) (*Nyso* Latreille, 1796, rejected by

- ICZN, 1979 (Opinion 1115); *Nyssonus* Rafinesque, 1815; *Synneurus* A. Costa, 1859; *Synneurus* Gerstaecker, 1867). Type species *Crabro spinosus* Fabricius, 1775 (= *Sphex spinosus* J. Forster, 1771). Number of species: World – 101, Palaearctic – 59, Russia – 19.
- Nysson amurensis** Nemkov, 199. Russia: **FE** (AM, PR).
- Nysson decemmaculatus** Spinola, 1807 (*N. variolatus* A. Costa, 1867; *N. variolatus* A. Costa, 1869; *N. curtulus* F. Morawitz, 1892; *N. mopsus* Handlirsch, 1898). Russia: **EP** (CR). – Europe (WE, SE, EE), Azerbaijan, Turkey, Jordan, Israel, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.
- Nysson dimidiatus** Jurine, 1807 (*N. wesmaeli* Lepeletier de Saint Fargeau, 1845; *N. distinguendus* Chevrier, 1867; *N. nigrina* Barovskij, 1929; *N. susterai* Zavadil in Zavadil et Šnoflák, 1948; *N. susterai* Zavadil, 1951). Russia: **EP** (N, NW, C, S, NC), **WS** (AL), **ES** (IR, ZB), **FE** (AM, SA, MG). – Europe (WE, NE, SE, EE), Armenia, Central Asia, Kazakhstan, Mongolia.
- Nysson epeoliformis** epeoliformis F. Smith, 1856 (*Synneurus procerus* A. Costa, 1859; *Nysson notabilis* Handlirsch, 1895). Russia: **EP** (CR). – Europe (SE), Armenia, Azerbaijan, Turkey, Turkmenistan.
- Nysson fulvipes** A. Costa, 1859. Russia: **EP** (C, S, NC). – Europe (WE, SE, EE), N Africa, Turkey, Israel, Abkhazia, Kazakhstan.
- Nysson fulviventris** Tsuneki, 1971. Russia: **ES** (ZB). – Mongolia.
- Nysson guichardi** de Beaumont, 1967. Russia: **EP** (S, CR). – Europe (SE), Turkey.
- Nysson harveyi** de Beaumont, 1967. Russia: **EP** (NC). – Turkey, Uzbekistan.
- Nysson interruptus** (Fabricius, 1798) [*Mellinus*] (*Ceropales spinosus* Fabricius, 1804; *N. panzeri* Lepeletier de Saint Fargeau, 1845; *N. shukardi* Wesmael, 1852). Russia: **EP** (N, NW, C, E, CR), **UR**, **WS** (NS). – Europe (WE, NE, SE, EE), N Africa, Turkey, Israel, Armenia, Kazakhstan.
- Nysson maculosus** (Gmelin, 1790) [Sphex] (*Sphex maculatus* Fabricius, 1787; *Mellinus dissectus* Panzer, 1800; *N. guttatus* Olivier, 1812; *N. omissus* Dahlbom, 1845; *N. lineolatus* Schenck, 1857; *N. dubius* A. Costa, 1859). Russia: **EP** (N, NW, C, E, CR), **UR**, **WS** (TM, KM, AL), **ES** (KR, IR, YA, ZB), **FE** (AM, KH, PR, SA). – Europe (WE, NE, SE, EE), N Africa, Georgia, Armenia, Turkey, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NE, NC), Korean Peninsula.
- Nysson militaris** Gerstaecker, 1867. Russia: **EP** (S, NC, CR). – Europe (SE, EE), Turkey, Israel.
- Nysson mimulus** Valkeila, 1964 (*N. handlirschi* Westerlund, 1896, nom. nud.; *N. handlirschi* Hellén, 1955). Russia: **EP** (NW). – Europe (WE, NE, EE).
- Nysson niger niger** Chevrier, 1868. Russia: **EP** (N, NW, C, E), **WS** (TM). – Europe (WE, NE, SE, EE), Tajikistan, Kazakhstan.
- Nysson quadriguttatus** Spinola, 1807 (?*N. quadriguttatus* Olivier, 1812). Russia: **EP** (C, E, S), **WS** (AL). – Europe (WE, SE, EE), Uzbekistan, Kyrgyzstan, Kazakhstan.
- Nysson roubali** Zavadil, 1937. Russia: **EP** (S). – Europe (SE, EE).
- Nysson spinosus** (J. Forster, 1771) [Sphex] (?*Vespa bidens* Linnaeus, 1767; *Sphex spinosus* J. Forster, 1770, nom. nud.; *Crabro spinosus* Fabricius, 1775; *Mellinus tricinctus* Fabricius, 1793; *Vespa trilineata* Turton, 1802; *N. geniculatus* Lepeletier de Saint Fargeau, 1845; *N. malaisei* Gussakovskij, 1932). Russia: **EP** (N, NW, C, E, NC), **WS** (TM, NS, AL), **ES** (KR, IR), **FE** (AM, KH, PR, SA, KU). – Europe (WE, NE, SE, EE), Azerbaijan, Turkey, Iran, Kazakhstan, Japan (Hon).
- Nysson tridens** Gerstaecker, 1867 (*N. melas* Müller, 1918; *N. fraterculus* Gussakovskij, 1932; *N. boemicus* Zavadil in Zavadil et Šnoflák, 1948; *N. boemicus* Zavadil, 1951). Russia: **EP** (C, S), **WS** (TM, AL), **FE** (AM, PR). – Europe (WE, NE, SE, EE), Turkey, Tajikistan, Uzbekistan, Kazakhstan, Mongolia.
- Nysson trimaculatus trimaculatus** (Rossi, 1790) [*Crabro*] (*N. nigripes* Spinola, 1807; *N. geniculatus* Olivier, 1812). Russia: **EP** (C, E, S), **WS** (AL), **ES** (KR, IR), **FE** (PR, KU). – Europe (WE, NE, SE, EE), Abkhazia, Azerbaijan, Kazakhstan.
- Nysson variabilis** Chevrier, 1867 (?*N. friesei* Handlirsch, 1887). Russia: **EP** (S). – Europe (WE, SE, EE), Turkey, Kazakhstan.

Tribe BEMBICINI

- AMMATOMUS** A. Costa, 1859. Type species *Gorytes coarctatus* Spinola, 1808. Number of species: World – 30, Palaearctic – 5, Russia – 3.
- Ammatomus coarctatus** (Spinola, 1808) [Gorytes] (*Gorytes handlirschi* F. Morawitz, 1890; *G. mavromoustakisi* Balthasar, 1954; *G. mitjaevi* Kazenas, 1972). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), N Africa, Armenia, Azerbaijan, Turkey, Syria,

Israel, Yemen, Iran, Afghanistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Ammatomus mesostenus (Handlirsch, 1888) [Gorytes] (*Gorytes rhopalocerus* Handlirsch, 1895; *G. nikolajevskii* Gussakovskij, 1928). Russia: **EP** (S, NC). – N Africa, Israel, United Arab Emirates, Oman, Iraq, Iran, Afghanistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, China (NW).

Ammatomus rogenhoferi (Handlirsch, 1888) [Gorytes] Russia: **EP** (E, S, CR). – Europe (SE, EE), Georgia, Armenia, Azerbaijan, Turkey, Jordan, Israel, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, China (NW).

ARGOGORYTES Ashmead, 1899 (*Archarpactus* Pate, 1937, suppressed by the ICN, 1939 (Opinion 135); *Malaygorytes* Nemkov, 1999). Type species: *Gorytes carbonarius* F. Smith, 1856. Number of species: World – 31, Palaearctic – 5, Russia – 3.

Argogorytes fargeii (Shuckard, 1837) [Gorytes] (*Gorytes croceipes* Eversmann, 1849; *A. mongolicus* Tsuneki, 1971; *A. przewalskyi* Kazenas, 1971; *A. przewalskyi* Kazenas, 1972). Russia: **EP** (N, C, E, S, CR), **UR**, **WS** (NS, AL), **ES** (KR, IR, BR, YA, ZB). – Europe (WE, NE, SE, EE), Armenia, Turkey, Israel, Kazakhstan, Mongolia, China (NE).

Argogorytes mystaceus grandis (Gussakovskij, 1932) [Gorytes]. Russia: **ES** (ZB), **FE** (AM, KH, PR, SA, KU). – China (NE), Korean Peninsula, Japan (Hok, Hon).

Argogorytes mystaceus mystaceus (Linnaeus, 1761) [Sphecius] (*Vespa campestris* Linnaeus, 1761; *V. inimicus* Harris, 1782; *Sphex longicornis* Rossi, 1790; *Crabro bicinctus* Fabricius, 1793; *Mellinus arpactus* Fabricius, 1804; *Vespa flavigaster* Donovan, 1808; *Gorytes tonsus* (Bondroit, 1933)). Russia: **EP** (N, C, E), **UR**, **WS** (AL), **ES** (KR, IR). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan.

Argogorytes nipponis Tsuneki, 1963. Russia: **FE** (PR, KU). – Japan (Hok, Hon).

BEMBECINUS A. Costa, 1859 (*Stizomorphus* A. Costa, 1859; *Gorystizus* Minkiewicz, 1934, not available (ICZN, 1999, Article 13.3); *Gorystizus* Pate, 1937; *Lavia* Rayment, 1953, nomen nudum). Type species: *Bembecinus meridionalis* A. Costa, 1859. Number of species: World – 186, Palaearctic – 49, Russia – 2.

Bembecinus hungaricus hungaricus (Frivaldszky, 1876) [Larra]. Russia: **EP** (E, S, NC, CR). – Europe (WE, SE, EE), Abkhazia, Turkey, Kazakhstan.

Bembecinus hungaricus sibiricus (Mocsáry in Mocsáry et Szépligeti, 1901) [Stizus]. Russia: **WS** (TK, AL), **ES** (KR, BR, ZB), **FE** (AM, KH, PR). – Mongolia, China (NC, NW).

Bembecinus tridens tridens (Fabricius, 1781) [Vespa] (*Crabro cinctus* Rossi, 1790; *Mellinus repandus* Panzer, 1800; *Stizus sinuatus* Latreille, 1805; *S. errans* Kohl: Ferton, 1912, nomen nudum; *S. tridens errans* de Beaumont, 1950). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (AL). – Europe (WE, SE, EE), N Africa, Abkhazia, Georgia, Armenia, Azerbaijan, Turkey, Syria, Jordan, Lebanon, Israel, Oman.

BEMBIX Fabricius, 1775 (*Bembyx* Fabricius, 1775; *Bembex* Fabricius, 1777; *Apobembex* Pate, 1937; *Epibembex* Pate, 1937). Type species: *Bembex rostrata* Fabricius, 1781 (= *Apis rostrata* Linnaeus, 1758). Number of species: World – 328, Palaearctic – 74, Russia – 16.

Bembix bicolor Radoszkowski, 1877 (*B. femoralis* Radoszkowski, 1877; *B. bipunctata* Radoszkowski, 1877; *B. barbiventris* F. Morawitz, 1889). Russia: **EP** (C). – Europe (SE, EE), Turkey, Israel, Oman, Iraq, Iran, Afghanistan, Central Asia, Kazakhstan, Mongolia, China (NC, NW).

Bembix bidentata Vander Linden, 1829 (*B. dalmatica* Kriechbaumer, 1869). Russia: **EP** (C, S, NC, CR), **WS** (AL). – Europe (WE, SE, EE), Abkhazia, Azerbaijan, Turkey, Iran, Kazakhstan.

Bembix canescens (Gmelin, 1790) [Apis]. Russia: **UR**.

Bembix cinctella cinctella Handlirsch, 1893. Russia: **EP** (CR). – Europe (SE), Turkey,

Bembix diversipes F. Morawitz, 1889. Russia: **WS** (AL), **ES** (KR, IR, BR), **FE** (PR). – Turkey, Iran, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NC, NW).

Bembix eburnea Radoszkowski, 1877 (*B. weberi* Handlirsch, 1893; *B. subeburnea* Tsuneki, 1971; *B. lama* Tsuneki, 1971). Russia: **EP** (S, CR). – Turkey, Iran, Afghanistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NE, NC).

Bembix integra Panzer, 1801, nom. praeocc., nec *B. integra* Fabricius, 1793 (now in *Stizus*) (*B. cristata* Mocsáry, 1883; *B. dubia* Gussakovskij, 1933; *B. iliensis* Kazenas, 1978; *B. iliensis* Kazenas, 1980). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), Iran, Turkmenistan, Kazakhstan.

Bembix lutescens Radoszkowski, 1877 (*B. gracilis* Handlirsch, 1893). Russia: **EP** (S, CR). – Europe

(EE), Armenia, Azerbaijan, Turkey, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Bembix megerlei Dahlbom, 1845 (*?B. undata* Dahlbom, 1845; *B. sarafschanii* Radoszkowski, 1877). Russia: **EP** (C, S, NC, CR). – Europe (C, EE), Turkey, Uzbekistan, Kazakhstan.

Bembix niponica F. Smith, 1873 (*B. picticollis* F. Mörner, 1889; *B. miserabilis* J. Parker, 1929). Russia: **ES** (KR, BR), **FE** (AM, KH, PR). – Turkmenistan, Uzbekistan, Kazakhstan, Mongolia, China (NE, NC, NW), Korean Peninsula, Japan (Hon, Kyu, Ryu).

Bembix oculata oculata Panzer, 1801 (*?Vespa pallescens* Giorna, 1791; *B. latreillii* Lepeletier de Saint Fargeau, 1845; *B. neglecta* Dahlbom, 1845; *B. ocellata* Dahlbom, 1845; *?B. rossii* Dahlbom, 1845; *B. panzeri* Handlirsch, 1893; *B. hispanica* Mercet, 1904; *B. ebusiana* Giner Marí, 1934; *B. oculata mongolica* Tsuneki, 1971; *B. oculata gegen* Tsuneki, 1971). Russia: **EP** (C, S, NC, CR), **UR**, **WS** (AL). – Europe (WE, SE, EE), N Africa, Abkhazia, Azerbaijan, Syria, Jordan, Lebanon, Israel, Saudi Arabia, United Arab Emirates, Oman, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NC).

Bembix olivacea olivacea Fabricius, 1787 (*?Vespa clipeata* Christ, 1791; *?V. apilinguaria* Christ, 1791; *B. senilis* Fabricius, 1804; *?B. octopunctata* Donovan, 1810; *B. notata* Dahlbom, 1845; *B. mediterranea* Handlirsch, 1893; *B. maroccana* Mercet, 1905; *B. pardoi* Giner Marí, 1945). Russia: **EP** (C, S, NC, CR). – Europe (WE, SE, EE), N Africa, Armenia, Azerbaijan, Turkey, Israel, Saudi Arabia, Yemen, Iran, Turkmenistan, Kazakhstan, Africa, Madagascar.

Bembix pallida Radoszkowski, 1877. Russia: **EP** (C, S, CR). – Europe (SE, EE), Turkey, Israel, Tajikistan, Uzbekistan.

Bembix portchinskii Radoszkowski, 1884 (*B. seminigra* F. Morawitz, 1889; *B. kirgisica* F. Morawitz, 1891; *B. ganglbaueri* Handlirsch, 1893; *B. trimaculata* Kazenas, 1978; *B. trimaculata* Kazenas, 1980). Russia: **EP** (S). – N Africa, Turkey, Iran, Afghanistan, Central Asia, Kazakhstan, Mongolia, China (NC, NW).

Bembix rostrata rostrata (Linnaeus, 1758) [*Apis*] (*Vespa armata* Sulzer, 1776; *Apis rostrata* Gmelin, 1790, nom. praeocc., nec *A. rostrata* Linnaeus, 1758; *B. vidua* Lepeletier de Saint Fargeau, 1845; *B. dissecata* Dahlbom, 1845; *B. gallica* Mocsáry, 1883; *B. paradoxa* Giner Marí, 1943). Russia: **EP** (NW,

C, E, S, NC), **UR**, **WS** (TK, NS, AL), **ES** (KR, IR). – Europe (WE, NE, SE, EE), Azerbaijan, Turkey, Iran, Afghanistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NC).

Bembix turca Dahlbom, 1845 (*B. melaena* F. Smith, 1856; *B. melanura* F. Morawitz, 1889; *B. asiatica* Radoszkowski, 1893; *B. gobiensis* Tsuneki, 1971; *B. atra* Kazenas, 1978; *B. atra* Kazenas, 1980). Russia: **EP** (S, CR). – Europe (SE, EE), N Africa, Turkey, Iran, Afghanistan, Turkmenistan, Tajikistan, Kazakhstan, Mongolia, China (NW).

GORYTES Latreille, 1805 (*Arpactus* Panzer, 1805; *Arpactus* Panzer, 1806, nom. praeocc., nec *Arpac-tus* Panzer, 1805; *Euzonia* Stephens, 1829; *Hoplis-sus* Lepeletier de Saint Fargeau, 1832; *Euspon-gus* Lepeletier de Saint Fargeau, 1832; *Pseudopli-sus* Ashmead, 1899; *Laevigorytes* Zavadil in Zavadil et Šnoblák, 1948; *Leiogorytes* Bohart, 2000). Type species *Mellinus quinquecinctus* Fabricius, 1793. Number of species: World – 73, Palaearctic – 25, Russia – 18.

Gorytes aino Tsuneki, 1963. Russia: **ES** (BR), **FE** (PR, SA, KU). – China (NC), Japan (Hok, Hon).

Gorytes albidulus (Lepeletier de Saint Fargeau, 1832) [Hoplisus] (*Mellinus dissectus* Panzer, 1801; *Euspon-gus albilabris* Lepeletier de Saint Fargeau, 1832; *G. elegans* F. Smith, 1856). Russia: **EP** (C, S, CR), **WS** (AL). – Europe (WE, SE, EE), N Africa, Azerbaijan, Turkey, Iran, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia.

Gorytes ambiguus Handlirsch, 1888 (*G. sogdianus* Gussakovskij, 1952; *G. verhoeffi* Tsuneki, 1967; *G. heptapotamensis* Kazenas, 1972; *G. zaysanicus* Kazenas, 1987; *G. nobilis* Nesterov, 1993). Russia: **WS** (AL), **ES** (KR, IR), **FE** (KH, PR). – Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NE).

Gorytes fallax Handlirsch, 1888. Russia: **EP** (C, E, CR), **WS** (AL), **ES** (KR, IR). – Europe (WE, SE, EE), Kazakhstan.

Gorytes foveolatus foveolatus Handlirsch, 1888. Russia: **EP** (NC, CR). – Europe (SE, EE), Turkey, Syria, Jordan.

Gorytes kohlii Handlirsch, 1888 (*G. kozlovi* Nesterov, 1993). Russia: **EP** (NC, CR). – Europe (SE, EE), Turkey, Syria, Turkmenistan, Kazakhstan.

Gorytes laticinctus koreanus Handlirsch, 1888 (*G. radoszkowskyi* Handlirsch, 1888). Russia: **ES** (ZB), **FE** (KH, PR, SA, KU). – Korean Peninsula, Japan (Hok).

Gorytes laticinctus laticinctus (Lepeletier de Saint Fargeau, 1832) [Euspongus]. Russia: **EP** (N, NW, C, E, CR), **WS** (AL). – Europe (WE, NE, SE, EE), N Africa, Turkey, Uzbekistan, Kazakhstan.

Gorytes maculicornis (F. Morawitz, 1889) [Hoplisus] (*G. eous* Gussakovskij, 1932; *G. harbinensis* Tsuneki, 1967). Russia: **ES** (ZB), **FE** (AM, KH, PR). – China (NE, NC), Korean Peninsula, Japan (Hok, Hon).

Gorytes neglectus Handlirsch, 1895 (*G. hakutozanus* Tsuneki, 1963). Russia: **EP** (N, NW, C), **WS** (TM, NS, KM, AL), **ES** (KR, IR, BR), **FE** (KH, PR, SA). – Europe (NE), Turkey, Kazakhstan, Korean Peninsula.

Gorytes nigrifacies (Mocsáry, 1879) [Hoplisus]. Russia: **EP** (NC, CR). – Europe (WE, SE, EE), Georgia, Azerbaijan, Turkey, Jordan, Israel.

Gorytes pieli Yasumatsu, 1943 (*G. takeuchii* Tsuneki, 1963). Russia: **ES** (ZB), **FE** (KH, PR). – Mongolia, China (NE, NC), Korean Peninsula.

Gorytes pleuripunctatus pleuripunctatus (A. Costa, 1859) [Hoplisus] (*Hoplisus tirolensis* Kohl, 1880). Russia: **EP** (C, S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Georgia, Armenia, Azerbaijan, Turkey, Syria, Israel, Kazakhstan.

Gorytes procrustes Handlirsch, 1888. Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (AL). – Europe (WE, SE, EE), Georgia, Turkey, Kazakhstan.

Gorytes quadrifasciatus (Fabricius, 1804) [*Mellinus*] (*Euspongus vicinus* Lepeletier de Saint Fargeau, 1832; *Hoplisus montivagus* Mocsáry, 1878). Russia: **EP** (N, C, E, S, NC, CR), **UR**, **WS** (TM, NS, KM, AL), **ES** (KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, MG). – Europe (WE, NE, SE, EE), N Africa, Turkey, Kyrgyzstan, Kazakhstan, Mongolia, Korean Peninsula.

Gorytes quinquecinctus quinquecinctus (Fabricius, 1793) [*Mellinus*] (*Crabro calceatus* Rossi, 1794; *Mellinus arenarius* Panzer, 1798; *G. ruficornis* Latreille, 1805; *G. proximus* Handlirsch, 1893). Russia: **EP** (N, C, E, S, NC, CR), **UR**, **WS** (TM, NS, KM, AL), **ES** (TU, KR, IR, ZB). – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey, Syria, Iran, Uzbekistan, Kyrgyzstan, Kazakhstan.

Gorytes quinquefasciatus quinquefasciatus (Panzer, 1798) [*Mellinus*] (*Hoplisus lacordairei* Lepeletier de Saint Fargeau, 1832; *H. anceps* Mocsáry, 1879; *H. geminatus* A. Costa, 1867; *H. geminatus* A. Costa, 1869). Russia: **EP** (C, E, NC, CR), **UR**, **WS** (AL), **ES** (KR, IR, BR), **FE** (PR). – Europe (WE, SE, EE).

Gorytes schmiedeknechti Handlirsch, 1888. Russia: **EP** (NC). – Europe (SE, EE), Azerbaijan, Turkey, Syria.

Gorytes sulcifrons mongolicus Tsuneki, 1971. Russia: **WS** (AL), **ES** (KR, IR, BR, ZB). – Azerbaijan, Kazakhstan, Mongolia, China (NC).

Gorytes sulcifrons sulcifrons (A. Costa, 1867) [Hoplisus] (*Hoplisus sulcifrons* A. Costa, 1869). Russia: **EP** (E, S, CR). – Europe (WE, SE, EE), N Africa, Armenia, Turkey, Lebanon, Central Asia, Kazakhstan.

HARPACTUS Shuckard, 1837 (*Arpactus* Jurine, 1807, nom. praeocc., nec *Arpactus* Panzer, 1805; *Harpactes* Dahlbom, 1843; *Dienoplus* W. Fox, 1894). Type species: *Arpactus formosus* Jurine, 1807. Number of species: World – 72, Palaearctic – 54, Russia – 11.

Harpactus affinis (Spinola, 1808) [Gorytes] (*Arpactus carceli* Lepeletier de Saint Fargeau, 1832; *A. tricolor* Lepeletier de Saint Fargeau, 1832). Russia: **EP** (C, S, NC), **WS** (AL). – Europe (WE, SE, EE), Turkey, Kazakhstan.

Harpactus annulatus Eversmann, 1849 (*Dienoplus varipes* Tsuneki, 1971). Russia: **EP** (S), **UR**. – Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia.

Harpactus elegans elegans (Lepeletier de Saint Fargeau, 1832) [Arpactus] (*H. vedicus* Nesterov, 1994). Russia: **EP** (C, E, S, NC, CR), **WS** (AL), **ES** (KR). – Europe (WE, NE, SE, EE), Armenia, Azerbaijan, Kazakhstan.

Harpactus formosus formosus (Jurine, 1807) [Arpactus]. Russia: **EP** (C, S, NC, CR). – Europe (WE, SE, EE), N Africa, Turkey, Israel, Saudi Arabia, Kuwait, Oman, Iran, Kyrgyzstan, Kazakhstan.

Harpactus laevis laevis (Latreille, 1792) [Mutilla] (*Sphex cruentus* Fabricius, 1798; *Evania ruficollis* Fabricius, 1798; *H. caucasicus* Radoszkowski, 1884; *Dienoplus dzhingis* Tsuneki, 1971). Russia: **EP** (C, S, CR), **WS** (AL), **ES** (IR, BR, ZB), **FE** (AM). – Europe (WE, SE, EE), Turkey, Iran, Kazakhstan, Mongolia, China (NE, NC).

Harpactus lunatus (Dahlbom, 1832) [Larra] (*Arpactus belgicus* Wesmael, 1839). Russia: **EP** (N, C, E), **WS** (AL), **ES** (KR, IR, YA). – Europe (WE, NE, SE, EE), Kazakhstan.

Harpactus morawitzi Radoszkowski, 1884 (*Gorytes moravicus* Šnoflák, 1943). Russia: **EP** (C, S, NC), **ES** (IR, BR, ZB). – Europe (WE, SE, EE), Turkey, Turkmenistan, Kazakhstan.

Harpactus tauricus Radoszkowski, 1884 (*Gorytes sareptanus* Handlirsch, 1888; *H. turcmenicus* Radoszkowski, 1893). Russia: **EP** (S, CR). – Europe (SE, EE), Turkey, Iran, Turkmenistan, Tajikistan, Kazakhstan.

Harpactus transbaicalicus Nemkov, 1994. Russia: **ES** (BR).

Harpactus transiens A. Costa, 1887 (*H. transiens* A. Costa, 1888; *Gorytes consanguineus* Handlirsch, 1888; *H. chosrovicus* Nesterov, 1994). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), Armenia, Turkey, Kazakhstan.

Harpactus tumidus tumidus (Panzer, 1801) [Pom-pilus] (*Dienoplus japonensis* Tsuneki, 1963; *H. transiliensis* Kazenas, 1989). Russia: **EP** (N, C, E), **WS** (TK, AL), **ES** (IR, BR, ZB), **FE** (AM, KH, PR, SA, KA, MG). – Europe (WE, NE, SE, EE), N Africa, Kyrgyzstan, Kazakhstan, Japan (Hon).

HOPLISOIDES Gribodo, 1884 (*Icuma* Cameron, 1905). Type species *Hoplisoides intricans* Gribodo, 1884. Number of species: World – 79, Palaearctic – 6, Russia – 4.

Hoplisoides craverii craverii (A. Costa, 1867) [Hoplisus] (*Hoplisus craverii* A. Costa, 1869; *H. ottomanus* Mocsáry, 1879). Russia: **EP** (S, NC), **UR**. – Europe (WE, SE, EE), Armenia, Azerbaijan, Turkey, Turkmenistan, Uzbekistan, Kazakhstan, Mongolia.

Hoplisoides gazagnairei distinguendus (Yasumatsu, 1939) [Gorytes]. Russia: **FE** (PR). – Kazakhstan, Mongolia, China (NE, NC), Korean Peninsula.

Hoplisoides latifrons (Spinola, 1808) [Gorytes] (*Hoplissus pulchellus* Wesmael, 1852; *H. minutus* Mocsáry, 1879). Russia: **EP** (E, S, CR), **UR**. – Europe (WE, SE, EE), N Africa, Turkey, Tajikistan, Uzbekistan, Kazakhstan.

Hoplisoides punctuosus punctuosus (Eversmann, 1849) [Hoplisus] (*Hoplisus punctatus* Kirschbaum, 1853; *H. crassicornis* A. Costa, 1859; *H. maculipennis* Giraud in von Frauenfeld, 1861; *Gorytes ibericus* Mercet, 1906). Russia: **EP** (C, S, NC, CR), **UR**, **WS** (NS, KM, AL). – Europe (WE, SE, EE), N Africa, Armenia, Azerbaijan, Turkey, Syria, Israel, Central Asia, Kazakhstan, Mongolia.

LESTIPHORUS Lepeletier de Saint Fargeau, 1832 (*Lestophorus* Agassiz, 1847; *Hypomellinus* Ashmead, 1899; *Mellinogastra* Ashmead, 1899). Type species *Crabro bicinctus* Rossi, 1794. Number of species: World – 18, Palaearctic – 8, Russia – 3.

Lestiphorus bicinctus (Rossi, 1794) [Crabro], validated by ICZN, 1963 (Opinion 675). Russia: **EP** (C, CR), **WS** (AL). – Europe (WE, SE, EE), Turkey.

Lestiphorus bilunulatus bilunulatus A. Costa, 1867 (*L. bilunulatus* A. Costa, 1869; *Gorytes semistriatus* Schmiedeknecht, 1881). Russia: **WS** (KM, AL), **ES** (KR, ZB), **FE** (AM, KH, PR, SA). – Europe (WE, SE, EE), Kazakhstan.

Lestiphorus pacificus (Gussakovskij, 1932) [Gorytes]. Russia: **FE** (PR).

OLGIA Radoszkowski, 1877 (*Kaufmannia* Radoszkowski, 1877). Type species *Olgia modesta* Radoszkowski, 1877. Number of species: World – 6, Palaearctic – 6, Russia – 1.

Olga helena de Beaumont, 1953. Russia: **EP** (CR). – Europe (SE), Turkey.

ORYTTUS Spinola, 1836 (*Agraptus* Wesmael, 1852; *Arcesillas* Pate, 1938). Type species *Sphex concinnus* Rossi, 1790. Number of species: World – 14, Palaearctic – 5, Russia – 2.

Oryttus concinnus concinnus (Rossi, 1790) [Sphex]. Russia: **EP** (C, CR). – Europe (WE, SE, EE).

Oryttus kaszabi Tsuneki, 1971 (*Gorytes kostjuki* Nesterov, 1993). Russia: **WS** (AL), **ES** (TU, ZB). – Kyrgyzstan, Kazakhstan, Mongolia.

PSAMMAECIUS Lepeletier de Saint Fargeau, 1832. Type species *Gorytes punctulatus* Vander Linden, 1829. Number of species: World – 5, Palaearctic – 5, Russia – 1.

Psammaecius punctulatus (Vander Linden, 1829) [Gorytes]. Russia: **EP** (C, S, NC, CR). – Europe (WE, SE, EE), N Africa, Armenia, Azerbaijan, Israel, Central Asia, Kazakhstan.

SPHECIUS Dahlbom, 1843 (*Hogardia* Dufour, 1841, not available (ICZN, 1999, Art. 11.5); *Hogardia* Lepeletier de Saint Fargeau, 1845; *Sphecienus* Patton, 1879; *Nothosphecius* Pate, 1936). Type species *Sphecius speciosus* Dahlbom, 1843 (= *Sphex speciosus* Drury, 1773). Number of species: World – 21, Palaearctic – 10, Russia – 2.

Sphecius antennatus (Klug, 1845) [Larra] (*Stizus aberrans* Eversmann, 1849; *Sphecius impressus* Kokujev, 1909). Russia: **EP** (S, NC, CR), **UR**, **WS** (AL). – Europe (WE, SE, EE), Azerbaijan, Turkey, Syria, Israel, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NC).

Sphecius conicus conicus (Germar, 1817) [Stizus] (*Stizus luniger* Eversmann, 1849). Russia: **EP** (S, CR), **UR**, **WS** (AL). – Europe (SE, EE), Turkey, Kazakhstan.

STIZOIDES Guérin-Méneville, 1844 (*Omphalius* Vachal, 1900, nom. praeocc., nec *Omphalius* Philippi, 1847 (Gastropoda); *Scotomphales* Vachal, 1900; *Tachystizus* Minkiewicz, 1934, not available (ICZN, 1999, Art. 13.3); *Tachystizus* Pate, 1937). Type species *Larra fasciata* Fabricius, 1798 (= *Sphex assimilis* Fabricius, 1787). Number of species: World – 30, Palaearctic – 10, Russia – 3.

Stizoides crassicornis (Fabricius, 1787) [Tiphia] (*Bembex rufipes* Olivier, 1789; *Pompilus alienus* Fischer de Waldheim, 1843; *Stizus fulvipes* Eversmann, 1846). Russia: **EP** (C, E, S, NC, CR). – Europe (WE, SE, EE), N Africa, Azerbaijan, Turkey, Israel, Iran, Central Asia, Kazakhstan.

Stizoides melanopterus (Dahlbom, 1845) [Stizus] (*Stizus concolor* Eversmann, 1849). Russia: **EP** (S, CR). – Europe (SE), Armenia, Turkey, Lebanon, Israel, Iraq, Central Asia, Kazakhstan.

Stizoides tridentatus tridentatus (Fabricius, 1775) [Crabro] (*Larra bifasciata* Fabricius, 1798). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Armenia, Azerbaijan, Turkey, Syria, Israel, Kazakhstan, Mongolia.

STIZUS Latreille, 1802 (*Megastizus* Patton, 1879; *Stizolarra* de Saussure, 1887; *Megalostizus* W. Schulz, 1906). Type species *Bembix ruficornis* Fabricius, 1787 (= *Vespa ruficornis* J. Forster, 1771). Number of species: World – 105, Palaearctic – 54, Russia – 6.

Stizus annulatus (Klug, 1845) [Larra] (*Larra subapicalis* Walker, 1871; *S. eversmanni* Radoszkowski, 1877; *S. kizilkumii* Radoszkowski, 1877; *S. picticornis* F. Morawitz, 1891; *S. pygidialis* Handlirsch, 1892; *S. gracilipes* Handlirsch, 1892; *S. hebraeus* Balthasar, 1954; *S. gobiensis* Tsuneki, 1971). Russia: **EP** (S). – Europe (SE), N Africa, Armenia, Turkey, Syria, Israel, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia.

Stizus bipunctatus (F. Smith, 1856) [Larra] (*S. eugeniae* Gussakovskij, 1935). Russia: **EP** (CR). – Europe (SE, EE), Turkey, Iran, Afghanistan, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan.

Stizus fasciatus (Fabricius, 1781) [Bembex] (*Bembex integra* Fabricius, 1793; *S. terminalis* Dahlbom, 1845;

S. mongolicus Tsuneki, 1971). Russia: **EP** (C, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Turkey, Israel, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NW).

Stizus perrisi perrisi Dufour, 1838 (*S. perrisi* Lepeletier de Saint Fargeau, 1845; *Larra pulcherrima* F. Smith, 1856; *S. pluschtschewskii* Radoszkowski, 1888). Russia: **EP** (C, E, S, CR), **UR**, **WS** (AL), **ES** (ZB), **FE** (PR). – Europe (WE, NE, SE, EE), Armenia, Turkey, Uzbekistan, Kyrgyzstan, Kazakhstan, China (NE, NC, NW), Korean Peninsula, Japan (Hon).

Stizus ruficornis ruficornis (J. Forster, 1771) [Vespa] (*Bembex ruficornis* Fabricius, 1787, nom. praeocc., nec *Vespa ruficornis* J. Forster, 1771; *Vespa gaditana* Gmelin, 1790; *S. ornatus* Lepeletier de Saint Fargeau, 1845; *Larra ferruginea* F. Smith, 1856; *S. komarovi* Radoszkowski, 1888; ?*S. ritsemae* Handlirsch, 1895; *S. ritsemae* Handlirsch, 1896; *S. scutellaris* W.F. Kirby, 1900; *S. distinguendus* Handlirsch, 1901). Russia: **EP** (S, CR), **UR**. – Europe (WE, SE, EE), N Africa, Armenia, Azerbaijan, Turkey, Jordan, Saudi Arabia, United Arab Emirates, Yemen, Oman, Iran, Central Asia, Kazakhstan, Mongolia, Africa.

Stizus rufiventris rufiventris Radoszkowski, 1877 (*S. sarmaticus* F. Morawitz, 1891; *S. kaszabi* Tsuneki, 1971). Russia: **EP** (S, CR). – N Africa, Turkey, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NW).

Subfamily CRABRONINAE

A.V. ANTROPOV

Tribe CRABRONINI

CРАБРО Fabricius, 1775 (*Crabro* Fabricius, 1775, nom. praeocc., nec *Crabro* Geoffroi de Saint Hilaire, 1762, which was suppressed by ICZN (1943, Opinion 144; 1994, Opinion 1754); *Carabro* Say, 1823, lapsus for *Crabro* Fabricius, 1775; *Pemphilis* Risso, 1826, nom. nud.; *Thyreopus* Lepeletier de Saint Fargeau et Brullé, 1835, objective synonym of *Crabro* Fabricius, 1775; *Thyreocnemus* A. Costa, 1867; *Paranothyreus* Kohl, 1897, unavailable under ICZN, 1961; available under ICZN, 1985, 1999; *Paranothyreus* Ashmead, 1899; *Dyscolocrabro* Kohl, 1915, unavailable under ICZN, 1961; available under ICZN, 1985, 1999; *Paranothyreus* Kohl, 1915, unavailable under ICZN, 1961; available under ICZN, 1985, 1999; *Parathyreopus* Kohl, 1915, unavailable under ICZN, 1961; available under ICZN, 1985, 1999; *Agnosicrabro*

Pate, 1944, validation of *Agnosicrabro* Kohl, 1915; *Dyscolocrabro* Pate, 1944, validation of *Dyscolocrabro* Kohl, 1915; *Paranothyreus* Pate, 1944, validation of *Paranothyreus* Kohl, 1915; *Parathyreopus* Pate, 1944, validation of *Parathyreopus* Kohl, 1915; *Pemphilis* Pate, 1944, validation of *Pemphilis* Riss, 1826, objective synonym of *Crabro* Fabricius, 1775). Type species *Vespa cibraria* Linnaeus, 1758. The genus includes eight subgenera: *Crabro* Fabricius, 1775, *Anothyreus* Dahlbom, 1845, *Synothyreopus* Ashmead, 1899, *Dyscolocrabro* Pate, 1944, *Hemithyreopus* Pate, 1944, *Paranothyreus* Pate, 1944, *Norumbega* Pate, 1947, and *Othyreus* Marshakov, 1977. Number of species: World – 88, Palaearctic – 26, Russia – 18.

Crabro (Anothyreus) lapponicus Zetterstedt, 1838 (*C. biguttatus* F. Morawitz, 1892). Russia: **EP** (N, NW), **ES** (BR, KR, IR), **FE** (AM, PR). – Europe (WE, NE, EE).

Crabro (Anothyreus) maeklini A. Morawitz, 1866 (*C. Mäklini* A. Morawitz, 1866a; *C. Mäklini* A. Morawitz, 1866b, objective synonym of A. Morawitz, 1866a). Russia: **EP** (N, NW), **UR**, **ES** (IR, YA, ZB), **FE** (KH, PR). – Europe (NE, EE), Kazakhstan, Mongolia.

Crabro (Crabro) alpinus Imhoff, 1863 (*Thyreopus interruptus* Lepeletier de Saint Fargeau et Brullé, 1835, junior secondary homonym of *Crabro interruptus* Fabricius, 1787; *Th. lactarius* Chevrier, 1867; *C. interruptulus* Dalla Torre, 1897, substitute name for *C. interruptus* Lepeletier de Saint Fargeau et Brullé, 1835). Russia: **EP** (S, NC). – Europe (WE, SE, EE), Turkey.

Crabro (Crabro) altaicus F. Morawitz, 1892. Russia: **WS** (AL), **ES** (TU). – Central Asia, Kazakhstan, Mongolia, China (NW).

Crabro (Crabro) cibrarius (Linnaeus, 1758) [*Vespa*] (*Vespa cibraria* Linnaeus, 1758; *Sphex patellarius* Schreber, 1784; *S. Cribaria Argus* Christ, 1791; *S. cibraria longa* Christ, 1791; *S. cibraria lunata* Christ, 1791, nom. praeocc., nec *S. lunatus* Fabricius, 1775; *C. palmatus* Panzer, 1797; *C. interruptus* Stephens, 1829, nom. praeocc., nec Fabricius, 1787 (now in *Palarus*), substitute name for *C. clypeatus* Panzer, 1804; *C. cibrarius inornatus* Mocsáry in Mocsáry et Szépligeti, 1901; *C. cibrarius hypotheticus* Kokujev, 1927). Russia: **EP** (N, NW, C, E, NC, CR), **UR**, **WS** (NS, AL), **ES** (KM, TM, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA). – Europe (WE,

NE, SE, EE), N Africa, Turkey, Iran, Kazakhstan, Mongolia, Korean Peninsula.

Crabro (Crabro) ingricus (F. Morawitz, 1888) [*Thyreopus*]. Russia: **EP** (N, NW, C), **UR**, **WS** (AL), **FE** (PR, KA). – Europe (WE, NE, SE, EE).

Crabro (Crabro) mocsaryi Kohl, 1915 (*C. Mocsáryi* Kohl, 1915). Russia: **WS** (AL). – Tadzhikistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NW).

Crabro (Crabro) peltarius (Schreber, 1784) [*Sphex*] (*Sphex peltaria* Schreber, 1784; *C. patellatus* Panzer, 1797; *C. dentipes* Panzer, 1797; *C. mediatus* Fabricius, 1798). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (OM, NS, AL), **ES** (BR, KR, IR), **FE** (AM, KH, PR, SA). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan, Mongolia, Korean Peninsula.

Crabro (Crabro) scutellatus (von Scheven, 1781) [*Sphex*] (*Sphex scutellata* von Scheven, 1781; *S. scutularia* Schreber, 1784; *S. scutellata quatuormaculata* Christ, 1791; *C. pterotus* Panzer, 1801; *C. geniculatus* Stephens, 1829, nom. nud.; *C. phaeopterus* Stephens, 1829, nom. nud.). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, NS, AL), **ES** (KR, BR, IR, YA), **FE** (AM, KH, PR, SA). – Europe (WE, NE, SE, EE), Kazakhstan, Mongolia, China (CE).

Crabro (Crabro) sibiricus A. Morawitz, 1866. Russia: **WS** (OM, NS, KM, AL), **ES** (TU, KR, IR, BR, ZB), **FE** (AM, PR). – Kazakhstan, Mongolia, China (NE).

Crabro (Crabro) signaticrus F. Morawitz, 1893. Russia: **WS** (AL). – Tajikistan, Kyrgyzstan, Kazakhstan, Mongolia.

Crabro (Crabro) ussuriensis Gussakovskij, 1932 (*C. koreanus* Tsuneki, 1947). Russia: **ES** (BR, ZB), **FE** (AM, KH, PR). – China (NE), Korean Peninsula.

Crabro (Crabro) werestschagini Gussakovskij, 1932 (*C. okabei* Yasumatsu, 1944). Russia: **ES** (BR, ZB), **FE** (AM, KH, PR). – China (NE).

Crabro (Hemithyreopus) caspicus (F. Morawitz, 1888) [*Blepharipus*]. Russia: **UR**. – Kazakhstan, Turkmenistan.

Crabro (Hemithyreopus) femoralis F. Morawitz, 1892. Russia: **EP** (S, NC), **ES** (BR, IR, KR, BR, ZB), **FE** (PR). – Mongolia, China (NE).

Crabro (Hemithyreopus) loewi Dahlbom, 1845 (*C. Loewi* Dahlbom, 1845; *C. Jaroszewskyi* F. Morawitz, 1892). Russia: **EP** (C, S), **UR**, **WS** (AL), **ES** (IR, ZB). – Europe (WE, SE, NE, EE), Turkmenistan, Kazakhstan.

Crabro (Hemithyreopus) malyshevi L. Ahrens, 1933. Russia: **EP** (C), **ES** (ZB). – Kazakhstan.

Crabro (Synothyreopus) pubens Marshakov, 1976.
Russia: EP (S). – Turkmenistan.

CROSSOCERUS Lepeletier de Saint Fargeau et Brullé, 1835 (*Coelocrabro* Thomson, 1874; *Dolichocrabro* Ashmead, 1899; *Stenocrabro* Ashmead, 1899; *Synorhopalum* Ashmead, 1899; *Ischnolynthus* Holmberg, 1903; *Yuchiha* Pate, 1944; *Fentis* Tsuneki, 1971; *Corenocrabro* Tsuneki, 1991, nom. nud., nom. praeocc., nec Tsuneki, 1974). Type species *Crabro scutatus* Fabricius, 1787 (= *Sphex palmipes* Linnaeus, 1767). The genus includes 23 subgenera: *Crossocerus* Lepeletier de Saint Fargeau et Brullé, 1835, *Blepharipus* Lepeletier de Saint Fargeau et Brullé, 1835, *Cuphopterus* A. Morawitz, 1866, *Hoplocrabro* Thomson, 1874, *Microcrabro* de Saussure, 1892, *Epicrossocerus* Ashmead, 1899, *Ablepharipus* Perkins, 1913, *Acanthocrabro* Perkins, 1913, *Apocrabro* Pate, 1944, *Nothocrabro* Pate, 1944, *Stictoptila* Pate, 1944, *Ainocrabro* Tsuneki, 1954, *Oxycrabro* Leclercq, 1961, *Paroxycrabro* Leclercq, 1963, *Alicrabro* Tsuneki, 1968, *Apoides* Tsuneki, 1968, *Neoblepharipus* Leclercq, 1968, *Towada* Tsuneki, 1970, *Bnunius* Tsuneki, 1971, *Ornicrabro* Leclercq, 1973, *Thao* Tsuneki, 1982, *Orthocrabro* Tsuneki, 1990, and *Yambal* Tsuneki, 1990. Number of species: World – 249, Palaearctic – 78, Russia – 39.

Crossocerus (Ablepharipus) assimilis (F. Smith, 1856) [Crabro] (*Crabro affinis* Wesmael, 1852, nom. praeocc., nec Rossi, 1792, non *Crossocerus affinis* Lepeletier de Saint Fargeau et Brullé, 1835; *C. socius* Thomson, 1870; *C. Tirolensis* Kohl, 1878). Russia: EP (C, E, S, NC), UR, WS (AL), ES (IR, ZB), FE (AM, KH, PR, SA, KU). – Europe (WE, NE, EE), Turkey, Kazakhstan, Mongolia, Korean Peninsula, Japan (Hon).

Crossocerus (Ablepharipus) congener (Dahlbom, 1844) [Crabro]. Russia: EP (C, E, NC), UR, WS (AL), FE (PR, KU). – Europe (WE, NE, SE, EE), Kazakhstan, Mongolia, Japan (Hon).

Crossocerus (Ablepharipus) podagricus (Vander Linden, 1829) [Crabro] (*Crabro tibialis* Stephens, 1829, nom. nud., nom. praeocc., nec *Vespa* (*Crabro*) *tibialis* Gmelin, 1790; *Crossocerus vicinus* Dahlbom, 1842; *Crabro Snoflaki* Zavadil in Zavadil et Šnoflak, 1948; *C. podagricus punctata* Šnoflak in Zavadil et Šnoflak, 1948). Russia: EP (N, C, E, NC, CR), UR, WS (AL), FE (PR, SA, KU). – Europe (WE, NE, SE, EE), N Africa, Turkey, Iran, Kazakhstan, Mongolia, Korean Peninsula, Japan (Hok, Hon).

Crossocerus (Ablepharipus) shibuyai (Iwata, 1934) [Crabro]. Russia: FE (PR, SA, KU). – Korean Peninsula, Japan (Hok, Hon).

Crossocerus (Acanthocrabro) vagabundus (Panzer, 1799) [Crabro] (*Crabro bojus* Schrank, 1802; *Blepharipus 5 maculatus* Lepeletier de Saint Fargeau et Brullé, 1835; *Crossocerus lefebvrei* Lepeletier de Saint Fargeau et Brullé, 1835; *Crabro vagabundus fasciatus* A. Costa, 1871; *C. vagabundus melanotica* Bischoff, 1921; *Corenocrabro ectemiformis* Tsuneki, 1974; *C. ectemiformis* Tsuneki, 1991, nom. nud., nom. praeocc., nec Tsuneki, 1974). Russia: EP (N, NW, C, E, NC, CR), UR, WS (TK, AL), ES (KR, IR, ZB), FE (AM, KH, PR, SA, KA). – Europe (WE, NE, SE, EE), N Africa, Turkey, Uzbekistan, Kazakhstan, Mongolia, China (NE, NC, NW, SW, CE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu), India.

Crossocerus (Ainocrabro) malaisei (Gussakovskij, 1932) [Crabro] (*Crabro aino* Tsuneki, 1947). Russia: FE (AM, PR, SA, KU). – Mongolia, Korean Peninsula, Japan (Hok, Hon, Shi).

Crossocerus (Blepharipus) annulipes annulipes (Lepeletier de Saint Fargeau et Brullé, 1835) [Blepharipus] (*C. gonager* Lepeletier de Saint Fargeau et Brullé, 1835; *Crabro nigritus* Gimmerthal, 1836, nom. praeocc., nec *Blepharipus nigritus* Lepeletier de Saint Fargeau et Brullé, 1835; *C. ambiguus* Dahlbom, 1842; *C. Capito* Dahlbom, 1845; *B. parkeri* Banks, 1921; *Crabro davidsoni* Sandhouse, 1938). Russia: EP (N, NW, C, E, NC), UR, WS (AL), ES (IR), FE (AM, PR). – Europe (WE, NE, SE, EE), N Africa, Abkhazia, Turkey, Israel, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, N America.

Crossocerus (Blepharipus) annulipes hokkaidoensis Tsuneki, 1954. Russia: FE (SA, KU). – China (NE, CE), Korean Peninsula, Japan (Hok, Hon).

Crossocerus (Blepharipus) barbipes Dahlbom, 1845 (*Crabro ater* Cresson, 1865, nom. praeocc., nec Olivier, 1792; *C. hirtipes* A. Morawitz, 1866; *C. hirtipes* A. Morawitz, 1866, nom. praeocc., nec A. Morawitz, 1866; *Dolichocrabro Wickhamii* Ashmead, 1899; *D. Wickhamii* Ashmead, 1902, nom. praeocc., nec Ashmead, 1899; *D. wickhami* Ashmead, 1910, nom. praeocc., nec Ashmead, 1899; *Crossocerus pammelas* Pate, 1944, substitute name for *Crabro ater* Cresson, 1865). Russia: EP (NW, C, E), UR, WS (TK, AL), ES (KR, IR, ZB), FE (KH, PR, SA, KU). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan, Mongolia, China (SW), Korean Peninsula, Japan (Hok, Hon, Yakushima, Ryu), N America.

Crossocerus (Blepharipus) capitosus (Shuckard, 1837) [Crabro] (*Crabro annulus* Dahlbom, 1839; *Crossocerus capitosus yezo* Tsuneki, 1960). Russia: **EP** (NW, C, E, S, NC, CR), **WS** (AL), **FE** (PR, SA, KU). – Europe (WE, NE, SE, EE), Georgia, Kazakhstan, Mongolia, China (CE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Crossocerus (Blepharipus) cetratus (Shuckard, 1837) [Crabro] (*Crabro van der Lindenii* Dahlbom, 1839; *C. dilatatus* Herrich-Schaeffer, 1841; *C. inornatus* Matsumura, 1912; *Crossocerus dentsukamus* Tsuneki, 1976). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (IR), **FE** (PR, SA, KU). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan, Mongolia, China (SW), Korean Peninsula, Japan (Hok, Hon).

Crossocerus (Blepharipus) cinxius (Dahlbom, 1839) [Crabro] (*Crossocerus capitosus yamato* Tsuneki, 1960). Russia: **EP** (N, C), **UR**, **WS** (AL), **ES** (IR), **FE** (PR, SA, KU, MG). – Europe (WE, NE, SE, EE), Turkey, Tajikistan, Kazakhstan, Mongolia, China (SW), Korean Peninsula, Japan (Hok, Hon).

Crossocerus (Blepharipus) heydeni Kohl, 1880 (*Crabro sutshanicus* Gussakovskij, 1932; *C. hedgreni* Kjellander, 1954; *Crossocerus heydeni nipponis* Tsuneki, 1966; *C. murotai* Tsuneki, 1992). Russia: **EP** (CR), **UR**, **WS** (AL), **FE** (PR, SA). – Europe (SE, NE, EE), Turkey, Kazakhstan, Mongolia, Korean Peninsula, Japan (Hon).

Crossocerus (Blepharipus) leucostoma (Linnaeus, 1758) [Sphecius] (*Crabro carbonarius* Dahlbom, 1839; *Crossocerus Rugosus* Herrich-Schaeffer, 1840, nom. nud.; *Crabro rugosus* Herrich-Schaeffer, 1841; *C. melanarius* Wesmael, 1852; *Blepharipus cinctipes* Provancher, 1882; *B. cinctipes* Provancher, 1883, objective synonym of Provancher, 1882; *Crabro niger* Provancher, 1888, nom. praeocc., nec *Crossocerus niger* Lepeletier de Saint Fargeau et Brullé, 1835; *Crabro nigror* W. Fox, 1895; *C. nigror* W. Fox, 1896, emendation of *C. nigror* W. Fox, 1895; *C. servus* Dalla Torre, 1897, substitute name for *C. niger* Provancher, 1888; *Stenocrabro cinctitarsis* Ashmead, 1901; *B. columbiae* Bradley, 1906; *Thyreopus stygius* Mickel, 1916; *Th. utensis* Mickel, 1916). Russia: **EP** (N, NW, C, E), **UR**, **WS** (AL), **ES** (IR), **FE** (KH, PR, SA, KA, MG). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan, Mongolia, Korean Peninsula, Japan (Hok, Hon), N America.

Crossocerus (Blepharipus) megacephalus (Rossi, 1790) [Crabro] (*Crabro aterrimus* Stephens, 1829, nom. nud.; *C. spiniceps* Stephens, 1829, nom. nud.;

C. tarsalis Stephens, 1829, nom. nud.; *Crabro bidens* Haliday, 1833, nom. praeocc., nec Schrank, 1802; *Crossocerus niger* Lepeletier de Saint Fargeau et Brullé, 1835; *C. rufipes* Lepeletier de Saint Fargeau et Brullé, 1835; *Crabro laeviceps* F. Smith, 1856; *C. bison* A. Costa, 1884; *Crossocerus bison* A. Costa, 1884, objective synonym of *Crabro bison* A. Costa, 1884; *?Crabro zaidamensis* Radoszkowski, 1887; *Coelocrabro leucostomoides* Richards, 1935, substitute name for *Crabro leucostoma* of Kohl, 1915 and authors, nec Linnaeus, 1758). Russia: **EP** (N, C, E, S, NC, CR), **UR**, **WS** (TK, AL), **ES** (IR), **FE** (AM, PR). – Europe (WE, NE, SE, EE), N Africa, Turkey, Iran, Tajikistan, Kazakhstan, Mongolia, China (NW), Japan (Hok), India.

Crossocerus (Blepharipus) minamikawai Tsuneki, 1966. Russia: **FE** (PR). – Japan (Hon).

Crossocerus (Blepharipus) nigritus (Lepeletier de Saint Fargeau et Brullé, 1835) [*Blepharipus*] (*Blepharipus nigrita* Lepeletier de Saint Fargeau et Brullé, 1835; *Crabro pubescens* Shuckard, 1837; *Crossocerus diversipes* Herrich-Schaeffer, 1841; *Crabro tischbeini* Dahlbom: Tischbein, 1850, nom. nud.; *C. inermis* Thomson, 1870; *Crossocerus melanogaster* Kohl, 1880; *B. nigricornis* Provancher, 1888; *Crabro sambucicola* C. Verhoeff, 1891; *C. pubescens daisetsuzanus* Tsuneki, 1947; *Crossocerus verhoeffi* Tsuneki, 1967; *C. verhoeffi sudai* Tsuneki, 1976; *C. babai* Tsuneki, 1979). Russia: **EP** (N, NW, C, E), **UR**, **WS** (AL), **ES** (IR), **FE** (PR, SA, KU, KA). – Europe (WE, NE, SE, EE), N Africa, Kazakhstan, Mongolia, China (NE), Korean Peninsula, Japan (Hok, Hon), N America.

Crossocerus (Blepharipus) styrius (Kohl, 1892) [Crabro] (*Crabro pauxillus* Gussakovskij, 1932; *C. suiglihai* Iwata, 1938; *Crossocerus pilicornis* Tsuneki, 1977). Russia: **EP** (E), **FE** (PR, SA, KU). – Europe (WE, NE, SE, EE), Korean Peninsula, Japan (Hok, Hon, Kyu).

Crossocerus (Blepharipus) takeuchii Tsuneki, 1957 (*C. etizenensis* Tsuneki, 1992). Russia: **FE** (PR). – Korean Peninsula, Japan (Hon, Kyu).

Crossocerus (Blepharipus) tanakai Tsuneki, 1954. Russia: **FE** (KH, PR, SA, KU). – Korean Peninsula, Japan (Hok, Hon).

Crossocerus (Blepharipus) tyuzendzianus Tsuneki, 1954. Russia: **EP** (C), **FE** (PR). – Korean Peninsula, Japan (Hok, Hon).

Crossocerus (Blepharipus) walkeri (Shuckard, 1837) [Crabro] (*Crabro Walkeri* Shuckard, 1837;

C. geniculatus Shuckard, 1837, nom. praeocc., nec Olivier, 1792, non Fabricius, 1793; *Crossocerus scaposus* Zetterstedt: Dahlbom, 1844, nom. nud.; *Crabro clypearis* Schenck, 1857; *C. clypearis* Schenck, 1860, nom. nud.; *C. cloëvorax* Nielson, 1901). Russia: **EP** (C, E, S, CR), **UR**, **WS** (AL), **ES** (IR, ZB), **FE** (PR, SA, KU). – Europe (WE, NE, EE), Turkey, Kazakhstan, Mongolia, Korean Peninsula, Japan (Hok, Hon).

Crossocerus (Crossocerus) denticoxa (Bischoff, 1932) [Crabro] (*Crabro denticornis* Gussakovskij, 1932, nom. praeocc., nec F. Smith, 1879; *Crossocerus republicus* Leclercq, 1954, substitute name for *Crabro denticornis* Gussakovskij, 1932). Russia: **EP** (NC), **WS** (AL), **ES** (IR), **FE** (AM, PR). – Europe (WE, SE, EE), Kazakhstan, Mongolia, China (CE), Korean Peninsula.

Crossocerus (Crossocerus) denticrus Herrich-Schaeffer, 1841 (*C. Denticrus* Herrich-Schaeffer, 1840, nom. nud.). Russia: **WS** (AL), **FE** (PR, SA, KU). – Europe (WE, NE, EE), Mongolia, Korean Peninsula, Japan (Hok, Hon).

Crossocerus (Crossocerus) distinguendus (A. Morawitz, 1866) [Crabro] (*Crabro mucronatus* Thomson, 1870, nom. praeocc., nec Fabricius, 1793). Russia: **EP** (C, E, NC, CR), **WS** (AL). – Europe (WE, NE, SE, EE), Kazakhstan, Mongolia.

Crossocerus (Crossocerus) elongatulus annulatus Lepeletier de Saint Fargeau et Brullé, 1835 (*C. annulatus* Lepeletier de Saint Fargeau et Brullé, 1835; *Crabro proximus* Shuckard, 1837; *C. hyalinus* Shuckard, 1837; *C. transversalis* Shuckard, 1837; *C. obliquus* Shuckard, 1837; *C. propinquus* Shuckard, 1837). Russia: **EP** (S), **UR**. – Europe (WE, SE, EE), Turkey.

Crossocerus (Crossocerus) elongatulus elongatulus (Vander Linden, 1829) [Crabro] (*Crabro Stigma* Stephens, 1829, nom. nud.; *Crossocerus varipes* Lepeletier de Saint Fargeau et Brullé, 1835; *C. affinis* Lepeletier de Saint Fargeau et Brullé, 1835; *C. luteipalpis* Lepeletier de Saint Fargeau et Brullé, 1835; *C. morio* Lepeletier de Saint Fargeau et Brullé, 1835; *C. pallidipalpis* Lepeletier de Saint Fargeau et Brullé, 1835; *Crabro brevis* Eversmann, 1849, nom. praeocc., nec Vander Linden, 1829; *C. scutellaris* F. Smith, 1851, nom. praeocc., nec Gimmerthal, 1836; *C. sulcus* W. Fox, 1895; *Stenocrabro plesius* Rohwer, 1912; *Crabro elongatulus Berlandi* Richards, 1928). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (KR, IR), **FE** (PR, SA, KU). – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey, Iran,

Turkmenistan, Tajikistan, Kyrgyzstan, Kazakhstan, Mongolia, Japan (Hok), N America, S America.

Crossocerus (Crossocerus) emarginatus (Kohl, 1899) [Crabro] (*Crabro pacificus* Gussakovskij, 1932). Russia: **ES** (IR, BR, ZB), **FE** (AM, KH, PR, SA, KU). – Kazakhstan, Mongolia, China (CE), Korean Peninsula, Japan (Hok, Hon, Shi, Yakushima, Ryu).

Crossocerus (Crossocerus) exiguum (Vander Linden, 1829) [Crabro] (*C. aphidum* Lepeletier de Saint Fargeau et Brullé, 1835). Russia: **EP** (N, NW, C, E), **UR**, **WS** (NS, AL), **ES** (IR, YA, ZB), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), N Africa, Turkmenistan, Kazakhstan, Mongolia, China (NE), Korean Peninsula.

Crossocerus (Crossocerus) kohli (Bischoff, 1921) [Crabro] (*Crabro diacanthus* Gussakovskij, 1930). Russia: **WS** (AL), **ES** (ZB), **FE** (PR). – Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NW), India.

Crossocerus (Crossocerus) lundbladi (Kjellander, 1954a) [Crabro]. Russia: **EP** (N). – Europe (NE).

Crossocerus (Crossocerus) ovalis Lepeletier de Saint Fargeau et Brullé, 1835 (*Crabro punctum* Zetterstedt, 1838; *C. anxius* Wesmael, 1852; *C. Shuckardi* F. Smith, 1856, nom. praeocc., nec Dahlbom, 1838; *C. ovatus* W. Schulz, 1906, emendation of Lepeletier de Saint Fargeau et Brullé, 1835). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), Kazakhstan, Mongolia.

Crossocerus (Crossocerus) palmipes chosensis Tsuneki, 1957. Russia: **ES** (ZB), **FE** (AM, KH, PR). – Korean Peninsula.

Crossocerus (Crossocerus) palmipes palmipes (Linnaeus, 1767) [Sphex] (*Sphex palmaria* Schreber, 1784; *Crabro scutatus* Fabricius, 1787; *Crossocerus ornatus* Lepeletier de Saint Fargeau et Brullé, 1835; *Crabro scutellaris* Gimmerthal, 1836; *C. gracilis* Eversmann, 1849). Russia: **EP** (N, NW, C, E), **UR**, **WS** (NS, AL), **ES** (KR, IR, BR). – Europe (WE, NE, SE, EE), N Africa, Turkey, Kazakhstan, Mongolia.

Crossocerus (Crossocerus) pullulus (A. Morawitz, 1866) [Crabro] (*Crabro imitans* Kohl, 1915). Russia: **EP** (C, E, S), **WS** (AL), **ES** (YA, IR, BR, ZB), **FE** (AM, KH, PR, SA, KA, MG). – Europe (WE, NE, SE, EE), Kazakhstan, Mongolia.

Crossocerus (Crossocerus) tarsatus (Shuckard, 1837) [Crabro] (*C. palmatus* De Stefani Perez, 1884, nom. praeocc., nec Panzer, 1797). Russia: **EP** (C, E, NC, CR), **WS** (AL), **FE** (PR, KA). – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, Iran, Kazakhstan, Mongolia, Korean Peninsula, N America.

Crossocerus (Crossocerus) ungyo Terayama, 2017

(*C. ungyo* Terayama in Terayama et Takahashi, 2017, substitute name for *Crabro uchidai* Tsuneki, 1947, nom. praeocc., nec Takeuchi, 1931 = *Cimbex femorata* (Linnaeus, 1758)). Russia: **ES** (IR, YA, ZB), **FE** (AM, KH, PR, KU, KA, MG, CH). – Mongolia, Japan (Hok).

Crossocerus (Crossocerus) varus Lepeletier de Saint Fargeau et Brullé, 1835, nom. praeocc., nec *Crabro varus* Panzer, 1799 (now in *Tracheliodes* A. Morawitz); (*C. pusillus* Lepeletier de Saint Fargeau et Brullé, 1835; *C. striatulus* Lepeletier de Saint Fargeau et Brullé, 1835; *Crabro scutellatus* Stephens, 1829, nom. nud., nom. praeocc., nec *Sphex scutellata* von Scheven, 1781 (now in *Crabro*); *Crabro spinipennis* Shuckard, 1837; *Crossocerus striatus* Lepeletier de Saint Fargeau, 1845, lapsus for or emendation of *C. striatulus* Lepeletier de Saint Fargeau et Brullé, 1835; *C. varius* Lepeletier de Saint Fargeau, 1845, emendation of *C. varus* Lepeletier de Saint Fargeau et Brullé, 1835; *Crabro intricatus* F. Smith, 1856, substitute name for *C. striatus* Lepeletier de Saint Fargeau, 1845; *C. Lepeletieri* F. Smith, 1856, substitute name for *C. striatulus* Lepeletier de Saint Fargeau et Brullé, 1835). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (AL), **ES** (KR, IR), **FE** (PR, SA, KU, KA). – Europe (WE, NE, SE, EE), N Africa, Turkey, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (SW), Korean Peninsula, Japan (Hok, Hon).
Crossocerus (Crossocerus) wesmaeli parvicorpus

Tsuneki, 1974 (*C. wesmaeli parvicorpus* Tsuneki, 1991, nom. nud.). Russia: **FE** (PR). – China (NE), Korean Peninsula, Japan (Hon).

Crossocerus (Crossocerus) wesmaeli wesmaeli

(Vander Linden, 1829) [Crabro] (*Ceratocolus maurus* Lepeletier de Saint Fargeau et Brullé, 1835; *C. ziegleri* Lepeletier de Saint Fargeau et Brullé, 1835). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TM, AL), **ES** (IR), **FE** (AM, PR, SA, KU, KA). – Europe (WE, NE, EE), N Africa, Turkey, Tajikistan, Uzbekistan, Kazakhstan, N America.

Crossocerus (Crossocerus) yasumatsui (Tsuneki, 1947) [Crabro] (*Crabro yasumatsui mongolensis* Tsuneki, 1972; *C. nambui* Tsuneki, 1982). Russia: **WS** (AL), **ES** (IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, MG, CH). – Kyrgyzstan, Kazakhstan, Mongolia, Japan (Hok, Hon, Shi), SE Asia.
Crossocerus (Cuphopterus) binotatus Lepeletier de Saint Fargeau et Brullé, 1835 (*Crabro signatus*

Panzer, 1798, nom. praeocc., nec Olivier, 1792; *C. monstrosus* Dahlbom, 1845; *C. confusus* Schulz, 1906, substitute name for *C. signatus* Panzer, 1798). Russia: **EP** (N, C, E, NC, CR), **UR**, **FE** (PR). – Europe (WE, NE, EE), N Africa, Iran, Kazakhstan, India.

Crossocerus (Cuphopterus) dimidiatus dimidiatus

(Fabricius, 1781) [Crabro] (*Crabro subpunctatus* Rossi, 1790; *C. signatus* Olivier, 1792; *C. sexmaculatus* Olivier, 1792; *C. serripes* Panzer, 1797; *C. notatus* Illiger in Rossi, 1807; *C. analis* Stephens, 1829, nom. nud.; *C. consobrinus* Stephens, 1829, nom. nud.; *Blepharipus pauperatus* Lepeletier de Saint Fargeau et Brullé, 1835; *Crabro armipes* von Siebold, 1844). Russia: **EP** (N, C, E, CR), **UR**, **WS** (AL), **ES** (TU, IR), **FE** (AM, PR, KA, MG, CH). – Europe (WE, NE, SE, EE), N Africa, Turkey, Kyrgyzstan, Kazakhstan, Mongolia, China (SW).

Crossocerus (Cuphopterus) dimidiatus agyo Terayama et Takahashi, 2017, substitute name for *Crabro dimidiatus sapporoensis* Kohl, 1915 (*Crabro serripes sapporoensis* Kohl, 1915, nom. praeocc., nec *C. sapporoensis* Matsumura, 1911 (= *Ectemnius fossorius* (Linnaeus, 1758))). Russia: **FE** (SA, KU). – Korean Peninsula, Japan (Hok, Hon).

Crossocerus (Cuphopterus) hakusanus Tsuneki, 1954. Russia: **FE** (PR, SA, KU). – Korean Peninsula, Japan (Hok, Hon, Kyu, Ryu), SE Asia.

Crossocerus (Cuphopterus) subulatus (Dahlbom, 1845) [Crabro]. Russia: **EP** (N, NW, C, E), **UR**, **WS** (AL), **FE** (AM, PR, SA, KU). – Europe (WE, NE, EE), Kazakhstan, Mongolia, Japan (Hok, Hon, Shi).

Crossocerus (Cuphopterus) suzukii (Matsumura, 1912) [Crabro] (*Crabro monstrosus suzukii* Matsumura, 1912). Russia: **FE** (PR, SA, KU). – Korean Peninsula, Japan (Hok, Hon, Shi).

Crossocerus (Cuphopterus) yanoi (Tsuneki, 1947) [Crabro]. Russia: **FE** (SA, PR, KU). – Korean Peninsula, Japan (Hok, Hon, Shi).

Crossocerus (Hoplocrabro) pseudopalmarius (Gusakovskij, 1932) [Crabro] (*C. asiaticus* Tsuneki, 1967). Russia: **FE** (AM, KH, PR). – China (NE), Korean Peninsula.

Crossocerus (Hoplocrabro) quadrimaculatus (Fabricius, 1793) [Crabro] (*Crabro 4punctatus* Fabricius, 1793; *C. murorum* Latreille, 1805; *C. levipes* Vander Linden, 1829; *Crossocerus bimaculatus* Lepeletier de Saint Fargeau et Brullé, 1835; *Crabro 5-maculatus* [sic] Dahlbom, 1838; *C. rotundarius* Dahlbom, 1838). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (TM, AL), **ES** (TU, KR, IR, BR, ZB), **FE** (AM, PR). – Europe

(WE, NE, SE, EE), N Africa, Turkey, Iran, Tajikistan, Uzbekistan, Kazakhstan, Mongolia.

Crossocerus (Neoblepharipus) amurensis (Kohl, 1892) [Crabro]. Russia: **FE** (AM, KH, PR, SA, KU). – Mongolia, China (CE), Korean Peninsula, Japan (Hok, Hon).

Crossocerus (Ornicrabro) flavissimus Leclercq, 1973. Russia: **FE** (PR). – SE Asia.

Crossocerus (Oxycrabro) acanthophorus (Kohl, 1892) [Crabro] (*Crabro pavlovskii* Gussakovskij, 1952). Russia: **EP** (S, NC), **WS** (AL). – Europe (WE, NE, SE, EE), N Africa, Turkey, Syria, Israel, Tadzhikistan, Kazakhstan, Mongolia.

Crossocerus (Towada) flavitarsus (Tsuneki, 1947) [Crabro] (*C. leclercqi* Tsuneki, 1959). Russia: **FE** (PR, SA, KU). – Korean Peninsula, Japan (Hok, Hon, Kyu).

ECTEMNIUS Dahlbom, 1845 (*Mesocrabro* C. Verhoeff, 1892; *Pseudocrabro* Ashmead, 1899; *Xestocrabro* Ashmead, 1899; *Xylocrabro* Ashmead, 1899; *Hylocrabro* R. Perkins, 1902; *Melanocrabro* R. Perkins, 1902; *Xenocrabro* R. Perkins, 1902; *Lophocrabro* Rohwer, 1916; *Ceratocrabro* Tsuneki, 1970; *Papuacrabro* Tsuneki, 1983; *Metactemnius* Tsuneki, 1984; *Spadicocrabro* Tsuneki, 1990). Type species “*E. guttatus*, Dahlb.”: Ashmead, 1899 (= *Crabro guttatus*: Dahlb., 1845 = *C. guttatus* Vander Linden, 1829). The genus includes 16 subgenera: *Ectemnius* Dahlbom, 1845, *Clytochrysus* A. Morawitz, 1864, *Thyreocerus* A. Costa, 1871, *Hypocrabro* Ashmead, 1899, *Metacrabro* Ashmead, 1899, *Protothyreopus* Ashmead, 1899, *Nesocrabro* R. Perkins, 1899, *Oreocrabro* R. Perkins, 1902, *Merospis* Pate, 1941, *Apoctemnius* Leclercq, 1950, *Cameronitus* Leclercq, 1950, *Protoctemnius* Leclercq, 1951, *Yanonus* Tsuneki, 1956, *Policrabro* Leclercq, 1958, *Iwataia* Tsuneki, 1959, and *Leocrabro* Leclercq, 1968. Number of species: World – 188; Palaearctic – 44; Russia – 27.

Ectemnius (Cameronitus) nigritarsus (Herrich-Schaeffer, 1841) [Crabro] (*Crabro Nigritarsus* Herrich-Schaeffer, 1840, nom. nud.; *C. trinotatus* A. Costa, 1867). Russia: **EP** (C, E, CR), **UR**, **WS** (KM, AL), **FE** (AM, PR). – Europe (WE, NE, SE, EE), Georgia, Turkey, Korean Peninsula, Japan (Hok, Hon), India, Sri Lanka, SE Asia, Australia.

Ectemnius (Cameronitus) radiatus (Pérez, 1905) [Crabro] (*Crabro nigritarsus mizuhō* Tsuneki, 1947; *E. mizuhō* Tsuneki, 1952). Russia: **FE** (PR). – Korean Peninsula, Japan (Hok, Hon, Kyu).

Ectemnius (Clytochrysus) cavifrons (Thomson, 1870)

[Crabro]. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (KM, AL), **ES** (KR, IR, ZB), **FE** (AM, KH, PR, SA, KU). – Europe (WE, NE, SE, EE), N Africa, Tajikistan, Kazakhstan, Mongolia, Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu), SE Asia.

Ectemnius (Clytochrysus) lapidarius (Panzer, 1803)

[Crabro] (*Crabro sinuatus* Fabricius, 1804; *C. chrysostomus* Lepeletier de Saint Fargeau et Brullé, 1835, nom. praeocc., nec Gmelin, 1790; *C. comptus* Lepeletier de Saint Fargeau et Brullé, 1835; *C. duodecimguttatus* Stephens, 1829, nom. nud.; *C. xylourgus* Shuckard, 1837; *C. interstinctus* F. Smith, 1851; *C. obscurus* F. Smith, 1856; *C. gracilissimus* Packard, 1866; *C. denticulatus* Packard, 1866; *C. effosus* Packard, 1866; *C. papagorum* Viereck, 1908). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (NS, KM, AL), **ES** (IR, BR, ZB), **FE** (AM, KH, PR, SA, KU). – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, Iran, Kyrgyzstan, Kazakhstan, Mongolia, China (NE), Korean Peninsula, Japan (Hok, Hon), Africa, N America.

Ectemnius (Clytochrysus) ruficornis (Zetterstedt, 1838) [Crabro] (*Crabro aurilabris* Herrich-Schaeffer, 1841; *C. nigrifrons* Cresson, 1865; *C. contiguus* Cresson, 1865; *C. septentrionalis* Packard, 1866; *C. planifrons* Thomson, 1870; *C. longipalpis* C. Verhoeff, 1892; *C. lineatotarsis* Matsumura, 1911; *C. chipsani* Matsumura, 1912). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (KM, AL), **ES** (KR, IR, BR, ZB), **FE** (AM, KH, PR, SA, KU, KA). – Europe (WE, NE, EE), Turkey, Kazakhstan, Mongolia, Korean Peninsula, Japan (Hok, Hon, Ryu), SE Asia, N America.

Ectemnius (Clytochrysus) sexcinctus (Fabricius, 1775) [Crabro] (*Crabro quadricinctus* Fabricius, 1787; *Sphex interrupte-fasciata* Retzius, 1783; *S. cribaria quatuorcincta* Christ, 1791; *Crabro tibialis* Olivier, 1792; *C. 8 maculata* Preyssler, 1793; *C. zonatus* Panzer, 1797; *C. vespiformis* Panzer, 1798; *C. cinctus* Spinola, 1806, nom. praeocc., nec Rossi, 1790; *C. flavipes* Lepeletier de Saint Fargeau et Brullé, 1835, nom. praeocc., nec Fabricius, 1781; *C. tetraedrus* Blanchard, 1840; *C. saundersi* R. Perkins, 1899; *E. yosemite* Pate, 1946). Russia: **EP** (NW, C, CR), **WS** (KM, AL), **ES** (TU, BR), **FE** (PR). – Europe (WE, NE, SE, EE), N Africa, Turkey, Iran, Afghanistan, Pakistan, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan, China (CE), Mongolia, India, N America.

Ectemnius (Ectemnius) borealis (Zetterstedt, 1838) [Crabro] (*Crabro borealis* Zetterstedt, 1838; *C. bipunctatus* Zetterstedt, 1838, nom. praeocc., nec Fabricius, 1787; *C. Nigrinus* Herrich-Schaeffer, 1840, nom. nud.; *C. nigrinus* Herrich-Schaeffer, 1841; *C. parvulus* Packard, 1866, nom. praeocc., nec Herrich-Schaeffer, 1841; *Lindenius Gredleri* Kohl, 1878; *Crabro proletarius* Mickel, 1916). Russia: **EP** (N, C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (KR, IR, BR, ZB), **FE** (AM, PR, SA, KU, KA). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan, Mongolia, Korean Peninsula, Japan (Hok), N America.

Ectemnius (Ectemnius) dives (Lepeletier de Saint Fargeau et Brullé, 1835) [Solenius] (*S. 8-notatus* Lepeletier de Saint Fargeau et Brullé, 1835; *S. octavonotatus* Lepeletier de Saint Fargeau et Brullé, 1835; *Crabro alatulus* Dahlbom, 1839; *C. Pictipes* Herrich-Schaeffer, 1840, nom. nud.; *C. pictipes* Herrich-Schaeffer, 1841; *C. auratus* F. Smith, 1856, substitute name for *Solenius dives* Lepeletier de Saint Fargeau et Brullé, 1835; *C. montanus* Cresson, 1865, nom. praeocc., nec Gistel, 1857; *C. cristatus* Packard, 1866; *C. cubiceps* Packard, 1866; *C. heraclei* Rohwer, 1908; *C. montivagans* Strand, 1916). Russia: **EP** (N, C, E, S, NC, CR), **UR**, **WS** (NS, KM, AL), **ES** (IR, BR, ZB), **FE** (AM, KH, PR, SA, KU). – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, Israel, Kazakhstan, China (SW), Korean Peninsula, Japan (Hok, Hon) N America.

Ectemnius (Ectemnius) guttatus (Vander Linden, 1829) [Crabro] (*Crabro laportei* Lepeletier de Saint Fargeau et Brullé, 1835; *C. Spinicollis* Herrich-Schaeffer, 1840, nom. nud.; *C. spinicollis* Herrich-Schaeffer, 1841; *C. parvulus* Herrich-Schaeffer, 1841; *C. pictus* Schenck, 1857, nom. praeocc., nec Fabricius, 1793 (now in *Dinetus*); *C. pictus* Schenck, 1860, nom. nud.; *C. divitoides* C. Verhoeff, 1892; *C. verhoeffii* Dalla Torre, 1897, substitute name for *C. parvulus* of C. Verhoeff, 1892; *C. dives schencki* Cockerell, 1907, substitute name for *C. pictus* Schenck, 1857). Russia: **EP** (N, NW, C, E, NC, CR), **UR**, **WS** (AL), **ES** (IR, ZB). – Europe (WE, NE, SE, EE), Kazakhstan.

Ectemnius (Ectemnius) rugifer (Dahlbom, 1845) [Crabro]. Russia: **EP** (C, E, S, CR), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), Turkey.

Ectemnius (Hypocrabro) confinis (Walker, 1871) [Crabro] (*E. laevigatus* De Stefani Perez, 1884; *Crabro flavidollis* F. Morawitz, 1892; *C. Hannonis* Gribodo, 1896; *C. subtilis* Pérez: Antiga, Bofill, 1904, nom. nud.; *C. subtilis* Pérez, 1905). Russia: **EP** (C, S,

CR), **UR**. – Europe (WE, NE, SE, EE), N Africa, Turkey, Jordan, Israel, Iraq, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, China (NE), Korean Peninsula, Japan (Ryu), India.

Ectemnius (Hypocrabro) continuus (Fabricius, 1804) [Crabro] (*Crabro 6-maculatus* Say, 1824, nom. praeocc., nec Olivier, 1792; *C. agrestis* Stephens, 1829, nom. nud.; *C. divisus* Stephens, 1829, nom. nud.; *C. Fuscitarsus* Herrich-Schaeffer, 1840, nom. nud.; *C. fuscitarsis* Herrich-Schaeffer, 1841; *C. vagatus* F. Smith, 1869; *C. granulatus* Walker, 1871; *C. rugosopunctatus* Taschenberg, 1875; *C. validus* De Stefani Perez, 1884; *C. vagans* Fokker, 1887; *Xylocrabro slossonae* Ashmead: Slosson, 1902: 5, nom. nud.; *C. bisexmaculatus* Viereck in J. Smith, 1910, substitute name for *Crabro 6-maculatus* Say, 1824; *C. sayi* Cockerell, 1910, substitute name for *Crabro 6-maculatus* Say, 1824; *Solenius giffardi* Rohwer, 1917). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, TK, NS, AL), **ES** (KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KA). – Europe (WE, NE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Syria, Jordan, Israel, Iran, Iraq, Central Asia, Kazakhstan, Mongolia, China (NW, NE, SW), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu), N America, Cuba, Guatemala, Venezuela.

Ectemnius (Hypocrabro) hypsae (De Stefani Perez, 1884) [Crabro] (*Crabro Hypsae* De Stefani Perez, 1884; *C. serotinus* De-Stefani, 1894; *C. laetus* Pérez: Antiga et Bofill, 1904, nom. nud.; *C. loetus* Pérez, 1905). Russia: **EP** (S). – Europe (WE, SE, EE), N Africa, Turkey, Israel.

Ectemnius (Hypocrabro) meridionalis (A. Costa, 1867) [Crabro] (*Crabro finitimus* F. Morawitz, 1894). Russia: **EP** (S, CR). – Europe (WE, SE, EE), N Africa, Turkey, Syria, Jordan, Israel, Iran, Afghanistan, Central Asia, Kazakhstan.

Ectemnius (Hypocrabro) pedicularis (F. Morawitz, 1889) [Crabro] (*Crabro Nielseni* Kohl, 1915). Russia: **ES** (ZB), **FE** (AM, KH, PR). – Mongolia, China (NE, NC, CE), Korean Peninsula.

Ectemnius (Hypocrabro) rubicola (Dufour et Perris, 1840) [Solenius] (*Crabro Microstictus* Herrich-Schaeffer, 1840, nom. nud.; *C. microstictus* Herrich-Schaeffer, 1841; *C. larvatus* Wesmael, 1852; *C. pumilus* A. Costa, 1871). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (KM, IR), **FE** (AM, PR, SA, KU). – Europe (WE, NE, SE, EE), Turkey, Iran, Kazakhstan, Japan (Hok, Hon, Shi, Kyu).

Ectemnius (Hypocrabro) schlettereri (Kohl, 1888) [Crabro] (*Crabro Schlettereri* Kohl, 1888). Russia:

EP (S), UR, WS (AL), ES (KR, IR, BR, ZB), FE (AM, KH, PR, SA). – Europe (WE, SE, EE), Iran, Kazakhstan, China (NE, CE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu), India, SE Asia.

Ectemnius (Hypocrabro) varentzowi F. Morawitz, 1894 [Crabro] (*Crabro Varentzowi* F. Morawitz, 1894). Russia: **UR, WS (NS), ES (IR)**. – Turkmenistan, Uzbekistan, Kazakhstan.

Ectemnius (Hypocrabro) walteri (Kohl, 1889) [Crabro] (*Crabro Walteri* Kohl in Kohl et Handlirsch, 1889). Russia: **EP (S)**. – Turkey, Turkmenistan, Kazakhstan.

Ectemnius (Metacrabro) cephalotes (Olivier, 1792a) [Crabro] (*Crabro cephalotes* Olivier, 1792; *C. floralis* Olivier, 1792; *C. geniculatus* Olivier, 1792; *C. tibialis* Olivier, 1792; *C. cephalotes* Panzer, 1799, nom. praeocc., nec Olivier, 1792; *C. subinterruptus* Stephens, 1829, replacement name for *C. lituratus* of Panzer, 1804; *C. striatus* Lepeletier de Saint Fargeau et Brullé, 1835; *C. ornatus* Lepeletier de Saint Fargeau et Brullé, 1835; *Ceratocolus striatus* Lepeletier de Saint Fargeau et Brullé, 1835, nom. praeocc., nec *Crabro striatus* Lepeletier de Saint Fargeau et Brullé, 1835; *Blepharipus striatulus* Lepeletier de Saint Fargeau et Brullé, 1835; *Crabro Lindenius* Shuckard, 1837; *C. Shuckardi* Dahlbom, 1839; *C. interruptus* Dahlbom, 1845, nom. praeocc., nec Fabricius, 1787 (now in *Palarus*); *C. Fargeii* Smith, 1856, substitute name for *Ceratocolus striatus* Lepeletier de Saint Fargeau et Brullé, 1835; *Crabro aciculatus* Provancher, 1882; *C. aciculatus* Provancher, 1883, objective synonym of Provancher, 1882; *C. ruthenicus* F. Morawitz, 1892). Russia: **EP (N, NW, C, E, S, NC, CR), UR, WS (AL), FE (PR)**. – Europe (WE, NE, SE, EE), N Africa, Turkey, Syria, Kazakhstan, N America.

Ectemnius (Metacrabro) chrysites (Kohl, 1892) [Crabro] (*Crabro auricomus* Bingham, 1897; *C. khasianus* Cameron, 1902; *E. butuani* Leclercq, 1963). Russia: **FE (PR)**. – China (NE, CE, SW, SE), Korean Peninsula, Japan (Hon, Ryu), SE Asia, India.

Ectemnius (Metacrabro) fossorius (Linnaeus, 1758) [Sphecius] (*Sphecius fossorius* Linnaeus, 1758; *S. scutellata* *Bucephalus* Christ, 1791; *Crabro aspidiphorus* Schrank, 1802; *Solenius fuscipennis* Lepeletier de Saint Fargeau et Brullé, 1835: 714, nom. praeocc., nec Lepeletier de Saint Fargeau et Brullé, 1835: 710; *S. grandis* Lepeletier de Saint Fargeau et Brullé, 1835; *S. tetraedrus* Dahlbom, 1845; *C. fumipennis* F. Smith, 1856, substitute name for *Solenius*

fuscipennis Lepeletier de Saint Fargeau et Brullé, 1835:714, nec Lepeletier de Saint Fargeau et Brullé, 1835:710; *C. montanus* Gistel, 1857; *C. Konowi* Kohl, 1905; *C. sapporoensis* Matsumura, 1911; *C. rubropictus* Matsumura, 1912). Russia: **EP (N, NW, C, E, S, NC, CR), UR, WS (NS, KM, AL), ES (KR, IR, BR, YA, ZB), FE (AM, KH, PR, SA, KU)**. – Europe (WE, NE, SE, EE), N Africa, Turkey, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE), Korean Peninsula, Japan (Hon, Hon).

Ectemnius (Metacrabro) iridifrons (Pérez, 1905) [Crabro] (*E. iridifrons* [sic] *meridionalis* Tsuneki, 1984, nom. praeocc., nec *Crabro meridionalis* A. Costa, 1871). Russia: **FE (PR)**. – Korean Peninsula, Japan (Hon, Hon, Shi, Kyu, Ryu), SE Asia.

Ectemnius (Metacrabro) lituratus (Panzer, 1803) [Crabro] (*Solenius petiolatus* Lepeletier de Saint Fargeau et Brullé, 1835; *Ceratocolus fasciatus* Lepeletier de Saint Fargeau et Brullé, 1835; *Ceratoculus* [*lapsus calami*] *reticulatus* Lepeletier de Saint Fargeau et Brullé, 1835; *Crabro Kollari* Dahlbom, 1845; *C. argenteus* Schenck, 1857; *C. vestitus* F. Smith, 1858; *C. intermedius* A. Morawitz, 1866; *C. intermedius* A. Morawitz, 1866: 339, objective synonym of A. Morawitz, 1866: 264; *C. luxuriosus* A. Costa, 1871; *C. kollari* *Dallatorreanus* Kohl, 1880; *C. transiens* Kohl, 1915). Russia: **EP (C, E, S, NC, CR), UR**. – Europe (WE, NE, SE, EE), N Africa, Turkey, Iran, Kazakhstan.

Ectemnius (Metacrabro) spinipes (A. Morawitz, 1866) [Crabro] (*Crabro spinipes* A. Morawitz, 1866b:341, objective synonym of A. Morawitz, 1866a:265; *C. Bulsanensis* Kohl, 1879). Russia: **EP (C, E, NC, CR), UR, WS (TK, NS, AL), ES (KR, IR), FE (AM, KH, PR, SA)**. – Europe (WE, NE, C, EE), Azerbaijan, Turkey, Turkmenistan, Kazakhstan, Mongolia, China (NE), Korean Peninsula, Japan (Hon, Hon).

Ectemnius (Thyreocerus) crassicornis (Spinola, 1808) [Crabro] (*Crabro hybridus* Eversmann, 1849; *E. punctulatus* De-Stefani, 1884; *E. Siculus* De Stefani Perez, 1884). Russia: **EP (E, S, NC, CR), UR, WS (AL), ES (KR, IR), FE (PR, SA, KU)**. – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey, Syria, Jordan, Israel, Iraq, Iran, Central Asia, Kazakhstan.

Ectemnius (Yanonius) martjanowi (F. Morawitz, 1892) [Crabro] (*Crabro Martjanowi* F. Morawitz, 1892; *Clytochrysis dubiosus* Ashmead, 1904). Russia: **WS (NS, AL), ES (KR, IR), FE (AM, KH, PR, SA, KU)**. – Mongolia, China (SW), Korean Peninsula, Japan (Hon, Hon), India.

ENTOMOGNATHUS Dahlbom, 1844 (*Florkinus* Leclercq, 1956). Type species *Crabro brevis* Vander Linden, 1829. The genus includes four subgenera: *Entomognathus* Dahlbom, 1844, *Koxinga* Pate, 1944, *Mashona* Pate, 1944, and *Toncahua* Pate, 1944. Number of species: World – 64, Palaearctic – 11, Russia – 4.

Entomognathus brevis (Vander Linden, 1829) [Crabro] (*Lindenius apicalis* Lepeletier de Saint Fargeau et Brullé, 1835; *L. nasutus* Gribodo, 1884). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (IR, YA), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), NW Africa, Georgia, Turkey, Syria, Israel, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NE, NC), Japan (Hok, Hon).

Entomognathus dentifer (Noskiewicz, 1929) [Crabro] (*E. permixtus* Nouvel et Ribaut, 1956). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), Turkey.

Entomognathus sahlbergi (A. Morawitz, 1866) [Crabro] (*C. Sahlbergi* A. Morawitz, 1866). Russia: **FE** (KH). – Mongolia.

Entomognathus schmiedeknechti (Kohl, 1905) [Crabro] (*C. Schmiedeknechti* Kohl, 1905). Russia: **EP** (C, S, NC, CR). – Europe (SE, EE), Turkey.

LESTICA Billberg, 1820 (*Solenius* Lepeletier de Saint Fargeau et Brullé, 1835; *Ceratocolus* Lepeletier de Saint Fargeau et Brullé, 1835; *Thyreus* Lepeletier de Saint Fargeau et Brullé, 1835, nom. praeocc., nec Panzer, 1806, non Swainson, 1821; *Hypothyreus* Ashmead, 1899; *Clypeocrabro* Richards, 1935; *Ptyx* Pate, 1947). Type species *Crabro subterraneus* Fabricius, 1775. Number of species: World – 44, Palaearctic – 12, Russia – 9.

Lestica alata (Panzer, 1797) [Crabro] (*Crabro basalis* F. Smith, 1856; *C. alatus japonicus* Schulz, 1904; *C. alatus rubidus* Schirmer, 1915; *L. spinicollis* Tsuneki, 1976). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (NS, AL), **ES** (KR, IR, BR, YA, ZB), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), N Africa, Turkey, Kyrgyzstan, Kazakhstan, Mongolia, China (NE, NC, CE), Korean Peninsula, Japan (Hok), N India.

Lestica camelus (Eversmann, 1849) [Crabro] (*Crabro sapporensis* Matsumura, 1912). Russia: **EP** (C, E), **UR**, **WS** (NS, KM, AL), **ES** (KR, IR, BR, YA), **FE** (AM, KH, PR, SA). – Turkey, Kazakhstan, Mongolia, China (NE, SW), Korean Peninsula, Japan (Hok, Hon).

Lestica clypeata (Schreber, 1759) [Apis] (*Sphex clypearia* Schreber, 1784, misspelling or substitute name for *Apis clypeata* Schreber, 1759; *S. clypeata* ovata Christ, 1791; *Crabro vexillatus* Panzer, 1797; *C. lapidarius* Fabricius, 1804, nom. praeocc., nec Panzer, 1804; *Philanthes clypeatus* Thunberg, 1815, nom. praeocc., nec *Apis clypeata* Schreber, 1759; *Crabro Nigridens* Herrich-Schaeffer, 1840, nom. nud.; *C. nigridens* Herich-Schaeffer, 1841; *C. quadrifer* Dufour, 1841). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, KM, AL), **ES** (IR, BR), **FE** (PR). – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, Syria, Lebanon, Israel, Iraq, Iran, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan.

Lestica collaris (Matsumura, 1912) [Crabro] (*Crabro aberrans* Gussakovskij, 1932). Russia: **FE** (KH, PR). – Korean Peninsula, Japan (Hok, Hon, Kyu).

Lestica constricta Krombein, 1949. Russia: **FE** (PR). – China (SE), SE Asia.

Lestica heros (Kohl, 1915) [Crabro]. Russia: **FE** (PR). – China (SE), Korean Peninsula, Japan (Hok, Hon).

Lestica pluschtschevskyi (F. Morawitz, 1891) [Crabro] (*Crabro Pluschtschevskyi* F. Morawitz, 1891). Russia: **UR**, **WS** (AL). – Spain, Turkey, Kazakhstan.

Lestica reiteri (Kohl, 1915) [Crabro] (*Crabro Reiteri* Kohl, 1915; *C. reiteri kuramaensis* Iwata, 1938). Russia: **FE** (KH, PR). – Korean Peninsula, Japan (Hok, Hon).

Lestica subterranea ochotica (A. Morawitz, 1866) [Crabro] (*Crabro ochoticus* A. Morawitz, 1866a; *C. ochoticus* A. Morawitz, 1866b: 337, objective synonym of *C. ochoticus* A. Morawitz, 1866a: 128). Russia: **WS** (AL), **ES** (KR, BR, YA, ZB), **FE** (KH, PR). – Mongolia, China (CE).

Lestica subterranea subterranea (Fabricius, 1775) [Crabro] (*Crabro philanthoides* Panzer, 1801). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), N Africa, Turkey, Iran, Tajikistan, Kyrgyzstan, Kazakhstan, China (NC).

LINDENIUS Lepeletier de Saint Fargeau et Brullé, 1835 (*Chalcolamprus* Wesmael, 1852; *Trachelosimus* A. Morawitz, 1866). Type species *Crabro albilabris* Fabricius, 1793. Number of species: World – 64, Palaearctic – 53, Russia – 13.

Lindenius aegyptius (Kohl, 1888) [Crabro]. Russia: **EP** (S, NC). – Europe (WE), N Africa, Jordan, United Arab Emirates, Pakistan, Uzbekistan, Kazakhstan.

Lindenius albilabris (Fabricius, 1793) [Crabro] (*Crabro aenescens* Dahlbom, 1839). Russia: **EP** (N, NW, C,

- E, S, NC, CR), **UR**, **WS** (TK, KM, AL), **ES** (IR, BR, YA, ZB), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, Syria, Israel, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, Korean Peninsula.
- Lindenius altaicus** Mokrousov, 2016. Russia: **WS** (AL).
- Lindenius gussakovskii** Marshakov, 1973. Russia: **EP** (S). – Uzbekistan, Kyrgyzstan, Kazakhstan.
- Lindenius ibex** Kohl, 1883. Russia: **EP** (C), **WS** (AL). – Greece, Bulgaria, Turkey, Syria, Israel.
- Lindenius laevis** A. Costa, 1867 (*Crabro rhaibopus* Kohl, 1915). Russia: **EP** (C, S, NC), **UR**. – Europe (SE, EE), Turkey, Kazakhstan.
- Lindenius latitarsis** Marshakov, 1973. Russia: **EP** (S, CR), **UR**. – Europe (EE), Turkey, Kazakhstan.
- Lindenius mesopleuralis** (F. Morawitz, 1890) [Crabro]. Russia: **EP** (E, S, NC, CR), **UR**, **WS** (NS, AL), **ES** (ZB), **FE** (PR). – Europe (WE, SE, EE), Turkey, Turkmenistan, Kazakhstan, Mongolia, China (NC, NW).
- Lindenius panzeri** (Vander Linden, 1829) [Crabro] (*L. venustus* Lepeletier de Saint Fargeau et Brullé, 1835; *Crabro latebrosus* Kohl, 1905; *L. harbinensis* Tsuneki, 1967; *L. panzeri mongolicus* Tsuneki, 1972). Russia: **EP** (C, E, S, NC), **UR**, **WS** (TM, AL), **ES** (IR, ZB), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), N Africa, Turkey, Iran, Afghanistan, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE).
- Lindenius parkanensis** (Zavadil, 1948) [Crabro] (*Crabro parkanensis* Zavadil in Zavadil et Šnolflak, 1948; *L. ponticus* de Beaumont, 1956). Russia: **EP** (C, S), **UR**, **WS** (AL). – Europe (SE, EE), Kazakhstan.
- Lindenius pygmaeus** (Rossi, 1794) [Crabro]. Russia: **EP** (C, E, S, NC), **UR**. – Europe (WE, NE, SE, EE), N Africa, Turkey, Syria, Jordan, Israel, Iran, Afghanistan, Central Asia, Kazakhstan.
- Lindenius satschouanus** (Kohl, 1915) [Crabro] (*Crabro satschouanus* Kohl, 1915; *L. kaszabi* Tsuneki, 1972). Russia: **EP** (S). – Turkey, Afghanistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (CE).
- Lindenius subaeneus** Lepeletier de Saint Fargeau et Brullé, 1835. Russia: **EP** (S), **WS** (AL). – Europe (WE, SE, EE), N Africa, Kazakhstan.
- ODONTOCRABRO** Tsuneki, 1971. Type species *Odontocrabro abnormis* Tsuneki, 1971. Number of species: World – 8, Palaearctic – 6, Russia – 1.
- Odontocrabro antropovi** Leclercq, 1991. Russia: **FE** (PR).
- RHOPALUM** Stephens, 1829 (*Euplilis* Risso, 1826, suppressed by ICZN (1978, Opinion 1106); *Rhopalum* Stephens, 1829, conserved by ICZN (1978, Opinion 1106); *Physoscelus* Lepeletier de Saint Fargeau et Brullé, 1835; *Physoscelis* Westwood, 1839, lapsus or emendation of *Physoscelus* Lepeletier de Saint Fargeau et Brullé, 1835; *Dryphus* Herrich-Schaeffer, 1840; *Alliognathus* Ashmead, 1899). Type species *Crabro rufiventris* Panzer, 1799. The genus includes seven subgenera: *Rhopalum* Stephens, 1829, *Corynopus* Lepeletier de Saint Fargeau et Brullé, 1835, *Calceorhopalum* Tsuneki, 1952, *Latrorhopalum* Tsuneki, 1952, *Aporhopalum* Leclercq, 1955, *Zelorrhopalum* Leclercq, 1955, and *Notorhopalum* Leclercq, 1979. Number of species: World – 287, Palaearctic – 21, Russia – 12.
- Rhopalum (Calceorhopalum) pygidiale** Bohart in Bohart et Menke, 1976, (substitute name for *Crabro calceatus* Tsuneki, 1947, nom. praeocc., nec Rossi, 1794). Russia: **FE** (AM, PR, SA, KU). – Korean Peninsula, Japan (Hok, Hon).
- Rhopalum (Calceorhopalum) watanabei** Tsuneki, 1952 (*Rh. arasianum* Tsuneki, 1972). Russia: **FE** (PR, SA, KU). – Japan (Hok, Hon), SE Asia.
- Rhopalum (Corynopus) coarctatum** (Scopoli, 1763) [*Sphecius*] (*Crabro crassipes* Fabricius, 1798; *C. tibialis* Fabricius, 1798, nom. praeocc., nec Olivier, 1792; *Rh. modestum* Rohwer, 1908). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TK, NS, AL), **ES** (IR), **FE** (AM, KH, PR, KA, MG). – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, Iran, Kazakhstan, China (CE), Korean Peninsula, Japan (Hok, Hon), N America.
- Rhopalum (Corynopus) gracile** Wesmael, 1852 (*Rh. nigrinum* Kiesenwetter, 1849, nom. praeocc., nec *Crabro nigrinus* Herrich-Schaeffer, 1841; *Rh. Gracile* Wesmael, 1852; *Crabro Kiesenwetteri* A. Morawitz, 1866, substitute name for *C. nigrinus* (Kiesenwetter, 1849); *Corynopus simplices* F. Morawitz, 1888). Russia: **EP** (C, S, NC), **UR**, **WS** (AL), **FE** (KH, PR, SA, KU, MG, KA). – Europe (WE, NE, SE, EE), Turkey, Uzbekistan, Kazakhstan, Korean Peninsula, Japan (Hok, Hon), N America.
- Rhopalum (Corynopus) kawabatai** Marshakov, 1976. Russia: **FE** (AM, KH, PR, SA).
- Rhopalum (Latrorhopalum) laticorne** (Tsuneki, 1947) [Crabro]. Russia: **ES** (IR, ZB), **FE** (AM, KH, PR, SA). – China (NE), Korean Peninsula.

Rhopalum (Latrorhopalum) latronum (Kohl, 1915)
[Crabro]. Russia: **ES** (ZB), **FE** (KH, PR, SA, KU). – China (NC), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Rhopalum (Rhopalum) austriacum (Kohl, 1899)
[Crabro]. Russia: **EP** (CR), **FE** (PR, SA). – Europe (WE, SE, EE), N Africa, Abkhazia, Turkey, Israel.

Rhopalum (Rhopalum) clavipes (Linnaeus, 1758)
[Sphex] (*Crabro rufiventris* Panzer, 1799). Russia: **EP** (N, C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (IR), **FE** (PR, KU, KA). – Europe (WE, NE, EE), N Africa, Uzbekistan, Kazakhstan, Mongolia, Korean Peninsula, Japan (Hok), N America.

Rhopalum (Rhopalum) jessonicum (Bischoff, 1921)
[Crabro] (*Rh. ebetsuense* Tsuneki, 1952). Russia: **FE** (PR, SA, KU). – Korean Peninsula, Japan (Hok, Hon).

Rhopalum (Rhopalum) kuwayamai Tsuneki, 1952.
Russia: **FE** (PR, SA, KU). – China (CE), Korean Peninsula, Japan (Hok, Hon).

Rhopalum (Rhopalum) mushaense Tsuneki, 1971.
Russia: **FE** (PR). – SE Asia.

TRACHELIODES A. Morawitz, 1866 (*Brachymerus* Dahlbom, 1845, nom. praeocc., nec *Chevrolet* in Dejean, 1835; *Fertonius* Pérez in Ferton, 1892). Type species *Crabro megerlei* Dahlbom, 1845. Number of species: World – 17, Palaearctic – 5, Russia – 2.

Tracheliodes alinae Nemkov, 1988. Russia: **FE** (PR).
Tracheliodes ghilarovi Nemkov, 1988. Russia: **FE** (PR).

TSUNEKIOLA Antropov, 1986. Type species *Tsunekiola tracheliformis* Antropov, 1986. Monotypic Palaearctic genus.

Tsunekiola tracheliformis Antropov, 1986. Russia: **FE** (PR).

Tribe LARRINI

GASTROSERICUS Spinola, 1839 (*Eparamostethus* Kohl, 1907; *Paralellopsis* Maidl, 1914; *Dinetomorpha* Gussakovskij, 1931, unavailable under ICBN (1999, Article 13.3); *Gastrargyron* Gussakovskij, 1931, unavailable under ICBN (1999, Article 13.3); *Dinetomorpha* Pate, 1937, validation of *Dinetomorpha* Gussakovskij, 1931 under ICBN (1999, Article 13.1.2); *Gastrargyron* Pate, 1937:28, validation of *Gastrargyron* Gussakovskij, 1931 under ICBN (1999, Article 13.1.2)). Type species

Gastrosericus waltlii Spinola, 1839. Number of species: World – 61, Palaearctic – 14, Russia – 1.

Gastrosericus waltlii Spinola, 1839 (*Dinetus niger* Dufour, 1854; *G. Maracandicus* Radoszkowski, 1877; *G. rufiventris* F. Morawitz, 1889; *G. rufitarsis* Cameron, 1902; *G. lanuginosus* Arnold, 1922; *G. maracandicus dubius* Gussakovskij, 1931; *G. aiunensis* Giner Marí, 1945). Russia: **EP** (S, NC). – Europe (SE), N Africa, Armenia, Turkey, Cyprus, Syria, Jordan, Israel, Oman, Quatar, Saudi Arabia, United Arab Emirates, Yemen, Iraq, Iran, Pakistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, China (NC), India, Africa.

LARRA Fabricius, 1793 (*Larrana* Rafinesque, 1815, emendation or substitute name for *Larra* Fabricius, 1793; *Lara* Drapiez, 1819, lapsus or emendation of *Larra* Fabricius, 1793; *Monomatium* Shuckard, 1840; *Lyrops* Dahlbom, 1843, nom. praeocc., nec Illiger, 1807; *Larraxena* F. Smith, 1851; *Lara* LeConte, 1852, nom. praeocc., nec Drapiez, 1819; *Larrada* F. Smith, 1856). The genus includes two subgenera: *Larra* Fabricius, 1793 and *Cratolarra* Cameron, 1900. Type species *Larra ichneumoniformis* Fabricius, 1793 (= *Sphex anathema* Rossi, 1790). Number of species: World – 65, Palaearctic – 9, Russia – 1.

Larra anathema (Rossi, 1790) [Sphex] (*L. ichneumoniformis* Fabricius, 1793; *Pompilus teutonus* Fabricius, 1804; *Tachytes grandis* Chevrier, 1872). Russia: **EP** (C, E, S, NC, CR), **WS** (NS). – Europe (WE, SE, EE), N Africa, Georgia, Turkey, Syria, Iraq, Iran, Saudi Arabia, Bahrain, Israel, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, China (NE).

LARROPSIS Patton, 1892. Type species *Larrada tenuicornis* F. Smith, 1856. The genus includes two subgenera: *Larropsis* Patton, 1892 and *Ancistromma* Fox, 1893. Number of species: World – 40, Palaearctic – 4, Russia – 1.

Larropsis (Ancistromma) asiatica (Gussakovskij, 1935) [*Ancistromma*] (*Ancistromma shestakovi* Gussakovskij, 1935). Russia: **EP** (S). – Turkey, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

LIRIS Fabricius, 1804 (*Liris* Rafinesque, 1815, emendation or substitute name for *Liris* Fabricius, 1804; *Notogonia* A. Costa, 1867, nom. praeocc., nec Perty, 1850 [Crustaceal]; *Caenolarra* Cameron, 1900; *Spanolarra* Cameron, 1900; *Notogonius* Howard, 1901, lapsus for *Notogonia* A. Costa, 1867; *Chrysolarra*

Cameron, 1901; *Notogonidea* Rohwer, 1911; *Dociliris* Tsuneki, 1967; *Nigliris* Tsuneki, 1967; *Colloliris* Tsuneki, 1974). The genus includes four subgenera: *Liris* Fabricius, 1804, *Motes* Kohl, 1897, *Leptolarra* Cameron, 1900, and *Pitaliris* Tsuneki, 1982. Type species *Sphex auratus* Fabricius, 1787, nom. praeocc., nec Linnaeus, 1758 (= *Sphex aurulentus* Fabricius, 1787). Number of species: World – 314, Palaearctic – 17, Russia – 6.

Liris atratus (Spinola, 1805) [Larra] (*Larra micans* Spinola, 1806, unjustified emendation of *Larra atrata* Spinola, 1905; *Tachytes nigrita* Lepeletier de Saint Fargeau, 1845; *Notogonia pharaonum* Kohl, 1906). Russia: UR. – Europe (WE, SE), N Africa, United Arab Emirates, Kuwait, Turkey, Jordan, Israel, Iraq, Iran, Afghanistan, Pakistan, Kazakhstan.

Liris festinans preaetermissus (Richards, 1928) [*Notogonia*] (*Leptolarra schulthessi* Giner Marí, 1942). Russia: EP (NC). – Europe (WE, SE, EE), N Africa, Oman, United Arab Emirates, Saudi Arabia, Israel, Abkhasia, Turkey, Iran, Afghanistan, Uzbekistan.

Liris inopinatus de Beaumont, 1961. Russia: EP (S). – Europe (SE), Turkey, Syria, Israel.

Liris niger (Fabricius, 1775) [Sphe] (*Sphex nigra* Fabricius, 1775; *Larra pompiliformis* Panzer, [1807–1809], nom. praeocc., nec Panzer, 1804; *Anoplus unioellatus* Dufour, 1833; *Larra confusa* Radoszkowski, 1887; *L. nigriventris* Cameron, 1889; *Tachytes germabensis* Radoszkowski, 1893; *Larra nana* Bingham, 1897; *L. iridipennis* Cameron, 1900; *Notogonidea baguenai* Giner Marí, 1934). Russia: EP (S, NC, CR). – Europe (WE, SE, EE), N Africa, Yemen, Turkey, Jordan, Iran, Iraq, Afghanistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NE, CE), SE Asia, India.

Liris nigricans (Walker, 1871) [Larrada] (*Notogonia argyropyga* A. Costa, 1875; *N. sculpturata* Kohl, 1892; *N. reticulata* de Saussure, 1892; *Larra Jugurtha* Gribodo, 1894; *L. palumbula* Kohl, 1894; *N. mahensis* Cameron, 1907; *N. punctipleura* Cameron, 1908). Russia: EP (NC). – Europe (WE, SE, EE), N Africa, Yemen (Socotra), Oman, United Arab Emirates, Turkey, Israel, Iraq, Iran, Afghanistan, Tajikistan, Uzbekistan, Kazakhstan, China (CE), Japan (Kyu), India, Africa.

Liris opalipennis (Kohl, 1898) [*Notogonia*]. Russia: EP (S). – N Africa, United Arab Emirates, Turkmenistan, Tajikistan, Kazakhstan.

PARAPIAGETIA Kohl, 1897a (*Lirosphe* Brèthes, 1913; *Psammosphe* Gussakovskij, 1952). Type species *Piagetia odontostoma* Kohl, 1884. Number of species: World – 29, Palaearctic – 17, Russia – 2.

Parapiagetia kaszabi Tsuneki, 1972. Russia: EP (S). – Turkmenistan, Kazakhstan, Mongolia, China (NW).

Parapiagetia tridentata Tsuneki, 1972. Russia: EP (S). – Iran, Pakistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NW).

PROSOPIGASTRA A. Costa, 1867 (*Homogambrus* Kohl, 1889; *Hologambrus* Morice, 1897, lapsus or emendation of *Homogambrus* Kohl, 1889). Type species *Prosopigastra punctatissima* A. Costa, 1867. Number of species: World – 36, Palaearctic – 22, Russia – 5.

Prosopigastra bulgarica Pulawski, 1958. Russia: EP (S). – Europe (SE, EE), Turkey, Iran, Kazakhstan.

Prosopigastra globiceps (F. Morawitz, 1889) [*Tachysphex*]. Russia: EP (S). – N Africa, United Arab Emirates, Turkmenistan, Kazakhstan, Mongolia, China (NC).

Prosopigastra latifrons Gussakovskij, 1933. Russia: EP (S). – Turkmenistan, Kazakhstan.

Prosopigastra orientalis de Beaumont, 1954 (*P. punctatissima orientalis* de Beaumont, 1947). Russia: EP (S, NC, CR). – Europe (SE, EE), Turkey, Israel, Tajikistan, Uzbekistan, Kazakhstan.

Prosopigastra zalinda de Beaumont, 1955. Russia: EP (S, NC). – Europe (WE, SE, EE), N Africa, Turkey, Israel, Kazakhstan, Mongolia.

TACHYSPEX Kohl, 1883 (*Schistosphe* Arnold, 1922; *Atelosphe* Arnold, 1923). Type species *Tachysphex filicornis* Kohl, 1883. Number of species: World – 446, Palaearctic – 136, Russia – 38.

Tachysphex angustatus Pulawski, 1967. Russia: EP (S). – Europe (SE), Turkey, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NC).

Tachysphex argentatus Gussakovskij, 1952. Russia: EP (S). – N Africa, Israel, Saudi Arabia, Turkey, Syria, Turkmenistan, Tadzhikistan, Kazakhstan.

Tachysphex blattivorus Gussakovskij, 1952. Russia: EP (S). – Europe (WE), Tajikistan, Uzbekistan, Kazakhstan.

Tachysphex brevipennis Mercet, 1909 (*T. minutus* Nurse, 1909; *T. lilliputianus* R. Turner, 1917; *Atelosphe miscophoides* Arnold, 1923; *A. lugubris*

- Arnold, 1924; *T. imperfectus* de Beaumont, 1940; *T. rugosus* Gussakovskij, 1952; *T. heterochromus* de Beaumont, 1955; *T. rhodius* de Beaumont, 1960; *T. sinaiticus* Pulawski, 1964; *T. convexus* Pulawski, 1971). Russia: **EP** (S, CR). – Europe (WE, SE, EE), N Africa, Turkey, Syria, Jordan, Israel, Saudi Arabia, United Arab Emirates, Oman, Iran, Turkmenistan, Tajikistan, Kazakhstan, India, Sri Lanka, Africa.
- Tachysphex brullii** (F. Smith, 1856) [Tachytes] (*Tachytes bicolor* Brullé, 1833, nom. praeocc., nec *Larra bicolor* Fabricius, 1804; *T. Brullii* F. Smith, 1856, substitute name for *T. bicolor* Brullé, 1833; *T. spoliata* Giraud, 1863; *T. Brulléi* Kirchner, 1867, emendation of *T. brullii* F. Smith, 1856; *T. rufipes* von Aichinger, 1870; *T. montanus* Radoszkowski, 1886, nom. praeocc., nec *Larrada montana* Cresson, 1865; *T. monticola* Dalla Torre, 1897, substitute name for *T. montanus* Radoszkowski, 1886). Russia: **EP** (C, S, NC, CR). – Europe (WE, SE, EE), N Africa, Turkey, Israel, Turkmenistan, Kazakhstan.
- Tachysphex consocius** Kohl, 1892 (*T. Cabrerai* Mercet, 1909; *T. minutulus* Arnold, 1923; *T. grandii* de Beaumont, 1965). Russia: **EP** (C, S, NC, CR). – Europe (WE, SE, EE), N Africa, Abkhazia, Azerbaijan, Turkey, Syria, Iran, Afghanistan, Saudi Arabia, Oman, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, India, Africa.
- Tachysphex coriaceus** (A. Costa, 1867) [Tachytes] (*Tachytes coriacea* A. Costa, 1867b; *Tachysphex coriaceus* A. Costa, 1867c, objective synonym of *T. coriaceus* A. Costa, 1867b; *T. Reiseri* Kohl, 1901). Russia: **EP** (S). – Europe (WE, SE, EE), Turkey.
- Tachysphex costae** (De Stefani Perez, 1882) [Tachytes] (*Tachytes Costae* De Stefani Perez, 1882; *Tachysphex laticeps* Arnold, 1924; *T. zarudnyi* Gussakovskij, 1933; *T. tadzhikus* Gussakovskij, 1952; *T. costai canariensis* de Beaumont, 1968; *T. costai fertoni* Pulawski, 1971). Russia: **EP** (S, CR). – Europe (WE, SE, EE), N Africa, Turkey, Israel, Iran, Turkmenistan, Tajikistan, Kazakhstan, Africa.
- Tachysphex ctenophorus** Pulawski, 1971. Russia: **EP** (S), **ES** (BR), **FE** (PR). – Kazakhstan.
- Tachysphex erythropus** (Spinola, 1839) [Lyrops] (*Tachytes Maracandica* Radoszkowski, 1877, *T. flavogeniculatus* Taschenberg, 1880; *Tachysphex heliopolites* Morice, 1897; *T. inventus* Nurse, 1903; *T. mantivora* de Beaumont, 1940). Russia: **EP** (S, NC). – Europe (WE, SE, EE), N Africa, Saudi Arabia, United Arab Emirates, Oman, Turkey, Israel, Iran, Pakistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, China (NW).
- Tachysphex excelsus** R. Turner, 1917 (*T. mysticus* Pulawski, 1971). Russia: **WS** (AL). – Europe (WE, SE), Turkey, Pakistan, Kazakhstan, Mongolia, China (NW).
- Tachysphex ferrugineus** Pulawski, 1971. Russia: **EP** (C, NC). – Europe (EE), Turkey, Kazakhstan.
- Tachysphex fugax** (Radoszkowski, 1877) [Tachytes] (*T. filicornis* Kohl, 1883; *T. micromegas* de Saussure, 1892; *T. sikorae* de Saussure, 1892; *T. filicornis excerptus* R. Turner, 1917). Russia: **EP** (S, NC). – Europe (WE, SE, EE), N Africa, Turkey, Syria, Saudi Arabia, Oman, Yemen, Israel, Uzbekistan, Kazakhstan, Africa.
- Tachysphex fulvitarsis** (A. Costa, 1867) [Tachytes] (*Tachytes fulvitarsis* A. Costa, 1867c, objective synonym of A. Costa, 1867b; *T. acrobates* Kohl, 1878; *T. strigosus* Mocsáry, 1879; *T. erythrogastera* A. Costa, 1882; *T. erythrogastera* A. Costa, 1884, objective synonym of A. Costa, 1882; *T. dubius* Radoszkowski, 1886; *Tachysphex bipunctatus* F. Morawitz, 1891). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (NS, AL), **ES** (BR, YA, ZB), **FE** (AM, PR). – Europe (WE, NE, SE, EE), N Africa, Turkey, Syria, Jordan, Israel, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.
- Tachysphex gibbus** Kohl, 1885 (*T. lativalvis gibba* Kohl, 1885; *T. philippi* S. Saunders, 1910). Russia: **EP** (NC, CR). – Europe (WE, SE), N Africa.
- Tachysphex gussakovskii** Pulawski, 1971. Russia: **EP** (S). – Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.
- Tachysphex helveticus** Kohl, 1885. Russia: **EP** (C, E, S, NC), **UR**, **WS** (AL), **ES** (KR, IR, BR, ZB), **FE** (PR). – Europe (WE, NE, SE, EE), N Africa, Turkey, Jordan, Tajikistan, Kazakhstan, Mongolia.
- Tachysphex incertus** (Radoszkowski, 1877) [Tachytes] (*Tachytes incerta* Radoszkowski, 1877; *T. panzeri geniculata* A. Costa, 1867b, nom. praeocc., nec *Lyrops geniculata* Spinola, 1839; *T. panzeri geniculata* A. Costa, 1867c, objective synonym of A. Costa, 1867b; *Tachysphex pygidialis* Kohl, 1883; *T. rufiventralis* Ferton, 1905; *T. incertus kallipygus* Pulawski, 1971). Russia: **EP** (S, NC, CR), **UR**, **WS** (AL). – Europe (WE, SE, EE), Turkey, Oman, Yemen, Israel, Jordan, Lebanon, Iran, Afghanistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, China (NW).
- Tachysphex julliani** Kohl, 1883 (*T. Julliani* Kohl, 1883; *T. semenovi* Gussakovskij, 1933; *T. julliani africanus* Pulawski, 1971). Russia: **EP** (S, CR). – Europe (WE,

- SE, EE), N Africa, Turkey, Lebanon, Israel, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.
- Tachysphex latifrons** Kohl, 1884 (*T. cyrenaicus* de Beaumont, 1947). Russia: **EP** (S, CR), **WS** (AL). – Europe (SE, EE), N Africa, Azerbaijan, Turkey, Israel, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, China (NC).
- Tachysphex mediterraneus** Kohl, 1883 (*T. mediterranea* Kohl, 1883; *T. collaris* Kohl, 1898; *T. lanatus* Arnold, 1947). Russia: **EP** (C, S, NC, CR). – Europe (WE, SE, EE), Turkey, Israel, Kazakhstan, India, Sri Lanka, Africa.
- Tachysphex melas** Kohl, 1898. Russia: **EP** (S), **WS** (AL), **ES** (IR), **FE** (PR). – N Africa, Turkey, Jordan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia.
- Tachysphex micans** (Radoszkowski, 1877) [Tachytes]. Russia: **EP** (S). – N Africa, Oman, United Arab Emirates, Turkmenistan, Tajikistan, Kazakhstan, Mongolia.
- Tachysphex mocsaryi** Kohl, 1884 (*T. Mocsaryi* Kohl, 1884; *T. algira* Kohl, 1892; *T. mocsaryi maroccana* de Beaumont, 1947). Russia: **EP** (S, CR), **UR**. – Europe (WE, SE, EE), N Africa, Turkey, Israel, Jordan, Afghanistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.
- Tachysphex moczari** Tsuneki, 1972. Russia: **ES** (ZB), **FE** (PR). – Mongolia, China (NC).
- Tachysphex nigricolor** (Dalla Torre, 1897) [Larra] (*Larrada nigricans* F. Smith, 1873, nom. praeocc., nec Walker, 1871; *Larra nigricolor* Dalla Torre, 1897, substitute name for *Larrada nigricans* F. Smith, 1873; *T. japonicus* Iwata, 1933). Russia: **FE** (AM, KH, PR, KU). – China (NE, SE), Korean Peninsula, Japan (Hon, Kyu, Ryu), SE Asia.
- Tachysphex nitidior** de Beaumont, 1940. Russia: **EP** (C, S, NC, CR). – Europe (WE, SE, EE), N Africa, Turkey, Israel, Lebanon, Iran, Turkmenistan, Uzbekistan, Kazakhstan, Mongolia, SE Asia.
- Tachysphex nitidus** (Spinola, 1805) [Astata] (*Tachytes ibericus* de Saussure, 1867; *Tachysphex ibericus borealis* Pulawski, 1971). Russia: **EP** (N, C, E, S, NC, CR), **WS** (NS, AL). – Europe (WE, NE, SE, EE), N Africa, Turkey, Israel, Kuwait, Oman, Turkmenistan, Tajikistan, Kazakhstan, China (NC).
- Tachysphex obscuripennis** (Schenck, 1857) [Tachytes] (*Tachytes obscuripennis* Scheck, 1860, nom. nud.; *T. lativalvis* Thomson, 1870). Russia: **EP** (N, C, E, S, NC), **WS** (AL). – Europe (WE, NE, SE, EE), Morocco, Turkey, Lebanon.

- Tachysphex panzeri** (Vander Linden, 1829) [Tachytes] (*Lyrops rufiventris* Spinola, 1839; *L. haemorrhoa* Spinola, 1843, unavailable under ICZN (1999, Article 11.6); *Tachytes Oraniensis* Lepeletier de Saint Fargeau, 1845; *T. aurifrons* Lucas, 1849; *T. discolor* Frivaldszky, 1877; *T. pulverosus* Radoszkowski, 1886; *T. ceylonica* Cameron, 1900; *T. aurifrons* Cameron, 1900, junior primary homonym of Lucas, 1849; *Tachysphex ablatus* Nurse, 1909; *T. panzeri fortunatus* de Beaumont, 1968; *T. panzeri cyprius* Pulawski, 1971; *T. panzeri sareptanus* Pulawski, 1971). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, NE, SE, EE), N Africa, Turkey, Syria, Israel, Jordan, Saudi Arabia, Yemen, Bahrain, United Arab Emirates, Djibouti, Iran, Pakistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, India, Sri Lanka.
- Tachysphex pomphiliformis** (Panzer, 1804) [Larra] (*Larrada parvula* Cresson, 1865; *Larra Quebecensis* Provancher, 1882; *L. Quebecensis* Provancher, 1883, objective synonym of Provancher, 1882; *T. decorus* W. Fox, 1894; *T. tenuipunctus* W. Fox, 1894; *T. consimilis* W. Fox, 1894; *T. rufo-niger* Bingham, 1897; *T. projectus* Nurse, 1903; *T. argyrotrichus* Rohwer, 1911; *T. granulosus* Mickel, 1916; *T. erythraeus* Mickel, 1916; *T. angularis* Mickel, 1916). Russia: **EP** (N, C, E, S, NC, CR), **UR**, **WS** (NS, AL), **ES** (KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, MG, KA). – Europe (WE, NE, SE, EE), N Africa, Turkey, Iran, Pakistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NC), Korean Peninsula, India, N America.
- Tachysphex psammobius** (Kohl, 1880) [Tachytes] (*Tachytes psammobia* Kohl, 1880; *Tachysphex asperatus* W. Fox, 1894; *T. sculptilis* W. Fox, 1894; *T. nigrescens* Rohwer, 1908; *T. sphecodoides* Rohwer, 1911). Russia: **EP** (C, S, NC, CR), **UR**, **WS** (TM, AL), **ES** (IR, BR), **FE** (PR). – Europe (WE, SE, EE), N Africa, Turkey, Israel, Kazakhstan, China, N America.
- Tachysphex schmiedeknechti** Kohl, 1883 (*T. Schmiedeknechti* Kohl, 1883; *T. psilopus* Kohl, 1884; *T. helophilus* Nurse, 1909; *T. ornatipennis* Gussakovskij, 1933; *T. fasciipennis* Gussakovskij, 1933; *T. calopteryx* Gussakovskij, 1933; *T. schmiedeknechti satanas* Pulawski, 1971). Russia: **EP** (E, S). – Europe (WE, SE), Saudi Arabia, United Arab Emirates, Oman, Yemen, Turkey, Syria, Jordan, Israel, Iran, Afghanistan, Turkmenistan, Uzbekistan, Kazakhstan, India, Sri Lanka, Africa.

Tachysphex speciosissimus Morice, 1897 (*T. redivivus* Kohl, 1901). Russia: **EP** (S). – N Africa, Syria, Israel, Iran, Uzbekistan.

Tachysphex stachi de Beaumont, 1936. Russia: **EP** (S). – Georgia, Turkey, Israel, Turkmenistan, Uzbekistan, Kazakhstan.

Tachysphex subdentatus F. Morawitz, 1893. Russia: **EP** (S), **WS** (AL). – Europe (SE, EE), Turkey, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Tachysphex tarsinus (Lepeletier de Saint Fargeau, 1845) [Tachytes] (*Tachytes tarsina* Lepeletier de Saint Fargeau, 1845). Russia: **EP** (C, S, NC), **WS** (NS, AL). – Europe (WE, SE, EE), N Africa, Turkey, Saudi Arabia, Oman, Israel, Lebanon, Iran, Tajikistan, Uzbekistan, Kazakhstan, China (NC).

Tachysphex unicolor (Panzer, 1807-1809) [Larra] (*Larra jurinii* Drapiez, 1819). Russia: **EP** (N, C, E, S). – Europe (WE, NE, SE, EE), N Africa, Turkey, Israel, Lebanon, Afghanistan, Turkmenistan, Kazakhstan, Mongolia, China (NE).

TACHYTES Panzer, 1806 (*Lyrops* Illiger, 1807; *Tachyptera* Dahlbom, 1843, nom. praeocc., nec Berge, 1842; *Holotachytes* R. Turner, 1917; *Calotachytes* R. Turner, 1917; *Tachyoides* Banks, 1942; *Tachyplena* Banks, 1942, substitute name for *Tachyptera* Dahlbom, 1843; *Tachynana* Banks, 1942). Type species *Pompilus tricolor* of Fabricius, 1798 (= *Sphex tricolor* Fabricius, 1793, nom. praeocc., nec Schrank, 1781; = *Sphex tricoloratus* Turton, 1801). Number of species: World – 297, Palaearctic – 58, Russia – 8.

Tachytes ambidens Kohl, 1884 (*T. caucasicus* Radoszkowski, 1886). Russia: **EP** (S, NC). – Europe (SE), N Africa, Georgia, Armenia, Turkmenistan, Tajikistan, Kazakhstan, China (NW).

Tachytes argyreus (F. Smith, 1856) [Larrada] (*L. argyrea* F. Smith, 1856; *T. melanopyga* A. Costa, 1893a; *T. melanopyga* A. Costa, 1893b, objective synonym of A. Costa, 1893a; *Tachysphex debilis* Pérez, 1907). Russia: **EP** (S, NC). – N Africa, Turkey, Israel, Saudi Arabia, United Arab Emirates, Bahrain, Iraq, Iran, Pakistan, Turkmenistan, Tajikistan, India.

Tachytes bidens Gussakovskij, 1952. Russia: **EP** (S). – Iran, Turkmenistan, Uzbekistan, Kazakhstan.

Tachytes etruscus etruscus (Rossi, 1790) [Andrena (*A. Etrusca* Rossi, 1790; *T. argentata* Brullé, 1832)]. Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (AL). – Europe (WE, SE, EE), N Africa, Turkey, Syria, Israel, Saudi Arabia, Iran, Turkmenistan, Kazakhstan, Mongolia, Africa.

Tachytes etruscus sibiricus Gussakovskij, 1932 (*T. sibirica* Gussakovskij, 1932; *T. nipponicus* Tsuneki, 1964). Russia: **FE** (KH, PR). – China (NE), Korean Peninsula, Japan (Hon).

Tachytes freygessneri Kohl, 1881 (*Apis hispanica* Linnaeus in Kirby, 1802, nom. praeocc., nec Fabricius, 1787; *T. Frey-Gessneri* Kohl, 1881). Russia: **EP** (S). – Europe (WE, NE, SE, EE), N Africa, Turkey, Iraq, Iran, Tajikistan, Uzbekistan, Kazakhstan, China (NC, NW).

Tachytes latifrons Tsuneki, 1964. Russia: **FE** (PR). – Korean Peninsula, Japan (Hok, Hon).

Tachytes obsoletus (Rossi, 1792) [*Apis*] (*Apis Obsoleta* Rossi, 1792). Russia: **EP** (S, NC), **UR**, **WS** (NS, AL). – Europe (WE, SE, EE), N Africa, Turkey, Israel, Iraq, Tajikistan, Uzbekistan, Kazakhstan, China (NE, NC).

Tachytes panzeri orientis Pulawski, 1962. Russia: **FE** (AM, KH, PR). – China (NE, NC).

Tachytes panzeri panzeri (Dufour, 1841) [Lyrops] (*Lyrops Panzeri* Dufour, 1841, substitute name for *Pompilus tricolor* of Panzer, 1801, nom. praeocc., nec *Tachytes panzeri* Vander Linden, 1829 (now in *Tachysphex*); *Tachytes europaea* Kohl, 1883). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (NS, AL), **ES** (IR). – Europe (WE, SE, EE), N Africa, Turkey, Israel, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Tribe MISCOPHINI

LYRODA Say, 1837 (*Morphota* F. Smith, 1856; *Odon-tolarra* Cameron, 1900; *Lyrodon* Howard, 1901, lapsus pro *Lyroda* Say, 1837, nom. praeocc., nec *Lyrodon* Goldfuss, 1837). Type species *Lyroda subita* Say, 1837. Number of species: World – 25, Palaearctic – 2, Russia – 1.

Lyroda nigra japonica Iwata, 1933 (*L. japonica* Iwata, 1933). Russia: **FE** (KH, PR). – China (NE), Korean Peninsula, Japan (Hon, Kyu, Ryu).

MISCOPHUS Jurine, 1807 (*Nitelopterus* Ashmead, 1897; *Hypomiscophus* Cockerell, 1898; *Miscophinus* Ashmead, 1898). Type species *Miscophus bicolor* Jurine, 1807. Number of species: World – 184, Palaearctic – 131, Russia – 14.

Miscophus albuferiae de Andrade, 1952. Russia: **EP** (NC). – Europe (WE, SE), Turkey.

Miscophus ater Lepeletier de Saint Fargeau, 1845 (*M. maritimus* F. Smith, 1858). Russia: **EP** (C, S), **WS**

- (AL), **ES (IR)**, **FE (AM, PR)**. – Europe (WE, NE, EE), N Africa, Azerbaijan, Turkey, Iran, Kazakhstan.
- Miscophus bicolor** Jurine, 1807 (*Larra dubia* Panzer, 1807-1809; *M. metallicus* C. Verhoeff, 1890; *M. tsunekii* de Andrade, 1960). Russia: **EP (C, E, S, NC)**, **UR, WS (AL)**, **ES (IR, BR, YA, ZB)**, **FE (AM, KH, PR, SA)**. – Europe (WE, NE, SE, EE), N Africa, Abkhasia, Turkey, Syria, Iran, Kazakhstan, Mongolia, Korean Peninsula.
- Miscophus concolor** Dahlbom, 1844 (*M. moravicus* Balthasar, 1957). Russia: **EP (C, E)**. – Europe (WE, NE, SE, EE).
- Miscophus gegensumus** Tsuneki, 1971 (*M. angarensis* Nemkov, 1985). Russia: **ES (IR, BR)**. – China (NC).
- Miscophus gussakovskiji** de Andrade, 1954. Russia: **EP (S)**. – Uzbekistan.
- Miscophus helveticus** Kohl, 1883 (*M. gallicus* Kohl, 1884; *M. rubriventris* Ferton, 1896; *M. deserti* Berland, 1943; *M. gallicus rufus* Giner Marí, 1943; *M. cypriacus* de Andrade, 1953; *M. cypriacus obscurus* de Andrade, 1954). Russia: **EP (C, E)**. – Europe (WE, SE, EE), N Africa, United Arab Emirates, Turkey, Israel.
- Miscophus johni** Mokrousov, 2004. Russia: **EP (C)**. – Europe (SE).
- Miscophus niger** Dahlbom, 1844. Russia: **EP (N, C, E, S)**, **UR, ES (IR)**, **FE (AM, PR, SA)**. – Europe (WE, NE, SE, EE), N Africa, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.
- Miscophus percitus** Mokrousov, 2004. Russia: **EP (C)**.
- Miscophus rufiventris** Tsuneki, 1972. Russia: **ES (BR)**. – Mongolia.
- Miscophus spurius** (Dahlbom, 1832) [*Larra*] (*Larra spuria* Dahlbom, 1832). Russia: **EP (N, S)**, **UR, ES (IR, YA)**, **FE (AM, PR, SA)**. – Europe (WE, NE, SE), N Africa Kazakhstan.
- Miscophus transcaspicus** de Andrade, 1960. Russia: **ES (BR)**. – Turkmenistan, Kazakhstan.
- Miscophus tricolor** Kazenas, 1992. Russia: **EP (S)**. – Kazakhstan.
- NITELA** Latreille, 1809 (*Rhinonitela* F. Williams, 1928). Type species *Nitela spinolae* Latreille, 1809. The genus includes two subgenera: *Nitela* Latreille, 1809 and *Tenila* Brèthes, 1913. Number of species: World – 66, Palaearctic – 14, Russia – 5.
- Nitela (Nitela) borealis** Valkeila, 1974. Russia: **EP (C, E, S)**, **UR, WS (AL)**. – Europe (WE, NE, SE, EE), Georgia, Turkey, Kazakhstan.
- Nitela (Nitela) fallax** Kohl, 1884. Russia: **EP (C, E, S)**, **UR**. – Europe (WE, SE, EE), Turkey.
- Nitela (Nitela) koreana** Tsuneki, 1982. Russia: **FE (PR)**. – Korean Peninsula.
- Nitela (Nitela) lucens** Gayubo et Felton, 2000. Russia: **EP (C)**. – Europe (WE, SE), N Africa.
- Nitela (Nitela) spinolae** Latreille, 1809. Russia: **EP (N, C, E, NC)**. – Europe (WE, NE, SE, EE), N Africa, Turkey, Japan (Hok.).
- PLENOCULUS** W. Fox, 1893 (*Ptygosphex* Gussakovskij, 1928; *Ptigosphex* Gussakovskij, 1928, lapsus for *Ptygosphex* Gussakovskij, 1928; *Pavlovskia* Gussakovskij, 1935). Type species *Plenoculus davisi* W. Fox, 1893. Number of species: World – 21, Palaearctic – 2, Russia – 1.
- Plenoculus murgabensis** (Gussakovskij, 1928) [*Ptygosphex*] (*Pavlovskia tadzhika* Gussakovskij, 1935). Russia: **EP (S)**. – Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.
- SOLIERELLA** Spinola, 1851 (*Sylaon* Piccioli, 1869; *Silaon* Piccioli, 1869, lapsus or emendation of *Sylaon* Piccioli, 1869; *Niteliopsis* S. Saunders, 1873; *Ammosphecidium* Kohl, 1878; *Lautara* Herbst, 1920). Type species *Solierella miscophoides* Spinola, 1851. Number of species: World – 113, Palaearctic – 34, Russia – 5.
- Solierella bactriana** Gussakovskij, 1930. Russia: **EP (S)**. – Iran, Tadzhikistan, Turkmenistan, Uzbekistan.
- Solierella compedita** (Piccioli, 1869) [*Silaon*] (*Silaon compeditus* Piccioli, 1869; *Ammosphecidium Helleri* Kohl, 1877; *Sylaon Xambeui* Er. André, 1896). Russia: **EP (C, S, CR)**. – Europe (WE, SE, EE), N Africa, Turkey, Syria, Jordan, Israel, Kazakhstan.
- Solierella pisonoides** (S. Saunders, 1873) [*Niteliopsis*] (*Solierella paradoxa* Gussakovskij, 1930). Russia: **EP (C, CR)**. – Europe (WE, SE, EE), Turkey, Syria, Israel, Turkmenistan, Kazakhstan.
- Solierella seabrai** de Andrade, 1950. Russia: **EP (S)**. – Europe (WE, SE, EE), N Africa.
- Solierella zimini** Gussakovskij, 1928. Russia: **EP (S)**. – Uzbekistan, Turkmenistan.

Tribe OXYBELINI

BELOMICRUS A. Costa, 1871b (*Oxybeloides* Radoszowski, 1877). Type species *Belomicrus italicus*

A. Costa, 1817. Number of species: World – 101, Palaearctic – 61, Russia – 8.

Belomicrus antennalis Kohl, 1899. Russia: **EP** (S), **WS** (AL), **ES** (IR). – Europe (SE), Kazakhstan, Mongolia.

Belomicrus borealis Forsius, 1923. Russia: **EP** (NW, E). – Europe (NE).

Belomicrus corniger Kazenas, 1995. Russia: **EP** (S). – Kazakhstan.

Belomicrus odontophorus (Kohl, 1892) [Oxybelus] (*B. odontomorphus*: von Schulthess, 1926, lapsus pro *B. odontophorus* (Kohl, 1892)). Russia: **EP** (S). – Europe (EE), Armenia, Turkey, Turkmenistan, Kazakhstan, Mongolia.

Belomicrus parvulus (Radoszkowski, 1877) [Oxybelus] (*B. femoralis* Kohl, 1899). Russia: **UR**. – Turkmenistan, Uzbekistan, Kazakhstan.

Belomicrus schulthessii Kohl, 1924 (*B. Schulthesii* Kohl, 1924). Russia: **EP** (S). – N Africa, Saudi Arabia, United Arab Emirates, Iran, Turkmenistan, Kazakhstan.

Belomicrus tricolor Kazenas, 1991. Russia: **EP** (S). – Kazakhstan.

Belomicrus ussuriensis Antropov, 1993. Russia: **FE** (PR).

OXYBELUS Latreille, 1796 (*Notoglossa* Dahlbom, 1845; *Alepidaspis* A. Costa, 1882; *Anoxybelus* Kohl, 1924; *Gonioxybelus* Minkiewicz, 1934, unavailable under ICZN (1999, Article 13.3); *Orthoxybelus* Minkiewicz, 1934, unavailable under ICZN (1999, Article 13.3); *Gonioxybelus* Pate, 1937, validation of *Gonioxybelus* Minkiewicz, 1934; *Orthoxybelus* Pate, 1937, validation of *Orthoxybelus* Minkiewicz, 1934; *Euoxybelus* Noskiewicz et Chudoba, 1950, unavailable under ICZN (1999, Article 13.3); *Latroxybelus* Noskiewicz et Chudoba, 1950). Type species *Crabro uniglumis* of Fabricius, 1775 (= *Vespa uniglumis* Linnaeus, 1758). Number of species: World – 264, Palaearctic – 86, Russia – 16.

Oxybelus argentatus Curtis, 1833 (?*Vespa decimmaculata* Donovan, 1806; *O. ferox* Shuckard, 1837; *O. nigricornis* Shuckard, 1837). Russia: **EP** (N, C, E, S), **UR**, **WS** (AL), **ES** (IR, ZB), **FE** (PR). – Europe (WE, NE, SE, EE), Mongolia.

Oxybelus aurantiacus Mocsáry, 1883 (*O. rufitarsis* F. Morawitz, 1894). Russia: **EP** (S), **WS** (AL). – Europe (WE, SE, EE), N Africa, Yemen, Saudi Arabia, Turkey, Syria, Jordan, Israel, Pakistan, Central Asia, Kazakhstan, Mongolia.

Oxybelus bipunctatus Olivier, 1812 (*O. nigroaeneus* Shuckard, 1837; *O. laevigatus* Schilling, 1848). Russia: **EP** (N, C, E, S), **UR**, **WS** (AL), **ES** (IR, BR), **FE** (AM, PR, SA). – Europe (WE, NE, SE, EE), N Africa, Turkey, Syria, Iran, Turkmenistan, Uzbekistan, Kazakhstan, Mongolia, Korean Peninsula, N America.

Oxybelus dissectus dissectus Dahlbom, 1845 (*O. monachus* Gerstaecker, 1867). Russia: **EP** (C). – Europe (WE, NE, SE, EE), N Africa, Kuwait, Turkey, Syria, Iran, Turkmenistan, Tajikistan, Kyrgyzstan, Kazakhstan, Mongolia.

Oxybelus dissectus eburneofasciatus Dahlbom, 1845 (*O. eburneo-fasciatus* Dahlbom, 1845; *O. dufouri* Marquet, 1896). Russia: **UR**. – Europe (WE, SE).

Oxybelus dissectus elegans Mocsáry, 1879. Russia: **EP** (S). – Europe (WE, SE, EE), N Africa, Turkey, Israel, Mongolia.

Oxybelus haemorrhoidalis Olivier, 1812 (*O. victor* Lepeletier de Saint Fargeau, 1845; *O. dubius* Dahlbom, 1845; *O. fasciatus* Dahlbom, 1845; *O. simplex* Dahlbom, 1845; *O. bicolor* Schilling, 1848; *O. elegantulus* Gerstaecker, 1867; *O. incomptus* Gerstaecker, 1867; *O. analis* Gerstaecker, 1867; *O. melancholicus* Chevrier, 1868). Russia: **EP** (NW, C, E, S), **UR**, **WS** (NS, AL), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey, Syria, Israel, Iran, Afghanistan, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NC), Korean Peninsula, Japan (Hon).

Oxybelus lamellatus Olivier, 1811 (*O. Savignyi* Spilola, 1839; *Notoglossa sagittata* Dahlbom, 1845; *O. squamosus* F. Smith, 1875; *O. rufipes* Taschenberg, 1880; *O. forticarinatus* Cameron, 1908; *O. ceylonicus* Cameron, 1897). Russia: **EP** (S). – Europe (WE, SE), N Africa, Cyprus, Turkey, Syria, Jordan, Israel, Iraq, Iran, Saudi Arabia, Kuwait, Oman, United Arab Emirates, Afghanistan, Pakistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, China (SW, SE), SE Asia, India, Africa.

Oxybelus latro Olivier, 1812 (*O. armiger* Olivier, 1812; *O. affinis* Marquet, 1881; *O. opacus* Tournier, 1901; *O. latro rugulosus* Móczár, 1957). Russia: **EP** (NC, E, S, CR), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), N Africa, Turkey, Iran, Afghanistan, Central Asia, Kazakhstan, Mongolia, China (NC).

Oxybelus lineatus (Fabricius, 1787) [*Nomada*] (*Nomada lineata* Fabricius, 1787; *O. bellicosus* Olivier, 1812). Russia: **EP** (C, E, S), **UR**, **ES** (KR). – Europe (WE, NE, SE, EE), N Africa, Turkey.

Oxybelus maculipes F. Smith, 1856 (*O. Solskii* Radoszkowski, 1877; *O. acuticornis* F. Morawitz, 1891; *O. pictipes* F. Morawitz, 1891). Russia: **EP** (C, S), **UR**. – Europe (SE, EE), Turkey, Iran, Tajikistan, Kyrgyzstan, Kazakhstan, China (SW).

Oxybelus mandibularis Dahlbom, 1845 (*O. sericatus* Gerstaecker, 1867). Russia: **EP** (C, E, S), **UR**, **WS** (TM, AL), **ES** (IR, BR), **FE** (AM, PR, SA). – Europe (WE, NE, SE, EE), N Africa, Turkey, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia.

Oxybelus mucronatus (Fabricius, 1793) [Crabro] (*O. pugnax* Olivier, 1812; *O. ambiguus* Gerstaecker, 1867; *O. scutellaris* A. Costa, 1867; *O. meridionalis* Mocsáry, 1879; *O. maculiventris* Tournier, 1901; *O. nigritiventris* Tournier, 1901; *O. nigripes immaculatus* Guiglia, 1944; *O. pugnax bistillatus* Balthasar, 1972, unavailable under ICBN (1999, Articles 1.3.4, 15.2)). Russia: **EP** (NW, C, E, S, CR), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), N Africa, Turkey, Syria, Jordan, Lebanon, Israel, Iraq, Iran, Central Asia, Kazakhstan, China (NE).

Oxybelus quatuordecimnotatus Jurine, 1807 (*O. 14 notatus* Jurine, 1807; *Crabro crassipes* Walckenaer, 1817, nom. praeocc., nec Fabricius, 1798; *C. punctatus* Walckenaer, 1817; *C. parisinus* Kittel, 1828; *Oxybelus bellus* Dahlbom, 1844; *O. furcatus* Lepeletier de Saint Fargeau, 1845; *O. raptor* Lepeletier de Saint Fargeau, 1845; *O. timidus* Chevrier, 1868; *O. venustus* Sickmann, 1894; *O. maritimus* Marquet, 1896). Russia: **EP** (NW, C, E, S, CR), **UR**, **WS** (TM, AL), **ES** (KR, IR), **FE** (KH, PR). – Europe (WE, NE, SE, EE), N Africa, Saudi Arabia, Oman, Yemen, Turkey, Syria, Jordan, Israel, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NE, SW).

Oxybelus subspinosus Klug in Waltl, 1835 (*O. fissus* Lepeletier de Saint Fargeau, 1845; *O. latidens* Gerstaecker, 1867; *O. psammobius* Kohl, 1884). Russia: **EP** (C, E, S, CR), **WS** (AL), **ES** (IR), **FE** (PR). – Europe (WE, NE, SE, EE), N Africa, Armenia, Saudi Arabia, Turkey, Syria, Jordan, Israel, Iran, Afghanistan, Central Asia, Kazakhstan, China (NC).

Oxybelus trispinosus (Fabricius, 1787) [Apis] (*Apis trispinosa* Fabricius, 1787; *O. nigripes* Olivier, 1812). Russia: **EP** (C, NC), **UR**, **WS** (AL), **ES** (IR, BR), **FE** (KH, PR). – Europe (WE, NE, SE, EE), N Africa, Turkey, Kazakhstan.

Oxybelus uniglumis (Linnaeus, 1758) [Vespa] (*No-mada punctata* Fabricius, 1793; *Crabro tridens* Fabricius, 1798; *?Vespa decim-maculata* Donovan, 1806; *O. pygmaeus* Olivier, 1812; *O. 4-notatus* Say, 1824; *O. impatiens* F. Smith, 1856; *O. interruptus* Cresson, 1865; *O. fallax* Gerstaecker, 1867; *O. Bro-diei* Provancher, 1883a; *O. Brodiei* Provancher, 1883b, objective synonym of Provancher, 1883a; *O. quadrinotatus montanus* Robertson, 1889; *O. hispanicus* García Mercet: Dusmet y Alonso, 1935, nom. nud.; *O. hispanicus* Giner Marí, 1943). Russia: **EP** (N, NW, C, E, S, CR), **UR**, **WS** (NS, AL), **ES** (KR, IR, BR, ZB), **FE** (SA, KU, KA). – Europe (WE, NE, SE, EE), N Africa, Turkey, Tajikistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE), N America.

Oxybelus variegatus Wesmael, 1852 (*O. pulchellus* Gerstaecker, 1867). Russia: **EP** (C, E, S, CR), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), Turkey, Iran, Afghanistan, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan.

Tribe PALARINI

PALARUS Latreille, 1802 (*Gonius* Panzer, 1806; *Gonius* Jurine, 1807, nom. praeocc., nec Panzer, 1806). Type species *Tiphia flavipes* Fabricius, 1793. Number of species: World – 34, Palaearctic – 18, Russia – 2.

Palarus funerarius F. Morawitz, 1889 (*P. gracilis* Kohl in Kohl et Handlirsch, 1889; *P. seraxensis* Radoszkowski, 1893; *P. incertus* Radoszkowski, 1893; *P. quiescens* Nurse, 1903; *P. nursei* R. Turner, 1911; *P. beaumonti* Bytinski-Salz, 1957). Russia: **EP** (S, NC). – Greece (Rhodes), Armenia, Turkey, Iran, Pakistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NC), India.

Palarus variegatus variegatus (Fabricius, 1781) [*Tiphia*] (*Tiphia variegata* Fabricius, 1781; *Crabro flauipes* Fabricius, 1781; *P. auriginosus* Eversmann, 1849; *P. affinis* F. Morawitz, 1893; *P. fortistriolatus* Cameron, 1907). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (KR, ZB). – Europe (WE, SE, EE), N Africa, Georgia, Armenia, Turkey, Afghanistan, Pakistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NC, NW), India.

Palarus variegatus varius Sickmann, 1894 (*P. flavipes varius* Sickmann, 1894; *P. saishiuensis* Okamoto, 1924). Russia: **FE** (AM, PR). – Korean Peninsula, China (NE).

Tribe TRYPOXYLINI

PISON Jurine, 1808 (*Pison* Jurine in Spinola, 1808; *Tachybulus* Latreille, 1809; *Nephridia* Brullé, 1833; *Pisonitus* Shuckard, 1838; *Pisum* Agassiz, 1846, nom. praeocc., nec Megerle, 1811, unjustified emendation of *Pison* Jurine, 1808; *Pisonoides* F. Smith, 1858; *Parapison* F. Smith, 1869; *Pseudo-Nysson* Radoszkowski, 1876; *Taranga* W.F. Kirby, 1883; *Paraceramius* Radoszkowski, 1887, nom. praeocc., nec de Saussure, 1854; *Pisum* W. Schulz, 1906, nom. praeocc., nec Megerle, 1811, non Agassiz, 1846, unjustified emendation of *Pison* Jurine, 1808; *Krombeiniellum* Richards, 1962, substitute name for *Paraceramius* Radoszkowski, 1887). Type species *Pison Jurini* Spinola, 1808 (= *Alyson ater* Spinola, 1808). Number of species: World – 183, Palaearctic – 11, Russia – 6.

Pison atrum (Spinola, 1808) [Alyson] (*Alyson ater* Spinola, 1808; *P. Jurini* Spinola, 1808; *Tachybulus niger* Latreille, 1809; *P. montanus* Cameron, 1908). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), N Africa, Georgia, Turkey, Oman, Iran, Kazakhstan, Africa.

Pison fasciatum (Radoszkowski, 1876) [Pseudonyszon] (*P. suspicax* Kokujev, 1912). Russia: **EP** (NC). – Europe (SE), Azerbaijan, Turkey, Kazakhstan.

Pison insigne Sickmann, 1894. Russia: **FE** (PR). – China (NE, NC, CE), Korean Peninsula, Japan (Hok).

Pison koreense (Radoszkowski, 1887) [Paraceramius]. Russia: **FE** (PR). – China (CE), Korean Peninsula, Japan (Hon), N America, SE Asia.

Pison sericeum Kohl, 1888. Russia: **EP** (S, NC). – Europe (SE, EE), Turkey, Israel.

Pison strandi Yasumatsu, 1935 (*P. iwatai* Yasumatsu, 1935; *P. ussuricense* Gussakovskij, 1937). Russia: **FE** (PR). – Korean Peninsula, Japan (Hon, Kyu, Tsushima).

TRYPOXYLON Latreille, 1796 (*Tripxylon* Spinola, 1806, lapsus or emendation of *Trypxylon* Latreille, 1796; *Apis* Panzer, 1806; *Apis* Jurine, 1807, nom. praeocc., nec Panzer, 1806; *Trypxylon* Jurine, 1807, lapsus or emendation of *Trypxylon* Latreille, 1796; *Trypxylum* Agassiz, 1846, unjustified emendation of *Trypxylon* Latreille, 1796; *Trypxylum* W. Schulz, 1906, nom. praeocc., nec Agassiz, 1846, unjustified emendation of *Trypxylon* Latreille, 1796; *Asaconoton* Arnold, 1959). Type species

Sphex figulus Linnaeus, 1758. The genus includes two subgenera: *Trypxylon* Latreille, 1796 and *Trypargilum* Richards, 1934. Number of species: World – 631, Palaearctic – 54, Russia – 28.

Trypxylon (Trypxylon) ambiguum Tsuneki, 1956. Russia: **FE** (KU). – Japan (Hon).

Trypxylon (Trypxylon) attenuatum F. Smith, 1851 (*T. hannibalis* Gribodo, 1894). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), N Africa, Turkey, Israel, Afghanistan, Uzbekistan, Kazakhstan, ?China (NW), N America.

Trypxylon (Trypxylon) beaumonti Antropov, 1991. Russia: **EP** (NC). – Europe (WE, SE, EE).

Trypxylon (Trypxylon) clavicerum Lepeletier de Saint Fargeau et Audinet-Serville, 1828 (*T. tibiale* Zetterstedt, 1840; *T. batumicum* Antropov, 1985). Russia: **EP** (N, NW, C, E, S, NC, CR), **WS** (AL), **ES** (IR). – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, Israel, Iran, Kazakhstan, Japan (Hachijo-jima), N America.

Trypxylon (Trypxylon) deceptorium Antropov, 1991. Russia: **EP** (E, S, NC, CR), **WS** (AL). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Syria, Lebanon, Israel, Iran, Central Asia, Kazakhstan, Mongolia.

Trypxylon (Trypxylon) exiguum exiguum Tsuneki, 1956 (*T. clavicerum exiguum* Tsuneki, 1956). Russia: **FE** (SA, KU). – Japan (Hok, Hon, Shi, Kyu).

Trypxylon (Trypxylon) exiguum gussakovskii Tsuneki, 1974 (*T. pygmaeum* Gussakovskij, 1933, nom. praeocc. nec Cameron, 1900; *T. clavicerum gussakovskii* Tsuneki, 1974; *T. exiguum gussakovskii*: Antropov, 1986; *T. clavicerum gussakovskii* Tsuneki, 1991). Russia: **FE** (PR). – Korean Peninsula.

Trypxylon (Trypxylon) figulus (Linnaeus, 1758) [*Sphex*] (*Sphex figulus* Linnaeus, 1758; *T. figulus major* Kohl, 1883; *T. apicalis* W. Fox, 1891; *T. figulus yezo* Tsuneki, 1956; *T. figulus barbarum* de Beaumont, 1957; *T. fieuzeti* Giner Marí, 1959). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, TK, NS, KM, AL), **ES** (KS, KR, IR, BR, ZB), **FE** (PR). – Europe (WE, NE, SE, EE), N Africa, Turkey, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE, CE), Japan (Hok, Hon), N America.

Trypxylon (Trypxylon) frigidum cornutum Gussakovskij, 1932 (*T. cornutum* Gussakovskij, 1932; *T. frigidum chongar* Tsuneki, 1956; *T. frigidum yamatonis* Tsuneki, 1981). Russia: **WS** (AL), **ES** (BR, ZB), **FE** (AM, KH, PR, SA, KU, MG, KA). – Korean Peninsula, Japan (Hon, Kyu).

Trypoxylon (Trypoxylon) fronticorne fronticorne Gussakovskij, 1936. Russia: **EP** (C, E, NC), **ES** (KR, IR). – Europe (WE, NE, SE, EE), Armenia, Azerbaijan, Turkey, Kazakhstan, India.

Trypoxylon (Trypoxylon) fronticorne japonense Tsuneki, 1956 (*T. fronticorne japonense*: Nemkov, 2009, 2012, lapsus pro *T. fronticorne japonense* Tsuneki, 1956). Russia: **WS** (AL), **ES** (BR, ZB), **FE** (AM, KH, PR, SA, KU, KA). – China (NE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Trypoxylon (Trypoxylon) imayoshii Yasumatsu, 1938. Russia: **FE** (PR). – Japan (Hon, Kyu).

Trypoxylon (Trypoxylon) inopinatum Antropov, 1986. Russia: **EP** (NC).

Trypoxylon (Trypoxylon) koikense Tsuneki, 1956. Russia: **FE** (PR, KU). – Japan (Hon, Shi).

Trypoxylon (Trypoxylon) kolazyi Kohl, 1893a (*T. Kolazyi* Kohl, 1893; *T. backi* Sandhouse, 1940). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), Georgia, Turkey, Uzbekistan, Kazakhstan, N America.

Trypoxylon (Trypoxylon) koma Tsuneki, 1956g (*T. figulus koma* Tsuneki, 1956). Russia: **FE** (AM, KH, PR). – Korean Peninsula.

Trypoxylon (Trypoxylon) konosuense Tsuneki, 1968. Russia: **FE** (KH, PR). – Japan (Hon).

Trypoxylon (Trypoxylon) koreanum Tsuneki, 1956 (*T. mowchowense* Tsuneki, 1981; *T. okeanskayanum* Tsuneki, 1981). Russia: **EP** (C), **FE** (PR, SA). – China (NC, SW), Korean Peninsula.

Trypoxylon (Trypoxylon) kostylevi Antropov, 1985. Russia: **EP** (NC). – Europe (SE, NE, EE), Abkhazia, Turkey.

Trypoxylon (Trypoxylon) malaisei Gussakovskij, 1932. Russia: **FE** (PR). – China (CE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu), SE Asia, N America.

Trypoxylon (Trypoxylon) medium de Beaumont, 1945 (*T. figulus minus rubi* Wolf, 1959). Russia: **EP** (C, E, S, NC), **WS** (AL), **ES** (KR). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan.

Trypoxylon (Trypoxylon) minus de Beaumont, 1945 (*T. figulus minor* de Beaumont, 1945; *T. figulus koma* Tsuneki, 1956). Russia: **EP** (C, E, S, NC, CR), **WS** (AL). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan.

Trypoxylon (Trypoxylon) pacificum Gussakovskij, 1932. Russia: **FE** (PR). – China (CE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Trypoxylon (Trypoxylon) regium regium Gussakovskij, 1932. Russia: **FE** (PR).

Trypoxylon (Trypoxylon) rubiginosum Gussakovskij, 1936 (*T. adzharicum* Antropov, 1984). Russia: **EP** (NC, CR). – Abkhazia, Georgia, Azerbaijan.

Trypoxylon (Trypoxylon) rufimaculatum Antropov, 1987 (*T. rufomaculatum*: Haneda et al., lapsus pro *T. rufimaculatum* Antropov, 1987). Russia: **FE** (KH, PR). – China (NE, NC, SW), Korean Peninsula, Japan (Hon, Kyu, Ryu).

Trypoxylon (Trypoxylon) scutatum scutatum Chevrier, 1867 (*T. ammophiloides* A. Costa, 1867; *T. quartinae* Gribodo, 1884; *T. scutigerum* Taschenberg, 1881). Russia: **EP** (S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Turkey, Syria, Israel, Iran, Afghanistan, Turkmenistan, Tajikistan, Kazakhstan.

Trypoxylon (Trypoxylon) shimoyamai Tsuneki, 1958. Russia: **FE** (PR). – China (SW), Japan (Hon).

Trypoxylon (Trypoxylon) ussuricense Kazenas, 1980. Russia: **FE** (PR).

Trypoxylon (Trypoxylon) varipes Pérez, 1905 (*T. monticola* Tsuneki, 1956, *T. varipes nasutum* Tsuneki, 1974, *T. varipes nasutum* Tsuneki, 1991, nom. nud.). Russia: **FE** (PR, KU). – Korean Peninsula, Japan (Hon, Kyu, Ryu).

Subfamily DINETINAE

P.G. NEMKOV

DINETUS Panzer, 1806 (*Dinetus* Jurine, 1807, nom. praeocc., nec *Dinetus* Panzer, 1806). Type species *Crabro pictus* Panzer, 1794 (= *Crabro pictus* Fabricius, 1793). Number of species: World – 12, Palaearctic – 11, Russia – 1.

Dinetus pictus Fabricius, 1793 [Crabro] (*Sphex guttatus* Fabricius, 1793, nom. praeocc., nec *S. guttatus* Gmelin, 1790; *Crabro ceraunius* Rossi, 1794). Russia: **EP** (N, C, E, S, NC), **WS** (AL). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan.

Subfamily EREMIASPHECIINAE

M.V. MOKROUSOV

EREMIASPHECIUM Kohl, 1897 (*Shestakovia* Gussakovskij, 1930; *Xanthospheciun* Simon Thomas, 1994). Type species *Eremiasphecium schmiedeknechtii* Kohl, 1897. Number of species: World – 16; Palaearctic – 16; Russia – 2.

Eremiasphecium bicolor (Gussakovskij, 1930) [Shestakovia]. Russia: **EP** (S). – Turkmenistan, Uzbekistan, Kazakhstan.

Eremiasphecium desertorum (Gussakovskij, 1930) [Shestakovia]. Russia: **EP** (S). – Turkmenistan, Uzbekistan.

Subfamily MELLININAE

P.G. NEMKOV

MELLINUS Fabricius, 1790 (*Trachogorytes* R. Bohart, 2000). Type species *Vespa arvensis* Linnaeus, 1758. Number of species: World – 16; Palaearctic – 3; Russia – 3.

Mellinus arvensis (Linnaeus, 1758) [Vespa] (*Sphex vagus* Linnaeus, 1758; *Crabro bipunctatus* Fabricius, 1787; *Philanthus quinquemaculatus* Fabricius, 1793; *M. pratensis* Jurine, 1807; *M. compactus* Handlirsch, 1888). Russia: **EP** (NW, C, E, NC, CR), **UR**, **WS** (AL), **ES** (IR). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan.

Mellinus crabroneus (Thunberg, 1791) [*Sphex*] (*Crabro sabulosus* Fabricius, 1787, nom. praecocc., nec *Sphex sabulosus* Linnaeus, 1758; *M. ruficornis* Fabricius, 1793; *M. fulvicornis* Fabricius, 1804). Russia: **EP** (NW, C, E, S), **UR**, **WS** (AL), **ES** (IR, BR, ZB), **FE** (AM, PR). – Europe (WE, NE, SE, EE), Turkey, Kyrgyzstan, Kazakhstan, Mongolia, China (NE, NC), Korean Peninsula.

Mellinus obscurus Handlirsch, 1888 (*M. tristis* Pérez, 1905). Russia: **FE** (AM, KH, PR, SA, KU). – Mongolia, China (NC, NE, SE), Korean Peninsula, Japan (Hon), Nepal.

Subfamily PEMPHREDONINAE

M.V. MOKROUSOV

The subfamily is distributed worldwide, mainly in humid regions. Number of taxa: World 43 genera, more than 1100 species, Palaearctic – 22/310, Russia – 17/127.

Tribe ENTOMOSERICINI

ENTOMOSERICUS Dahlbom, 1845. Type species *Entomericus* [sic] *concinus* Dahlbom, 1845. Number of species: World – 3; Palaearctic – 3; Russia – 1.

Entomosericus concinnus Dahlbom, 1845. Russia: **EP** (S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Georgia, Armenia, Turkey, Syria, Israel, Turkmenistan, Kazakhstan.

Tribe PEMPHREDONINI

AMMOPLANELLUS Gussakovskij, 1931. Type species *Ammoplanus chorasmius* Gussakovskij, 1931. Number of species: World – 17; Palaearctic – 6; Russia – 2.

Ammoplanellus chorasmius (Gussakovskij, 1931) [*Ammoplanus*]. Russia: **EP** (S). – Turkmenistan, Uzbekistan.

Ammoplanellus simplex (Gussakovskij, 1951) [*Ammoplanus*] (*Ammoplanus orbiculatus* Tsuneki, 1972). Russia: **EP** (S). – Europe (SE), N Africa, Israel, Iran, United Arab Emirates, Tajikistan, Kazakhstan, Mongolia, India.

AMMOPLANUS Giraud, 1869 (*Hoplocrabron* De Stefani Perez, 1887; *Ceballosia* Giner Marí, 1943, nom. praeocc., nec Mercet, 1921). Type species *Ammoplanus perrisi* Giraud, 1869. Number of species: World – 49; Palaearctic – 32; Russia – 6.

Ammoplanus gegen Tsuneki, 1972. Russia: **EP** (S). – Europe (WE, EE), Cyprus, Jordan, Turkmenistan, Tajikistan, Mongolia.

Ammoplanus kaszabi Tsuneki, 1972 (*A. crudelis* Marshakov, 1976). Russia: **ES** (S), **UR**, **ES** (IR), **WS** (AL), **FE** (KH). – Europe (WE SE), Turkey, Kazakhstan, Mongolia.

Ammoplanus marathroicus (De Stefani Perez, 1887) [*Hoplocrabron*] (*A. handlirschi* Gussakovskij, 1931; *A. metatarsalis* Gussakovskij, 1931; *A. sibiricus* Gussakovskij, 1931; *A. mongolensis* Tsuneki, 1972). Russia: **EP** (C, E, NC, CR), **UR**, **WS** (AL), **ES** (KR, IR). – Europe (WE, SE, EE), N Africa, Turkey, Syria, Jordan, Israel, Iraq, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NC).

Ammoplanus perrisi Giraud, 1869 (*A. wesmaeli* Giraud, 1869; *A. maidli* Gussakovskij, 1931). Russia: **EP** (C, NC). – Europe (WE, SE, EE), N Africa, Turkey, Jordan, Israel, Pakistan.

Ammoplanus rjabovi Gussakovskij, 1931. Russia: **EP** (NC). – Europe (SE), Israel.

Ammoplanus serratus Tsuneki, 1972. Russia: **WS** (AL). – Mongolia.

CARINOSTIGMUS Tsuneki, 1954 (*Perissostigmus* Krombein, 1984). Type species *Stigmus*

congruus Walker, 1860. Number of species: World – 35; Palaearctic – 5; Russia – 1.

Carinostigmus filippovi (Gussakovskij, 1934) [Stigmus]. Russia: **FE** (KH, PR). – China (SW), Korean Peninsula, Japan (Hok, Hon, Shi).

CERATOPHORUS Shuckard, 1837 (*Pemphredon* auct.). Type species *Pemphredon morio* Vander Linden, 1829. Number of species: World – 2; Palaearctic – 2; Russia – 2.

Ceratophorus clypeata (Thomson, 1870) [Pemphredon] (*Pemphredon morio* auct.). Russia: **EP** (N, C). – Europe (WE, NE, EE).

Ceratophorus morio (Vander Linden, 1829) [Pemphredon] (*C. anthracinus* F. Smith, 1851; *Pemphredon carinata* Thomson, 1870; *P. intermedia* Tsuneki, 1951). Russia: **EP** (N, NW, C, E, NC), **UR**, **FE** (PR, MG). – Europe (WE, NE, SE, EE), Turkey, Iran, Uzbekistan, Kazakhstan, Korean Peninsula, Japan (Hok), N America.

DIODONTUS Curtis, 1834 (*Xylocelia* Rohwer, 1915; *Neodiodontus* Tsuneki, 1972; *Corenius* Tsuneki, 1974; *Corenius* Tsuneki, 1991). Type species *Pemphredon tristis* Vander Linden, 1829. Number of species: World – 76; Palaearctic – 47; Russia – 18.

Diodontus brevilabris de Beaumont, 1967. Russia: **EP** (C, S, E, NC). – Europe (SE, EE), Turkey, Kyrgyzstan.

Diodontus changaiensis Tsuneki, 1972. Russia: **WS** (AL). – Mongolia.

Diodontus collaris Tsuneki, 1972. Russia: **WS** (AL), **ES** (KR, IR, ZB). – Kazakhstan, Mongolia.

Diodontus dziuroo Tsuneki, 1972 (*D. pygmaeus* Tsuneki, 1972). Russia: **ES** (ZB), **FE** (PR). – Kazakhstan, Mongolia.

Diodontus gegen Tsuneki, 1972. Russia: **ES** (BR, ZB). – Mongolia.

Diodontus handlirschi Kohl, 1888. Russia: **WS** (AL), **FE** (MG). – Europe (WE, SE), Kyrgyzstan, Kazakhstan, Mongolia.

Diodontus hyalipennis Kohl, 1892 (*D. friesei* Kohl, 1901; *D. major gobiensis* Tsuneki, 1972; *D. mongolicus* Tsuneki, 1972; *D. insidiosus* auct.). Russia: **EP** (S, NC), **WS** (AL), **ES** (TU). – Europe (SE, EE), N Africa, Azerbaijan, Iran, Kazakhstan, Mongolia.

Diodontus insidiosus Spooner, 1938. Russia: **EP** (C, S), **UR**. – Europe (WE, NE, SE, EE).

Diodontus kaszabi Tsuneki, 1972 (*D. monticola* Tsuneki, 1972; *D. obo* Tsuneki, 1972). Russia: **WS** (AL), **ES** (IR, ZB), **FE** (AM, KA, MG). – Mongolia.

Diodontus kohli Tsuneki, 1972. Russia: **EP** (S). – Kazakhstan, Mongolia.

Diodontus kuroo Tsuneki, 1972. Russia: **EP** (S). – Kazakhstan, Mongolia.

Diodontus luperus Shuckard, 1837. Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (KR, IR, YA, ZB). – Europe (C, SE, EE), N Africa, Turkey, Kazakhstan, Mongolia.

Diodontus major Kohl, 1901. Russia: **EP** (C, S). – Europe (C, SE, EE).

Diodontus mediuss Dahlbom, 1844 (*D. dahlbomi* A. Morawitz, 1864). Russia: **EP** (N, C, E), **UR**, **WS** (TM, AL), **ES** (IR), **FE** (AM, PR, SA, KA, MG). – Europe (C, NE, SE, EE), Turkey.

Diodontus minutus (Fabricius, 1793) [Crabro] (*Xylocelia franclemonti* Krombein, 1939). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, AL), **ES** (KR, IR, ZB), **FE** (PR). – Europe (WE, NE, SE, EE), N Africa, Turkey, Syria, Israel, Iraq, Iran, Kuwait, Afghanistan, Central Asia, Kazakhstan, Mongolia, China (NC), Korean Peninsula, N America.

Diodontus parvulus (Radoszkowski, 1877) [Passaloecus]. Russia: **EP** (S). – Central Asia, Kazakhstan.

Diodontus spinicollis Gussakovskij, 1933. Russia: **EP** (S). – Iran.

Diodontus tristis (Vander Linden, 1829) [Pemphredon]. Russia: **EP** (N, NW, C, E, S), **UR**, **WS** (AL), **ES** (IR, YA, ZB), **FE** (AM). – Europe (WE, NE, SE, EE), Turkey, Iran, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia.

PASSALOECUS Shuckard, 1837 (*Xyloecus* Shuckard, 1837, nom. praeocc., nec Déjean, 1833; *Coeloeucus* C. Verhoeff, 1890; *Heroecus* C. Verhoeff, 1890). Type species *Pemphredon insignis* Vander Linden, 1829. Number of species: World – 40; Palaearctic – 24; Russia – 17.

Passaloecus borealis Dahlbom, 1844. Russia: **EP** (N, NW), **WS** (AL), **ES** (IR), **FE** (KH, PR, SA, MG). – Europe (WE, NE, SE, EE), Kazakhstan, N America. Records in **EP** (C, E) and **UR** erroneous.

Passaloecus brevilabris Wolf, 1958. Russia: **EP** (C, CR). – Europe (WE, NE, SE, EE).

Passaloecus clypealis Faester, 1947 (*P. angustus* Gussakovskij, 1952). Russia: **EP** (N, C, E, NC), **UR**, **WS** (AL), **ES** (IR), **FE** (AM, KH, PR, SA, KU, KA). – Europe (WE, NE, SE, EE), Tajikistan, Uzbekistan,

- Kyrgyzstan, Kazakhstan, China (NC, CC, SE), Korean Peninsula, Japan (Hok, Hon).
- Passaloecus corniger** Shuckard, 1837. Russia: **EP** (N, C, E, S, NC, CR). – Europe (WE, NE, SE, EE), N Africa, Turkey, Kazakhstan, Japan (Hon).
- Passaloecus eremita** Kohl, 1893. Russia: **EP** (N, C, E). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan.
- Passaloecus gracilis** (Curtis, 1834) [*Diodontus* (*P. brevicornis* A. Morawitz, 1864)]. Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), N Africa, Abkhazia, Turkey, Iran, Uzbekistan, Kazakhstan, N America.
- Passaloecus hinganicus** Merisuo, 1976. Russia: **ES** (ZB).
- Passaloecus insignis** (Vander Linden, 1829) [Pemphredon] (*P. roettgeni* C. Verhoeff, 1890; *P. shuckardi* Yasumatsu, 1934). Russia: **EP** (NC, CR), **WS** (AL), **FE** (AM, PR). – Europe (WE, NE, SE, EE), Abkhazia, Turkey, Kazakhstan, China (CC, SW), Korean Peninsula, Japan (Hok, Hon). Records in **EP** (N, C, E) and **UR** doubtful.
- Passaloecus koreanus** Tsuneki, 1974 (*P. iwatai* Merisuo, 1976). Russia: **FE** (PR). – Korean Peninsula, Japan (Hok, Hon).
- Passaloecus longiceps** Merisuo, 1973. Russia: **EP** (N).
- Passaloecus mongolicus** Tsuneki, 1972 (*P. altaicus* Merisuo, 1976). Russia: **WS** (AL), **ES** (IR, ZB), **FE** (KH, PR, SA). – Kazakhstan, Mongolia, Japan (Hok, Hon).
- Passaloecus monilicornis** Dahlbom, 1842. Russia: **EP** (N, NW, C, E), **UR**, **WS** (TK, AL), **ES** (KR, IR, ZB), **FE** (AM, KH, PR, SA, KU, KA, MG). – Europe (WE, NE, SE, EE), Kazakhstan, Mongolia, China (NE, SW, WP, SE), Korean Peninsula, Japan (Hok, Hon, Kyu), N America.
- Passaloecus nipponicola** Tsuneki, 1955. Russia: **FE** (KH, PR). – Japan (Hon).
- Passaloecus pictus** Ribaut, 1952. Russia: **EP** (NC, CR). – Europe (WE, SE, EE), N Africa, Abkhazia, Georgia, Turkey, Syria, Iran, S America.
- Passaloecus ribauti** Merisuo, 1974. Russia: **EP** (CR). – Europe (WE, SE), N Africa, Turkey, Syria.
- Passaloecus singularis** Dahlbom, 1844 (*P. tenuis* A. Morawitz, 1864; *P. gertrudis* Krombein, 1938; *P. gracilis* auct.). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **FE** (PR). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan, Japan (Hon), N America.
- Passaloecus turionum** Dahlbom, 1844 (*P. borealis* auct.). Russia: **EP** (NW, C, E, NC), **UR**. – Europe (WE, NE, SE, EE), Turkey, Tajikistan, Kyrgyzstan, Kazakhstan.
- PEMPHREDON** Latreille, 1796 (*Cemonus* Panzer, 1806; *Cemonus* Jurine, 1807; *Dineurus* Westwood, 1837; *Diphlebus* Westwood, 1840; *Chevrieria* Kohl, 1883; *Susanowo* Tsuneki, 1972; *Sinostigma* Hong, 1984). Type species *Crabro lugubris* Fabricius, 1793. Number of species: World – 44; Palaearctic – 28; Russia – 16.
- Pemphredon austriaca** (Kohl, 1888) [Diphlebius (*P. coracina* Valkeila, 1972; *P. tenera* Valkeila, 1972; *P. nescia* Merisuo, 1972)]. Russia: **EP** (C, S, NC, CR). – Europe (WE, SE, EE), N Africa, Abkhazia, Azerbaijan, Turkey.
- Pemphredon baltica** Merisuo, 1972 (*P. fennica* Merisuo, 1972). Russia: **EP** (N, C), **UR**, **WS** (AL), **FE** (KH, PR, KA). – Europe (WE, SE, NE), N America.
- Pemphredon enslini** (Wagner, 1932) [Dineurus] (*P. austriaca* auct.). Russia: **EP** (C, E). – Europe (WE, SE).
- Pemphredon fabricii** (M. Müller, 1911) [Cemonus] (*P. lethifer* auct.). Russia: **EP** (C, S, NC). – Europe (WE, SE), Turkey, Kazakhstan, Japan (Hon).
- Pemphredon flavistigma** Thomson, 1874. Russia: **EP** (NW, C, E), **UR**, **ES** (KR, IR, BR), **FE** (KH, PR, SA). – Europe (WE, NE, SE, EE), Korean Peninsula, Japan (Hok, Hon).
- Pemphredon inornata** Say, 1824 (*Cemonus shuckardi* A. Morawitz, 1864; *C. dentatus* Puton, 1871; *P. tenax* W. Fox, 1892). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TK, KM, AL), **ES** (KR, IR, BR, ZB), **FE** (AM, KH, PR, SA, KU, KA, MG). – Europe (WE, NE, SE, EE), N Africa, Turkey, Abkhazia, Central Asia, Kazakhstan, Mongolia, China (NE, NC, CC), Korean Peninsula, Japan (Hok, Hon), N America.
- Pemphredon japonica** Matsumura, 1912. Russia: **FE** (PR, SA, KU). – Nepal, Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Pemphredon krombeini** Tsuneki, 1960 (*P. mandibularis* Tsuneki, 1951, nom. praeocc., nec Cresson, 1865). Russia: **FE** (SA). – Japan (Hok, Hon).
- Pemphredon lethifer** (Shuckard, 1837) [Cemonus] (*Cemonus strigatus* Chevrier, 1870; *Diphlebus fuscatus* A. Wagner, 1918; *D. littoralis* A. Wagner, 1918; *D. neglectus* A. Wagner, 1918; *D. minutus* A. Wagner, 1918; *P. brevipetiolata* A. Wagner, 1932; *P. gemina* Valkeila, 1972; *P. trichogastor* Valkeila, 1972). Russia: **EP** (N, C, E, S, NC, CR), **UR**, **WS** (NS, AL), **FE** (AM, KH, PR, SA, KU). – Europe (WE, NE,

SE, EE), N Africa, Abkhazia, Azerbaijan, Turkey, Israel, Iraq, Iran, Tajikistan, Uzbekistan, Kyrgyzstan, Afghanistan, Kazakhstan, Mongolia, China (NE, NC, CC, SE), Korean Peninsula, Japan (Hok, Hon), N America.

Pemphredon lugens Dahlbom, 1842. Russia: **EP** (N, NW, C, E), **UR**, **WS** (AL), **ES** (KR, IR), **FE** (MG). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan.

Pemphredon lugubris (Fabricius, 1793) [Crabro]
(*P. concolor* Say, 1824; *P. fuscipennis* Stephens, 1829; *P. nitida* Stephens, 1829; *P. ocellaris* Gimmerthal, 1836; *P. luctuosa* Shuckard, 1837; *P. morio* Cresson, 1865, nom. praeocc., nec Vander Linden, 1829; *P. cressoni* Dalla Torre, 1897; *P. provancheri* Dalla Torre, 1897; *P. provancheri* Ashmead, 1899, nom. praeocc., nec Dalla Torre, 1897; *P. tinctipennis* Cameron, 1908; *P. shawii* Rohwer, 1917; *P. pacifica* Gussakovskij, 1932). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (TK, AL), **ES** (IR, ZB), **FE** (KH, PR, SA, KU, KA, MG). – Europe (WE, NE, SE, EE), Turkey, Iran, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE, CC), Korean Peninsula, Japan (Hok, Hon), N America.

Pemphredon montana Dahlbom, 1844 (*P. angularis* W. Fox, 1892). Russia: **EP** (N, NW, C, E), **UR**, **ES** (KR, IR, BR, YA), **FE** (AM, KH, PR, SA, KU, KA, MG). – Europe (WE, NE, SE, EE), Kazakhstan, China (NE), Korean Peninsula, Japan (Hok, Hon), N America.

Pemphredon mortifer Valkeila, 1972 (*P. bipartitor* W. Fox, 1892; *P. harbecki* Rohwer, 1910; *P. rugifer* auct.). Russia: **EP** (NW, C, E, S, NC), **UR**, **ES** (IR), **FE** (KH, PR, SA). – Europe (WE, NE, SE, EE), Azerbaijan, Turkey, Korean Peninsula, Japan (Hok), N America.

Pemphredon podagrica Chevrier, 1870 (*P. laeviceps* Gussakovskij, 1932). Russia: **FE** (PR, SA). – Europe (WE, SE, EE), Japan (Hok, Hon).

Pemphredon rugifer (Dahlbom, 1844) [Cemonus] (*Sphex unicolor* Panzer, 1798, nom. praeocc., nec Fabricius, 1787; *Cenomus* [sic] *pilosus* Gimmerthal, 1836, nom. oblitum; *Cemonus solivagus* Bondroit, 1932; *P. cythica* Valkeila, 1972; *P. punctifer* Merisuo, 1972). Russia: **EP** (C, E, S, NC), **UR**, **WS** (NS, AL), **ES** (KR), **FE** (SA). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan, China (NE, SW), Korean Peninsula, Japan (Hon), N America.

Pemphredon wesmaeli (A. Morawitz, 1864) [Cemonus] (*P. scotica* Perkins, 1929; *P. rugifer* auct.).

Russia: **EP** (N, NW, C, E), **UR**. – Europe (WE, NE, SE, EE), Kazakhstan.

POLEMISTUS de Saussure, 1892. Type species *Polemistus macilentus* de Saussure, 1892. Number of species: World – 37; Palaearctic – 2; Russia – 1.

Polemistus abnormis (Kohl, 1888) [Passaloecus]. Russia: **EP** (C), **WS** (AL), **FE** (KH, PR). – Europe (WE, SE), Kazakhstan, China (CC), Korean Peninsula, Japan (Hok, Hon), SE Asia.

SPILOMENA Shuckard, 1838 (*Celia* Shuckard, 1837, nom. praeocc., nec Zimmermann, 1832; *Microglossa* Rayment, 1930, nom. praeocc., nec Voight, 1831; *Microglossella* Rayment, 1935; *Taialia* Tsuneki, 1971). Type species *Stigmus troglodytes* Vander Linden, 1829. Number of species: World – 87; Palaearctic – 22; Russia – 7.

Spilomena beata Blüthgen, 1953 (*S. exspectata* Valkeila, 1957). Russia: **EP** (C, S), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), Georgia, Kazakhstan.

Spilomena curruca (Dahlbom, 1844) [Celia] (*Spilomena pulawskii* Dollfuss, 1983; *S. nikkoensis* Tsuneki, 1971). Russia: **EP** (N, E), **UR**, **WS** (AL), **FE** (AM, PR, KU). – Europe (WE, NE, SE, EE), Japan (Hon).

Spilomena dedzeli Tsuneki, 1971. Russia: **FE** (KU). – Japan (Hon).

Spilomena differens Blüthgen, 1953. Russia: **EP** (N). – Europe (WE, NE, EE), Estonia.

Spilomena enslini Blüthgen, 1953. Russia: **EP** (E), **UR**, **WS** (AL). – Europe (WE, NE, EE), Kazakhstan.

Spilomena mocsaryi Kohl, 1898 (*S. capitata* Gussakovskij, 1931; *S. zavadili* Šnoflák, 1942). Russia: **EP** (C, S, CR), **UR**. – Europe (C, SE, EE), Georgia, Cyprus, Turkey, Syria, Jordan, Israel, United Arab Emirates, Central Asia.

Spilomena troglodytes (Vander Linden, 1829) [Stigmus] (*Stigmus minutissimus* Radoszkowski, 1877; *Spilomena vagans* Blüthgen, 1953). Russia: **EP** (NW, C, NC, CR), **UR**, **FE** (PR). – Europe (WE, NE, SE, EE), N Africa, Abkhazia, Georgia, Turkey, Israel, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan, N America.

STIGMUS Panzer, 1804 (*Antronius* Zetterstedt, 1838; *Gonostigmus* Rohwer, 1911; *Atopostigmus* Krombein, 1973). Type species *Stigmus pendulus* Panzer, 1804. Number of species: World – 25; Palaearctic – 9; Russia – 7.

Stigmus convergens Tsuneki, 1954 (*S. solskyi*: Nemkov, 1986; *S. pendulus*: Mokrousov, 2009). Russia: **FE** (KU). – Japan (Hok, Hon).

Stigmus eurasaticus Mokrousov, 2017. Russia: **EP** (N, C, E), **UR**, **WS** (AL), **ES** (IR), **FE** (KH, PR). – China (SE), Korean Peninsula.

Stigmus japonicus Tsuneki, 1954. Russia: **FE** (AM, KH, PR, SA, KU). – Korean Peninsula, Japan (Hok, Hon).

Stigmus munakatai Tsuneki, 1954. Russia: **FE** (PR). – Japan (Hok, Hon, Kyu).

Stigmus pendulus Panzer, 1804 (*S. ater* Jurine, 1807). Russia: **EP** (C, E, NC), **UR**. – Europe (WE, NE, SE, EE).

Stigmus quadriceps Tsuneki, 1954. Russia: **FE** (KU). – Japan (Hok, Hon).

Stigmus solskyi A. Morawitz, 1864 (*S. europaeus* Tsuneki, 1954; *S. verhoeffi* Tsuneki, 1954). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), N Africa, Turkey, Kazakhstan.

TZUSTIGMUS Finnimore, 1995. Type species *Tzustigmus syam* Finnimore, 1995. Number of species: World – 8; Palaearctic – 3; Russia – 1.

Tzustigmus rhinocerus (Budrys, 1987) [Carinostigmus]. Russia: **FE** (KH, PR). – Korean Peninsula.

Tribe PSENINI

MIMESA Shuckard, 1937 (*Aporia* Wesmael, 1852, nom. praeocc., nec Hübner, 1819; *Aporina* Gussakovskij, 1937, nom. praeocc., nec Fuhrmann, 1902). Type species *Trypoxyton equestre* Fabricius, 1804. Number of species: World – 76; Palaearctic – 40; Russia – 15.

Mimesa beckeri Tournier, 1889. Russia: **EP** (S). – Kazakhstan, Mongolia.

Mimesa bicolor (Jurine, 1807) [Psen]. Russia: **EP** (N, NW, C, E, S, CR), **UR**, **WS** (NS, AL), **ES** (KR, IR, ZB), **FE** (PR, SA). – Europe (WE, NE, SE, EE), Israel, Kyrgyzstan, Kazakhstan, China (NC).

Mimesa breviventris F. Morawitz, 1891. Russia: **EP** (S). – Kazakhstan, China (NE).

Mimesa bruxellensis Bondroit, 1934 (*Psen rossicus* Gussakovskij, 1937). Russia: **EP** (NW, C, E), **UR**. – Europe (WE, NE, SE, EE), Kazakhstan.

Mimesa caucasica Maidl, 1914. Russia: **EP** (C, S, NC), **UR**. – Europe (EE), Georgia, Azerbaijan, Iran, Central Asia, Kazakhstan, Mongolia, China (NC).

Mimesa crassipes A. Costa, 1871 (*M. carbonaria*

A. Costa, 1867, nom. praeocc., nec F. Smith, 1865; *M. ochroptera* A. Costa, 1871; *M. costae* Ed. André, 1888; *M. helvetica* Tournier, 1889; *M. lixivia* Tournier, 1889; *M. pannonica* Maidl, 1914). Russia: **EP** (C, E, S, NC, CR), **WS** (AL). – Europe (WE, SE, EE), Georgia, Turkey, Iran, Kyrgyzstan, Kazakhstan, China (NW).

Mimesa equestris (Fabricius, 1804) [Trypoxylon] (*Psen basalis* Stephens, 1829). Russia: **EP** (N, NW, C, E), **UR**, **WS** (NS, AL), **ES** (KR, IR, YA, ZB), **FE** (KH, PR, SA, KA, MG). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan, Korean Peninsula.

Mimesa grandii Maidl, 1933 (*Psen carinatus* Gussakovskij, 1937; *P. sciaphilus* Arcidiacono, 1965). Russia: **EP** (S). – Europe (WE, SE, EE), N Africa, Turkey, Iran, Turkmenistan, Kazakhstan, China (NC).

Mimesa gussakovskiji (de Beaumont, 1941) [Psen] (*Psen costae*: Gussakovskij, 1937). Russia: **EP** (E, S). – Europe (SE, EE), Kazakhstan.

Mimesa kurzenkoi Budrys, 1988. Russia: **FE** (PR, KU).

Mimesa lutaria (Fabricius, 1787) [Sphecius] (*M. shuckardi* Wesmael, 1852; *M. basirufa* Packard, 1867; *M. nebrascensis* H. Smith, 1908; *Psen dispar* Gussakovskij, 1937; *M. mallochi* Finnimore, 1980). Russia: **EP** (NW, C, E), **UR**, **WS** (AL), **ES** (KR, IR, YA), **FE** (AM, KH, PR, SA, KU, KA, MG). – Europe (WE, NE, SE, EE), Turkey, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE), Korean Peninsula, Japan (Hok, Hon, Shi), N America.

Mimesa nigrita Eversmann, 1849 (*M. brevis* Maidl, 1914). Russia: **EP** (S), **UR**, **WS** (AL), **ES** (KR, IR). – Europe (WE, SE, EE), Turkey, Syria, Uzbekistan, Kazakhstan, Mongolia.

Mimesa punctipleuris (Gussakovskij, 1937) [Psen] (*Psen angulicollis* Tsuneki, 1972). Russia: **WS** (AL), **ES** (KR, ZB), **FE** (PR). – Kyrgyzstan, Kazakhstan, Mongolia.

Mimesa sibirica (de Beaumont, 1937) [Psen] (*Psen albopilosus* Tsuneki, 1972). Russia: **ES** (IR, BR, ZB). – Kazakhstan, Mongolia, China (NC).

Mimesa vindobonensis Maidl, 1914 (*Psen bidentatus* Gussakovskij, 1937, nom. praeocc., nec Cameron, 1908). Russia: **EP** (C, S), **WS** (AL), **ES** (KR, BR, ZB). – Europe (WE, EE), Turkey, Kazakhstan, Mongolia, China (NC).

MIMUMESA Malloch, 1933. Type species *Psen niger* Packard, 1867. Number of species: World – 32; Palaearctic – 13 (1 introduced); Russia – 7.

Mimumesa atratina (F. Morawitz, 1891) [Mimesa]

(*Mimesa carbonaria* Tournier, 1889, nec F. Smith, 1865; *M. belgica* Bondroit, 1932; *M. longula* Gussakovskij, 1932; *Psen sameshima* Yasumatsu, 1937). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TM, AL), **ES** (KR, BR, ZB), **FE** (KH, PR, SA, KU, KA, MG). – Europe (WE, NE, SE, EE), Turkey, Kyrgyzstan, Kazakhstan, Mongolia, China (NE, NC, NW), Korean Peninsula, Japan (Hon, Kyu), N America.

Mimumesa beaumonti (van Lith, 1949) [Psen]. Russia:

EP (N, NW, C, E), **UR**, **WS** (TK), **ES** (IR), **FE** (PR). – Europe (WE, NE, EE), Kazakhstan.

Mimumesa dahlbomi (Wesmael, 1852) [Mimesa]. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (IR), **FE** (AM, KH, PR, SA, KU, KA, MG). – Europe (WE, NE, SE, EE), Turkey, Kyrgyzstan, Kazakhstan, Mongolia, China (NE, NC, NW, CC, WP), Korean Peninsula, Japan (Hon).
Mimumesa littoralis (Bondroit, 1934) [Mimesa]

(*Psen fulvitarsis* Gussakovskij, 1934; *Mimesa celtica* Spooner, 1948). Russia: **EP** (N, NW, C, S, NC), **UR**, **WS** (AL), **ES** (TU, IR, ZB), **FE** (AM, KH, PR, SA, MG). – Europe (WE, NE, SE, EE), Tajikistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE), Korean Peninsula, Japan (Hok, Hon), India.

Mimumesa sibiricana R. Bohart, 1976 (*Psen sibiricus*

Gussakovskij, 1937, nom. praeocc., nec de Beaumont, 1937). Russia: **WS** (AL), **ES** (IR, YA, ZB), **FE** (KA). – Europe (WE, Mongolia. Record in **UR** doubtful.

Mimumesa spooneri (Richards, 1948) [Mimesa]. Russia: **EP** (C). – Europe (WE, NE, EE).**Mimumesa unicolor** (Vander Linden, 1829) [Psen]

(*Mimesa borealis* Dahlbom, 1842; *M. palliditarsis* E. Saunders, 1904; *Psen oresterus* van Lith, 1976). Russia: **EP** (N, C, E, S, NC), **UR**, **WS** (NS, AL), **ES** (KR, IR, YA, ZB). – Europe (WE, NE, SE, EE), Azerbaijan, Turkey, Israel, Iraq, Afghanistan, Pakistan, Central Asia, Kazakhstan, China (NE, NC, NW, WP).

PSEN Latreille, 1796 (*Psenus* Rafinesque, 1815; *Psenia* Stephens, 1829; *Dahlbomia* Wissmann, 1849; *Mesopora* Wesmael, 1852; *Caenopsen* Cameron, 1899). Type species: *Crabro ater* Olivier, 1792. Number of species: World – 95; Palaearctic – 25; Russia – 9.**Psen affinis** Gussakovskij, 1937 (*P. hakusanus* Tsuneki, 1959). Russia: **FE** (AM, PR, SA, KU). – China (SW, SE), Korean Peninsula, Japan (Hok, Hon, Shi).

Psen ater (Olivier, 1792) [Crabro] (*Sphex ater* Fabricius, 1794; *Pelopoeus unicolor* Fabricius, 1804; *Pelopoeus compressicornis* Fabricius, 1804; *Psen atratus* Jurine, 1807; *Psen serraticornis* Jurine, 1807). Russia: **EP** (C, S, CR), **UR**, **WS** (AL), **ES** (KR, IR, BR, ZB), **FE** (AM, KH, PR, KU). – Europe (WE, NE, SE, EE), Turkey, Uzbekistan, Kazakhstan, Mongolia, China (NE, NW), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Psen aurifrons Tsuneki, 1959 (secondary homonym of *P. aurifrons* (Taschenberg, 1875), now in *Pseneo* Malloch; *P. caocinnus* Tsuneki, 1973). Russia: **FE** (AM, KH, PR, SA, KU). – China (NC), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu).

Psen bettoh Tsuneki, 1977. Russia: **FE** (KU). – China (SE), Japan (Hok, Hon).

Psen foveolatus Budrys, 1986. Russia: **FE** (PR).

Psen koreanus Tsuneki, 1959 (*P. ohnonis* Tsuneki, 1973). Russia: **FE** (PR). – China (SE), Korean Peninsula, Japan (Hon, Ryu).

Psen miyagino Tsuneki, 1983 (*P. exorientis* Budrys, 1988). Russia: **FE** (PR). – Japan (Hon).

Psen seminitidus van Lith, 1965 (*Mimesa kohli* Gussakovskij, 1934, secondary homonym of *Psen kohli* W. Fox, 1898; *P. alticola* Tsuneki, 1977; *P. takanensis* Tsuneki, 1978). Russia: **ES** (ZB), **FE** (AM, PR, SA, KU). – China (NC, NW, WP, SW, SE), Korean Peninsula, Japan (Hok, Hon).

Psen ussuriensis van Lith, 1959 (*Mimesa orientalis* Gussakovskij, 1932, secondary homonym of *Psen orientalis* Cameron, 1890; *P. mandibularis* Tsuneki, 1959; *P. tsunekii* van Lith, 1965). Russia: **FE** (AM, PR). – China (NE), Korean Peninsula, Japan (Hok, Hon, Shi).

PSENEO Malloch, 1933 (*Punctipsen* van Lith, 1968). Type species: *Psen kohlii* W. Fox, 1898. Number of species: World – 26; Palaearctic – 1.

Pseneo exaratus (Eversmann, 1849) [Mimesa] (*Mimesa superba* Tournier, 1889; *M. picicornis* F. Morawitz, 1892). Russia: **EP** (C, E, NC), **UR**, **WS** (TK, KM, AL), **ES** (IR), **FE** (AM, PR). – Europe (WE, SE, EE), Abkhazia, Georgia, Kazakhstan, China (SE), Korean Peninsula, Japan (Hok, Hon, Shi, Ryu), India.

PSENULUS Kohl, 1897 (*Neoxoxia* Viereck, 1901; *Stenomellinus* W. Schulz, 1911; *Eopsenulus* Gussakovskij, 1934; *Nipponopsen* Yasumatsu, 1938). Type species: *Psen fuscipennis* Dahlbom, 1843.

Number of species: World – 160; Palaearctic – 34; Russia – 15.

Psenulus anomoneurae (Yasumatsu, 1938) [Nipponopsen] (*P. mandibularis* Tsuneki, 1959). Russia: **FE** (KH, PR, SA, KU). – Korean Peninsula, Japan (Hon, Kyu).

Psenulus carinifrons (Cameron, 1902) [Psen]. Russia: **EP** (NC). – Europe (SE), Turkey, Pakistan, China (SW), Korean Peninsula, Japan (Hon, Ryu), SE Asia, India, Australia.

Psenulus chevrieri (Tournier, 1889) [Psen] (*P. brevitarvis* Merisuo, 1937; *P. pallipes* auct.). Russia: **EP** (C, E, S, CR), **UR**. – Europe (WE, NE, SE, EE), Armenia.

Psenulus concolor (Dahlbom, 1843) [Psen] (*Psen intermedius* Schenck, 1857; *P. ambiguus* Schenck, 1857). Russia: **EP** (N, C, E, NC, CR). – Europe (WE, NE, SE, EE), Turkey. Records in **UR** and Japan (Hok, Hon) doubtful.

Psenulus fulvicornis (Schenck, 1857) [Psen] (*P. schencki* auct.). Russia: **EP** (NC). – Europe (WE, C, SE, EE), N Africa, Turkey, Syria.

Psenulus fuscipennis (Dahlbom, 1843) [Psen] (*Psen nylanderi* Dahlbom, 1845; *P. dufouri* Dahlbom, 1845; *P. migratus* Brischke, 1862; *P. procerus* A. Costa, 1867). Russia: **EP** (N, C, E, S, NC, CR), **UR**, **FE** (PR). – Europe (WE, NE, SE, EE), Georgia, Turkey, Korean Peninsula, Japan (Hok, Hon), India.

Psenulus ghilarovi Budrys, 1988. Russia: **FE** (KH, PR).

Psenulus laevigatus (Schenck, 1857) [Psen] (*Psen distinctus* Chevrier, 1870; *Psenulus fuscipes* Tsuneki, 1959). Russia: **EP** (C, E, NC, CR), **ES** (ZB), **FE** (PR, SA, KU). – Europe (WE, SE, EE), Turkey, Korean Peninsula, Japan (Hok, Hon).

Psenulus lubricus (Pérez, 1905) [Psen]. Russia: **FE** (SA, KU). – China (SW), Japan (Hok, Hon, Shi).

Psenulus maculipes Tsuneki, 1959. Russia: **FE** (KU). – Japan (Hon).

Psenulus meridionalis de Beaumont, 1937. Russia: **EP** (C, E, S, NC). – Europe (WE, SE, EE), Armenia, Turkey, Syria, Jordan, Iraq, Iran, Turkmenistan.

Psenulus nikkoensis Tsuneki, 1959. Russia: **FE** (KU). – Japan (Hok, Hon, Kyu).

Psenulus pallipes (Panzer, 1798) [Sphex] (*Trypoxyylon atratum* Fabricius, 1804; *Psen montanus* A. Costa, 1861; *P. haemorrhoidalis* A. Costa, 1867; *P. minutus* Tournier, 1889; *P. nigricornis* Tournier, 1889; *P. pygmaeus* Tournier, 1889; *Psenulus rubicola* Harttig, 1931; *P. puncticeps* Gussakovskij, 1932, secondary

homonym of *Psen puncticeps* Cameron, 1907; *P. gussakovskij* van Lith, 1973). Russia: **EP** (N, C, E, S, NC, CR), **UR**, **WS** (TK, AL), **ES** (IR), **FE** (AM, KH, PR, SA, KU, KA). – Europe (WE, NE, SE, EE), Abkhazia, Armenia, Turkey, Iran, Kazakhstan, China (NC, SW), Korean Peninsula, Japan (Hok, Hon), N America.

Psenulus schencki (Tournier, 1889) [Psen] (*Psen simplex* Tournier, 1889; *P. longulus* Tournier, 1889). Russia: **EP** (C, E, S, NC, CR). – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey. Records in **WS** (AL) doubtful.

Psenulus tanakai Tsuneki, 1959. Russia: **FE** (PR). – Japan (Hon).

Subfamily PHILANTHINAE

YU.N. DANILOV AND M.V. MOKROUSOV

Tribe CERCERINI

CERCERIS Latreille, 1802 (*Apiraptrix* Shestakov, 1923; *Bucerceris* Minkiewicz, 1934; *Stercobata* Gussakovskij, 1935; *Apicerceris* Pate, 1937). Type species *Philanthus ornatus* Fabricius, 1790 [= *Sphex rybyensis* Linnaeus, 1771]. Number of species: World – about 860; Palaearctic – about 270; Russia – 62.

Cerceris adelpha Kohl in Schletterer, 1887 (*C. associa* Kohl, 1898). Russia: **ES** (IR, BR, YA, ZB), **FE** (AM, KH, PR). – Mongolia, China (NE, NC), Korean Peninsula.

Cerceris albicolor Shestakov, 1918. Russia: **EP** (S), **WS** (AL). – Europe (EE), Azerbaijan, Turkey, Turkmenistan, Kazakhstan.

Cerceris albofasciata Rossi, 1790 [*Vespa*] (*C. luctuosa* A. Costa, 1867; *C. navitatis* F. Smith, 1873; *C. cibrata* Mocsáry, 1879). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (NS, AL), **ES** (KS, KR, BR, ZB), **FE** (PR). – Europe (WE, SE, EE), Turkey, Iran, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE, NC, SE), Korean Peninsula, Japan (Hok, Hon, Kyu, Ryu).

Cerceris angelica Kazenas, 1977. Russia: **EP** (S). – Turkmenistan, Uzbekistan, Kazakhstan.

Cerceris angustirostris Shestakov, 1918. Russia: **EP** (CR), **WS** (AL). – Europe (EE), Kazakhstan.

Cerceris apakensis Tsuneki, 1961. Russia: **ES** (ZB). – Mongolia, China (NC).

Cerceris arenaria (Linnaeus, 1758) [Sphex] (*C. striolata* Schletterer, 1887; *C. flavesrens* Schletterer, 1889; *C. stecki* Schletterer, 1889). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, ZB), **FE** (AM, PR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran, Central Asia, Kazakhstan, Mongolia, China (NE), Korean Peninsula, Japan (Hon, Ryu).

Cerceris bicincta Klug in Waltl, 1835 (*C. quadrimaculata* Dufour, 1849; *C. leucozonica* Schletterer, 1887; *C. schulthessi* Schletterer, 1889). Russia: **EP** (C, S, CR), **UR**, **WS** (OM, NS, AL), **ES** (KS), **FE** (KH, PR). – Europe (WE, SE, EE), N Africa, Turkey, Israel, Afghanistan, Pakistan, Central Asia Kazakhstan, Mongolia, China (NE, NC), Korean Peninsula.

Cerceris bracteata Eversmann, 1849 (*C. penicillata* Mocsáry, 1879; *C. mirabilis* Shestakov, 1927). Russia: **EP** (C, E, S, CR), **UR**, **WS** (OM, NS, AL). – Europe (WE, SE, EE), Turkey, Iran, Turkmenistan, Uzbekistan, Kazakhstan, China (NW).

Cerceris bupresticida Dufour, 1841 (*C. argentifrons* Lepeletier de Saint Fargeau, 1845; *C. mixta* Radoszkowski, 1877; *C. quadripunctata* Radoszkowski, 1877). Russia: **EP** (E, S, NC, CR), **UR**, **WS** (AL). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Jordan, Lebanon, Israel, Iraq, Iran, Afghanistan, Central Asia, Kazakhstan, Mongolia, China (NE, NC).

Cerceris circularis (Fabricius, 1804) [*Philanthus*] (*C. dacica* Schletterer, 1887). Russia: **EP** (C, S, NC, CR), **UR**, **WS** (AL). – Europe (WE, SE, EE), N Africa, Armenia, Turkey, Syria, Israel, Iran, Turkmenistan, Uzbekistan, Kazakhstan, Mongolia.

Cerceris conica Shestakov, 1918. Russia: **EP** (NC). – Armenia, Turkey, Turkmenistan.

Cerceris coreensis Tsuneki, 1961. Russia: **FE** (PR). – Korean Peninsula.

Cerceris cupes Shestakov, 1918 (*C. pulawskii* Kazenas, 1972, nom. praeocc., nec Tsuneki, 1971; *C. kazenasi* Pulawski, 1976). Russia: **WS** (AL). – Turkey, Central Asia, Kazakhstan.

Cerceris dorsalis Eversmann, 1849 (*C. solskii* Radoszkowski, 1877; *C. nobilis* Radoszkowski, 1877; *C. excavata* Schletterer, 1889; *C. caspica* F. Morawitz, 1891). Russia: **EP** (C, S, NC), **UR**, **WS** (AL). – Europe (EE), Armenia, Turkey, Iran, Afghanistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NC).

Cerceris elegans Evermann, 1849. Russia: **UR**. – Kazakhstan.

Cerceris errata Shestakov, 1918. Russia: **EP** (S). – Turkmenistan, Uzbekistan, Kazakhstan.

Cerceris eryngii Marquet, 1875 (*C. haueri* Schletterer, 1887; *C. robusta* Shestakov, 1915; *C. prahenda* Kazenas, 1978). Russia: **EP** (S, CR), **UR**, **WS** (AL). – Europe (WE, SE, EE), N Africa, Turkey, Israel, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Cerceris eversmanni W. Schulz, 1912 (*C. cornuta* Eversmann, 1849, secondary homonym of *C. cornuta* (Fabricius, 1787); *C. caucasica* Shestakov, 1915). Russia: **EP** (NC, CR), **UR**, **WS** (TK, NS, KM, AL), **ES** (TU, KR, IR). – Europe (WE, SE, EE), Georgia, Azerbaijan, Turkey, Kazakhstan, Mongolia, China (NE).

Cerceris falcifera Tsuneki, 1961. Russia: **ES** (ZB). – China (NC).

Cerceris fimbriata (Rossi, 1790) [Crabro] (*C. pallidopicta* Radoszkowski, 1877; *C. polita* Schletterer, 1889; *C. cogens* Kohl, 1916). Russia: **EP** (C, S), **UR**. – Europe (WE, SE, EE), N Africa, Georgia, Armenia, Turkey, Syria, Israel, Afghanistan, Pakistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia.

Cerceris flavicornis Brullé, 1833 (*C. conigera* Dahlbom, 1845). Russia: **EP** (S, NC, CR), **UR**. – Europe (WE, SE, EE), Georgia, Turkey, Israel, Central Asia, Kazakhstan, Mongolia, China (NW).

Cerceris flavilabris (Fabricius, 1793) [*Hylaeus*] (*C. aurita* Latreille, 1805; *C. ferreri* Vander Linden, 1829; *C. laminata* Eversmann, 1849). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (OM, NS, AL), **ES** (KS). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, China (NC).

Cerceris fodiens Eversmann, 1849 (*C. charusini* F. Morawitz, 1891; *C. sareptana* Schletterer, 1887; *C. opalipennis* Kohl, 1888; *C. shur* Shestakov, 1918). Russia: **EP** (S, NC, CR), **UR**, **WS** (AL). – Europe (EE), Azerbaijan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Cerceris fulvipes Eversmann, 1849 (*C. shaman* Shestakov, 1918). Russia: **ES** (KR, IR, ZB). – Kazakhstan, China (WP).

Cerceris hohlbecki Shestakov, 1914 (*C. fragosa* Kohl, 1916). Russia: **EP** (S, NC). – Armenia, Central Asia, Kazakhstan.

Cerceris hortivaga Kohl, 1880. Russia: **FE** (AM, PR, KU). – Europe (WE, SE, EE), China (NE, NC), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu).

Cerceris impercepta de Beaumont, 1950. Russia: **EP** (S), **WS** (AL). – Europe (C, E), Kazakhstan.

Cerceris interrupta (Panzer, 1799) [Philanthus] (*C. labiata* (Fabricius, 1793), secondary homonym of *C. labiata* (Olivier, 1792); *C. peninsularis* Mercet, 1903). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (NS, AL), **ES** (KR). – Europe (WE, SE, EE), Turkey, Kazakhstan.

Cerceris jakowleffi Kohl, 1898 (*C. pulawskii* Tsuneki, 1971). Russia: **ES** (IR, BR, ZB). – Mongolia, China (NC).

Cerceris kaszabi Tsuneki, 1971. Russia: **ES** (KR, IR, BR, YA, ZB), **FE** (AM, PR, SA). – Mongolia, China (NE).

Cerceris kuznetzovi Kazenas, 1984. Russia: **EP** (S). – Uzbekistan, Kazakhstan.

Cerceris lunata A. Costa, 1867 (*C. dorsalis* Dufour, 1849, nom. praeocc., nec Eversmann, 1849; *C. funerea* A. Costa, 1867). Russia: **EP** (C, S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Armenia, Turkey, Syria, Israel, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Cerceris manchuriana Tsuneki, 1961. Russia: **FE** (PR). – China (NE).

Cerceris manflava Tsuneki, 1971 (*C. flava* Tsuneki, 1961, nom. praeocc., nec Mochi, 1939). Russia: **FE** (AM, PR). – Mongolia, China (NC).

Cerceris media Klug, 1835 (*C. capitata* F. Smith, 1856). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), Turkey, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Cerceris micropunctata Shestakov, 1918. Russia: **WS** (AL), **ES** (KS, KR, IR, BR, YA, ZB).

Cerceris odontophora Schletterer, 1887. Russia: **EP** (S, NC). – Europe (SE, EE), Azerbaijan, Turkey, Iran, Tajikistan, Kazakhstan, China (NW).

Cerceris pedetes Kohl, 1887 (*C. bicornuta* F. Smith, 1856, nom. praeocc., nec Guérin-Méneville, 1844; *C. smithii* Dalla Torre, 1897). Russia: **ES** (ZB), **FE** (AM, PR). – China (NE), Korean Peninsula, Japan (Hon).

Cerceris pucilii Radoszkowski, 1869 (*C. poutziloi* W. Schulz, 1906; *C. nupta* Shestakov, 1918). Russia: **ES** (TU, BR, ZB), **FE** (AM, KH, PR). – Mongolia, China (NE, NC, NW).

Cerceris quadricincta (Panzer, 1799) [Philanthus]. Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (AL). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iraq, Iran, Afghanistan, Central Asia, Kazakhstan.

Cerceris quadricolor F. Morawitz, 1889. Russia: **ES** (BR). – Mongolia, China (NE, NC).

Cerceris quadrifasciata (Panzer, 1799) [Philanthus] (*C. truncatula* Dahlbom, 1844). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (NS, AL), **ES** (KS, KR, IR, ZB), **FE** (AM, KH, PR, SA). – Europe (WE, NE, SE, EE), Abkhazia, Georgia, Armenia, Turkey, Kazakhstan, China (NE), Korean Peninsula.

Cerceris quinquefasciata quinquefasciata (Rossi, 1792) [Crabro] (*C. nasuta* Dahlbom, 1844). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (OM, NS, KM, AL), **ES** (KS, TU, KR). – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey, Kazakhstan.

Cerceris quinquefasciata seoulensis Tsuneki, 1961. Russia: **ES** (ZB), **FE** (AM, PR). – China (NE), Korean Peninsula.

Cerceris rossica Shestakov, 1914. Russia: **EP** (S, CR). – Europe (EE), Kazakhstan.

Cerceris rubida (Jurine, 1807) [Philanthus] (*C. albonotata* Vander Linden, 1829). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (NS, AL), **FE** (PR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Syria, Jordan, Israel, Iraq, Iran, Afghanistan, Central Asia, Kazakhstan, China (NE, NC, NW), Korean Peninsula.

Cerceris ruficornis (Fabricius, 1793) [Philanthus] (*C. cunicularia* (Schrank, 1802); *C. labiata* auct.). Russia: **EP** (N, C, E, S, CR), **UR**, **WS** (TM, NS, AL), **ES** (IR, BR, YA), **FE** (KH, SA). – Europe (WE, NE, SE, EE), N Africa, Turkey, Iran, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan, China (NE, NW), Korean Peninsula.

Cerceris rybyensis (Linnaeus, 1771) [Sphex] (*C. ornata* (Fabricius, 1790); *C. variabilis* (Schrank, 1802); *C. jacobsoni* Shestakov, 1923). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, KR, IR, BR, YA, ZB), **FE** (PR, SA). – Europe (WE, NE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Syria, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Korean Peninsula, Japan (Hon), India.

Cerceris sabulosa (Panzer, 1799) [Philanthus] (*C. emarginata* Panzer, 1799; *C. subgibbosa* Yasumatsu, 1935). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (OM, NS, AL). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Lebanon, Israel, Iraq, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NE, NC, NW, WP), Korean Peninsula.

Cerceris scutifera Shestakov, 1914 (*C. tadzhika* Gussakovskij, 1952). Russia: **EP** (NC), **WS** (AL). – Turkmenistan, Tadzhikistan, Uzbekistan, Kazakhstan.

Cerceris semilunata Radoszkowski, 1870. Russia: **ES** (KR, IR, BR, ZB), **FE** (AM, PR). – Mongolia, China (NE), Korean Peninsula.

Cerceris shestakoviiana Gussakovskij, 1952. Russia: **EP** (S). – Turkmenistan, Tadzhikistan, Uzbekistan, Kazakhstan.

Cerceris sibirica F. Morawitz, 1892. Russia: **ES** (KR, BR, ZB), **FE** (AM). – China (NE).

Cerceris somotorensis Balthasar, 1956. Russia: **EP** (S, NC, CR), **UR**. – Europe (WE, EE), Kazakhstan.

Cerceris specularis A. Costa, 1867. Russia: **EP** (S, CR), **UR**. – Europe (WE, SE, EE), N Africa, Georgia, Turkey, Syria, Jordan, Israel, Iran, Afghanistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Cerceris spinipectus F. Smith, 1856 (*C. prisca* Schletterer, 1887; *C. accola* Kohl, 1916). Russia: **EP** (S). – Europe (SE, EE), N Africa, Georgia, Armenia, Turkey, Jordan, Israel, Saudi Arabia, Yemen, Iraq, Iran, Afghanistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Cerceris stratiotes Schletterer, 1887. Russia: **EP** (NC, CR), **UR**, **WS** (AL). – Europe (SE, EE), Armenia, Azerbaijan, Turkey, Kyrgyzstan.

Cerceris tenuivittata Dufour, 1849. Russia: **EP** (S, CR). – Europe (WE, SE, EE), Turkey, Syria, Israel, Central Asia, Kazakhstan.

Cerceris tuberculata (de Villers, 1789) [Crabro] (*C. rufipes* (Fabricius, 1787); *C. caspia* (Gmelin, 1790); *C. semirufa* F. Smith, 1856; *C. evecta* Shestakov, 1918). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (OM, NS, AL), **ES** (KR, BR, ZB). – Europe (WE, SE, EE), Georgia, Armenia, Azerbaijan, Turkey, Israel, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NE, NC), Korean Peninsula.

Cerceris vagans Radoszkowski, 1877 (*C. rufonodis* Radoszkowski, 1877, nom. praeocc., nec *C. rufinodis* F. Smith, 1856; *C. turkestanica* Radoszkowski, 1893). Russia: **EP** (S, NC). – Turkey, United Arab Emirates, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan.

Cerceris verhoeffi Tsuneki, 1961. Russia: **FE** (AM, PR). – China (NE).

Cerceris vitticollis F. Morawitz, 1894. Russia: **EP** (S, CR). – Turkmenistan, Tadzhikistan, Uzbekistan, Kazakhstan.

Tribe PHILANTHINI

PHILANTHINUS de Beaumont, 1949 (*Shestakoviel-la* Gussakovskij, 1952). Type species *Philanthus integer* de Beaumont, 1949. Number of species: World – 4; Palaearctic – 4; Russia – 1.

Philanthinus quattuordecimpunctatus (F. Morawitz, 1888) [Anthophilus] (*Anthophilus elegans* F. Morawitz, 1888; *Philanthus eximius* F. Morawitz, 1894). Russia: **EP** (C, CR), **WS** (AL). – Turkey, Turkmenistan, Kazakhstan, China (NW).

PHILANTHUS Fabricius, 1790 (*Anthophilus Dahlbom*, 1844). Type species *Philanthus coronatus* Fabricius, 1790 (= *Sphex coronatus* Thunberg, 1784). Number of species: World – 133; Palaearctic – 32; Russia – 8.

Philanthus coronatus (Thunberg, 1784) [Sphex] (*Ph. coronatus* Fabricius, 1790, secondary homonym of *Ph. coronatus* (Thunberg, 1784)). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (NS, AL), **ES** (KR). – Europe (WE, SE, EE), N Africa, Turkey, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE, NC, NW), Korean Peninsula.

Philanthus decemmaculatus Eversmann, 1849 (*Ph. angustifrons* Kohl, 1891). Russia: **EP** (C, CR), **UR**, **WS** (TK). – Europe (WE), N Africa, Armenia, Turkey, Syria, Kyrgyzstan, Kazakhstan.

Philanthus desertorum (F. Morawitz, 1890). Russia: **EP** (S). – Turkmenistan, Uzbekistan.

Philanthus glaber Kohl, 1891 (*Ph. rubriventris* Kazenas, 1970; *Ph. kaszabi* Tsuneki, 1971). Russia: **WS** (AL), **ES** (BR, ZB). – Kazakhstan, Mongolia, China (NC).

Philanthus hellmanni (Eversmann, 1849) [Anthophilus] (*Ph. sibiricus* Radoszkowski, 1888). Russia: **WS** (AL), **ES** (KR, IR, BR, ZB), **FE** (AM, PR). – Tajikistan, Kazakhstan, Mongolia, China (NC).

Philanthus triangulum (Fabricius, 1775) [Vespa]. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (NS, KM, AL). – Europe (WE, NE, SE, EE), N Africa, Abkhazia, Georgia, Azerbaijan, Turkey, Syria, Jordan, Israel, Saudi Arabia, Yemen, Iraq, Iran, Afghanistan, Central Asia, Kazakhstan, Africa.

Philanthus variegatus Spinola, 1839 (*Ph. kizilkumi* Radoszkowski, 1877). Russia: **EP** (S). – N Africa, Armenia, Azerbaijan, Turkey, Syria, Israel, Saudi Arabia, Kuwait, United Arab Emirates, Oman, Iraq, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, China (NC).

Philanthus venustus (Rossi, 1790) [Crabro]. Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Turkey, Syria, Israel, Kazakhstan, Mongolia.

Tribe PSEUDOSCOLIINI

PSEUDOSCOLIA Radoszkowski, 1876 (*Philoponus* Kohl, 1889; *Acolpus* Vachal, 1893; *Philoponidea* Pate, 1937; *Philoponoides* Giner Marí, 1945). Type species *Pseudoscolia maculata* Radoszkowski, 1876. Number of species: World – 49; Palaeartic – 48; Russia – 5.

Pseudoscolia corporica Kazenas, 1994. Russia: **EP** (S). – Turkmenistan.

Pseudoscolia diversicornis (F. Morawitz, 1894) [*Philanthus*]. Russia: **EP** (CR). – Turkmenistan, Tajikistan, Uzbekistan.

Pseudoscolia frontalis Kazenas, 1994. Russia: **EP** (S). – Turkmenistan, Kazakhstan.

Pseudoscolia marshakovi Kazenas, 1994. Russia: **EP** (S). – Turkmenistan.

Pseudoscolia simplicicornis (F. Morawitz, 1894) [*Philanthus*] (*Philanthus variegatus* (F. Morawitz, 1889), secondary homonym of *Ph. variegatus* Spinola, 1839). Russia: **EP** (S, NC), **ES** (ZB). – Europe (EE), Turkmenistan, Uzbekistan, Kazakhstan, Mongolia.

SECTION APIFORMES – BEES

73. FAMILY COLLETIDAE

M.YU. PROSHCHALYKIN

The family contains perhaps 2600 species around the world. There are five subfamilies: Colletinae, Hylaeinae, Diphaglossinae, Xeromelissinae, and Euryglossinae. The subfamily Hylaeinae is found on all continents except Antarctica but is most abundant and diverse in Australia. It is represented in Palaearctic region by the genus *Hylaeus*, which contains small, slender, black species with two submarginal cells, usually with yellow markings at least on the face. Females lack external pollen-carrying apparatus (scopa), they carry pollen in their crops. Most species nest in dead hollow or pithy stems. The subfamily Colletinae occurs around the world but it is diverse only in the southern continents. It is represented in Palaearctic region only by genus *Colletes*. Females carry pollen externally, on the well-developed scopa of the hind legs (trochanter to tibia). Nests are in burrows in the ground. Number of taxa: World – 86 genera, 2630 species; Palaearctic – 2/432; Russia – 2/100.

References. Eversmann, 1852; Morawitz, 1874; Radoszkowski, 1891; Noskiewicz, 1936; Nikiforuk, 1957; Osytshnjuk, 1970; Blagoveshchenskaya, 1970; Warncke, 1972, 1978; Sedykh, 1974; Osytshnjuk et al., 1978; Dathe, 1980a, b; Sittikov, 1986; Ikudome, 1989; Osytshnjuk, Romankova, 1995; Proshchalykin, 2004, 2007a, 2008a, 2010, 2012a, 2014a, 2017; Humala, 2006; Kuhlmann, Quest, 2006; Levchenko, 2009; Kuhlmann, Proshchalykin, 2011, 2013a, b, 2014, 2016; Proshchalykin, Kuhlmann, 2012, 2015a, b; Proshchalykin, Dathe, 2012, 2016a, b, 2017; Shibaev, Polomordvinov, 2012; Levchenko, Yuferev, 2013; Proshchalykin, Lelej, 2013; Paukkunen, Kozlov, 2015; Dathe, Proshchalykin, 2016, 2017; Proshchalykin, Astafurova, 2016; Ascher, Pickering, 2017; Kuhlmann et al., 2017.

Subfamily COLLETINAE

COLLETES Latreille, 1802 (*Evodia* Panzer, 1806; *Monia* Westwood, 1875; *Monidia* Cockerell, 1905). Type species *Apis succincta* Linnaeus, 1758. Number of species: World – 500; Palaearctic – 212; Russia – 43.

Colletes albomaculatus (Lucas, 1849) [Halictus] (*C. spectabilis* Morawitz, 1868; *C. niveofasciatus*

Dours, 1872). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Syria, Jordan, Iran, Tajikistan, Kyrgyzstan.

Colletes alini Kuhlmann, 2000. Russia: **ES** (KS, TU, KR, BR, ZB), **FE** (PR). – Mongolia, China (NE, NC, NW).

Colletes anceps Radoszkowski, 1891 (*C. caspicus anceps*: Warncke, 1978). Russia: **EP** (NC). – Turkey, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, China (NW).

Colletes arsenjevi Kuhlmann, 2006. Russia: **FE** (PR). – Kazakhstan, Mongolia, China (NC, NW).

Colletes caspicus Morawitz, 1874 (*C. balticus* Alfken, 1912). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (OM, AL), **ES** (TU, KR). – Europe (WE, NE, EE), Georgia, Azerbaijan, Central Asia, Kazakhstan.

Colletes chengtehensis Yasumatsu, 1935 (*C. pallescens* Noskiewicz, 1936; *C. tsunekii* Yasumatsu, 1948; *C. marginatus pallescens*: Warncke, 1978). Russia: **EP** (NW, S, NC, CR), **UR**, **WS** (OM, NS, KM, AL), **ES** (KS, TU, KR, BR, ZB). – Europe (WE, SE, EE), Georgia, Azerbaijan, Iran, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE, NC, NW).

Colletes cinerascens Morawitz, 1893. Russia: **ES** (TU). – Tajikistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NW, WP).

Colletes collaris Dours, 1872 (*C. incerta* Radoszkowski, 1891; *C. frigidus* Pérez, 1903; *Andrena riukiensis* Matsumura, 1926). Russia: **EP** (C, E), **WS** (KM, AL), **ES** (TU, KR, IR, BR, ZB), **FE** (KH, PR, SA, KU). – Europe (WE, SE, EE), Kazakhstan, Mongolia, China (NE, NC, CC, SW, WP).

Colletes cunicularius (Linnaeus, 1761) [*Apis*] (*Apis cinerea* Geoffroy, 1785; *Colletes hirtus* Lepeletier de Saint Fargeau, 1825; *C. pilosus* Spinola, 1838; *C. cunicularius infuscatus* Noskiewicz, 1936; *C. c. celticus* O'Tolle, 1974; *C. c. tauricus* Warncke, 1978; *C. c. khasanensis* Osytshnjuk, 1995). Russia: **EP** (NW, C, E, S, CR), **UR**, **WS** (TK, KM, AL), **ES** (KR, IR, BR), **FE** (KH, PR). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan, Mongolia, China (NE, NC).

Colletes daviesanus Smith, 1846 (?*Apis canus* Gmelin, 1790; *Colletes daviesanus signatus* Verhoeff, 1890). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (OM, TK, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH). – Europe (WE, NE, SE, EE), Georgia, Armenia, Turkey, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE, WP).

Colletes ebmeri Kuhlmann, 2002. Russia: **ES** (BU). – Kazakhstan, Mongolia.

- Colletes eous** Morice, 1904 (*C. cecrops* Morice, 1904; *C. porosicus* Strand, 1921; *C. illyricus* Noskiewicz, 1936; *C. dimidiatus eous*: Warncke, 1978). Russia: **EP** (C, S, NC, CR). – Europe (WE, SE, EE), N Africa, Azerbaijan, Georgia, Turkey, Lebanon, Iran, Central Asia, Kazakhstan, India.
- Colletes floralis** Eversmann, 1852 (*C. alpinus* Morawitz, 1872; *C. montana* Morawitz, 1876; *C. suecica* Aurivillius, 1903; *C. kudiensis* Cockerell, 1924; *C. yasumatsui* Hirashima et Ikudome, 1989). Russia: **EP** (N, C, E, S, NC), **UR**, **WS** (TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU). – Europe (WE, NE, SE, EE), Georgia, Turkey, Iran, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, Japan (Hok, Hon), India.
- Colletes fodiens** (Fourcroy, 1785) [*Apis*] (?*Apis fosiarius* Razoumowsky, 1789; *A. palidicincta* Kirby, 1802; *Colletes kirgisica* Radoszkowski, 1867; *C. fodiens hispanicus* Noskiewicz, 1936). Russia: **EP** (NW, C, S, NC, CR), **UR**, **WS** (OM, KM, AL), **ES** (KS, TU, KR). – Europe (WE, NE, SE, EE), Azerbaijan, Turkey, Iran, Pakistan, Kyrgyzstan, Kazakhstan, Mongolia.
- Colletes friesei** Cockerell, 1918 (*C. mongolicus* Friese, 1913, nom. praeocc., nec Pérez, 1903). Russia: **ES** (BU). – Tajikistan, China (NC).
- Colletes fulvicornis** Noskiewicz, 1936. Russia: **WS** (AL), **ES** (TU, ZB). – Mongolia, China (NW, WP).
- Colletes hakkari** Kuhlmann, 2002. Russia: **EP** (NC). – Turkey, Iran.
- Colletes hederae** Schmidt et Westrich, 1993. Russia: **EP** (CR). – Europe (WE, SE).
- Colletes hylaeiformis** Eversmann, 1852 (*C. chobauti* Pérez, 1903). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), Georgia, Azerbaijan, Tajikistan, Uzbekistan, Kazakhstan.
- Colletes impunctatus** Nylander, 1852. Russia: **EP** (N, NW, E), **WS** (AL), **ES** (KS, TU, IR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KA). – Europe (WE, NE, SE, EE), Kazakhstan, Mongolia, China (NC).
- Colletes inexpectatus** Noskiewicz, 1936. Russia: **EP** (C, S), **UR**. – Europe (WE, SE, EE), Turkey, Central Asia, Kazakhstan, Mongolia.
- Colletes jankowskyi** Radoszkowski, 1891 (*C. ventralis* Pérez, 1903; *C. ventraliformis* Cockerell, 1924). Russia: **WS** (AL), **ES** (KS, TU, KR, IR, BR, ZB), **FE** (AM, KH, PR). – Tajikistan, Mongolia, China (NE, NC, SW).
- Colletes kaszabi** Kuhlmann, 2002. Russia: **ES** (TU, BR). – Mongolia, China (NC).
- Colletes kozlovi** Friese, 1913. Russia: **EP** (S, NC), **ES** (TU, BR). – Azerbaijan, Central Asia, Kazakhstan, Mongolia, China (NW).
- Colletes laevifrons** Morawitz, 1893. Russia: **WS** (AL). – Iran, Tajikistan, Kyrgyzstan, Kazakhstan.
- Colletes maidli** Noskiewicz, 1936 (*C. lebedewi* Noskiewicz, 1936). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Syria, Israel, Iran, Kazakhstan.
- Colletes marginatus** Smith, 1846 (*C. thomsoni* Alfken, 1899). Russia: **EP** (NW, C, E, S), **UR**, **WS** (OM, TK, KM, AL), **ES** (TU, KR). – Europe (WE, NE, SE, EE), Georgia, Turkey.
- Colletes mlokosewiczi** Radoszkowski, 1891 (*C. brevicornis* Pérez, 1903, nom. praeocc., nec Robertson, 1897; *C. brachycerus* Swenk, 1906). Russia: **EP** (N, C, S, NC, CR), **UR**. – Europe (WE, SE, EE), Georgia, Azerbaijan, Turkey, Central Asia, Kazakhstan.
- Colletes nasutus** Smith, 1853. Russia: **EP** (S, NC). – Europe (WE, SE, EE), Armenia, Azerbaijan, Turkey, Syria, Iran.
- Colletes patellatus** Pérez, 1905 (*C. kobensis* Cockerell, 1926). Russia: **ES** (ZB). – China (NC, CC, SW, SE), Korean Peninsula, Japan (Hon, Shi, Kyu).
- Colletes perforator** Smith, 1869 (*C. vogti* Pérez, 1905). Russia: **ES** (ZB), **FE** (AM, PR). – Mongolia, China (NE, NC, CC), Japan (Hok, Hon, Shi, Kyu).
- Colletes pseudocinerascens** Noskiewicz, 1936. Russia: **WS** (AL), **ES** (TU, YA). – Mongolia.
- Colletes ravulus** Noskiewicz, 1936. Russia: **ES** (TU, BR). – Mongolia, China (NC, NW).
- Colletes roborovskyi** Friese, 1913. Russia: **EP** (E, S), **ES** (BR). – Georgia, Azerbaijan, Turkey, Iran, Pakistan, Turkmenistan, Tajikistan, Kyrgyzstan, Kazakhstan, Mongolia.
- Colletes senilis** (Eversmann, 1852) [*Andrena*] (*C. juvenilis* Eversmann, 1852). Russia: **EP** (E, S), **UR**. – Europe (SE, EE), Azerbaijan, Turkey, Iran, Kazakhstan.
- Colletes sidemii** Radoszkowski, 1891 (*C. mongolicus* Pérez, 1903). Russia: **EP** (E), **UR**, **WS** (AL), **ES** (KS, TU, KR, IR, BR, ZB), **FE** (AM, PR). – Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE, NC, NW, SW, WP), Nepal.
- Colletes similis** Schenck, 1853 (*C. picistigma* Thomson, 1872). Russia: **EP** (C, S, NC, CR), **UR**, **WS** (TM, NS, KM, AL), **ES** (KS, KR, IR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Syria, Jordan, Israel, Central Asia, Kazakhstan, China (NW).

Colletes subnitens Noskiewicz, 1936. Russia: **EP** (S). – Iran, Kazakhstan.

Colletes succinctus (Linnaeus, 1758) [Apis] (*Apis glutinans* Cuvier, 1798; *A. calendarum* Panzer, 1806; *Colletes balteatus* Nylander, 1852; *Andrena xanthothorax* Eversmann, 1852). Russia: **EP** (N, NW, C, E, NC), **UR**. – Europe (WE, NE, SE, EE), Georgia, Kazakhstan.

Colletes tuberculatus Morawitz, 1893 (*C. tuberculatus siculus* Noskiewicz, 1959; *C. t. anatolicus* Noskiewicz, 1959; *C. daviesanus tuberculatus*: Warncke, 1978). Russia: **EP** (C, S, CR). – Europe (SE, EE), N Africa, Georgia, Armenia, Turkey, Jordan, Israel, Iran, Pakistan, Central Asia, Kazakhstan.

Colletes ulrikae Kuhlmann, 2002. Russia: **FE** (MG).

Colletes wacki Kuhlmann, 2002. Russia: **EP** (S), **ES** (TU). – Kazakhstan, Mongolia, China (NC).

Colletes warnekei Kuhlmann, 2002. Russia: **EP** (NC). – Turkey, Iran.

Subfamily HYLAEINAE

HYLAEUS Fabricius, 1793 (*Prosopis* Fabricius, 1804).

Type species *Prosopis annulata* Fabricius, 1804. Number of species: World – 760; Palaearctic – 220; Russia – 57.

Hylaeus (Abrupta) cornutus Curtis, 1831 (*H. cornutus* Smith, 1842; *H. plantaris* Smith, 1842; *H. pluminicornis* Costa, 1884). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, NE, SE, EE), N Africa, Armenia, Azerbaijan, Turkey, Syria, Israel, Iran.

Hylaeus (Dentigera) brevicornis Nylander, 1852 (*Prosopis pygmaea* Schenck, 1853; *P. rubicola* Smith, 1869, nom. praeocc., nec Saunders, 1850; *Hylaeus discretus* Förster, 1871; *H. atratulus* Förster, 1871, nom. praeocc., nec Schenck, 1853; *H. suspectus* Förster, 1871; *Prosopis brevicornis sardoa* Alfken, 1934; *Hylaeus parvulus* Janvier, 1972, nom. praeocc., nec Schenck, 1853). Russia: **EP** (NW, C, E, S, NC, CR), **WS** (KM, AL), **ES** (KR). – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey, Israel, Iran, Central Asia.

Hylaeus (Dentigera) gredleri Förster, 1871 (*H. fumipennis* Förster, 1871; *H. brevipalpis* Förster, 1871; *H. confinis* Förster, 1871; *H. rotundatus* Förster, 1871). Russia: **EP** (C, S, NC), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey, Lebanon.

Hylaeus (Dentigera) pallidicornis Morawitz, 1876 (*Prosopis breviceps* auct., nec Morawitz, 1876).

Russia: **EP** (NC, CR). – Europe (EE), Uzbekistan, Kyrgyzstan.

Hylaeus (Hylaeus) aborigensis Dathe, 1994 (*Prosopis chasanensis* Romankova, 1995; *Hylaeus chasanensis negoroi* Ikudome, 2004). Russia: **WS** (AL), **ES** (KS, TU, YA), **FE** (AM, KH, PR, SA, KU, MG). – Mongolia, Japan (Hon).

Hylaeus (Hylaeus) altaicus Dathe, 1986. Russia: **WS** (AL), **ES** (TU, IR, ZB). – Mongolia.

Hylaeus (Hylaeus) angustatus (Schenck, 1861) [*Prosopis*] (*H. subexcisus* Förster, 1871; *H. subtilis* Förster, 1871; *H. subpunctatus* Förster, 1871; *H. submarginatus* Thomson, 1872; *Prosopis punctifrons* Pérez, 1903; *Hylaeus pauperculus* Cockerell, 1931; *Prosopis curviscapa* Benoist, 1960). Russia: **EP** (NW, C, S, NC, CR), **WS** (AL), **ES** (KS, TU). – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey, Lebanon, Iran, Mongolia.

Hylaeus (Hylaeus) annulatus (Linnaeus, 1758) [Apis] (*Apis xanthometopa* Preyssler, 1793; *Prosopis elliptica* Kirby, 1827; *P. patellata* Eversmann, 1852; *P. varifrons* Cresson, 1869; *P. antennata* Cresson, 1869; *Hylaeus barbatus* Förster, 1871; *Prosopis tamanukii* Yasumatsu, 1939). Russia: **EP** (NW, C, E, CR), **UR**, **WS** (TM, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, MG, CH). – Europe (WE, NE, SE, EE), Armenia, Azerbaijan, Mongolia, N America.

Hylaeus (Hylaeus) cardioscapus Cockerell, 1924 (*Prosopis vallei* Niemelä, 1947). Russia: **EP** (C, E), **WS** (TM, KM, AL), **ES** (KS, TU, KR, IR, ZB), **FE** (AM, KH, PR, SA, KA, MG). – Europe (WE, NE, SE, EE), Azerbaijan, Mongolia.

Hylaeus (Hylaeus) communis Nylander, 1852 (?*Vespa minuta* Rossi, 1790, nom. praeocc., nec Linné, 1767; ?*Hylaeus triangulus* Fabricius, 1793; *H. communis* Nylander, 1852; *Prosopis ciliata* Eversmann, 1852; *Hylaeus alienatus* Förster, 1871; *H. marginatus* Förster, 1871, nom. praeocc., nec Thomson, 1870; *H. ebeninus* Förster, 1871; *H. nigricornis* Förster, 1871, nom. praeocc., nec Schenck, 1853; *H. nitidulus* Förster, 1871, nom. praeocc., nec Fabricius, 1804; *Prosopis rupestris* Smith, 1872; *P. annulata* var. *tristis* Dalla Torre, 1877; *P. foersteri* Dalla Torre et Friese, 1895; *P. annulata* var. *nigrifacies* Alfken, 1913; *P. communis* var. *reiecta* Stoeckhert, 1954). Russia: **EP** (NW, C, E, NC, CR), **UR**, **WS** (TM, KM, AL), **ES** (IR, ZB), **FE** (AM, KH, PR, SA, KA, MG). – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey, Lebanon, Iran, Central Asia, Kazakhstan.

- Hylaeus (Hylaeus) dolichocephalus** Morawitz, 1876
[*Prosopis plurifasciata* Alfken, 1931; *P. heliaca* Warncke, 1992]. Russia: **ES** (TU). – Turkey, Afghanistan, Tajikistan, Kazakhstan, Mongolia, China (NW).
- Hylaeus (Hylaeus) dorni** Dathe, 1986. Russia: **ES** (TU). – Mongolia.
- Hylaeus (Hylaeus) gracilicornis** (Morawitz, 1867)
[*Prosopis*] (*H. nigrolineatus* Cockerell, 1924). Russia: **EP** (NW, C, E, NW, NC), **WS** (TM, NS, KM, AL), **ES** (KS, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KM, MG). – Europe (WE, NE, SE, EE), Azerbaijan, Turkey, Syria, Iran, Mongolia, China (SW).
- Hylaeus (Hylaeus) hungaricus** (Alfken, 1905) [*Prosopis*] (*H. affinis* Morawitz, 1876, nom. praeocc., nec Smith, 1853; *Prosopis morawitzi* Dalla Torre, 1896, nom. praeocc., nec Radoszkowski, 1893). Russia: **WS** (AL). – Iran, Afghanistan, Turkmenistan, Tajikistan, Mongolia, China (NW).
- Hylaeus (Hylaeus) leleji** Proshchalykin et Dathe, 2016. Russia: **WS** (AL).
- Hylaeus (Hylaeus) leptocephalus** (Morawitz, 1871)
[*Prosopis*] (*H. bisinuatus* Förster, 1871; *Prosopis discrepans* Schenck, 1875; *P. incompleta* Alfken, 1937; *P. stevensi* Crawford, 1913). Russia: **EP** (C, E, S, NC), **UR**, **WS** (TM, KM), **ES** (TU), **FE** (KH). – Europe (WE, NE, SE, EE), Armenia, Azerbaijan, Iran.
- Hylaeus (Hylaeus) montivagus** Dathe, 1986. Russia: **WS** (AL), **ES** (TU), **FE** (KH, PR). – Mongolia.
- Hylaeus (Hylaeus) moricei** (Fries, 1898) [*Prosopis*] (*Prosopis gracilicornis* var. *luteifrons* Strand, 1909; *P. moricella* Bischoff, 1954; *P. m. f. atricallosa* Bischhoff, 1954; *P. nigrifacies rhenana* Warncke, 1986). Russia: **EP** (C, S, NC), **ES** (KS). – Europe (WE, SE, EE), N Africa, Armenia, Azerbaijan, Turkey, Iran.
- Hylaeus (Hylaeus) nigritus** (Fabricius, 1798) [Melittinus] (*Prosopis atrata* Fabricius, 1804; *P. laevigata* Eversmann, 1852; *Hylaeus propinquus* Nylander, 1852; *Prosopis nitidiuscula* Schenck, 1853; *P. tuberculata* Smith, 1853). Russia: **EP** (C, E), **UR**. – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey, Iran.
- Hylaeus (Hylaeus) nimbatus** Dathe, 1986. Russia: **WS** (AL), **ES** (TU). – Mongolia.
- Hylaeus (Hylaeus) oehlkei** Dathe, 2010. Russia: **WS** (AL), **ES** (TU, KR). – Mongolia.
- Hylaeus (Hylaeus) paulus** Bridwell, 1919 (*H. lepidulus* Cockerell, 1924). Russia: **EP** (C), **WS** (AL), **ES** (KS, TU, YA), **FE** (AM, KH, PR, SA, KU, KA). – Europe (WE, NE, SE, EE), Azerbaijan, Mongolia, China (NE).
- (WE, NE, SE, EE), Azerbaijan, Mongolia, China (NE, NC), Japan (Hok, Hon, Kyu).
- Hylaeus (Hylaeus) pesenkoi** Proshchalykin et Dathe, 2016. Russia: **ES** (TU). – Mongolia.
- Hylaeus (Hylaeus) scutellaris** Morawitz, 1874 (*H. flavigipes* Morawitz, 1876, nom. praeocc., nec Smith, 1853; *Prosopis morawitzi* Radoszkowski, 1893; *P. turcestanica* Dalla Torre, 1896). Russia: **EP** (S, NC). – Azerbaijan, Afghanistan, Turkmenistan, Uzbekistan.
- Hylaeus (Hylaeus) sibiricus** (Strand, 1909) [*Prosopis*] (*Prosopis sjostedti* Alfken, 1936; *Hylaeus communis excurrens* Cockerell, 1937). Russia: **WS** (AL), **ES** (TU, IR, BR), **FE** (KH, PR). – Mongolia, China (NE, NC).
- Hylaeus (Hylaeus) stubbei** Dathe, 1986 (*H. peregrinus* Dathe, 1986). Russia: **WS** (AL). – Mongolia.
- Hylaeus (Hylaeus) telmenicus** Dathe, 1986. Russia: **WS** (AL). – Mongolia.
- Hylaeus (Hylaeus) tsingtauensis** (Strand, 1915)
[*Prosopis*] (*H. pallescens* Cockerell, 1924; *H. stenoriscapus* Dathe, 1986). Russia: **WS** (TM, KM, AL), **ES** (KS, TU, KR, IR, BR, ZB), **FE** (AM, KH, PR). – Mongolia.
- Hylaeus (Koptogaster) punctulatissimus** Smith, 1842 (*Prosopis obscurata* Schenck, 1853; *P. quadrimaculata* Schenck, 1861). Russia: **EP** (C, S, NC, CR). – Europe (WE, SE, EE), N Africa, Azerbaijan, Turkey, Israel.
- Hylaeus (Lambdopsis) dilatatus** (Kirby, 1802) [Melitta] (*Melitta annularis* Kirby, 1802; *Prosopis varians* Spinola, 1806; *Hylaeus distans* Eversmann, 1852; *Prosopis cervicornis* Costa, 1858; *Hylaeus compar* Förster, 1871; *H. smithii* Förster, 1871; *H. oculatus* Förster, 1871; *H. schenckii* Förster, 1871; *Prosopis consobrina* Schenck, 1874; *Hylaeus strigulosus* Costa, 1884; *Prosopis annularis elbursa* Warncke, 1981). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (TM, KM, AL), **ES** (KS, TU, KR, IR). – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey, Kazakhstan.
- Hylaeus (Lambdopsis) euruscapus** Förster, 1871 (*H. rimosus* Förster, 1871). Russia: **EP** (CR). – Europe (WE, SE, EE), N Africa, Turkey.
- Hylaeus (Lambdopsis) pfankuchi** (Alfken, 1919)
[*Prosopis*] (*H. nipponicus* Bridwell, 1919; *H. polevoiae* Cockerell, 1924; *Prosopis nigrifacies* Gussakovskij, 1932). Russia: **WS** (KM, AL), **ES** (KS, TU, KR), **FE** (KH, PR, SA, KU). – Europe (WE, NE, SE, EE), Azerbaijan, Mongolia, China (NE).
- Hylaeus (Lambdopsis) rinki** (Gorski, 1852) [*Prosopis*] (*Prosopis gerstaeckeri* Hensel, 1870; *Hylaeus*

- clathratus* Thomson, 1870; *H. immaculatus* Förster, 1871; *H. atromicans* Cockerell, 1924; *H. sinicola* Cockerell, 1924). Russia: **EP** (NW, C, E), **WS** (TM, NS, KM, AL), **ES** (KS, TU, KR, IR, YA, ZB), **FE** (AM, KH, PR, SA, KU). – Europe (WE, SE, EE), Azerbaijan, Central Asia, Mongolia.
- Hylaeus (Nesohylaeus) niger** Bridwell, 1919. Russia: **FE** (KU). – China (CC), Japan (Hok, Hon).
- Hylaeus (Nesoprosopis) floralis** (Smith, 1873) [*Prosopis*]. Russia: **FE** (SA). – China (CC, SW, SE), Japan (Hok, Hon, Shi, Kyu).
- Hylaeus (Nesoprosopis) globulus** (Vachal, 1903) [*Prosopis*]. Russia: **FE** (PR, SA). – Japan (Hok, Hon, Shi, Kyu).
- Hylaeus (Nesoprosopis) noomen** Hirashima, 1977. Russia: **FE** (KH). – Japan (Hok, Hon, Shi, Kyu).
- Hylaeus (Nesoprosopis) pectoralis** Förster, 1871 (*H. kriechbaumeri* Förster, 1871; *Prosopis palustris* Perkins, 1900). Russia: **EP** (NW), **WS** (TM, KM, AL), **ES** (KS), **FE** (PR, SA, KU). – Europe (WE, NE, SE, EE), Azerbaijan, Turkey, Mongolia, Japan (Hok, Hon).
- Hylaeus (Nesoprosopis) transversalis** Cockerell, 1924 (*Prosopis transversalis* Gussakovskij, 1932; *P. sericata* Warncke, 1972; *Hylaeus nippon* Hirashima, 1977). Russia: **FE** (PR, KU). – Japan (Hok, Hon, Shi, Kyu, Tsu).
- Hylaeus (Paraprosopis) clypearis** (Schenck, 1853) [*Prosopis*] (*Prosopis angulata* Smith, 1853; *Hylaeus floridula* Förster, 1871; *H. siculus* Förster, 1871; *H. blandus* Förster, 1871; *H. opacus* Förster, 1871; *H. moniliatus* Förster, 1871; *Prosopis subopaca* Pérez, 1905; *P. clypearis* var. *genimaculata* Strand, 1909; *Hylaeus lineolellus* Cockerell, 1938). Russia: **EP** (E, CR), **UR**. – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey.
- Hylaeus (Paraprosopis) concinnus** Cockerell, 1924 (*H. monostictus* Cockerell, 1924). Russia: **FE** (PR). – China (NE, NC).
- Hylaeus (Paraprosopis) lineolatus** (Schenck, 1861) [*Prosopis*] (*H. inaequalis* Förster, 1871; *Prosopis lineolata* var. *nigrifacialis* Alfken, 1913; *P. creticola* Strand, 1915; *P. lineolata* *rudis* Pittioni, 1950; *P. l. ferina* Warncke, 1992). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), N Africa, Azerbaijan, Turkey, Jordan, Lebanon, Israel.
- Hylaeus (Paraprosopis) pictipes** Nylander, 1852 (*Sphex annulata* Panzer, 1798, nom. praecocc., nec Linnaeus, 1758; *Prosopis varipes* Smith, 1853; *P. excisa* Schenck, 1853; *Hylaeus vicinus* Förster, 1871, nom. praecocc., nec Sichel, 1867; *H. melanarius* Förster, 1871; *H. claripennis* Förster, 1871; *H. panzeri* Förster, 1871; *Prosopis trimaculata* Schenck, 1875; *P. aquisgranensis* Dalla Torre, 1896; *P. pictipes* var. *nigriscutum* Alfken, 1913). Russia: **EP** (CR). – Europe (WE, NE, SE, EE), Azerbaijan.
- Hylaeus (Paraprosopis) sinuatus** (Schenck, 1853) [*Prosopis*] (*Prosopis minuta* var. *nana* Alfken, 1927, nom. praecocc., nec Friese, 1924). Russia: **EP** (C, E, NC, CR). – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey, Iraq, Iran.
- Hylaeus (Paraprosopis) styriacus** Förster, 1871 (*Prosopis tristis* Frey-Gessner, 1900, nom. praecocc., nec Dalla Torre, 1870; *P. styriaca* var. *nigroelypeata* Alfken, 1913). Russia: **EP** (C, S, NC, CR), **WS** (AL). – Europe (WE, NE, SE, EE), Armenia, Azerbaijan, Turkey, Lebanon, Iran.
- Hylaeus (Patagiata) difformis** (Eversmann, 1852) [*Prosopis*] (*Prosopis subfasciata* Schenck, 1868, nom. praecocc., nec Imhoff, 1832, nec Schenck, 1861; *Hylaeus marginatus* Thomson, 1870; *Prosopis difformis hispanica* Warncke, 1972). Russia: **EP** (C, E, NC, CR), **UR**, **WS** (TM, KM), **FE** (PR). – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey, Iran, China (NW, SW).
- Hylaeus (Patagiata) nigrocuneatus** Cockerell, 1924 (*H. nigrocuneatus* var. *rufipennis* Cockerell, 1924; *H. brevicuneatus* Cockerell, 1924; *H. laevifrons* Cockerell, 1924; *H. paradiformis* Ikudome, 1989). Russia: **WS** (KM, AL), **ES** (TU, KR, IR, YA, ZB), **FE** (AM, KH, PR, SA). – Mongolia, China (NE, CC), Japan (Hok, Hon, Kyu).
- Hylaeus (Prosopis) confusus** Nylander, 1852 (*Prosopis sublaevis* Schenck, 1853; *Hylaeus xanthocnemis* Förster, 1871; *H. nigriceps* Förster, 1871; *H. similis* Förster, 1871; *Prosopis pallididens* Dalla Torre, 1896; *Hylaeus monticola* Bridwell, 1919; *H. wilmatiae* Cockerell, 1924). Russia: **EP** (NW, C, E, NC, CR), **WS** (TM, KM, AL), **ES** (KS, TU, KR, YA, ZB), **FE** (AM, KH, PR, SA, KU). – Europe (WE, NE, SE, EE), Armenia, Azerbaijan, Turkey, Central Asia, Mongolia, China (NE, NW, WP), Japan (Hok, Hon).
- Hylaeus (Prosopis) gibbus** Saunders, 1850 (*Prosopis mixta* Schenck, 1861, nom. praecocc., nec Smith, 1852; *Hylaeus kirschbaumi* Förster, 1871; *H. aemulus* Förster, 1871; *H. genalis* Thomson, 1872). Russia: **EP** (C, E, NC, CR). – Europe (WE, SE, EE), N Africa, Azerbaijan, Turkey, Iraq, Iran.
- Hylaeus (Prosopis) hyrcanius** Dathe, 1980. Russia: **EP** (S, NC). – Azerbaijan, Iran.

Hylaeus (Prosopis) incongruus Förster, 1871

(*H. genalis* Thomson, 1872; *H. biareolatus* Morawitz, 1876). Russia: **EP** (S, NC), **ES** (BR). – Europe (WE, SE, EE), Uzbekistan, Kazakhstan.

Hylaeus (Prosopis) meridionalis Förster, 1871 (*Prosopis variegata* var. *brevimaculata* Strand, 1909; *P. maculiscutum* Alfken, 1928; *P. effasciata* Alfken, 1930). Russia: **EP** (S, NC). – Europe (WE, SE), N Africa, Azerbaijan, Turkey, Jordan, Israel.**Hylaeus (Prosopis) rugicollis** Morawitz, 1874. Russia: **EP** (NC). – Europe (SE), Caucasus, Turkey, Lebanon, Israel, Iran.**Hylaeus (Prosopis) signatus** (Panzer, 1798) [Sphex]
(*Vespa pratensis* Geoffroy, 1785, nom. praeocc., nec Miller, 1759; *Mellinus bipunctatus* Fabricius, 1798; *?Philanthes bilineatus* Gravenhorst, 1807; *Prosopis albinasata* Strand, 1912; *P. pratensis* var. *nigrifacies* Alfken, 1913; *P. berlandi* Benoist, 1943). Russia: **EP** (C, E, CR), **UR**. – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey, Iran, Uzbekistan.**Hylaeus (Prosopis) variegatus** (Fabricius, 1798) [*Mellinus*] (*Mellinus labiatus* Fabricius, 1798; *Philanthes analis* Fabricius, 1804; *Prosopis colorata* Panzer, 1805; *P. coriacea* Pérez, 1895; *P. variegata* var. *obscura* Höppner, 1901; *Hylaeus hookeri* Cockerell, 1931;

Prosopis impressifrons Gussakovskij, 1932; *P. rufithorax* Alfken, 1933; *P. variegata relicta* Warncke, 1981). Russia: **EP** (C, E, NC, CR), **UR**, **WS** (OM, TM, KM, AL), **ES** (KS, TU, KR, IR, BR, ZB), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), N Africa, Armenia, Azerbaijan, Turkey, Syria, Lebanon, Iran, Central Asia, Mongolia, China (NC).

Hylaeus (Spatulariella) hyalinatus Smith, 1842 (*Vespa bipunctata* Fadricius, 1798, nom. praeocc., nec Fabricius, 1798; *Hylaeus hyalinatus* Smith, 1842; *H. ormillatus* Nylander, 1848; *Prosopis longicornis* Schenck, 1853; *P. similis* Schenck, 1853, nom. praeocc., nec Smith, 1853, nec Fabricius, 1793; *Hylaeus decipiens* Förster, 1871; *H. tricuspidis* Förster, 1871; *H. miscellus* Förster, 1871; *Prosopis hyalinata* var. *lugubris* Dalla Torre, 1877; *Spatulariella hyalinata montebovensis* Pittioni, 1950; *S. h. aragonensis* Pittioni, 1950; *Prosopis hyalinata minos* Warncke, 1981). Russia: **EP** (C, NC, CR). – Europe (WE, NE, SE, EE), N Africa, Armenia, Azerbaijan, Turkey, Iran.**Hylaeus (Spatulariella) punctatus** (Brüllle, 1832)
[*Prosopis*] (*H. subquadratus* Förster, 1871; *H. insignis* Förster, 1871; *H. corvinus* Förster, 1871; *Prosopis sinuata* var. *atrifacialis* Strand, 1909). Russia: **EP** (CR). – Europe (WE, SE, EE), Azerbaijan, Turkey.

74. FAMILY ANDRENIDAE

M.YU. PROSHCHALYKIN, YU.V. ASTAFUROVA
AND D.A. SIDOROV

The family Andrenidae is one of the largest bee families, colonizing all continents except Australia. They are also almost absent from the tropical Asian region. Similarly, in sub-Saharan Africa there are only a few genera and species. The Andrenidae are divided into four subfamilies, two of them are widely distributed, in the Palaearctic (Andreninae and Panurginae), and two Neotropical (Alocandreninae and Oxaeinae). Most species belong to the genus *Andrena*, which is ubiquitous in the north temperate areas. All species nest in the ground and most of them are strictly solitary, although they often nest in large aggregations (Michener, 2007; Osytshnjuk et al., 2005). Number of taxa: World – 77 genera, 2917 species; Palaearctic – 13/1100; Russia – 5/244.

R e f e r e n c e s. Lebedev, 1932; Popov, 1940, 1949, 1958b; Warncke, 1967; Osytshnjuk, 1977, 1982, 1983, 1984, 1986a,b, 1993, a,b, 1994a,b, 1995; Tadauchi, Xu, 1995, 1999, 2002, 2003; Xu, Tadauchi, 1999, 2005, 2006, 2009a,b, 2011, 2012; Xu et al., 2000; Gusenleitner, Schwarz, 2002; Davydova, Pesenko, 2002; Gusenleitner et al., 2005; Osytshnjuk et al., 2005, 2008; Schmid-Egger, 2005; Tadauchi, 2006, 2008; Proshchalykin, 2007b, 2009; 2010, 2012a; Shebl, Tadauchi, 2009; 2011; Scheuchl, 2010; Romankova, Astafurova, 2011; Astafurova, Romankova, 2012; Proshchalykin, Lelej, 2013; Shebl, Li, 2013; Proshchalykin, Astafurova, 2017; Aliyev et al., 2017; Ascher, Pickering, 2017; Kuhlmann et al., 2017; Levchenko, 2017; Sidorov, Proshchalykin, 2017a, b; Sidorov et al., 2017.

Subfamily ANDRENINAE

ANDRENA Fabricius, 1775. Type species *Apis helvola* Linnaeus, 1758. Genus is distributed throughout the Holarctic region, south in the Western Hemisphere to Panama, where one species occurs in the tropical lowlands, in Africa through the East African highlands and south to the Cape of Good Hope, and in Asia to the mountains of southern India and of Malaysia. Number of species: World – 1500; Palaearctic – 950; Russia – 225.

Andrena (Aciandrena) aciculata Morawitz, 1886 (*A. tenuis* sensu E. Stoeckhert, 1930, nom. praeocc., nec Morawitz, 1878; *A. tenuicula* E. Stoeckhert,

1950; *A. strymonia* Pittioni, 1950). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), Azerbaijan, Turkey, Iran.

Andrena (Aciandrena) chersona Warncke, 1972. Russia: **EP** (CR). – Europe (EE), Azerbaijan.

Andrena (Aciandrena) tenuis Morawitz, 1877. Russia: **EP** (S). – Europe (EE), Georgia, Azerbaijan, Turkey, Iran, Turkmenistan.

Andrena (Aciandrena) volgensis Osytshnjuk, 1994. Russia: **EP** (S). – Kazakhstan.

Andrena (Aenandrena) aeneiventris Morawitz, 1872 (*A. locularoides* Strand, 1915; *A. aeneiventris* var. *punctata* Jaeger, 1934). Russia: **EP** (S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Jordan, Israel, Iraq, Iran, Turkmenistan, Tajikistan, Kazakhstan.

Andrena (Aenandrena) bisulcata Morawitz, 1877 (*A. echinulata* Pérez, 1903; *A. doderoi* Jaeger, 1934). Russia: **EP** (CR). – Europe (WE, SE, EE), Georgia, Azerbaijan, Turkey, Lebanon, Iran.

Andrena (Aenandrena) hedikae Jaeger, 1934 (*A. hedikae* ssp. *inexpectata* Warncke, 1975). Russia: **EP** (S). – Europe (WE, SE, EE), N Africa, Azerbaijan, Turkey, Iran.

Andrena (Aenandrena) hystrix Schmiedeknecht, 1883 (*A. opaciventris* Friese, 1921, nom. praeocc., nec Cockerell, 1916; *A. hystrix* var. *lateritia* Benoist, 1961; *A. h.* ssp. *rufilata* Warncke, 1975). Russia: **EP** (CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Turkmenistan.

Andrena (Andrena) aino Tadauchi, Hirashima et Matsumura, 1987. Russia: **ES** (BR, YA, ZB), **FE** (AM, KH, PR, SA, MG). – Japan (Hok).

Andrena (Andrena) apicata Smith, 1847 (*A. batava* Pérez, 1902; *A. apicata* var. *tristis* Alfken, 1904; *A. a.* var. *kamtschatica* Alfken, 1929). Russia: **EP** (NW, C), **UR**, **FE** (KA). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan.

Andrena (Andrena) benefica Hirashima, 1962. Russia: **FE** (KH, PR). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Andrena (Andrena) brevihirtiscopa Hirashima, 1962. Russia: **FE** (KH, PR). – Korean Peninsula, Japan (Hok, Hon, Kyu).

Andrena (Andrena) bulgariensis Warncke, 1965. Russia: **EP** (CR). – Europe (EE).

Andrena (Andrena) clarkella (Kirby, 1802) [Melitta]. Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (TM, NS, KM), **ES** (KR, BR, YA, ZB), **FE** (AM, KH, PR, SA, MG, CH). – Europe (WE, NE, SE, EE), N America.

Andrena (Andrena) fucata Smith, 1847 (*A. clypearis* Nylander, 1848; *A. norvegica* Strand, 1910). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (TK, KM, AL), **ES** (KS, TU, KR). – Europe (WE, NE, SE, EE), Armenia, Turkey, Kazakhstan.

Andrena (Andrena) helvola (Linnaeus, 1758) [Apis] (*Melitta subdentata* Kirby, 1802; ?*Andrena incerta* Eversmann, 1852; *A. helvola* var. *concolor* Blüthgen, 1919, nom. praeocc., nec Robertson, 1898, nec Alfken, 1914). Russia: **EP** (C, E, S), **UR**, **WS** (TK, NS, KM). – Europe (WE, NE, SE, EE), Armenia, Turkey, Kazakhstan, China (NE).

Andrena (Andrena) hondoica Hirashima, 1962. Russia: **FE** (KH, PR, SA). – Korean Peninsula, Japan (Hok, Hon).

Andrena (Andrena) ishiharai Hirashima, 1962. Russia: **ES** (BR), **FE** (PR). – Korean Peninsula, Japan (Hok, Hon, Kyu).

Andrena (Andrena) kamtschatkaensis Friese, 1914. Russia: **FE** (KA).

Andrena (Andrena) lapponica Zetterstedt, 1838 (*A. alpina* Zetterstedt, 1838, nom. praeocc., nec Morawitz, 1872; *A. shirozui* Hirashima, 1962; *A. lapponica* ssp. *sumizome* Tadauchi, Hirashima et Matsumura, 1987). Russia: **EP** (N, NW, C), **UR**, **WS** (TM, TK), **ES** (IR, BR, YA, ZB), **FE** (PR, SA, KU, KA, MG). – Europe (WE, NE, SE, EE), Georgia, Kyrgyzstan, Kazakhstan, Japan (Hok, Hon).

Andrena (Andrena) maukensis Matsumura, 1911. Russia: **WS** (KM), **ES** (IR, YA, ZB), **FE** (KH, PR, SA, KU, KA). – Japan (Hok, Hon, Kyu).

Andrena (Andrena) mitis Schmiedeknecht, 1883 (*A. palumboi* Destefani, 1889). Russia: **EP** (C, CR). – Europe (WE, SE, EE), Azerbaijan, Turkey.

Andrena (Andrena) nawai Cockerell, 1913. Russia: **FE** (KH, PR). – China (NC), Japan (Hok, Hon, Kyu).

Andrena (Andrena) nycthemera Imhoff, 1868 (*A. nycthemera* var. *tergestensis* Alfken, 1904). Russia: **EP** (C, E), **UR**. – Europe (WE, NE, SE, EE), Kazakhstan.

Andrena (Andrena) praecox (Scopoli, 1763) [Apis] (*Melitta smithella* Kirby, 1802; *Andrena flavescens* Schenck, 1853; *A. clypeata* Smith, 1855, nom. praeocc., nec Illiger, 1806, nec Brullé, 1832, nec Smith, 1853, nec Schenck, 1853; *A. torkai* Alfken, 1913). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TK, NS, KM), **ES** (IR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Kazakhstan.

Andrena (Andrena) sakagamii Tadauchi, Hirashima et Matsumura, 1987. Russia: **ES** (BR), **FE** (AM, KH, SA). – Korean Peninsula, Japan (Hok, Hon).

Andrena (Andrena) varians (Kirby, 1802) [Melitta] (*Melitta angulosa* Kirby, 1802; *Andrena mesoxantha* Imhoff, 1834; *A. longula* Eversmann, 1852; *A. mixta* Schenck, 1853; *A. rectangula* Schenck, 1853; *A. varians* var. *conjungens* Blüthgen, 1916). Russia: **EP** (N, NW, C, E, S, CR), **UR**. – Europe (WE, NE, SE, EE), Kazakhstan.

Andrena (Brachyandrena) colletiformis Morawitz, 1874 (*Colletes parvula* Morawitz, 1871, nom. praeocc., nec Kirby, 1802; *Andrena dissidens* Schmiedeknecht, 1884; *A. colletiformis* ab. *gandzhensis* Lebedev, 1933; *A. c.* ssp. *insulana* Pittioni, 1950; *A. c.* var. *sardinica* Strand, 1915; *A. nanana* Strand, 1915; *A. subsquamularis* Noskiewicz, 1960). Russia: **EP** (S, NC, CR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran, Turkmenistan, Kazakhstan.

Andrena (Brachyandrena) limonii Osytshnjuk, 1983. Russia: **EP** (S). – Kazakhstan.

Andrena (Calomelissa) tsukubana Hirashima, 1957. Russia: **FE** (AM, KH, PR). – Korean Peninsula, Japan (Hon, Shi, Kyu).

Andrena (Campylogaster) erberi Morawitz, 1871 (*Campylogaster fulvo-crustatus* Dours, 1873; *Andrena amasia* Radoszkowski, 1876; *A. squamigera* Bramson, 1879, nom. praeocc., nec Schenck, 1874; ?*A. kozarowi* Nedalkov, 1914; *A. erberi* var. *sanguiniventris* Friese, 1921; *A. e.* ssp. *migrans* Warncke, 1967). Russia: **EP** (CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Syria, Iran.

Andrena (Campylogaster) incisa Eversmann, 1852 (*A. gracilis* Eversmann, 1852, nom. praeocc., nec Schenck, 1869; *A. punctatissima* Morawitz, 1866, nom. praeocc., nec Kriechbaumer, 1873). Russia: **EP** (S, CR), **UR**, **WS** (AL), **ES** (KS, KR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Tajikistan, Kyrgyzstan, Kazakhstan.

Andrena (Campylogaster) nova Popov, 1940. Russia: **ES** (BR, ZB), **FE** (AM, PR).

Andrena (Campylogaster) phaneroleuca Cockerell, 1929. Russia: **ES** (IR).

Andrena (Carandrena) semiflava Lebedev, 1932. Russia: **EP** (S, NC). – Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Andrena (Charitandrena) hattorfiana (Fabricius, 1775) [Nomada] (*Apis splendida* Müller, 1776; *A. marginata* Müller, 1776, nom. praeocc., nec Fabricius, 1776, nec Torka, 1913; *Andrena rubida* Olivier, 1789; *Apis fuscicornis* Gmelin, 1790; *A. plumosa* Christ, 1791, nom. praeocc., nec Kim et Kim, 1989,

nec Christ, 1791; *Andrena equestris* Panzer, 1797; *Apis flororum* Cuvier, 1798; *Melitta lathamana* Kirby, 1802; *Andrena quadripunctata* Fabricius, 1804; *A. clypeata* Illiger, 1806, nom. praeocc., nec Brullé, 1832, nec Smith, 1853, nec Smith, 1855; *A. labialis* Gravenhorst, 1807, nom. praeocc., nec Kirby, 1802; *A. dimidiata* Brullé, 1832; *A. elongata* Imhoff, 1834, nom. praeocc., nec Radoszkowski, 1871, nec Radoszkowski, 1876; *A. gravida* Eversmann, 1852, nom. praeocc., nec Imhoff, 1832, nec Dours, 1872; *A. clypeata* Schenck, 1853, nom. praeocc., nec Illiger, 1806, nec Brullé, 1832, nec Smith, 1853, nec Smith, 1855; *A. fischeri* Verhoeff, 1890; *A. hattorfiana* var. *rufa* Torka, 1913; *A. h.* var. *punctulata* Torka, 1913, nom. praeocc., nec Schenck, 1853; *A. h.* var. *marginalata* Torka, 1913, nom. praeocc., nec Müller, 1776, nec Fabricius, 1776; *A. haemorrhoidula* Viereck, 1916; *A. hattorfiana* ab. *deandrenalis* Strand, 1922). Russia: EP (NW, C, E, S, NC, CR), UR. – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey.

Andrena (Chlorandrena) cinereophila Warncke, 1965. Russia: EP (NC). – Europe (SE, EE), Turkey, Israel, Afghanistan.

Andrena (Chlorandrena) humilis Imhoff, 1832 (*A. fulvescens* Smith, 1847; *A. cinerascens* Nylander, 1848, nom. praeocc., nec Eversmann, 1852; *A. scabrosa* Morawitz, 1866; *A. nasalis* Thomson, 1870; *A. nudigastra* Alfken, 1914; ?*A. nudigastra* ssp. *nudigastra stroides* Yasumatsu, 1935; *A. humilis* ssp. *ardeola* Warncke, 1975; *A. h.* ssp. *cucullata* Warncke, 1975; *A. h.* ssp. *hiaticula* Warncke, 1975; *A. h.* ssp. *indigena* Warncke, 1975). Russia: EP (N, NW, C, E, S, NC, CR), UR, WS (TK, KM, AL), FE (PR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Iran, Kyrgyzstan, Kazakhstan.

Andrena (Chlorandrena) panurgimorpha Mavromoustakis, 1957. Russia: EP (CR). – Europe (WE, SE, EE), Georgia, Turkey, Israel, Iran.

Andrena (Chlorandrena) taraxaci Giraud, 1861 (*A. curtivalvis* Morice, 1899; *A. truncata* Pérez, 1903, nom. praeocc., nec Viereck, 1903; *A. humilis* var. *orienticola* Strand, 1915; *A. truncatiformis* Cockerell, 1922; *A. taraxaci* ssp. *chikuzenensis* Hirashima, 1957). Russia: EP (S, NC, CR), UR, FE (PR). – Europe (WE, SE, EE), N Africa, Georgia, Turkey, Lebanon, Israel, Iran, Turkmenistan, Tajikistan, Kazakhstan, China (NE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Andrena (Chlorandrena) tricuspidata Scheuchl, 2010. Russia: EP (CR).

Andrena (Chrysandrena) colonialis Morawitz, 1886. Russia: EP (S), WS (NS, AL). – Europe (EE), Azerbaijan.

Andrena (Chrysandrena) fulvago (Christ, 1791) [Apis] (*A. constrictus* Smith, 1849). Russia: EP (C, E, CR), UR, WS (KM, AL), ES (BR). – Europe (WE, NE, SE, EE), Georgia, Armenia, Turkey.

Andrena (Chrysandrena) hesperia Smith, 1853 (*A. opacicollis* Morawitz, 1886; *A. marcescens* Pérez, 1895; *A. freya* Strand, 1915; *A. asunica* Strand, 1921). Russia: EP (CR). – Europe (WE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Israel, Iran, Turkmenistan, Tajikistan.

Andrena (Chrysandrena) khankensis Osytshnjuk, 1995. Russia: FE (AM, PR), ES (ZB).

Andrena (Cnemidandrena) albicaudata Hirashima, 1966. Russia: FE (PR). – Mongolia, Japan (Hon).

Andrena (Cnemidandrena) denticulata (Kirby, 1802) [Melitta] (*Melitta listerella* Kirby, 1802; *Andrena edentata* Friese, 1922; *A. archanensis* Cockerell, 1929; *A. baicaliella* Cockerell, 1929; *A. epimelaena* Cockerell, 1924; *A. seneciorum* Hirashima, 1964). Russia: EP (NW, C, E), UR, WS (TK, KM), ES (IR, ZB), FE (AM, KH, PR, SA, KU). – Europe (WE, NE, SE, EE), Georgia, Kazakhstan, Mongolia, China (NE, NC, NW), Korean Peninsula, Japan (Hok, Hon, Shi).

Andrena (Cnemidandrena) fuscipes (Kirby, 1802) [Melitta] (*Melitta pubescens* Kirby, 1802, nom. praeocc., nec Olivier, 1789; *Andrena cincta* Nylander, 1848, nom. praeocc., nec Fabricius, 1781; *A. germanica* Verhoeff, 1890). Russia: EP (NW, C, E), UR. – Europe (WE, NE, SE, EE), Kazakhstan.

Andrena (Cnemidandrena) maetai Hirashima, 1964. Russia: FE (PR). – Korean Peninsula, Japan (Kyu, Tsu).

Andrena (Cnemidandrena) nigriceps (Kirby, 1802) [Melitta] (*Melitta lanifrons* Kirby, 1802; *Andrena jagnobensis* Morawitz, 1894; *A. nigriceps* var. *hispanica* Friese, 1922, nom. praeocc., nec Alfken, 1927; *A. n.* ssp. *pamirensis* Alfken, 1931; *A. marchica* Alfken, 1939; *A. nigriceps* ssp. *comata* Warncke, 1974). Russia: EP (NW, C, E), UR, ES (YA). – Europe (WE, NE, SE, EE), Armenia, Azerbaijan, Turkey, Tajikistan, Uzbekistan.

Andrena (Cnemidandrena) rufoclypeata Alfken, 1936. Russia: WS (AL), ES (TU). – Mongolia, China (NW).

Andrena (Cnemidandrena) simillima Smith, 1851 (*Anthrena* (!) *baicalica* Kokujev, 1927; *Andrena*

- bremensis* Alfken, 1900; *A. simillima* ssp. *sischkai* Warncke, 1988). Russia: **EP** (NW, C, E, S), **UR**, **WS** (TK, KM, AL), **ES** (KS, BR, YA). – Europe (WE, NE, SE, EE), Azerbaijan, Iran, Mongolia.
- Andrena (Cnemidandrena) tridentata** Kirby, 1802
[*Melitta*] (*Melitta rufitarsis* Kirby, 1802, nom. praeocc., nec Zetterstedt, 1838). Russia: **EP** (NC), **UR**, **WS** (NS, KM, AL). – Europe (WE, SE, EE).
- Andrena (Cordandrena) cordialis** Morawitz, 1877.
Russia: **EP** (S, NC, CR), **UR**, **ES** (ZB). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Central Asia, Kazakhstan, China (NW).
- Andrena (Cryptandrena) ventricosa** Dours, 1873
(*A. ventricosa* ssp. *ridibundus* Warncke, 1975). Russia: **EP** (NC). – Europe (WE, SE, EE), N Africa, Armenia, Azerbaijan, Turkey, Israel.
- Andrena (Didonia) stepposa** Osytshnjuk, 1977. Russia: **EP** (C). – Europe (EE).
- Andrena (Euandrena) allosa** Warncke, 1975 (*A. allosa* ssp. *canigica* Warncke, 1975; *A. a.* ssp. *pileata* Warncke, 1975). Russia: **EP** (S, NC), **UR**. – Europe (SE), Caucasus, Kazakhstan.
- Andrena (Euandrena) asperula** Osytshnjuk, 1977.
Russia: **EP** (S, CR). – Europe (EE), Azerbaijan.
- Andrena (Euandrena) bicolor** Fabricius, 1775 (*Apis icterica* Christ, 1791; *A. fuliginosa* Christ, 1791, nom. praeocc., nec Scopoli, 1770, nec Panzer, 1798; *Andrena lucida* Panzer, 1798, nom. praeocc., nec Lepeletier de Saint-Fargeau, 1841; *Melitta picicornis* Kirby, 1802; *M. pilosula* Kirby, 1802; *M. gwynana* Kirby, 1802; *Apis ephippiata* Schrank, 1802; *Andrena oraniensis* Lepeletier de Saint-Fargeau, 1841; *A. proxima* Smith, 1847, nom. praeocc., nec Kirby, 1802; *A. aestiva* Smith, 1849; *A. consimilis* Smith, 1849, nom. praeocc., nec Smith, 1847, nec Alfken, 1900; *A. nigrifrons* Eversmann, 1852, nom. praeocc., nec Smith, 1853, nec Smith, 1855, nec Cresson, 1878; *A. laeviuscula* Schenck, 1853; *A. fuscohirta* Schenck, 1853; *A. distinguenda* Schenck, 1853, nom. praeocc., nec Schenck, 1871; *A. marginalis* Schenck, 1861; *A. violascens* Thomson, 1870; *A. beuthini* Schenck, 1876; *A. gwynana* var. *testacea* Dalla Torre, 1877, nom. praeocc., nec Warncke, 1975; *A. croatica* Friese, 1887; *A. lavandulae* Pérez, 1902; *A. nigrosterna* Pérez, 1902; *A. nigriventris* Pérez, 1902, nom. praeocc., nec Gmelin, 1790, nec Saunders, 1908; *A. fervida* Pérez, 1902; *A. heterodoxa* Pérez, 1903, nom. praeocc., nec Cockerell, 1930; *A. phaneromelas* Cockerell, 1929; *A. baleina* Cockerell, 1929; *A. bicolor* ssp. *agraria* Warncke, 1974; *A. b.* ssp. *apricaria* Warncke, 1975). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (TK, KM, AL), **ES** (KS, TU, IR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran, Mongolia, China (NW).
- Andrena (Euandrena) capillosa** Morawitz, 1876. Russia: **UR**. – Uzbekistan, Kazakhstan.
- Andrena (Euandrena) chrysopus** Pérez, 1903 (*A. dobowlanensis* Noskiewicz, 1923). Russia: **EP** (C, E, S, CR). – Europe (WE, SE, EE), Turkey, Kazakhstan.
- Andrena (Euandrena) fulvida** Schenck, 1853 (*A. fasciatella* Schenck, 1853; *A. jasnitzkii* Cockerell, 1929). Russia: **EP** (NW, C), **WS** (TK, KM, AL), **ES** (KR, IR, BR, YA), **FE** (AM, KH, PR, SA). – Europe (WE, NE, SE, EE).
- Andrena (Euandrena) glabriventris** Alfken, 1935.
Russia: **EP** (CR). – Caucasus, Turkey.
- Andrena (Euandrena) hebes** Pérez, 1905 (*A. praecociformis* Cockerell, 1911). Russia: **FE** (SA). – Japan (Hon, Shi, Kyu).
- Andrena (Euandrena) khabarovi** Osytshnjuk, 1986.
Russia: **FE** (AM, KH, PR).
- Andrena (Euandrena) kudiana** Cockerell, 1924. Russia: **FE** (PR).
- Andrena (Euandrena) mutini** Osytshnjuk, 1986. Russia: **FE** (AM, KH, PR).
- Andrena (Euandrena) orientaliella** Osytshnjuk, 1986.
Russia: **ES** (TU, IR, BR, YA, ZB), **FE** (AM, KH, PR, MG). – Mongolia.
- Andrena (Euandrena) roscipes** Alfken, 1933. Russia: **EP** (S, NC). – Europe (EE).
- Andrena (Euandrena) rudolfae** Osytshnjuk, 1986.
Russia: **UR**, **ES** (KS, BR, ZB). – Kazakhstan.
- Andrena (Euandrena) ruficrus** Nylander, 1848. Russia: **EP** (N, NW, C, E), **UR**, **WS** (TK, KM), **ES** (IR, BR, YA, ZB), **FE** (AM, KH, PR, SA). – Europe (WE, NE, SE, EE), Turkey, Kyrgyzstan, Kazakhstan, Korean Peninsula, Japan (Hok, Hon).
- Andrena (Euandrena) symphyti** Schmiedeknecht, 1883 (*A. bothriorhina* Pérez, 1902; *A. furcata* Friese, 1921; *A. furcata* var. *amanensis* Friese, 1921). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), Georgia, Azerbaijan, Turkey, Iran.
- Andrena (Euandrena) verae** Osytshnjuk, 1986 (*A. verae* ssp. *nikiforuki* Osytshnjuk, 1986). Russia: **EP** (S), **UR**. – Europe (EE), Kazakhstan.
- Andrena (Euandrena) vulpecula** Kriechbaumer, 1873.
Russia: **FE** (AM, PR). – Europe (WE, SE, EE), N Africa.
- Andrena (Graecandrena) hyemala** Warncke, 1973
(*A. hyemala* ssp. *represa* Warncke, 1975; *A. h.* ssp.

- kushkia* Osytshnjuk, 1994). Russia: **EP** (S, NC). – Europe (SE, EE), Turkey, Israel, Turkmenistan.
- Andrena (Graecandrena) impunctata** Pérez, 1895
(*A. contusa* Pérez, 1895; *A. paula* Noskiewicz, 1939). Russia: **EP** (S, CR). – Europe (WE, SE, EE), N Africa, Azerbaijan, Turkey, Israel, Iran.
- Andrena (Graecandrena) schwarzi** Warncke, 1975
(*A. schwarzi* ssp. *walishanovi* Osytshnjuk, 1994). Russia: **EP** (S, CR). – Europe (EE), Turkey, Kazakhstan.
- Andrena (Holandrena) decipiens** Schenck, 1861 (*Apis albifrons* Christ, 1791, nom. praeocc., nec Förster, 1771, nec Rossi, 1790; *Andrena strigosa* Dours, 1872, nom. praeocc., nec Panzer, 1799; *A. insolita* Dours, 1872; ?*A. flavilabris* Schenck, 1874; *A. deceptoria* Schmiedeknecht, 1883; *A. fimbriata paganettii* Alfken, 1904, nom. praeocc., nec Strand, 1921, nec Maidl, 1922; *A. decipiens* ssp. *capta* Warncke, 1974). Russia: **EP** (S, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Iran.
- Andrena (Holandrena) labialis** (Kirby, 1802) [Melitta] (*A. succinata* Imhoff, 1832; *A. separata* Smith, 1847; *A. albipes* Schenck, 1853; *A. fulvipes* Schenck, 1853; *A. lutescens* Schenck, 1853; *A. trachyderma* Dours, 1872; *A. nigro-stincta* Dours, 1872; *A. succinctula* Viereck, 1916; *A. labialis* ssp. *megala* Warncke, 1975; *A. l. ssp. stenura* Warncke, 1975). Russia: **EP** (NW, C, E, S, CR), **UR**, **WS** (OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Syria, Iran, Afghanistan, Kyrgyzstan, Tajikistan, Uzbekistan, Kazakhstan.
- Andrena (Holandrena) labiatula** Osytshnjuk, 1993. Russia: **EP** (CR).
- Andrena (Holandrena) valeriana** Hirashima, 1957. Russia: **ES** (BR), **FE** (AM, KH, PR). – Mongolia, China (NC, NW, WP), Korean Peninsula, Japan (Hok, Hon, Kyu).
- Andrena (Holandrena) variabilis** Smith, 1853 (?*A. kirbyi* Curtis, 1826; *A. calabra* Costa, 1863; *A. fons-colombii* Dours, 1872; *A. picciolii* Dours, 1872; *A. piceicornis* Dours, 1872; *A. pulcherrima* Schmiedeknecht, 1884; *A. macedonica* Strand, 1919, nom. praeocc., nec Friese, 1923). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), N Africa, Azerbaijan, Turkey, Lebanon, Israel, Iran, Turkmenistan, Kazakhstan.
- Andrena (Hoplodrena) carantonica** Pérez, 1902
(?*Apis sabulosa* Scopoli, 1763; ?*A. degener* Scopoli, 1770; ?*A. acris* Harris, 1776; ?*A. lutulenta* Gmelin, 1790; *Andrena carantonica* Pérez, 1902; *A. trimmerana* var. *scotica* Perkins, 1917; *A. jacobi* Perkins, 1921; *A. jacobi* var. *johsoni* Perkins, 1921). Russia: **EP** (N, NW, C, E, CR), **UR**. – Europe (WE, NE, SE, EE), Armenia, Azerbaijan, Iran.
- Andrena (Hoplodrena) dentata** Smith, 1879. Russia: **WS** (TK, KM), **FE** (AM, KH, PR, KU). – China (NE, NC, SW, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Andrena (Hoplodrena) ferox** Smith, 1847 (*A. ferox* Smith, 1847; *A. distincta* Smith, 1847, nom. praeocc., nec Lucas, 1849, nec Schenck, 1861; *A. obscurata* Smith, 1853). Russia: **EP** (CR). – Europe (WE, SE, EE), Armenia, Turkey.
- Andrena (Hoplodrena) miyamotoi** Hirashima, 1964. Russia: **FE** (PR, KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu).
- Andrena (Hoplodrena) romankovae** Osytshnjuk, 1995. Russia: **ES** (BR), **FE** (PR).
- Andrena (Hoplodrena) rosae** Panzer, 1801 (*Melitta zonalis* Kirby, 1802; *Andrena postica* Imhoff, 1832; *A. coarctata* Imhoff, 1832, nom. praeocc., nec Pérez, 1902; *A. austriaca* var. *incompta* Schmiedeknecht, 1883; *A. rosae* var. *alfkeni* Friese, 1914; *A. florea* ssp. *sachalinensis* Yasumatsu, 1939; *A. austriaca* auct., nec Panzer, 1798). Russia: **EP** (NW, C, E, S, CR), **UR**, **WS** (TK, NS, KM, AL), **ES** (KS, TU, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU). – Europe (WE, NE, SE, EE), Armenia, Azerbaijan, Turkey, Iran, Kyrgyzstan, Kazakhstan, Mongolia, China (NE), Korean Peninsula, Japan (Hok).
- Andrena (Hoplodrena) trimmerana** (Kirby, 1802) [Melitta] (*Melitta spinigera* Kirby, 1802; *Andrena fusca* Lepeletier de Saint-Fargeau, 1841; *A. arietina* Dours, 1872; *A. ustulata* Dours, 1872; *A. lombardica* Schmiedeknecht, 1883; *A. dragana* Friese, 1887; *A. anglica* Alfken, 1911). Russia: **EP** (C, NC, CR), **UR**, **ES** (KR). – Europe (WE, NE, SE, EE), N Africa, Turkey, Turkmenistan,
- Andrena (Larandrena) sericata** Imhoff, 1868 (*A. favosa* Morawitz, 1872). Russia: **EP** (E, S, NC, CR), **UR**. – Europe (WE, SE, EE), Georgia, Armenia, Turkey, Turkmenistan, Kazakhstan.
- Andrena (Larandrena) ventralis** Imhoff, 1832 (?*Apis tenella* Christ, 1791; ?*Andrena analis* Fabricius, 1804; *A. ventralis* Imhoff, 1832; *A. rufiventris* Eversmann, 1852; *A. mutabilis* Morawitz, 1866, nom. praeocc., nec Pérez, 1895; *A. fukuiana* Hirashima et Haneda, 1973). Russia: **EP** (N, C, E, S, CR), **UR**, **WS** (OM, TK, KM), **ES** (IR, BR). – Europe (WE, SE, EE), Caucasus, Turkey, Central Asia, Kazakhstan, China (NC), Korean Peninsula, Japan (Hon, Shi).

Andrena (Lepidandrena) caprimulga Warncke, 1975.
Russia: EP (CR). – Azerbaijan, Turkey.

Andrena (Lepidandrena) curvungula Thomson, 1870
(*A. pictipes* Giraud, 1861; *A. squamigera* Schenck, 1874, nom. praeocc., nec Bramson, 1879). Russia: EP (C, S, CR), UR, WS (OM, TK, KM, AL). – Europe (WE, SE, EE), Georgia, Azerbaijan, Turkey, Kazakhstan.

Andrena (Lepidandrena) florivaga Eversmann, 1852
(*A. molhusina* Blüthgen, 1914). Russia: EP (C, S, NC, CR), UR, WS (NS, KM). – Europe (WE, NE, SE, EE), Turkey, Iran, Kazakhstan.

Andrena (Lepidandrena) paucisquama Noskiewicz, 1924 (*A. paucisquama* ssp. *curonica* Warncke, 1975). Russia: EP (C, NC, CR), UR. – Europe (WE, SE, EE), Caucasus, Turkey.

Andrena (Lepidandrena) rufizona Imhoff, 1834
(*A. alpina* Morawitz, 1872, nom. praeocc., nec Zetterstedt, 1838; *A. pretiosa* Schmiedeknecht, 1880, nom. praeocc., nec Morawitz, 1894). Russia: EP (C, E, NC), UR, WS (TK, NS, KM, AL), ES (KS, KR). – Europe (WE, SE, EE), Caucasus, Kazakhstan.

Andrena (Leucandrena) argentata Smith, 1844. Russia: EP (NW, C, E), UR, WS (TK, NS, AL), ES (KS, BR), FE (AM, PR). – Europe (WE, NE, SE, EE), Georgia, Turkey, Kazakhstan.

Andrena (Leucandrena) barbilabris (Kirby, 1802)
[*Melitta*] (*Apis sericea* Christ, 1791, nom. praeocc., nec Förster, 1771; *Hylaeus similis* Fabricius, 1793, nom. praeocc., nec Smith, 1849; *Melitta albicrus* Kirby, 1802; *Andrena albocincta* Zetterstedt, 1838; *A. barbatula* Zetterstedt, 1838, nom. praeocc., nec Kirby, 1802; *A. parumpunctata* Schenck, 1853; *A. placida* Smith, 1853; *A. angustipes* Schenck, 1866; *A. ciliata* Schenck, 1869, nom. praeocc., nec Gmelin, 1790; *A. macilenta* Provancher, 1888, nom. praeocc., nec Pérez, 1895; *A. macgillivrayi* Cockerell, 1897; *A. salicacea* Robertson, 1900; *A. sapellonis* Cockerell, 1900; *A. pauperata* Pérez, 1902; *A. ciliatula* Viereck, 1916; *A. sericea* var. *malaisei* Alfken, 1929). Russia: EP (NW, C, E, S), UR, WS (TM, TK, AL), ES (BR, YA, ZB), FE (AM, PR, KA, MG). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan, Mongolia, China (NE, NC), Japan (Hon), N America.

Andrena (Leucandrena) parviceps Kriechbaumer, 1873 (*A. parviceps* var. *basalis* Kriechbaumer, 1873, nom. praeocc., nec Dours, 1872; *A. lederi* Morawitz, 1886; *A. basilaris* Viereck, 1918). Russia: EP (C, CR). – Europe (WE, SE, EE), Georgia, Azerbaijan, Turkey, Iran, Turkmenistan.

Andrena (Margandrena) marginata Fabricius, 1776 (?*Apis obscura* Linné, 1764, nom. praeocc., nec Müller, 1776, nec Robertson, 1902, nec Friese, 1914; *Nomada cingulata* Fabricius, 1775; *Apis cetii* Schrank, 1781; *Andrena suecica* Gmelin, 1790; *Apis marginella* Gmelin, 1790; *A. sphegoides* Panzer, 1798, nom. praeocc., nec Schrank, 1781; *Melitta schrankella* Kirby, 1802; *M. affinis* Kirby, 1802; *Andrena frontalis* Smith, 1849; *A. fulva* Eversmann, 1852, nom. praeocc., nec Müller, 1766, nec Schrank, 1781; *A. hyalinella* Costa, 1888; *A. marginata* var. *nigrescens* Aurivillius, 1903; *A. m.* var. *absoluta* Friese, 1914; *A. eversmanniana* Osytshnjuk, 1994). Russia: EP (NW, C, E, S, NC), UR, WS (TM, KM), ES (KS, KR, IR, BR, ZB), FE (AM, PR). – Europe (WE, NE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Kazakhstan, Turkmenistan.

Andrena (Melandrena) albopunctata (Rossi, 1792)
[*Apis*] (*A. funebris* Panzer, 1798; *A. lugubris* Lepeletier de Saint-Fargeau, 1841, nom. praeocc., nec Erichson, 1841; *A. nigrobarbata* Morawitz, 1871; *A. macularis* Kriechbaumer, 1873; *A. lugubrescens* Cockerell, 1917; *A. albopunctata* ssp. *melona* Warncke, 1967). Russia: EP (E, S, NC, CR), UR. – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iraq, Iran, Afghanistan, Pakistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Andrena (Melandrena) comta Eversmann, 1852
(*A. orenburgensis* Schmiedeknecht, 1884; *Anthrena* (!) *csikiana* Mocsáry, 1901; *Andrena wulungshanensis* Yasumatsu, 1935). Russia: EP (C), UR, WS (NS, KM, AL), ES (KS, TU, IR, BR, ZB), FE (AM, PR). – Europe (WE, EE), Kazakhstan, Mongolia, China (NE, NC, NW), Japan (Hon).

Andrena (Melandrena) cineraria (Linnaeus, 1758)
[*Apis*] (*Apis atra* Scopoli, 1763, nom. praeocc., nec Müller, 1776, nec Smith, 1847, nec Popov, 1940; *Lamprocolletes peregrinus* Smith, 1878; *Andrena cineraria* var. *nigrifacies* Alfken, 1913; *A. danuvia* E. Stoeckhert, 1950; *A. barbareae* auct., partim, nec Panzer, 1805). Russia: EP (N, NW, C, E), UR, WS (OM, TK, NS, KM, AL), ES (KS, TU, IR, BR, YA, ZB), FE (KH). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran, Afghanistan, Pakistan, Turkmenistan, Kazakhstan, Mongolia, China (NC, NW, WP).

Andrena (Melandrena) cussariensis Morawitz, 1886
(*A. cussariensis* var. *kohatensis* Cockerell, 1917). Russia: EP (S, NC, CR), UR. – Georgia, Azerbaijan, Turkey, Iran, Tajikistan, Kazakhstan, Mongolia.

- Andrena (Melandrena) gallica** Schmiedeknecht, 1883
(A. gallica ssp. *alpicola* Bischoff, 1922; *A. g.* ssp. *fulvitigularis* Bischoff, 1922; *A. g.* ssp. *taurica* Bischoff, 1922; *A. g.* ssp. *iberica* Bischoff, 1922). Russia: **EP** (C, NC, CR), **UR**, **WS** (OM, KM, AL), **ES** (KS, IR, BR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Kazakhstan.
- Andrena (Melandrena) gussakovskii** Lebedev, 1932.
 Russia: **WS** (AL). – Iran, Central Asia.
- Andrena (Melandrena) limata** Smith, 1853 (*A. lucida* Lepeletier de Saint-Fargeau, 1841, nom. praeocc., nec Panzer, 1798; *A. pectoralis* Schmiedeknecht, 1883; *A. batesiae* Cockerell, 1910; *A. bateiae* Cockerell, 1914; *A. nitida* var. *algeriensis* Alfken, 1914; *A. thoracica* var. *secunda* Friese, 1922; *A. nitida* var. *aestivialis* Friese, 1922; *A. pectoralis* var. *alemannia* Alfken, 1927; *A. nitida* var. *hispanica* Alfken, 1927, nom. praeocc., nec Friese, 1922; *A. nitida* ssp. *hispaniola* Warncke, 1967; *A. limata* ssp. *mixtura* Warncke, 1967). Russia: **EP** (C, E, S, CR). – Europe (WE, NE, SE, EE), N Africa, Georgia, Armenia, Turkey, Israel, Iran, Uzbekistan, Kyrgyzstan, Kazakhstan.
- Andrena (Melandrena) magna** Warncke, 1965. Russia: **EP** (CR). – Europe (SE, EE), Georgia, Armenia, Turkey.
- Andrena (Melandrena) metallescens** Cockerell, 1906
(A. metallica Radoszkowski, 1876, nom. praeocc., nec Fabricius, 1793). Russia: **EP** (S). – Azerbaijan, Kazakhstan.
- Andrena (Melandrena) morio** Brullé, 1832 (*A. lugubris* Erichson, 1841; *A. holomelana* Lepeletier de Saint-Fargeau, 1841; *A. collaris* Lepeletier de Saint-Fargeau, 1841; *A. violaceipennis* Dufour, 1841, nom. praeocc., nec Cameron, 1902; ?*A. phaeoptera* Gistel, 1857; *A. rhodia* Kriechbaumer, 1873; *A. occipitalis* Pérez, 1895, nom. praeocc., nec Gistel, 1857; *A. asterabadiæ* Strand, 1921; *A. ephippium* var. *scopipes* Friese, 1923, nom. praeocc., nec Friese, 1923; *A. e.* var. *albicollis* Friese, 1923; *A. morio* var. *scopipes* Friese, 1923, nom. praeocc., nec Friese, 1923; *A. m.* ssp. *athalassae* Pittioni, 1950). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Israel, Iran, Kazakhstan.
- Andrena (Melandrena) nigroaenea** (Kirby, 1802)
 [Melitta] (*Apis obscura* Müller, 1776, nom. praeocc., nec Linné, 1764, Robertson, 1902, nec Friese, 1914; *Andrena splendens* Imhoff, 1832; *A. bipartita* Lepeletier de Saint-Fargeau, 1841, nom. praeocc., nec Brullé, 1839; *A. aprilina* Smith, 1848;

- A. nigrosericea* Dours, 1872; *A. serotina* Destefani, 1889, nom. praeocc., nec Robertson, 1893; *A. lepeltieri* Dalla Torre, 1896, nom. praeocc., nec Lucas, 1849; *A. candiae* Strand, 1915; *A. nigroaenea* var. *fasciculata* Friese, 1922; *A. melandura* Cockerell, 1922; *A. aemula* Alfken, 1926, nom. praeocc., nec Alfken, 1926; *A. nigroaenea sarnia* Richards, 1979; *A. n. corsa* Tkalcú, 1984). Russia: **EP** (NC, CR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel, Syria, Iran, Afghanistan.
- Andrena (Melandrena) nitida** (Müller, 1776) [*Apis*] (?*Apis assiduus* Harris, 1776; ?*A. fortis* Harris, 1776; *Andrena pubescens* Olivier, 1789, nom. praeocc., nec Kirby, 1802; *Apis nigriventris* Gmelin, 1790, nom. praeocc., nec Pérez, 1902, nom. praeocc., nec Saunders, 1908; *Andrena velutina* Lepeletier de Saint-Fargeau, 1841; *A. consimilis* Smith, 1847, nom. praeocc., nec Smith, 1849, nec Alfken, 1900; *A. nitens* Schenck, 1869; *A. pectoraloides* Verhoeff, 1890; *A. denticornis* Verhoeff, 1890; *A. nitida baltica* Alfken, 1913; *A. pubescens* ssp. *omissa* Valkeila, 1954). Russia: **EP** (NW, C, NC, CR), **UR**, **WS** (TK, KM). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Iran, Central Asia, Kazakhstan.
- Andrena (Melandrena) parathoracica** Hirashima, 1957. Russia: **FE** (PR, KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Andrena (Melandrena) pyropygia** Kriechbaumer, 1873. Russia: **EP** (NC, CR). – Europe (SE), Georgia, Azerbaijan, Turkey, Syria, Israel, Iran.
- Andrena (Melandrena) senex** Eversmann, 1852
(A. ovinella Friese, 1914). Russia: **UR**, **ES** (TU, IR, BR). – Kazakhstan, Mongolia, China (NE).
- Andrena (Melandrena) stigmatica** Morawitz, 1895. Russia: **EP** (CR). – Turkmenistan.
- Andrena (Melandrena) thoracica** (Fabricius, 1775)
 [*Apis*] (*Melitta melanocephala* Kirby, 1802; *Andrena sinensis* Cockerell, 1910; *A. thoracica* ssp. *melanoptera* Hedicke, 1934; *A. th.* ssp. *kotschy* Mavromoustakis, 1953). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (OM, TK, NS, KM, AL), **ES** (KS, TU, IR, BR, ZB), **FE** (AM, KH, PR, SA, KA, MG). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel, Jordan, Lebanon, Iran, Afghanistan, Turkmenistan, Tajikistan, Uzbekistan, Mongolia, China (NE, NC, NW), Korean Peninsula.
- Andrena (Melandrena) vaga** Panzer, 1799 (*Apis pratensis* Müller, 1776; *Andrena ovina* Klug, 1810; *A. nitidiventris* Blanchard, 1840, nom. praeocc., nec Dufour, 1841; *A. nitidiventris* Dufour, 1841, nom.

- praeocc., nec Blanchard, 1840; *A. atricula* Bischhoff, 1922). Russia: **EP** (NW, C, E, S), **UR**, **WS** (OM, TM, KM), **ES** (IR, BU, YA, ZB). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Kyrgyzstan, Kazakhstan.
- Andrena (Melandrena) watasei** Cockerell, 1913 (*A. azalearum* Cockerell, 1930). Russia: **FE** (PR, SA). – Japan (Hok, Hon, Shi, Kyu).
- Andrena (Melanapis) fuscosa** Erichson, 1835 (?*Andrena aethiops* Imhoff, 1832; *A. rutila* Spinola, 1838, nom. praeocc., nec Pérez, 1895; *A. ephippium* Spinola, 1838; *A. xanthoscelis* Brullé, 1839; *A. dorsalis* Lepeletier de Saint-Fargeau, 1841, nom. praeocc., nec Brullé, 1832; *A. lepeletieri* Lucas, 1849, nom. praeocc., nec Dalla Torre, 1896; *A. ruficornis* Smith, 1853; *A. atrocoerulea* Giraud, 1863; *A. dagestanica* Radoszkowski, 1867; *A. turcestanica* Morawitz, 1876; *A. dilecta* Mocsáry, 1879; *A. lepeletieri* var. *aschabadensis* Radoszkowski, 1893, nom. praeocc., nec Friese, 1922; *A. jucunda* Pérez, 1895; *A. ephippium* var. *cleopatra* Friese, 1899; *Melanapis violaceipennis* Cameron, 1902, nom. praeocc., nec Dufour, 1841; *M. rufifrons* Nurse, 1904; *Andrena cyprica* Cockerell, 1910; *A. ephippium* var. *flavipennis* Friese, 1914; *A. e.* var. *pallipes* Friese, 1914; *A. e.* var. *rufocincta* Friese, 1914, nom. praeocc., nec Ashmead, 1890; *A. e.* var. *rubriventris* Friese, 1922; *A. dorsatula* Viereck, 1918; *A. rutila* ab. *carnea* Lebedev, 1933; *A. rutila* ab. *zonaria* Lebedev, 1933; *A. ankarae* Alfken, 1935; *A. rutila* ssp. *mauritanica* Patiny, 1997; *A. canariensis* Patiny, 1997). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Jordan, Lebanon, Israel, Iran, Afghanistan, Central Asia, Kazakhstan, India.
- Andrena (Micrandrena) alfkenella** Perkins, 1914 (*A. moricella* Perkins, 1914; *A. pillichi* Noskiewicz, 1939; *A. alfkenella* ssp. *sunna* Warncke, 1975). Russia: **EP** (C, E, S, CR). – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey.
- Andrena (Micrandrena) enslinella** Stoeckhert, 1924 (*A. chimaera* Blüthgen, 1949). Russia: **EP** (C, E, S, NC), **UR**, **WS** (KM). – Europe (WE, SE, EE), Azerbaijan, Turkey, Iran.
- Andrena (Micrandrena) falsifica** Perkins, 1915. Russia: **EP** (C, E, NC, CR), **UR**, **WS** (KM), **ES** (IR). – Europe (WE, NE, SE, EE), Georgia, Azerbaijan.
- Andrena (Micrandrena) falsificissima** Hirashima, 1966. Russia: **FE** (KH, PR). – Korean Peninsula, Japan (Hok, Hon).
- Andrena (Micrandrena) floricola** Eversmann, 1852 (*A. punctulata* Schenck, 1853, nom. praeocc., nec Torka, 1913; *A. ochropyga* Alfken, 1916). Russia: **EP** (C, E, NC, CR), **UR**, **WS** (TK). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran.
- Andrena (Micrandrena) hikosana** Hirashima, 1957. Russia: **FE** (PR). – Korean Peninsula, Japan (Hok, Hon, Kyu, Shi).
- Andrena (Micrandrena) lazoiana** Osytshnjuk, 1995. Russia: **FE** (PR).
- Andrena (Micrandrena) magunta** Warncke, 1965. Russia: **EP** (NC, CR). – Europe (SE), Georgia, Azerbaijan, Turkey.
- Andrena (Micrandrena) minutissima** Osytshnjuk, 1995. Russia: **FE** (AM, KH, PR, SA).
- Andrena (Micrandrena) minutula** (Kirby, 1802) [Melitta] (*Melitta parvula* Kirby, 1802, nom. praeocc., nec Morawitz, 1871; *Andrena nigrifrons* Smith, 1855, nom. praeocc., nec Eversmann, 1852, nec Smith, 1853, nec Cresson, 1878; *A. gilvifrons* Pérez, 1903; *A. lenis* Pérez, 1903; *A. minutula* ssp. *alga* Warncke, 1974). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, China (NE, NC, NW), Japan (Hok, Hon, Shi, Kyu).
- Andrena (Micrandrena) minutuloides** Perkins, 1914 (*A. parvuloides* Perkins, 1914; *A. sparsiciliata* Alfken, 1925). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (TK, KM, AL), **ES** (KS, KR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran.
- Andrena (Micrandrena) nana** (Kirby, 1802) [Melitta] (*A. schenkella* Pérez, 1903; *A. nana* var. *taorminae* Strand, 1921; *A. asuniensis* Strand, 1921). Russia: **EP** (C). – Europe (WE, NE, SE, EE), N Africa.
- Andrena (Micrandrena) nanaeformis** Noskiewicz, 1925. Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (KM, AL), **ES** (KS, IR). – Europe (WE, EE), Azerbaijan, Kazakhstan.
- Andrena (Micrandrena) nanula** Nylander, 1848. Russia: **EP** (NW, C, E), **UR**, **WS** (KM, AL), **ES** (KS, IR, BR, ZB), **FE** (PR). – Europe (WE, NE, SE, EE), Azerbaijan.
- Andrena (Micrandrena) niveata** Friese, 1887 (*A. niveata* ssp. *bubulca* Warncke, 1975; *A. n.* ssp. *haloga* Warncke, 1980; *A. n.* ssp. *lecania* Warncke, 1975). Russia: **EP** (C, CR), **UR**. – Europe (WE, NE, SE, EE), Turkey.
- Andrena (Micrandrena) roripae** Osytshnjuk, 1993. Russia: **EP** (S, CR).
- Andrena (Micrandrena) rugulosa** Stoeckhert, 1935. Russia: **EP** (CR). – Europe (WE, SE, EE), Turkey.

- Andrena (Micrandrena) rugulosella** Osytshnjuk, 1993. Russia: EP (S). – Kazakhstan.
- Andrena (Micrandrena) semilaevis** Pérez, 1903 (*A. saundersella* Perkins, 1914). Russia: EP (NW, C, E, S, CR). – Europe (WE, NE, SE, EE), Azerbaijan.
- Andrena (Micrandrena) semirugosa** Cockerell, 1924 (*A. brassicae* Hirashima, 1957, nom. praeocc., nec Mitchell, 1960). Russia: FE (AM, KH, PR, SA, KU). – China (NE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu).
- Andrena (Micrandrena) sillata** Warncke, 1975 (*A. sil-lata* ssp. *histrionica* Warncke, 1975). Russia: EP (NC, CR). – Europe (SE), Turkey.
- Andrena (Micrandrena) stoeckhertella** Pittioni, 1948. Russia: EP (S, NC, CR). – Azerbaijan, Turkey.
- Andrena (Micrandrena) subopaca** Nylander, 1848. Russia: EP (N, NW, C, E, S, NC, CR), UR, WS (TK, KM, AL), ES (KR, IR, BR, ZB), FE (AM, KH, PR, SA, KU, KA). – Europe (WE, NE, SE, EE), Georgia, Azerbaijan, Kazakhstan, China (NE, SW), Japan (Hok, Hon, Shi, Kyu).
- Andrena (Micrandrena) tringa** Warncke, 1973. Russia: EP (C, S, NC, CR). – Europe (EE), Azerbaijan, Turkey.
- Andrena (Nobandrena) anatolica** Alfken, 1935. Russia: EP (S). – Europe (SE, EE), Azerbaijan, Turkey, Syria, Lebanon.
- Andrena (Nobandrena) athenensis** Warncke, 1965. Russia: EP (NC, CR). – Europe (SE, EE), Caucasus, Turkey.
- Andrena (Nobandrena) flavobila** Warncke, 1965 (*A. flavobila* ssp. *albicilla* Warncke, 1975). Russia: EP (CR). – Europe (SE, EE), Turkey.
- Andrena (Nobandrena) fratercula** Warncke, 1975. Russia: EP (NC). – Turkey.
- Andrena (Nobandrena) nobilis** Morawitz, 1874 (*A. palouri* Morawitz, 1877). Russia: EP (S, NC, CR), UR. – Europe (WE, SE, EE), Caucasus, Turkey, Iran, Turkmenistan.
- Andrena (Notandrena) chrysosceles** (Kirby, 1802) [Melitta] (*Melitta connectens* Kirby, 1802; *Andrena fallax* Eversmann, 1852; *A. erythrocnemis* Morawitz, 1870, nec auct.). Russia: EP (C, E, S, NC), UR, WS (OM, TK, KM, AL). – Europe (WE, NE, SE, EE), Georgia, Turkey, Kazakhstan.
- Andrena (Notandrena) minor** (Radoszkowski, 1891) [Cilissa] (*A. kurdistanica* Engel, 2005). Russia: FE (PR).
- Andrena (Notandrena) nitidiuscula** Schenck, 1853 (?*A. xanthopyga* Illiger, 1806; *A. fulvicornis* Schenck, 1853; *A. fulvicornis* Schenck, 1861; *A. lucens* Imhoff, 1868; *A. nigellata* Pérez, 1895; *A. gaschetti* Pérez, 1903; *A. petroselini* Pérez, 1903; *A. divergens* Pérez, 1903; *A. rostellata* Pérez, 1903; *A. rubrosignata* Saunders, 1908; *A. lucens* var. *algira* Friese, 1922; *A. franconica* E. Stoeckhert, 1922). Russia: EP (C, E, S, NC, CR), WS (TK, KM), ES (TU, IR, BR, ZB), FE (KH). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Syria, Israel, Iran, Mongolia, China (NE, NW, WP), Japan (Hok, Hon, Kyu).
- Andrena (Notandrena) pallitarsis** Pérez, 1903. Russia: UR, WS (KM). – Europe (WE, SE, EE).
- Andrena (Opandrena) schencki** Morawitz, 1866 (*A. labiata* sensu E. Stoeckhert, 1930, nom. praeocc., nec Fabricius, 1781; *A. bipustulata* Friese, 1921). Russia: EP (NC), UR. – Europe (WE, NE, SE, EE), Caucasus, Turkey, Israel, Iran, Turkmenistan.
- Andrena (Orandrena) oralis** Morawitz, 1876 (*A. sog-diana* Morawitz, 1876; *A. sisymbrii* Friese, 1887). Russia: EP (S, NC, CR), UR. – Europe (WE, SE, EE), Turkey, Uzbekistan, Kazakhstan.
- Andrena (Oreomelissa) coitana** (Kirby, 1802) [Melitta] (*Melitta shawella* Kirby, 1802; *Andrena nylanderi* Morawitz, 1864; *A. amaguensis* Cockerell, 1924; *A. coitana* var. *pilosodorsata* Alfken, 1929; *A. c. ssp. xema* Warncke, 1975). Russia: EP (NW, C, E, NC, CR), UR, WS (TK, KM, AL), ES (KS, TU, IR, BR, YA, ZB), FE (AM, KH, PR, SA, KU, KA, MG). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Kyrgyzstan, Kazakhstan, China (NW), Korean Peninsula, Japan (Hok).
- Andrena (Oreomelissa) kamikochiana** Hirashima, 1963. Russia: FE (PR). – Mongolia, China (NE, NC), Korean Peninsula, Japan (Hon).
- Andrena (Oreomelissa) media** (Radoszkowski, 1891) [Cilissa] (*A. amurensis* Friese, 1922, nom. praeocc., nec Radoszkowski, 1891; *A. amurica* Engel, 2005). Russia: FE (PR). – China (NE), Korean Peninsula.
- Andrena (Oreomelissa) mitakensis** Hirashima, 1963. Russia: FE (SA, KU). – China (NC), Japan (Hok, Hon).
- Andrena (Pallandrena) braunsiana** Friese, 1887 (*A. braunsiana* ssp. *detorta* Warncke, 1975). Russia: EP (C, S, CR). – Europe (WE, SE, EE), Azerbaijan, Turkey.
- Andrena (Parandrenella) atrata** Friese, 1887 (*A. bicarinata* auct., nec Morawitz, 1876). Russia: EP (C, S, NC, CR), UR, WS (NS, KM), ES (KS). – Europe (WE, SE, EE), Azerbaijan, Kazakhstan.

Andrena (Parandrenella) dentiventris Morawitz, 1874. Russia: EP (S, NC). – Georgia, Azerbaijan, Turkey, Iran, Kazakhstan.

Andrena (Parandrenella) figurata Morawitz, 1866. Russia: EP (E, S, NC, CR), UR. – Europe (SE, EE), Caucasus, Turkey, Iran, Kazakhstan.

Andrena (Plastandrena) bimaculata (Kirby, 1802) [Melitta] (*A. decorata* Smith, 1847; *A. conjuncta* Smith, 1847; *A. articulata* Smith, 1847; *A. vitrea* Smith, 1847; *A. consobrina* Eversmann, 1852, nom. praecocc., nec Schenck, 1861; *A. vitreipennis* Costa, 1861; *A. mystacea* Dours, 1861; *A. intermedia* Morawitz, 1870, nom. praecocc., nec Thomson, 1870; *A. morawitzi* Thomson, 1872; *A. atro-rubricata* Dours, 1872; *A. aulica* Morawitz, 1876; *A. compara-ta* Morawitz, 1876; *A. melanura* Morawitz, 1877; *A. paveli* Schmiedeknecht, 1883; *A. magrettiana* Schmiedeknecht, 1884; *A. germabica* Radoszkowski, 1893; *A. vitrea* var. *tenebrosa* Gribodo, 1894; *A. atrorubricata* var. *concolor* Alfken, 1914, nom. praecocc., nec Robertson, 1898, nec Blüthgen, 1919; *A. bimaculata* var. *mondaensis* Friese, 1922; *A. bimaculata* var. *serotinella* Friese, 1922; *A. morawitzi* var. *hirtella* Friese, 1922; *A. tibialis* var. *tricolorata* Friese, 1922; *A. florea* var. *clavipes* Friese, 1924; *A. blüthge-ni* Stoeckhert, 1930; *A. bimaculata* ssp. *oligotricha* Mavromoustakis, 1952; *A. gaetula* Benoist, 1961; *A. bimaculata* ssp. *lichata* Warncke, 1967). Russia: EP (NW, C, E, S, NC, CR), UR, WS (OM, TK, KM), ES (KR, IR, BR). – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, Syria, Israel, Iraq, Iran, Afghanistan, Central Asia, Kazakhstan, Mongolia.

Andrena (Plastandrena) dzynnanica Popov, 1949. Russia: ES (TU, IR, BR, ZB), FE (KH, PR). – Mongolia, China (NE, NC).

Andrena (Plastandrena) eversmanni Radoszkowski, 1867 (*A. fasciata* Radoszkowski, 1876, nom. praecocc., nec Fabricius, 1775, nec Imhoff, 1832, nec Nylander, 1852; *A. radoszkowskyi* Schmiedeknecht, 1884, nom. praecocc., nec Dalla Torre, 1896; *A. eversmanni* ssp. *ciscaspica* Popov, 1949). Russia: EP (NC). – Azerbaijan, Kazakhstan.

Andrena (Plastandrena) khasania Osytshnjuk, 1995. Russia: FE (AM, PR).

Andrena (Plastandrena) pilipes Fabricius, 1781 (?*Apis riparia* Scopoli, 1763; *A. atra* Müller, 1776, nom. praecocc., nec Scopoli, 1763, nec Smith, 1847, nec Popov, 1940; *A. ciliata* Gmelin, 1790, nom. praecocc., nec Schenck, 1869; *Andrena aterrima* Panzer, 1799; *Nomada nigrita* Panzer, 1800, nom. praecocc., nec

Fabricius, 1775, nec Morawitz, 1876; *Andrena atra* Smith, 1847, nom. praecocc., nec Müller, 1776, Scopoli, 1763, nec Popov, 1940; *A. spectabilis* Smith, 1853; *A. elongata* Radoszkowski, 1871, nom. praecocc., nec Imhoff, 1834, nec Radoszkowski, 1876; *A. nigrospina* Thomson, 1872; *A. praetexta* Smith, 1872; *A. doloro-sa* Nurse, 1904; *A. carbonaria* var. *autumnalis* Friese, 1922, nom. praecocc., nec Viereck et Cockerell, 1914; *A. c. var. claripennis* Friese, 1922; *A. c. ssp. iliensis* Alfken, 1938; *A. c. ssp. imamiana* Osytshnjuk, 1994; *Apis carbonaria* auct., nec Linné, 1967). Russia: EP (N, NW, C, E, S, CR), UR, WS (NS, KM, AL), ES (KS, TU, IR, BR, ZB), FE (AM, PR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Syria, Is-rael, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NC, NW, WP), India.

Andrena (Plastandrena) tibialis (Kirby, 1802) [Melitta] (*Melitta mouffetella* Kirby, 1802; *M. atriceps* Kirby, 1802; *Andrena fulvitarsis* Eversmann, 1852, nom. praecocc., nec Brullé, 1832; *A. ambiguia* Eversmann, 1852, nom. praecocc., nec Perkins, 1895; *A. subfascia-ta* Schenck, 1853; *A. indobonensis* Stoeckhert, 1950; *A. tibialis* ssp. *concreta* Warncke, 1975; *A. t. ssp. cor-vina* Warncke, 1967; *A. t. ssp. porzana* Warncke, 1975). Russia: EP (NW, C, E, S, CR), UR, WS (OM, TK, NS, KM, AL), ES (KS, TU, KR, IR, BR), FE (PR). – Europe (WE, NE, SE, EE), Caucasus, Tur-key, Iran, Kyrgyzstan, Kazakhstan, China (NC).

Andrena (Plastandrena) transbaicalica Popov, 1949 (*A. astragalina* Hirashima, 1957). Russia: WS (TK, NS, KM), ES (IR, BR, YA, ZB), FE (AM, KH, PR). – Korean Peninsula, Japan (Hok, Hon, Kyu).

Andrena (Poecilandrena) fukuokensis Hirashima, 1952 (*A. leleji* Osytshnjuk, 1982). Russia: ES (ZB), FE (PR). – China (NE, NC, CC), Korean Peninsula, Japan (Hon, Shi, Kyu, Tsu).

Andrena (Poecilandrena) hybrida Warncke, 1975 (*A. hybrida* ssp. *tauriensis* Warncke, 1975; *A. h. ssp. uralskia* Warncke, 1975). Russia: EP (S, CR). – Europe (SE), Turkey.

Andrena (Poecilandrena) labiata Fabricius, 1781 (*Apis fusciventris* Gmelin, 1790; *A. maculosa* Gmelin, 1790; *A. albilabris* Panzer, 1798; *Andrena divisa* Imhoff, 1834; *A. ferruginata* Schenck, 1853; *A. regina* Friese, 1921; *A. labiata* ssp. *bellina* Warncke, 1967; *A. cin-gulata* auct., nec Fabricius). Russia: EP (NW, C, E, S, NC, CR), UR, WS (TK, KM, AL), ES (KS, IR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Kyrgyzstan, Kazakhstan.

- Andrena (Poecilandrena) potentillae** Panzer, 1809
(*A. genevensis* Schmiedeknecht, 1884). Russia: **EP** (E, S), **UR**. – Europe (WE, SE, EE).
- Andrena (Poecilandrena) seminuda** Friese, 1896
(*A. setigera* Alfken, 1911). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), Georgia, Azerbaijan, Turkey.
- Andrena (Poecilandrena) semirubra** Morawitz, 1876.
Russia: **EP** (S, NC, CR). – Caucasus, Turkey, Israel.
- Andrena (Poecilandrena) sphecodimorpha** Hedicke, 1942 (*A. potentillae* ssp. *pendelica* Mavromoustakis, 1958). Russia: **EP** (S, CR). – Europe (SE, EE), Israel.
- Andrena (Poliandrena) altaica** Lebedev, 1932. Russia: **WS** (NS). – Kazakhstan.
- Andrena (Poliandrena) caspica** Morawitz, 1886
(*A. unifasciata* Friese, 1899). Russia: **EP** (NC). – Europe (SE), Azerbaijan, Turkey, Lebanon.
- Andrena (Poliandrena) florea** Fabricius, 1793 (*A. austriaca* Panzer, 1798; *A. rubricata* Smith, 1847; *A. globulilabris* Pérez, 1895; *A. florea* var. *obscura* Friese, 1914, nom. praeocc., nec Linné, 1764, nec Müller, 1776, Robertson, 1902; *A. f. var. rubra* Friese, 1914; *A. f. f. haslauica* Strand, 1921; *A. f. ssp. anhinga* Warncke, 1975; *A. f. ssp. corsica* Warncke, 1975). Russia: **EP** (C, S, CR), **UR**. – Europe (WE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Iran, Turkmenistan.
- Andrena (Poliandrena) limbata** Eversmann, 1852
(*A. squamea* Giraud, 1863; *A. limbata* ssp. *dusmeti* Warncke, 1967). Russia: **EP** (S, NC, CR), **UR**. – Europe (WE, NE, EE), Georgia, Azerbaijan, Turkey, Syria, Iran, Turkmenistan.
- Andrena (Poliandrena) ornata** Morawitz, 1866. Russia: **EP** (E, S, CR), **UR**. – Azerbaijan, Kazakhstan.
- Andrena (Poliandrena) polita** Smith, 1847 (*A. ochracea* Morawitz, 1872; *A. pseudopolita* Alfken, 1939). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Iran, Kazakhstan.
- Andrena (Poliandrena) tatjanae** Osytschnjuk, 1995.
Russia: **FE** (PR).
- Andrena (Proxiandrena) alutacea** E. Stoeckhert, 1942. Russia: **EP** (C, E). – Europe (WE, NE, SE, EE), Azerbaijan, Turkey, Iran, Turkmenistan.
- Andrena (Proxiandrena) proxima** (Kirby, 1802) [Melitta] (*Melitta digitalis* Kirby, 1802). Russia: **EP** (C, E, NC, CR), **UR**, **WS** (TK, KM). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Central Asia.
- Andrena (Ptilandrena) vetula** Lepetletier de Saint-Fargeau, 1841 (*A. megacephala* Smith, 1853; *A. opaca* Morawitz, 1868, *A. tyrrhena* Alfken, 1938. Russia: **EP** (NC). – Europe (WE, SE), N Africa, Caucasus, Turkey, Lebanon, Israel, Iraq, Iran.
- Andrena (Scitandrena) scita** Eversmann, 1852 (*A. obscuricanda* Costa, 1861; *A. transcaspica* Radoszkowski, 1886, nom. praeocc., nec Radoszkowski, 1893; *A. mesopyrrha* Dours, 1872; *A. scita* var. *nigra* Friese, 1914, nom. praeocc., nec Provancher, 1895, nec Saunders, 1908, nec Wu, 1982; *A. scita* var. *nigrofasciata* Friese, 1914). Russia: **EP** (E, S, NC, CR). – Europe (WE, SE, EE), Caucasus, Turkey, Iran, Afghanistan, Turkmenistan, Kazakhstan.
- Andrena (Simandrena) combinata** (Christ, 1791)
[*Apis*] (*A. candens* Eversmann, 1852; *A. albibarbis* Schenck, 1853; *A. sahlbergi* Morawitz, 1888; *A. mehelyi* Alfken, 1936; *A. combinata* ssp. *crudelis* Warncke, 1967). Russia: **EP** (C, E, CR), **UR**, **WS** (TK, NS, KM, AL), **ES** (KS, IR, BR, YA, ZB), **FE** (MG). – Europe (WE, SE, EE), N Africa, Georgia, Armenia, Turkey, Syria, Kyrgyzstan, Kazakhstan, Mongolia, China (NW).
- Andrena (Simandrena) congruens** Schmiedeknecht, 1884 (*A. dallatorrei* Clément, 1922; *A. confinis* E. Stoeckhert, 1930). Russia: **EP** (C, E), **UR**. – Europe (WE, NE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Syria.
- Andrena (Simandrena) dorsata** (Kirby, 1802) [Melitta] (?*Apis transfuga* Christ, 1791; *Melitta lewinella* Kirby, 1802; *M. collinsonana* Kirby, 1802; *M. nudiuscula* Kirby, 1802; *M. subincana* Kirby, 1802; *Andrena propinquua* Schenck, 1853; *A. cognata* Schenck, 1853; *A. griseola* Schenck, 1861; *A. dubitata* Schenck, 1870; *A. connexa* Pérez, 1895; *A. alliaria* Pérez, 1903; *A. thomsoni* Aurivillius, 1903, nom. praeocc., nec Ducke, 1898). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (TK). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel, Lebanon, Iran, Turkmenistan, Kazakhstan.
- Andrena (Simandrena) kerriæ** Hirashima, 1965. Russia: **WS** (KM), **ES** (BR, ZB), **FE** (AM, PR). – China (NE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Andrena (Simandrena) lepida** Schenck, 1861 (*A. separanda* Schmiedeknecht, 1884; *A. ardens* Pérez, 1895; *A. lepida* ssp. *eleonorae* Warncke, 1967). Russia: **EP** (C, NC, CR), **UR**. – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, Syria, Israel, Iran.
- Andrena (Simandrena) nippon** Tadauchi et Hirashima, 1983. Russia: **WS** (TK, KM), **ES** (KS, BR, ZB), **FE** (PR). – China (NE), Korean Peninsula, Japan (Hon),

Andrena (Simandrena) susterai Alfken, 1914. Russia: EP (NC, CR). – Europe (WE, SE, EE).

Andrena (Simandrena) opacifovea Hirashima, 1952. Russia: FE (KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu).

Andrena (Simandrena) transitoria Morawitz, 1871 (*A. basalis* Dours, 1872, nom. praeocc., nec Kriechbaumer, 1873; *A. aurora* Friese, 1922; *A. transitoria* ssp. *schmidti* Alfken, 1936). Russia: EP (S, NC, CR). – Europe (WE, SE, EE), Georgia, Azerbaijan, Turkey, Israel, Jordan, Syria, Iran, Afghanistan, Central Asia.

Andrena (Stenomelissa) halictoides Smith, 1869. Russia: FE (PR). – China (NE), Korean Peninsula, Japan (Hok, Hon).

Andrena (Taeniandrena) aberrans Eversmann, 1852 (*A. ratisbonensis* Stoeckhert, 1924). Russia: EP (S). – Europe (WE, SE, EE), Azerbaijan.

Andrena (Taeniandrena) callopyrrha Cockerell, 1929 (*A. callopyrrha* ssp. *kozlovi* Osytshnjuk, 1994). Russia: WS (AL). – Kazakhstan, Mongolia, China (NC).

Andrena (Taeniandrena) ezoensis Hirashima, 1965. Russia: ES (YA), FE (SA, KU). – Kazakhstan, China (NE, NC), Korean Peninsula, Japan (Hok, Hon).

Andrena (Taeniandrena) gelriae van der Vecht, 1927 (*A. podolica* Noskiewicz, 1930; *A. gelriae karellica* Niemel, 1949; *A. g. k. f. fuscata* Niemel, 1949; *A. g. greedana* Warncke, 1975; *A. g. vocifera* Warncke, 1975). Russia: EP (N, NW, C, E, NC, CR), UR, WS (OM, TK, KM, AL), ES (KS, IR, YA, ZB), FE (AM, KH, PR). – Europe (WE, NE, SE, EE), Caucasus, Turkey.

Andrena (Taeniandrena) lathyri Alfken, 1899. Russia: EP (C, E, S, CR), WS (TK, NS, KM), ES (IR), FE (AM, KH, PR). – Europe (WE, NE, SE, EE), Azerbaijan, Turkey.

Andrena (Taeniandrena) intermedia Thomson, 1870. Russia: EP (NW, C, E). – Europe (WE, NE, SE, EE), Turkey, Iran.

Andrena (Taeniandrena) similis Smith, 1849 (?*Apis ocreata* Christ, 1791; *Andrena croceiventris* Morawitz, 1871; *A. rufohispida* Dours, 1872; *A. stefanii* Pérez, 1895; *A. ocreata* ssp. *cypriolina* Warncke, 1975; *A. similis* ssp. *caraimica* Osytshnjuk, 1994). Russia: EP (CR). – Europe (WE, NE, SE, EE), N Africa, Armenia, Turkey, Iran, Afghanistan, Kazakhstan, China (NW).

Andrena (Taeniandrena) ovatula (Kirby, 1802) [Melitta] (*Melitta afzeliella* Kirby, 1802; *M. fuscata* Kirby, 1802, nom. praeocc., nec Niemalä, 1949; *M. picipes* Kirby, 1802; *M. barbata* Kirby, 1802; *Andrena hirticeps* Eversmann, 1852, nom. praeocc., nec Smith,

1853; *A. interrupta* Eversmann, 1852, nom. praeocc., nec Schenck, 1869; *A. octostrigata* Schenck, 1853; *A. gibba* Schenck, 1853; *A. ovata* Schenck, 1853; *A. plantaris* Schenck, 1853; *A. canescens* Schenck, 1853; *A. albofimbriata* Schenck, 1853; *A. distincta* Schenck, 1861, nom. praeocc., nec Smith, 1847, nec Lucas, 1849; *A. albofasciata* Thomson, 1870; ?*A. turbida* Walker, 1871; *A. poupillieri* Dours, 1872; *A. meliloti* Verhoeff, 1890; *A. afzeliella* var. *heliopolis* Friese, 1914; *A. pseudovatula* Alfken, 1926; *A. lecerfi* Benoist, 1961; *A. poupillieri* ssp. *incana* Warncke, 1975; *A. ovatula* ssp. *transcaspiensis* Osytshnjuk, 1994). Russia: EP (C, E, S, NC, CR), UR, WS (OM, TK, NS, KM, AL), ES (KS, IR, BR, YA, ZB), FE (AM, KH, PR, KA). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel, Syria, Iraq, Iran, Afghanistan, Central Asia, Kazakhstan, China (NW).

Andrena (Taeniandrena) wilkella (Kirby, 1802) [Melitta] (*Melitta xanthura* Kirby, 1802; *M. convexicula* Kirby, 1802; *M. barbatula* Kirby, 1802, nom. praeocc., nec Zetterstedt, 1838; *Andrena convexa* Schenck, 1853, nom. praeocc., nec Schrank, 1781, nec Provancher, 1888; *A. quadristrigata* Schenck, 1853; *A. winkleyi* Viereck, 1907; *A. meraca* Cockerell, 1924; *A. beaumonti* Benoist, 1961). Russia: EP (NW, C, S, NC, CR), UR, WS (TK, NS, KM), ES (IR, BR, YA, ZB), FE (AM, PR). – Europe (WE, NE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Kyrgyzstan, Kazakhstan, China (NC, NW, CC), N America.

Andrena (Tarsandrena) angarensis Cockerell, 1929. Russia: WS (TK, KM, AL), ES (KS, TU, IR, BR, ZB), FE (AM, KH, PR). – Kyrgyzstan, Mongolia.

Andrena (Tarsandrena) bonivuri Osytshnjuk, 1984. Russia: ES (ZB), FE (PR).

Andrena (Tarsandrena) ehnbergi Morawitz, 1888 (*A. belicovi* Cockerell, 1929). Russia: UR, WS (OM, KM, AL), ES (KS, TU, KR, IR, BR, ZB), FE (AM, PR). – Mongolia, China (NC, NW).

Andrena (Tarsandrena) tarsata Nylander, 1848 (*A. basilinea* Kriechbaumer, 1873; *A. gentianae* Vachal, 1906; *A. baicalensis* Cockerell, 1929; *A. universitatis* Cockerell, 1929). Russia: EP (NW, C, NC), UR, WS (KM, AL), ES (KS, TU, IR, BR, YA, ZB), FE (MG). – Europe (WE, NE, SE, EE), Georgia, Azerbaijan, Turkey, Mongolia, China (NW, SW).

Andrena (Thysandrena) hypopolia Schmiedeknecht, 1884 (*A. consobrina* Schenck, 1861, nom. praeocc., nec Eversmann, 1852; *A. inconspicua* Morawitz, 1871; ?*A. holosericea* Bramson, 1879; *A. interrogationis* Dalla Torre, 1884, nom. praeocc., nec Viereck

et Cockerell, 1914; *A. setosa* Pérez, 1903). Russia: **EP** (S, NC, CR), **WS** (NS). – Europe (WE, SE, EE).

Andrena (Thysandrena) ranunculorum Morawitz, 1877. Russia: **EP** (S, NC, CR), **UR**, **WS** (NS, KM, AL), **ES** (TU, BR, ZB). – Europe (SE), Georgia, Azerbaijan, Turkey, Iran, Turkmenistan.

Andrena (Trachandrena) haemorrhoa (Fabricius, 1781) [*Apis*] (?*Apis sordida* Scopoli, 1763, nom. praeocc., nec Gmelin, 1790, nec Morawitz, 1876; ?*A. tranquilla* Müller, 1776; ?*A. hiberus* Harris, 1776; *A. haemorrhoa* Fabricius, 1781; *A. convexa* Schrank, 1781, nom. praeocc., nec Schenck, 1853, nec Provancher, 1888; *A. nebulosa* Gmelin, 1790; *A. haemorrhoidalis* Christ, 1791, nom. praeocc., nec Fabricius, 1775; *Andrena salicis* Verhoeff, 1890, nom. praeocc., nec Robertson, 1891; *A. haemorrhoa* ssp. *japonibia* Hirashima, 1957; *Apis albicans* auct., nec Müller, 1776). Russia: **EP** (N, NW, C, E, S, CR), **UR**, **WS** (TK, NS, KM, AL), **ES** (KS, TU, KR, IR, YA, ZB), **FE** (AM, KH, PR, KA). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Syria, Iran, Turkmenistan, Kazakhstan, Mongolia, China (NE), Korean Peninsula, Japan (Hok, Hon, Kyu, Shi, Tsu).

Andrena (Truncandrena) derbentina Morawitz, 1886. Russia: **EP** (NC). – N Africa, Azerbaijan, Turkey, Israel, Iran.

Andrena (Truncandrena) rufomaculata Friese, 1921 (*A. paramythensis* Mavromoustakis, 1957; *A. rufomaculata* ssp. *optata* Warncke, 1975). Russia: **EP** (CR). – Europe (SE, EE), Caucasus, Turkey, Syria, Israel.

Andrena (Truncandrena) truncatilabris Morawitz, 1877 (*A. truncatilabris* ssp. *espanola* Warncke, 1967; *A. t.* ssp. *nigropilosa* Warncke, 1967). Russia: **EP** (S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Israel, Syria, Iran, Kazakhstan.

Andrena (Ulandrena) combaella Warncke, 1966 (*A. leucorhina* auct., part., nec Morawitz, 1876). Russia: **EP** (S). – Europe (WE, SE, EE), Georgia, Azerbaijan, Turkey, Israel.

Andrena (Ulandrena) elegans Giraud, 1863 (*Apis grisea* Fabricius, 1798, nom. praeocc., nec Gmelin, 1790). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), Turkey, Kazakhstan.

Andrena (Ulandrena) fulvitarsis Brullé, 1832 (*Camptogaster subglobosus* Dours, 1873; *Andrena carinula* Pérez, 1895). Russia: **EP** (CR). – Europe (SE, EE), Turkey, Israel.

Andrena (Ulandrena) osychniukae Osytshnjuk, 1977 (*A. abbreviata* ssp. *osychniuki* Warncke, 1973;

A. osychniuki Osytshnjuk, 1977; *A. abbreviata* ssp. *osychniukae* Osytshnjuk, 1977). Russia: **EP** (CR). – Europe (SE, EE), Caucasus, Turkey.

Andrena (Zonandrena) chrysopyga Schenck, 1853 (*A. integra* Thomson, 1870, nom. praeocc., nec Smith, 1853; *A. chrysopyga* var. *stefaniana* Schmiedeknecht, 1884). Russia: **EP** (C, E, S, NC), **UR**, **WS** (OM, TK, KM, AL), **ES** (KS, KR, IR). – Europe (WE, NE, SE, EE), Georgia, Azerbaijan, Turkey, Kyrgyzstan, Kazakhstan.

Andrena (Zonandrena) flavipes Panzer, 1799 (*Apis sordida* Gmelin, 1790, nom. praeocc., nec Scopoli, 1763, nec Morawitz, 1876; *Melitta fulvicrus* Kirby, 1802, nom. praeocc., nec Dufour, 1841; *Melitta continua* Kirby, 1802; *Andrena quadricincta* Brullé, 1832, nom. praeocc., nec Eversmann, 1852; *A. tricincta* Brullé, 1832; *A. fasciata* Imhoff, 1832, nom. praeocc., nec Fabricius, 1775, nec Nylander, 1852, nec Radoszkowski, 1876; *A. puber* Erichson, 1835; *A. mactae* Lepeletier de Saint-Fargeau, 1841; *A. fulvicrus* Dufour, 1841, nom. praeocc., nec Kirby, 1802; *A. cinerascens* Eversmann, 1852, nom. praeocc., nec Nylander, 1848; *A. capitalis* Smith, 1853; *A. interrupta* Schenck, 1869, nom. praeocc., nec Eversmann, 1852; *A. problematica* Pérez, 1903; *A. levilabris* Cameron, 1908; *A. punjaubensis* Cameron, 1909; *A. interruptula* Viereck, 1916; *A. quadricinctula* Viereck, 1916; *A. kraussei* Strand, 1921; *A. kengracensis* Cockerell, 1930; *A. flavipes* ssp. *alexandrina* Warncke, 1966; *A. flavipes* ssp. *ibicensis* Warncke, 1984). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel, Syria, Jordan, Iraq, Iran, Afghanistan, Central Asia, Kazakhstan, China (NW, WP), Nepal, India, N America.

Andrena (Zonandrena) sibirica Morawitz, 1888. Russia: **UR**, **WS** (NS, KM, AL), **ES** (KS, KR, IR, BR, ZB), **FE** (AM, KH, PR). – Kazakhstan, Mongolia, China (NE, NC).

Andrena (?) pseudothoracica Engel, 2005 (*Ciliessa thoracica* Radoszkowski, 1891, nom. praeocc., nec Fabricius, 1775). Russia: **FE** (PR).

Subfamily PANURGINAE

Tribe PANURGINI

CAMPTOPOEUM Spinola, 1843. Type species *Prosopis frontalis* Fabricius, 1804. Genus is distributed only in the Palaearctic region. Number of species: World – 30; Palaearctic – 30; Russia – 2.

Camptopoeum (Camptopoeum) friesei Mocsáry, 1894 (*C. friesei* var. *disparilis* Alfken, 1934; *C. f.* var. *intermedius* Alfken, 1934; *C. f.* var. *subinterruptus* Alfken, 1934). Russia: **EP** (S, CR), **UR**. – Europe (WE, SE, EE), Georgia, Turkey, Iran, Turkmenistan.

Camptopoeum (Camptopoeum) frontale (Fabricius, 1804) [Prosopis] (*Panurgus fasciatus* Giraud, 1857; *Camptopoeum altaicum* Morawitz, 1892). Russia: **EP** (C, S, CR). – Europe (WE, SE, EE), Azerbaijan, Turkey, Israel, Iran, Pakistan, Turkmenistan, Kazakhstan.

PANURGINUS Nylander, 1848 (*Scapteroides* Griswold, 1894; *Greeleyella* Cockerell, 1904; *Birkmania* Viereck, 1909). Type species *Panurginus niger* Nylander, 1848. Genus is distributed in the Holarctic region. Number of species: World – 53; Palaearctic – 31; Russia – 13.

Panurginus alpotanini Romankova et Astafurova, 2011. Russia: **FE** (KH). – Mongolia.

Panurginus alticolus Morawitz, 1876. Russia: **EP** (E, S, NC). – Georgia, Kazakhstan.

Panurginus arsenievi Romankova et Astafurova, 2011. Russia: **FE** (PR).

Panurginus crawfordi Cockerell, 1914. Russia: **FE** (PR). – Korean Peninsula, Japan (Hon, Shi, Kyu).

Panurginus herzi Morawitz, 1891 (*P. tyrolensis* Richards, 1932; *Panurgus montanus tyrolensis* Warncke, 1972; *P. m. herzi* Warncke, 1972). Russia: **WS** (KM, AL), **ES** (TU, YA). – Europe (NE), Mongolia.

Panurginus kropotkini Romankova et Astafurova, 2011. Russia: **FE** (PR).

Panurginus labiatus (Eversmann, 1852) [Panurgus] (*Panurgus clypeatus* Eversmann, 1852; *Panurginus arenarius* Schenck, 1861). Russia: **EP** (C, E, S), **UR**, **ES** (IR). – Europe (WE, SE, EE), Georgia, Azerbaijan, Kazakhstan.

Panurginus lactipennis Friese, 1897. Russia: **EP** (C, S, NC, CR). – Europe (SE), Azerbaijan, Turkey, Syria, Jordan.

Panurginus mikhno Romankova et Astafurova, 2011. Russia: **WS** (AL), **ES** (ZB).

Panurginus muraviovi Romankova et Astafurova, 2011. Russia: **WS** (KM, AL). – Mongolia.

Panurginus niger Nylander, 1848 (*P. dubius* Osytshnuk, 1995). Russia: **ES** (IR, BR, ZB), **FE** (AM, KH, PR). – Mongolia, China (NC).

Panurginus romani Aurivillius, 1914. Russia: **EP** (N), **WS** (KM, AL), **ES** (TU, KR, IR, BR, ZB), **FE** (AM, KH, PR, SA, KA, MG). – Europe (NE).

Panurginus sculpturatus Morawitz, 1872. Russia: **EP** (E, S, CR), **UR**. – Europe (EE).

PANURGUS Panzer, 1806 (*Eriops* Klug, 1807). Type species *Andrena lobata* Panzer, 1799 = *Apis calcarata* Scopoli, 1763. Genus is distributed only in the Palaearctic region. Number of species: World – 35; Palaearctic – 35; Russia – 1.

Panurgus calcaratus (Scopoli, 1763) [*Apis*] (*Apis calcarata* Scopoli, 1763; *A. denticulata* Gmelin, 1790; *A. ursina* Gmelin, 1790; *Philanthus ater* Fabricius, 1793; *Andrena lobata* Panzer, 1799; *Apis linnaeella* Kirby, 1802; *Panurgus unicolor* Spinola, 1808; *P. calcaratus* var. *nigricornis* Dalla Torre, 1877; *P. c.* var. *macrocephalus* Friese, 1897). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (TM, KM, AL), **ES** (KS, KR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Kazakhstan.

Tribe MELITTURGINI

MELITTURGA Latreille, 1809. Type species *Eucera clavicornis* Latreille, 1806. Genus is distributed in the Palaearctic and the Afrotropical regions. Number of species: World – 17; Palaearctic – 14; Russia – 3.

Melitturga clavicornis (Latreille, 1806) [*Eucera*] (*Macrocera nasalis* Eversmann, 1852; *Melitturga clavicornis* var. *thuringiaca* Friese, 1895). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (KS, TU, KR, IR, BR). – Europe (WE, SE, EE), Caucasus, Turkey, Iran, Central Asia, Kazakhstan, China (NW).

Melitturga mongolica Alfken, 1936. Russia: **ES** (BR, ZB), **FE** (AM, PR). – Turkey, Armenia, Iran, Afghanistan, Kyrgyzstan, Mongolia, China (NE, NC, NW, WP).

Melitturga praestans Giraud, 1861 (*M. caucasica* Morawitz, 1877). Russia: **UR**. – Europe (WE, SE, EE), Georgia, Turkey, Syria, Lebanon, Israel, Kazakhstan.

75. FAMILY HALICTIDAE

YU.V. ASTAFUROVA AND M.YU. PROSHCHALYKIN

The family Halictidae is a very large and nearly cosmopolitan group of bees, often dominate other bees in numbers of individuals. This family is monophyletic unit and belongs to generalised group conditionally named short-tongued bees. Number of taxa: World – the number of halictid genera varies in classifications by different authors, 4000 species; Palaearctic – 23 genera, 950 species; Russia – 13 genera, 263 species.

The classification of Halictini and Nomioidinae follows that of Michener (2007) and Nomiinae and Rophitinae that of Astafurova (2014).

R e f e r e n c e s. Morawitz, 1865, 1869; Mocsáry, Szépligeti, 1901; Blüthgen, 1923, 1924, 1925, 1931, 1935, 1955; Popov, 1935; Nikiforuk, 1957; Elfving, 1968; Ebmer, 1972, 1974, 1975, 1978, 1980, 1986, 1988a, b, 1995; 1998, 2000, 2004, 2005; Pesenko, 1972, 1985, 1986, 2005a, b, 2006a, b, 2007a, b; Bytinski-Salz, Ebmer, 1974; Sedykh, 1974; Warncke, 1975, 1982a, 1992; Dathe, 1980b; Ebmer, Sakagami, 1990; Fan, 1991; Rasmont et al., 1995; Pesenko et al., 2000; Niu et al., 2004; Murao, Tadauchi, 2007; Kuznetsova, 2009; Proshchalykin, 2009b, 2012a, 2014b; Bogusch, Straka, 2012; Proshchalykin, Astafurova, 2012, 2016; Levchenko, Yuferev, 2013; Mitai, Tadauchi, 2013; Astafurova, 2014; Astafurova, Proshchalykin, 2014, 2015, 2016, 2017a, b, c; Levchenko, 2014, 2015; Murao, 2015, 2017; Sidorov et al., 2016; Astafurova et al., 2017.

Subfamily HALICTINAE

This is an enormous and nearly cosmopolitan subfamily. Halictinae is subdivided by into two tribes: Augochlorini and Halictini. The tribe Augochlorini with more than 500 species is distributed in Western Hemisphere. The tribe Halictini is cosmopolitan group and includes more than 2500 species. Most of Halictini nest in burrows in banks or flat soil, except for a few groups that nest in rotting wood. The Halictinae exhibit both solitary behaviour and nearly every degree of sociality, ranging from subsocial to eusocial behaviour. Some augochlorini genera and halictini subtribe Sphecodina are cleptoparasites in nests of halictids and other bees. The majority of species are polilectic.

HALICTUS Latreille, 1804. Type species *Apis quadricincta* Fabricius, 1776. Widely distributed in the World (except Australia), but most of species

occur in northern hemisphere. Number of taxa: World – about 200 species and 15 subgenera, Palaearctic – more 150 species, Russia – 48.

Halictus (Argalictus) cochlearitarsis (Dours, 1872) [Lucasius] (*H. cochleareitarsis* Dalla Torre, 1896; *H. anomalipes* Lebedev, 1911). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), Armenia, Azerbaijan, Turkey.

Halictus (Argalictus) luganicus Blüthgen, 1936. Russia: **EP** (S, NC, CR). – Europe (EE), Georgia, Azerbaijan, Turkey, Kazakhstan.

Halictus (Argalictus) senilis (Eversmann, 1852) [*Hylaeus*] (*H. fucus* Morawitz, 1876; *H. albarius* Pérez, 1895; *H. albarius* Pérez, 1895; *H. bicinctus* Vachal, 1902; *H. aegypticola* Strand, 1909). Russia: **EP** (S, NC), **UR**. – Europe (SE, EE), N Africa, Caucasus, Turkey, Near East, Iraq, Iran, Afghanistan, Central Asia, Kazakhstan, Mongolia, China (NC, NW, SW).

Halictus (Halictus) brunnescens (Eversmann, 1852) [*Hylaeus*] (*H. quadricinctus maximus* Friese, 1916; *H. quadricinctus aegyptiacus* Friese, 1916). Russia: **EP** (S, NC, CR), **UR**, **ES** (BR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Syria, Israel, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NC, NW, WP), N India.

Halictus (Halictus) quadricinctus (Fabricius, 1776) [*Apis*] (*Apis hortensis* Geoffroy, 1785; *Halictus quadriguttatus* Latreille, 1805; *Hylaeus grandis* Illiger, 1806; *Halictus ecaphosus* Walckenaer, 1817; *H. charensis* Yasumatsu, 1940). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (OM, TK, NS, AL), **ES** (KR, IR, BR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Israel, Iran, Afghanistan, Central Asia, Kazakhstan, Mongolia, China (NE, NC, NW, WP).

Halictus (Hexataenites) resurgens Nurse, 1903 (*H. turkomannus* Pérez, 1903; *H. holtzi* Schulz, 1906; *H. asiaeminoris* Strand, 1921). Russia: **EP** (S, NC, CR). – Europe (SE, EE), N Africa, Caucasus, Turkey, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, China (NW), N India.

Halictus (Hexataenites) sexcinctus sexcinctus (Fabricius, 1775) [*Apis*] (*Apis ichneumonea* Christ, 1791; *Hylaeus arbustorum* Panzer, 1797; *Andrena rufipes* Spinola, 1806). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE).

Halictus (Monilapis) adjikenticus Blüthgen, 1923 (*H. scardicus* Blüthgen, 1936). Russia: **EP** (S, NC). – Europe (WE, SE), Armenia, Azerbaijan, Turkey.

Halictus (Monilapis) compressus compressus

(Walckenaer, 1802) [Andrena] (*Apis flavipes* Panzer, 1798; *Hylaeus tomentosus* Herrich-Schäffer, 1840; *H. senex* Förster, 1860; *Halictus eurygnathus* Blüthgen, 1931; *H. eurygnathopsis* Blüthgen, 1936; *H. veneticus* Ebmer, 1969). Russia: EP (N, NW, C, E, S, NC, CR). – Europe (WE, NE, SE, EE), Georgia, Azerbaijan, Turkey, Near East.

Halictus (Monilapis) compressus transvolgensis

Pesenko, 1985. Russia: EP (E), UR, WS (OM, TK, KM, AL), ES (KR, IR, BR, ZB). – Kyrgyzstan, Kazakhstan, China (NW).

Halictus (Monilapis) patellatus patellatus Morawitz, 1874 (*H. wagneri* Blüthgen, 1937). Russia: EP (NC). – Iran, Turkmenistan.**Halictus (Monilapis) patellatus taorminicus** Strand, 1921. Russia: EP (C, E, S, NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Syria, Iran.**Halictus (Monilapis) ponticus** Blüthgen, 1934. Russia: EP (NC). – Europe (EE), Turkey.**Halictus (Monilapis) rossicus** Ebmer, 1978 (*H. sarmensis* Blüthgen, 1936). Russia: EP (C, E, NC, CR), UR. – Europe (EE), Azerbaijan.**Halictus (Monilapis) sajoi** Blüthgen, 1923 (*H. veneticus* Móczár, 1967; *H. bifidus* Warncke, 1975). Russia: EP (C, S, CR), UR. – Europe (WE, SE, EE), Azerbaijan, Armenia, Turkey, Iran, Turkmenistan, Kyrgyzstan, Kazakhstan.**Halictus (Monilapis) simplex** Blüthgen, 1923 (*H. ibex* Warncke, 1973). Russia: EP (NW, C, E, S, NC, CR), UR. – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Kazakhstan.**Halictus (Monilapis) tetrazonianellus** Strand, 1909 (*H. leucognathus* Morice, 1921; *H. apatellatus* Strand, 1921; *H. gusenleitneri* Ebmer, 1975). Russia: EP (S, NC, CR). – Europe (SE, EE), Caucasus, Turkey, Iraq, Iran, Turkmenistan.**Halictus (Monilapis) tetrazonius** (Klug, 1817) [*Hylaeus*] (*H. furcatus* Blüthgen, 1925; *H. galileus* Blüthgen, 1955; *H. pannonicus* Ebmer, 1969). Russia: EP (S, E, CR). – Europe (WE, SE, EE), Georgia, Azerbaijan, Turkey, Israel, Iran, Turkmenistan.**Halictus (Monilapis) tsingtauensis** Strand, 1910 (*H. tsingtaensis* Blüthgen, 1923). Russia: WS (AL), ES (KR, IR, ZB), FE (AM, KH, PR, SA, KU). – China (NE, CC, SE), Korean Peninsula, Japan (Hok, Hon).**Halictus (Platyhalictus) alfkenellus cedens** Strand, 1909 (*H. posthumus* Blüthgen, 1925; *H. subalfkenellus* Blüthgen, 1936). Russia: EP (S, NC). – Europe

(WE, SE), Azerbaijan, Turkey, Lebanon, Israel, Iran, Kazakhstan.

Halictus (Platyhalictus) graecus Blüthgen, 1933. Russia: EP (CR). – Europe (SE, EE), Armenia, Azerbaijan, Turkey.**Halictus (Platyhalictus) lussinicus** Blüthgen, 1936. Russia: EP (S, NC). – Europe (SE).**Halictus (Platyhalictus) minor** Morawitz, 1876 (*H. altaicus* Pérez, 1903; *H. jarkandensis* Strand, 1909). Russia: WS (AL). – Azerbaijan, Iran, Pakistan, Central Asia, Kazakhstan, China (NE, NC), N India.**Halictus (Protohalictus) hedini hedini** Blüthgen, 1934. Russia: WS (KM, AL), ES (TU, KR, IR, BR, ZB), FE (AM, KH, PR, SA, KU). – Kazakhstan, Mongolia, China (NE, NC, NW), Korean Peninsula, Japan (Hok).**Halictus (Protohalictus) rubicundus** (Christ, 1791) [Apis] (*H. nidulans* Walckenaer, 1817; *H. lerouxi* Lepeletier de Saint Fargeau, 1841; *H. quadrisfuscatus* Smith, 1870; *H. lerouxi ruborum* Cockerell, 1898; *H. rubicundus nesiotes* Perkins, 1922; *H. r. laticincta* Blüthgen, 1923; *H. r. mongolensis* Blüthgen, 1936; *H. lupinelli* Cockerell, 1939; *H. frater* Pesenko, 1984). Russia: EP (N, NW, C, E, S, NC), UR, WS (OM, TK, KM, AL), ES (TU, KR, IR, BR, YA, ZB), FE (AM, KH, PR, SA, KA). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Kazakhstan, Mongolia, China (NE, NC, NW), Korean Peninsula, Japan (Hok), N America.**Halictus (Seladonia) aerarius** Smith, 1873 (*H. confluens* Morawitz, 1889; *H. alexoides* Strand, 1910; *H. pseudoconfluens* Strand, 1910; *H. nikkoensis* Cockerell, 1911; *H. leucopogon* Strand, 1914; *H. tussimae* Friese, 1916; *H. eruditus* Cockerell, 1924). Russia: FE (AM, KH, PR). – Mongolia, China (NE, NC, CC, SW, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Tsu, Yak).**Halictus (Seladonia) cephalicus** Morawitz, 1874 (*H. conjungens* Blüthgen, 1921; *H. neuter* Blüthgen, 1923). Russia: EP (S, NC, CR). – Europe (SE, EE), Caucasus, Turkey, Syria, Israel, Iraq, Iran, Afghanistan.**Halictus (Seladonia) confusus pelagi** Ebmer, 1996. Russia: WS (AL), ES (KS, KR, BR, YA), FE (AM, KH, PR, SA, KU, MG). – Mongolia, China (NC).**Halictus (Seladonia) confusus perkinsi** Blüthgen, 1926. Russia: EP (N, NW, C, E, S, NC, CR), UR. – Europe (WE, NE, SE, EE), Georgia, Turkey, Kazakhstan.

- Halictus (Seladonia) gavarnicus** Pérez, 1903 (*H. tataricus* Blüthgen, 1933; *H. gavarnicus delphinalis* Blüthgen, 1935; *H. g. longigenae* Warncke, 1988). Russia: **EP** (C, NC, CR), **UR**, **WS** (AL), **FE** (PR). – Europe (WE, SE, EE), Caucasus, Turkey, Mongolia, China (NC).
- Halictus (Seladonia) kessleri** Bramson, 1879 (*Osmia pannonica* Zilahí-Kiss, 1915; *Halictus kessleri nebulosus* Warncke, 1975; *H. k. hakkarius* Warncke, 1984). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (SE, EE), Caucasus, Turkey.
- Halictus (Seladonia) leleji** (Pesenko, 2006) [Seladonia]. Russia: **FE** (PR).
- Halictus (Seladonia) leucaheneus arenosus** Ebmer, 1976 (*H. fasciatus* auct.). Russia: **EP** (NW, C, E), **UR**. – Europe (WE, NE, SE, EE).
- Halictus (Seladonia) leucaheneus leucaheneus** Ebmer, 1972. Russia: **WS** (AL), **ES** (BR, IR, YA, ZB), **FE** (AM, KH, PR). – Kazakhstan, Kyrgyzstan, Mongolia, China (NE, NC, NW).
- Halictus (Seladonia) mondaensis** Blüthgen, 1923. Russia: **ES** (TU, BR, YA), **FE** (MG). – Mongolia.
- Halictus (Seladonia) mongolicus** Morawitz, 1880. Russia: **ES** (BR). – Mongolia, China (NW, WP).
- Halictus (Seladonia) nikolskayae** (Pesenko, 2006) [Seladonia]. Russia: **FE** (PR).
- Halictus (Seladonia) seladonius** (Fabricius, 1794) [*Apis*] (*H. geminatus* Pérez, 1903). Russia: **EP** (E, S), **UR**, **ES** (AL). – Europe (WE, SE, EE), Caucasus, Turkey, Israel, Iran, Afghanistan, Turkmenistan.
- Halictus (Seladonia) semitectus** Morawitz, 1874. Russia: **EP** (C, S, NC, CR), **UR**, **WS** (AL), **ES** (BR, ZB), **FE** (KH, PR). – Europe (WE, NE, SE, EE), Armenia, Afghanistan, Mongolia, China (NC, NW).
- Halictus (Seladonia) subauratus subauratus** (Rossi, 1792) [*Apis*] (*H. virescens* Lepeletier de Saint Fargeau, 1841; *H. gramineus* Smith, 1849; *H. meridionalis* Morawitz, 1874; *H. subauratus syrius* Blüthgen, 1933). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran, Central Asia, Pakistan, China (NW), N India.
- Halictus (Seladonia) submediterraneus** (Pauly, 2015) [Seladonia]. Russia: **EP** (S, NC, CR), **UR**. – Europe (WE, SE, EE), Turkey, Iran.
- Halictus (Seladonia) transbaikalensis** Blüthgen, 1933. Russia: **ES** (IR, ZB). – Mongolia, China (NC).
- Halictus (Seladonia) tumulorum ferripennis** Cockerell, 1929 (*H. tumulorum higashi* Sakagami et Ebmer, 1979). Russia: **WS** (AL), **ES** (KR, IR, BR, YA, ZB),

- FE** (AM, PR, SA, MG). – Mongolia, China (NE, NC), Korean Peninsula, Japan (Hok, Hon).
- Halictus (Seladonia) tumulorum tumulorum** (Linnaeus, 1758) [*Apis*] (*Apis flavipes* Fabricius, 1787; *Halictus fasciatus* Nylander, 1848; *H. tumulorum deviridatus* Strand, 1910). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TK, AL). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Syria, Israel, Iran, Central Asia.
- Halictus (Tytthaliictus) asperulus** Pérez, 1895 (*H. rugosulus* Pérez, 1895). Russia: **EP** (S, NC, CR). – Europe (SE, EE), Caucasus, Turkey, Syria, Israel, Iran.
- Halictus (Tytthaliictus) maculatus maculatus** Smith, 1848 (*H. interruptus* Lepeletier de Saint Fargeau, 1841). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (TM, TK, KM, AL), **ES** (KR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Afghanistan, Central Asia, Kazakhstan, China (NW).
- Halictus (Vestitohalictus) mucoreus** (Eversmann, 1852) [*Hylaeus*]. Russia: **EP** (S, NC), **UR**. – Georgia, Iran, Afghanistan, Central Asia, Kazakhstan.
- Halictus (Vestitohalictus) pollinosus cariniventris** Morawitz, 1876 (*H. carinaeventris* Fahringer et Friese, 1921; *H. c. creticola* Strand, 1921; *H. c. flavotectus* Cockerell, 1922; *H. pollinosus limmissicus* Blüthgen, 1937). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), Turkey, Israel, Iran, Afghanistan, Pakistan, Central Asia, Mongolia, China (NC, NW).
- Halictus (Vestitohalictus) pseudomucoreus** Ebmer, 1975. Russia: **EP** (S). – Azerbaijan, Turkey, Turkmenistan, Kazakhstan, China (NW).
- Halictus (Vestitohalictus) pulvereus** Morawitz, 1874 (*H. sogdianus* Morawitz, 1876; *Nomiooides aenescens* Radoszkowski, 1893). Russia: **EP** (S, NC, CR). – Cyprus, Turkey, Iran, Afghanistan, Central Asia, Mongolia, China (NW).
- Halictus (Vestitohalictus) radoszkovskii** Vachal, 1902 (*H. rufus* Friese, 1916). Russia: **EP** (NC). – Armenia, Turkmenistan, Uzbekistan, Iran, Afghanistan.
- Halictus (Vestitohalictus) tectus** Radoszkowski, 1876 (*H. vestitus* auct.). Russia: **EP** (S, CR), **UR**. – Europe (WE, SE, EE), Caucasus.
- Halictus (Vestitohalictus) tuberculatus** Blüthgen, 1925. Russia: **EP** (E, S, NC, CR). – Europe (EE), Armenia, Turkey, Syria, Israel.
- LASIOGLOSSUM** Curtis, 1833. Type species *Lasioglossum tricingulum* Curtis, 1833 (= *Melitta*

xanthopus Kirby, 1802). Widely distributed in the World, but most of species occur in Holarctic region. Number of taxa: World – about 1200 species, Palaearctic – about 450, Russia – 148.

Lasioglossum (Ctenonomia) blakistoni Sakagami et Munakata, 1990. Russia: **FE** (PR). – Japan (Hok).

Lasioglossum (Lasioglossum) aegyptiellum (Strand, 1909) [Halictus] (*Halictus divergens* Pérez, 1911; *H. orontis* Cockerell, 1918; *H. triopolitanus* Blüthgen in Schulthess, 1924; *Lasioglossum aegyptiellum dalmaticum* Ebmer, 1970). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Near East, Iraq, Iran, Turkmenistan.

Lasioglossum (Lasioglossum) agelastum Fan et Ebmer, 1992 (*L. koreanum*: Pesenko, 1986; *L. nipponicola* Sakagami et Tadauchi, 1995). Russia: **FE** (PR, KU). – China (NE, NC, CC, SW), Korean Peninsula, Japan (Hok, Hon).

Lasioglossum (Lasioglossum) alinense (Cockerell, 1924) [Halictus] (*Halictus lutzenkoi* Cockerell, 1925; *H. gorkiensis* Blüthgen, 1931). Russia: **WS** (AL), **ES** (IR), **FE** (AM, KH, PR). – Europe (NE, EE), China (NE, NC, CC, SW).

Lasioglossum (Lasioglossum) bicallosum (Morawitz, 1874) [Halictus] (*Halictus dmitrijewi* Morawitz, 1891; *H. dmitrijewii* Dalla Torre, 1896). Russia: **EP** (S, NC, CR). – Europe (SE, EE), Caucasus, Turkey, Israel, Iran.

Lasioglossum (Lasioglossum) caspicum (Morawitz, 1874) [Halictus] (*Halictus emesianus* Pérez, 1911). Russia: **EP** (NC). – Caucasus, Turkey, Near East, Iran, Afghanistan, Turkmenistan.

Lasioglossum (Lasioglossum) chloropus (Morawitz, 1894) [Halictus] (*L. aksuense* Ebmer, 1972). Russia: **ES** (IR). – Tajikistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NW).

Lasioglossum (Lasioglossum) costulatum (Kriechbaumer, 1873) [Halictus] (*Andrena campestris* Eversmann, 1852; *Halictus alpestris* Morawitz, 1877). Russia: **EP** (C, E), **UR**, **WS** (AL), **ES** (KR, IR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Syria, Iran, Central Asia, Kazakhstan.

Lasioglossum (Lasioglossum) cristula (Pérez, 1895) [Halictus] (*Halictus korbi* Blüthgen, 1929; *H. mesoleus* Cockerell, 1938; *H. cristula donatus* Warncke, 1975). Russia: **EP** (NC, CR). – Europe (WE, SE), N Africa, Caucasus, Turkey, Syria, Israel, Iran.

Lasioglossum (Lasioglossum) denticolle (Morawitz, 1891) [Halictus] (*Halictus glycybromifer* Strand,

1915; *H. laevifrons* Blüthgen, 1923 *H. morbillosus r. orientis* Cockerell, 1924). Russia: **ES** (KR, BR), **FE** (AM, KH, PR). – China (NE, NC, NW, CC, SW, WP), Korean Peninsula.

Lasioglossum (Lasioglossum) discum (Smith, 1853) [Halictus] (*Halictus morbillosus* Kriechbaumer, 1873; *H. major*: Morawitz, 1876; *H. fertoni* Vachal, 1895; *H. morbillosus r. glasunovi* Cockerell, 1924; *H. platycestus* auct.; *Lasioglossum pseudomorbillosum* Ebmer, 1970). Russia: **EP** (C, S, NC, CR), **UR**, **ES** (TU). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran, Afghanistan, Central Asia, Kazakhstan, China (NW).

Lasioglossum (Lasioglossum) eos Ebmer, 1978 (*L. kasparyani* Pesenko, 1986; *L. kerzhneri* Pesenko, 1986). Russia: **ES** (TU, IR, BR), **FE** (AM, PR). – Mongolia, China (NE, NC, NW).

Lasioglossum (Lasioglossum) exiliceps (Vachal, 1903) [Halictus]. Russia: **FE** (KH, PR). – Korean Peninsula, Japan (Hok, Hon, Kyu, Ryu).

Lasioglossum (Lasioglossum) euxinicum euxinicum Ebmer, 1972. Russia: **EP** (CR). – Azerbaijan, Iran.

Lasioglossum (Lasioglossum) fallax fallax (Morawitz, 1874) [Halictus]. Russia: **EP** (S, NC, CR), **UR**. – Georgia, Turkey, Iran, Turkmenistan.

Lasioglossum (Lasioglossum) harmandi (Vachal, 1903) [Halictus] (*L. nippонense*: Pesenko, 1986). Russia: **FE** (KU). – Japan (Hok, Hon, Ryu).

Lasioglossum (Lasioglossum) kansuense (Blüthgen, 1934) [Halictus] (*L. esoense* Hirashima et Sakagami, 1966). Russia: **ES** (BR), **FE** (AM, KH, PR, KU). – China (NE, NC, CC, SW, WP), Korean Peninsula, Japan (Hokk, Hon, Ryu).

Lasioglossum (Lasioglossum) korbi (Blüthgen, 1929) [Halictus] (*Halictus haesitans* Blüthgen, 1931). Russia: **EP** (NC). – Azerbaijan, Turkey.

Lasioglossum (Lasioglossum) kussariense (Blüthgen, 1925) [Halictus]. Russia: **EP** (NC, CR). – Europe (SE, EE), Caucasus, Turkey.

Lasioglossum (Lasioglossum) laevigatum (Kirby, 1802) [Melitta] (*Melitta lugubris* Kirby, 1802; *Halictus levigatus* Lepeletier de Saint Fargeau, 1841; *Hylaeus semicinctus* Förster, 1860). Russia: **EP** (C), **UR**. – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran.

Lasioglossum (Lasioglossum) lativentre (Schenck, 1853) [Hylaeus] (*Hylaeus bisbistrigatus* Schenck, 1853; *Halictus bicinctus* Schenck, 1874; *H. decipiens* Perkins, 1913; *H. lativentris sotschica* Blüthgen, 1931). Russia: **EP** (NW, C, E, NC, CR). – Europe

(WE, NE, SE, EE), Caucasus, Turkey, Near East, Iran.

Lasioglossum (Lasioglossum) leviventre (Pérez, 1905) [Halictus]. Russia: **FE** (PR, KU). – Japan (Hok, Hon, Izu, Shi, Ryu).

Lasioglossum (Lasioglossum) leucozonium (Schrank, 1781) [Apis] (*Apis leucostoma* Schrank, 1781; *Halictus similis* Smith, 1853; *H. bifasciatellus* Schenck, 1875; *H. leucozonius nigrotibialis* Dalla Torre, 1877; *H. deiphobus* Bingham, 1908; *H. tadschicus* Blüthgen, 1929; *H. satschauensis* Blüthgen, 1934; *H. leucozonius clusius* Warncke, 1975; *Lasioglossum leucozonium cedri* Ebmer, 1976; *L. zonulum elysium* Ebmer, 1979; *L. satschauense mandschuricum* Ebmer, 1978; *L. zonulum xylopedis* Ebmer, 1978). Russia: **EP** (N, NW, C, E, S, CR), **UR**, **ES** (TU, KS, IR, BR), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Near East, Afghanistan, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NE, NC, CC, SW, WP), N India, N America.

Lasioglossum (Lasioglossum) niveocinctum (Blüthgen, 1923) [Halictus]. **EP** (S, NC). – Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NC, NW).

Lasioglossum (Lasioglossum) occidens (Smith, 1873) [Halictus] (*Halictus quadraticollis* Vachal, 1903; *Lasioglossum koreanum* Ebmer, 1978). Russia: **FE** (SA, KU). – China (NC, CC, SW, WP, SE), Japan (Hok, Hon, Izu, Kyu).

Lasioglossum (Lasioglossum) pallens (Brullé, 1932) [Halictus] (*Halictus lineolatus* Lepoletier de Saint Fargeau, 1841; *H. cirrhozonius* Vachal, 1895; *H. pallens kantarae* Warncke, 1982). Russia: **EP** (NC). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Iran.

Lasioglossum (Lasioglossum) proximatum (Smith, 1879) [Halictus] (*Halictus discrepans* Pérez, 1905; *H. moltrechi* Cockerell, 1925; *H. kraloffi* Cockerell, 1925; *H. emelianoffi* Cockerell, 1925; *Lasioglossum acuticrista* Pesenko, 1986). Russia: **FE** (AM, KH, PR, KU). – China (NE, NC, CC, SW, WP), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu).

Lasioglossum (Lasioglossum) quadrinotatum (Kirby, 1802) [Melitta]. Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Kazakhstan.

Lasioglossum (Lasioglossum) rostratum (Eversmann, 1852) [Hylaeus] (*Halictus chlapovskii* Vachal, 1902). Russia: **UR**, **WS** (TM, TK, OM, AL), **ES**

(TU, KR, IR, ZB), **FE** (AM, KH, PR). – Kazakhstan, Mongolia, China (NE, NC, NW, WP).

Lasioglossum (Lasioglossum) scitulum (Smith, 1873) [Halictus] (*Halictus japonica* Strand, 1910; *H. basicirrus* Cockerell, 1919). Russia: **FE** (AM, KH, PR, SA, KU). – China (NE, NC, NW, CC, SW), Korean Peninsula, Japan (Hok, Hon, Ryu).

Lasioglossum (Lasioglossum) sexnotatum (Nylander, 1852) [Halictus]. Russia: **EP** (N, NW, C, E), **UR**, **WS** (OM). – Europe (WE, NE, SE, EE), Kazakhstan.

Lasioglossum (Lasioglossum) sexnotatum (Kirby, 1802) [Melitta] (*Andrena nitida* Panzer, 1798). Russia: **EP** (NW, C, E, NC, CR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Mongolia.

Lasioglossum (Lasioglossum) sutshanicum Pesenko, 1986. Russia: **FE** (PR). – China (SE), Korean Peninsula.

Lasioglossum (Lasioglossum) tungusicum Ebmer, 1978 (*Halictus tinnunculus* Warncke, 1982; *Lasioglossum sexmaculatum*: Pesenko, 1986). Russia: **EP** (NW, C, E, S), **UR**, **WS** (TM, AL), **ES** (IR, YA), **FE** (AM). – Caucasus, Turkey, Iran, Kazakhstan, Mongolia, China (NW, SW, WP).

Lasioglossum (Lasioglossum) upinense (Morawitz, 1889) [Halictus] (*Halictus carbonarius* Blüthgen, 1923; *H. tacitus* Cockerell, 1924; *H. carbonatus* Blüthgen, 1925; *H. wittenbourgi* Cockerell, 1925). Russia: **FE** (AM, KH, PR). – China (NE, NC, NW, SW, WP), Korean Peninsula.

Lasioglossum (Lasioglossum) xanthopus (Kirby, 1802) [Melitta] (*Apis emarginata* Christ, 1791; *Hylaeus derasus* Imhoff, 1832; *Lasioglossum tricinulum* Curtis, 1833; *Hylaeus fulvicrus* Eversmann, 1852; *Halictus soreli* Dours, 1872). Russia: **EP** (N, C, E, S, NC, CR), **UR**. – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NW).

Lasioglossum (Lasioglossum) zeyanense Pesenko, 1986 (*L. acervolum* Fan et Ebmer, 1992). Russia: **ES** (BR, YA, ZB), **FE** (AM, KH, PR). – China (NC, CC, SW, WP).

Lasioglossum (Lasioglossum) zonulum (Smith, 1848) [Halictus] (*Hylaeus trifasciatus* Schenck, 1853; *Halictus rhenanus* Verhoeff, 1890; *H. receptarius* Vachal, 1902; *H. craterus* Lovell, 1908; *H. zonulus sinistralis* Blüthgen, 1934; *H. z. dexter* Blüthgen, 1934; *Lasioglossum zonulum euronotum* Ebmer, 1998). Russia: **EP** (N, NW, C, E, S), **UR**, **WS** (TM), **ES** (KR, IR, BR). – Europe (WE, NE, SE, EE),

- Caucasus, Turkey, Iran, Central Asia, Kazakhstan, China (NC, NW, SW), N America.
- Lasioglossum (Evylaeus) aeratum** (Kirby, 1802) [Melitta] (*Halictus semiaeneus* Brullé, 1832; *H. viridiaeetus* Blüthgen, 1918; *H. aeratus caudatus* Warncke, 1982). Russia: **EP** (NW, C, E, S, NC). – Europe (WE, NE, SE, EE), Turkey, Syria, Iran, Central Asia, Kazakhstan.
- Lasioglossum (Evylaeus) affine** (Smith, 1853) [*Halictus*] (*Halictus leoninus* Vachal, 1903; *H. mandarinus* Strand, 1910; *H. nagasakiensis* Strand, 1910; *H. investigator* Strand, 1910; *H. investigatoris* Strand, 1915). Russia: **FE** (KH, PR). – China (NE, NC, CC, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu, Tsu).
- Lasioglossum (Evylaeus) albipes albipes** (Fabricius, 1781) [Apis] (*Hylaeus abdominalis* Panzer, 1798; *Halictus malachurellus* Strand, 1909; *H. albipes alpicola* Blüthgen, 1921; *H. a. rubelloides* Blüthgen, 1924). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (KM, AL), **ES** (BR, KR, IR, ZB), **FE** (KH, PR, SA, KU). – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, Iran, Central Asia, Mongolia, China (NE, NC), Korean Peninsula, Japan (Hok, Hon).
- Lasioglossum (Evylaeus) albipes villosum** Ebmer, 1995. Russia: **FE** (AM, PR). – Korean Peninsula.
- Lasioglossum (Evylaeus) allodatum** Ebmer et Sakagami, 1985. Russia: **FE** (AM, KH, PR). – China (NC, SW), Korean Peninsula, Japan (Hok, Hon).
- Lasioglossum (Evylaeus) amurense** (Vachal, 1902) [*Halictus*]. Russia: **WS** (KM), **FE** (KH, PR). – China (NE, NC).
- Lasioglossum (Evylaeus) anellum** (Vachal, 1905) [*Halictus*] (*Halictus schelkovnikovi* Kokujev, 1912). Russia: **EP** (S, NC). – Europe (SE, EE), Caucasus, Turkey, Syria, Jordan, Israel, Iran.
- Lasioglossum (Evylaeus) angaricum** (Cockerell, 1937) [*Halictus*]. Russia: **ES** (ZB), **FE** (AM, PR). – Mongolia.
- Lasioglossum (Evylaeus) annulipes** (Morawitz, 1876) [*Halictus*] (*Halictus metopias* Vachal in Kohl, 1903). Russia: **EP** (NC). – Armenia, Turkey, Afghanistan, Pakistan, Tajikistan, Uzbekistan, Kazakhstan.
- Lasioglossum (Evylaeus) apristum** (Vachal, 1903) [*Halictus*] (*Sphecodes pallidulus* Matsumura, 1912). Russia: **FE** (PR, KU). – China (SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu).
- Lasioglossum (Evylaeus) baleicum baleicum** (Cockerell, 1937) [*Halictus*]. Russia: **ES** (IR, ZB), **FE** (PR). – China (NE), Korean Peninsula.
- Lasioglossum (Evylaeus) baleicum insulicolum** (Pesenko, 2007) [*Evylaeus*]. Russia: **FE** (SA, KU). – Japan (Hok, Hon, Kyu, Ryu).
- Lasioglossum (Evylaeus) bavaricum olympicum** (Warncke, 1982) [*Halictus*]. Russia: **EP** (CR). – Europe (SE).
- Lasioglossum (Evylaeus) bluethgeni** (Ebmer, 1971) [*Halictus*] (*Halictus hirtiventris* Blüthgen, 1923). Russia: **EP** (NC). – Europe (WE, SE, EE), Georgia, Turkey.
- Lasioglossum (Evylaeus) brachycephalum** (Cockerell, 1925) [*Halictus*]. Russia: **FE** (PR).
- Lasioglossum (Evylaeus) briseis briseis** Ebmer, 2005. Russia: **ES** (TU). – Mongolia.
- Lasioglossum (Evylaeus) briseis eomontanum** Ebmer, 2006. Russia: **ES** (IR, YA), **FE** (PR, MG, CH).
- Lasioglossum (Evylaeus) brevicorne brevicorne** (Schenck, 1869) [*Halictus*] (*Halictus lucidicollis* Pérez, 1895; *H. analis* Pérez, 1903; *H. brevicornis aciculatus* Blüthgen in Schmiedeknecht, 1930; *H. optimellus* Cockerell, 1938). Russia: **EP** (C, E, S). – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey, Israel, Iran, Afghanistan.
- Lasioglossum (Evylaeus) buccale** (Pérez, 1903) [*Halictus*]. Russia: **EP** (S), **UR**. – Europe (WE, SE, EE), Turkey, Iran.
- Lasioglossum (Evylaeus) calceatum** (Scopoli, 1763) [Apis] (*Hylaeus cylindricus* Fabricius, 1793; *Melitta fulvocincta* Kirby, 1802; *M. obovata* Kirby, 1802; *Andrena vulpina* Fabricius, 1804; *Halictus terebrator* Walckenaer, 1817; *Hylaeus rubellus* Eversmann, 1852; *H. bipunctatus* Schenck, 1853; *Halictus rubens* Smith, 1854; *H. rufiventris* Giraud, 1861; *H. cylindricus rhodostoma* Dalla Torre, 1877; *H. calceatus ulterior* Cockerell, 1929). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (AL), **ES** (KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Iran, Central Asia, Kazakhstan, Mongolia, China (NE, NC), Korean Peninsula, Japan (Hok, Hon, Kyu).
- Lasioglossum (Evylaeus) caliginosum** Murao, Ebmer et Tadauchi, 2006 (*L. nemorae* Ebmer, 2006). Russia: **ES** (IR), **FE** (AM, KH, PR). – Mongolia, China (NE), Japan (Hok, Hon, Kyu).
- Lasioglossum (Evylaeus) clypeare** (Schenck, 1853) [*Hylaeus*]. Russia: **EP** (S, NC, CR), **UR**. – Europe (WE, SE, EE), Turkey, Iran, Kyrgyzstan, Kazakhstan.
- Lasioglossum (Evylaeus) convexiusculum** (Schenck, 1853) [*Hylaeus*] (*Halictus appropinquans* Schenck,

- 1869; *H. clypeatus* Schenck, 1869; *H. porcus* Morawitz, 1872; *H. puncticollis genevensis* Frey-Gessner, 1903). Russia: **EP** (C, S), **UR**. – Europe (WE, NE, SE, EE), Turkey, Iran.
- Lasioglossum (Evylaeus) corvinum** (Morawitz, 1877) [Halictus]. Russia: **EP** (S, NC, CR). – Europe (WE, EE, SE), N Africa, Caucasus, Turkey, Iran.
- Lasioglossum (Evylaeus) crassepunctatum** (Blüthgen, 1923) [Halictus]. Russia: **EP** (C, S, NC, CR). – Europe (WE, SE, EE), Azerbaijan, Turkey, Jordan, Israel, Iran.
- Lasioglossum (Evylaeus) damascenum** (Pérez, 1911) [Halictus] (*Halictus semitomentosus* Blüthgen, 1923). Russia: **EP** (CR). – Europe (SE, EE), Armenia, Turkey, Syria, Israel, Iran.
- Lasioglossum (Evylaeus) duckei duckei** (Alfken, 1909) [Halictus]. Russia: **EP** (S, CR). – Europe (SE, EE), Turkey.
- Lasioglossum (Evylaeus) dybowskii** (Radoszkowski, 1877) [Halictus] (*Halictus dubowスキi* Dalla Torre, 1896; *H. griseipennis* Cockerell, 1924). Russia: **FE** (AM, KH, PR, SA). – China (NE), Korean Peninsula.
- Lasioglossum (Evylaeus) elegans** (Lepeletier de Saint Fargeau, 1841) [Halictus] (*Halictus rubellus* Giraud, 1861; *H. giraudii* Dalla Torre et Friese, 1895). Russia: **EP** (S). – Europe (WE, SE, EE), N Africa, Georgia, Turkey, Iran, Turkmenistan.
- Lasioglossum (Evylaeus) ellipticeps** (Blüthgen, 1923) [Halictus] (*Halictus permicus* Blüthgen, 1923; *H. mayacensis* Cockerell, 1924). Russia: **UR**, **ES** (IR, YA), **FE** (KH, PR). – Mongolia, Korean Peninsula.
- Lasioglossum (Evylaeus) epiphron** Ebmer, 1982. Russia: **ES** (IR), **FE** (AM, KH, PR). – Mongolia.
- Lasioglossum (Evylaeus) eriphyle** Ebmer, 1996. Russia: **FE** (AM, KH, PR, KU). – China (CC).
- Lasioglossum (Evylaeus) euboeense** (Strand, 1909) [Halictus] (*Halictus furnasensis* Strand, 1909; *H. kirschbaumi* Blüthgen, 1919; *H. euboeensis anatolica* Blüthgen, 1931). Russia: **EP** (NW, E, S, NC). – Europe (WE, SE, EE), N Africa, Azerbaijan, Turkey, Iran.
- Lasioglossum (Evylaeus) fratellum betulae** Ebmer, 1978 (*Lasioglossum nupricola* Sakagami, 1988). Russia: **ES** (YA), **FE** (AM, KH, PR, KA, SA, KU). – Mongolia, Korean Peninsula, Japan (Hok, Hon).
- Lasioglossum (Evylaeus) fratellum fratellum** (Pérez, 1903) [Halictus] (*Halictus subfasciatus* Nylander, 1848; *H. freygessneri* Alfken, 1905; *H. norvegicus* Strand, 1910; *H. niger* auct.; *Lasioglossum nigrum* auct.). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (KM, AL), **ES** (KR, IR, ZB). – Europe (WE, NE, SE, EE).
- Lasioglossum (Evylaeus) fulvicorne fulvicorne** (Kirby, 1802) [Melitta] (*Hylaeus laeviusculus* Schenck, 1853; *H. albifrons* Schenck, 1853; *H. bimaculatus* Schenck, 1853; *Halictus opacifrons* Pérez, 1911). Russia: **EP** (N, NW, C, E, NC, CR), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Turkmenistan.
- Lasioglossum (Evylaeus) fulvicorne melanocorne** Ebmer, 1988. Russia: **WS** (AL), **ES** (KR, IR, YA), **FE** (AM, PR, SA, KU). – Mongolia.
- Lasioglossum (Evylaeus) gilanum** (Blüthgen, 1931) [Halictus]. Russia: **EP** (CR). – Iran.
- Lasioglossum (Evylaeus) gorge** Ebmer, 1982. Russia: **WS** (AL), **ES** (YA). – Mongolia.
- Lasioglossum (Evylaeus) glabriusculum glabriusculum** (Morawitz, 1872) [Halictus] (*Halictus leucopygus* Pérez, 1903; *H. truncatus* Alfken, 1905; *H. granulosus* Alfken, 1906). Russia: **EP** (S, NC). – Europe (WE, SE, EE), Georgia, Turkey, Syria, Jordan, Israel, Iran.
- Lasioglossum (Evylaeus) griseolum griseolum** (Morawitz, 1872) [Halictus] (*Halictus labrosus* Vachal, 1895; *H. misellus* Pérez, 1903; *H. dubitabilis* Saunders, 1904). Russia: **EP** (S, NC). – Europe (WE, SE), Georgia, Turkey, Israel, Iran, Afghanistan.
- Lasioglossum (Evylaeus) hoffmanni** (Strand, 1915) [Halictus] (*Halictus atropis* Strand, 1915; *H. shishkini* Cockerell, 1925; *H. suprafulgens* Cockerell, 1925; *H. speculicaudatus* Cockerell, 1931). Russia: **FE** (KH, AM, PR). – China (NE, NC), Korean Peninsula, Japan (Hon, Hon, Shi, Kyu, Tsu).
- Lasioglossum (Evylaeus) intermedium** (Schenck, 1869) [Halictus] (*Halictus melanoproctus* Pérez, 1903; *H. servulellus* Strand, 1909). Russia: **EP** (NC). – Europe (WE, NE, SE, EE), Turkey, Iran.
- Lasioglossum (Evylaeus) interruptum interruptum** (Panzer, 1798) [Hylaeus] (*Apis basimacula* Schrank, 1802; *Hylaeus geminus* Erichson, 1835; *H. quadrimaculatus* Schenck, 1853). Russia: **EP** (C, E, S). – Europe (SE, WE, NE, EE), N Africa.
- Lasioglossum (Evylaeus) interruptum trispinosus** (Alfken, 1907) [Halictus]. Russia: **EP** (CR). – Europe (EE), Georgia, Armenia, Turkey, Iran.
- Lasioglossum (Evylaeus) kankaucharis** (Strand, 1914) [Halictus]. Russia: **FE** (PR). – China (SE), Korean Peninsula, Japan (Hon).
- Lasioglossum (Evylaeus) kiautschouense** (Strand, 1910) [Halictus]. Russia: **FE** (KH, PR). – China (NE, NC), Korean Peninsula, Japan (Hok).

Lasioglossum (Evylaeus) kirgisicum Ebmer, 1972. Russia: **WS** (AL). – Europe (EE), Turkey, Kazakhstan.

Lasioglossum (Evylaeus) kozlovi (Friese, 1913) [Halictus]. Russia: **ES** (BR). – China (NC, NW).

Lasioglossum (Evylaeus) laeve (Kirby, 1802) [Melitta] (*Hylaeus convexus* Schenck, 1853; *Halictus nigritiventris* Arnold, 1894; *H. levis* Dalla Torre, 1896; *H. kriegeri* Alfken, 1897). Russia: **EP** (C, E), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), Georgia, Armenia, Turkey, Syria, Israel, Iran.

Lasioglossum (Evylaeus) laevoides Ebmer, 2005. Russia: **ES** (IR, BR). – Mongolia, China (NE), Korean Peninsula.

Lasioglossum (Evylaeus) laticeps laticeps (Schenck, 1869) [Halictus] (*Hylaeus affinis* Schenck, 1853; *H. nigricornis* Schenck, 1853; *Halictus mendax* Alfken, 1912). Russia: **EP** (C, E, NC, CR), **UR**. – Europe (WE, NE, SE, EE).

Lasioglossum (Evylaeus) leucopum (Kirby, 1802) [Melitta]. Russia: **EP** (N, NW, C, E, NC, CR), **UR**, **WS** (OM, KM), **ES** (KR, IR, BR, YA). – Europe (WE, NE, SE, EE), Turkey, Iran.

Lasioglossum (Evylaeus) lineare (Schenck, 1869) [Halictus] (*Halictus longuloides* Strand, 1909; *H. smyrnae* Strand, 1909; *H. aeginus* Strand, 1921; *Lasioglossum sublineare* Ebmer, 1974). Russia: **EP** (C, E, NC, CR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Syria, Israel, Iran, Turkmenistan.

Lasioglossum (Evylaeus) limbellum (Morawitz, 1876) [Halictus] (*Halictus limbellus dongarica* Blüthgen, 1934). Russia: **EP** (NC). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran, Afghanistan, Tajikistan, Uzbekistan, China (NC).

Lasioglossum (Evylaeus) lissonotum (Noskiewicz, 1926) [Halictus]. Russia: **EP** (C, E). – Europe (WE, SE, EE).

Lasioglossum (Evylaeus) littorale colchicum Ebmer, 1972. Russia: **EP** (CR). – Europe (SE), Turkey.

Lasioglossum (Evylaeus) lucidulum (Schenck, 1861) [Hylaeus] (*Hylaeus tenellus* Schenck, 1861; *Halictus gracilis* Morawitz, 1865; *H. unguinosus* Pérez, 1903; *H. chotanensis* Strand, 1909). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (KR). – Europe (WE, NE, SE, EE), Georgia, Turkey, Israel, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NC, NW).

Lasioglossum (Evylaeus) malachurum (Kirby, 1802) [Melitta] (*Halictus longulus* Smith, 1848; *Hylaeus apicalis* Schenck, 1853; *H. coriarius* Schenck, 1853

Halictus malachuroides Strand, 1909; *H. malachurus sharificus* Cockerell, 1937). Russia: **EP** (E, S, CR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran, Turkmenistan.

Lasioglossum (Evylaeus) mandibulare (Morawitz, 1866) [Halictus] (*Halictus coloratus* Morawitz, 1874; *H. czeckelii* Friese, 1916; *H. sareptanus* Blüthgen, 1923). Russia: **EP** (S, NC). – Europe (WE, SE, EE), Armenia, Turkey, Israel.

Lasioglossum (Evylaeus) marginatum (Brullé, 1832) [Halictus] (*Halictus fasciatellus* Schenck, 1869; *H. gribodoi* Kriechbaumer, 1873; *H. riparius* Morawitz, 1874; *H. vulgaris* Morawitz, 1876; *H. kervileanus* Pérez, 1910). Russia: **EP** (E, NC). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Syria, Jordan, Israel, Iraq, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, NW India, Nepal.

Lasioglossum (Evylaeus) marginellum (Schenck, 1853) [Halictus] (*Halictus marqueti* Pérez, 1903). Russia: **EP** (S, CR). – Europe (WE, SE, EE).

Lasioglossum (Evylaeus) matianense pluto Ebmer, 1980. Russia: **FE** (PR). – Uzbekistan, Kyrgyzstan, Kazakhstan.

Lasioglossum (Evylaeus) mesosclerum (Pérez, 1903) [Halictus] (*Halictus rhodosianus* Strand, 1909; *H. mesoclerus* Blüthgen, 1935; ?*Lasioglossum balneorum* Ebmer, 1974; *Halictus bubulcus* Warncke, 1982). Russia: **EP** (S, NC). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Syria, Jordan, Israel, Iran, Afghanistan, Central Asia, Kazakhstan.

Lasioglossum (Evylaeus) minutissimum (Kirby, 1802) [Melitta] (*Hylaeus exilis* Schenck, 1861; *Halictus hollandi* Saunders, 1904; *H. costiferellus* Strand, 1909; *H. kosensis* Strand, 1909; *H. xanthosensis* Strand, 1909; *H. arnoldi* Saunders, 1910; *H. lucidellus* Cockerell, 1937; *H. lilliput* Benoist, 1962). Russia: **EP** (C, E, CR), **UR**. – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel.

Lasioglossum (Evylaeus) minutuloides Ebmer, 1978. Russia: **FE** (PR). – China (NE).

Lasioglossum (Evylaeus) miyabei Murao, Ebmer et Tadauchi, 2006. Russia: **FE** (PR, KU). – Korean Peninsula, Japan (Hok, Hon, Kyu).

Lasioglossum (Evylaeus) monstrificum (Morawitz, 1891) [Halictus]. Russia: **ES** (IR).

Lasioglossum (Evylaeus) morio (Fabricius, 1793) [Hylaeus] (*Halictus morio basalis* Dalla Torre, 1877; *H. balticus* Blüthge, 1919). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Israel.

Lasioglossum (Evylaeus) nigripes (Lepeletier de Saint Fargeau, 1841) [Halictus] (*Halictus vulpinus* Nylander, 1852; *H. vulpinus nigroclypeata* Dalla Torre, 1877; *H. cylindricus orientalis* Magretti, 1890; *H. vulpinus nigriclypeatus* Dalla Torre, 1896; *H. nylanderi* Pérez, 1903; *H. pharaonis* Strand, 1909; *H. syriacus* Pérez, 1910). Russia: EP (S, NC, CR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Jordan, Israel, Iran, Turkmenistan.

Lasioglossum (Evylaeus) nipponense (Hirashima, 1953) [Halictus]. Russia: FE (AM, KH, PR, SA, KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu).

Lasioglossum (Evylaeus) nitidiusculum nitidiusculum (Kirby, 1802) [Melitta] (*Hylaeus rugosula* Schenck, 1853; *H. rugosula* Schenck, 1853; *H. pusillus* Schenck, 1853; *Halictus nitidulus* Pérez, 1903; *H. arnoldi* Saunders, 1910; *H. obsoletus* Warncke, 1975). Russia: EP (N, NW, C, E, NC), UR. – Europe (WE, NE, SE, EE), N Africa, Turkey, Iran.

Lasioglossum (Evylaeus) nitidulum nitidulum (Fabricius, 1804) [Prosopis] (*Halictus aeneidorsum* Alfken, 1920; *H. fudakowskii* Noskiewicz, 1926; *H. smeachmanellus* auct.; *H. continentalis* Blüthgen, 1944). Russia: EP (NW, C, S.), UR. – Europe (WE, NE, SE, EE), Turkey.

Lasioglossum (Evylaeus) nodicorne (Morawitz, 1889) [Halictus]. Russia: ES (IR, ZB). – Mongolia, China (NC).

Lasioglossum (Evylaeus) obscuratum obscuratum (Morawitz, 1876) [Halictus]. Russia: EP (NC). – Azerbaijan, Turkey, Syria, Jordan, Israel, Iran, Afghanistan, Central Asia, Kazakhstan.

Lasioglossum (Evylaeus) obscuratum acerbum (Warncke, 1975) [Halictus]. Russia: EP (S, CR). – Europe (WE, SE, EE).

Lasioglossum (Evylaeus) pallidum (Radoszkowski, 1888) [Nomioides] (*Halictus sahlbergi* Friese, 1916). Russia: EP (S). – Afghanistan, Turkmenistan.

Lasioglossum (Evylaeus) pallilolum (Strand, 1914) [Halictus]. Russia: FE (KH, PR). – China (NE, SE), Korean Peninsula, Japan (Hok, Hon, Shi).

Lasioglossum (Evylaeus) parvulum (Schenck, 1853) [Hylaeus] (*Melitta minuta*: Kirby, 1802; *Halictus striatus* Schenck, 1869; *H. striatus* Schenck, 1869; *H. ferrugineipes* Schenck, 1869; *H. minutus* auct., before 1974; *Lasioglossum minutum*: Ebmer, 1971). Russia: EP (N, NW, C, NC), UR. – Europe (WE, NE, SE, EE), Armenia, Turkey, Iran.

Lasioglossum (Evylaeus) pauxillum (Schenck, 1853) [Hylaeus] (*Hylaeus fulvicornis*: Schenck, 1851; *H. similis* Schenck, 1853; *H. immarginatus* Schenck, 1853; *H. flavidicornis* Schenck, 1853; *H. immarginatus fuscitarsis* Schenck, 1853; *Halictus delicatus* Pérez, 1903; *H. algericolellus* Strand, 1909; *H. luteostigmatellus* Strand, 1909; *H. pseudopunctulatus* Strand, 1909; *H. schulthessi* Blüthgen in Schulthess, 1924). Russia: EP (C, E, S, NC, CR), UR, WS (KM). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Turkmenistan.

Lasioglossum (Evylaeus) peregrinum (Blüthgen, 1923) [Halictus]. Russia: EP (CR). – Europe (WE, SE, EE), Caucasus, Turkey, Israel, Iran.

Lasioglossum (Evylaeus) perplexans (Cockerell, 1925) [Halictus]. Russia: FE (KH, PR).

Lasioglossum (Evylaeus) podolicum podolicum (Noskiewicz, 1925) [Halictus] (*Halictus smeachmanellus nucleolus* Warncke, 1982; *H. smeachmanellus petrosus* Warncke, 1975). Russia: EP (C, CR). – Europe (WE, SE, EE), Turkey.

Lasioglossum (Evylaeus) politum politum (Schenck, 1853) [Hylaeus] (*Hylaeus nanulus* Schenck, 1853; *Halictus granulosus* Alfken, 1906). Russia: EP (C, E, S, NC, CR), UR. – Europe (WE, SE, EE).

Lasioglossum (Evylaeus) problematicum (Blüthgen, 1923) [Hylaeus]. Russia: FE (SA, KU). – Korean Peninsula, Japan (Hok, Hon).

Lasioglossum (Evylaeus) punctatissimum punctatissimum (Schenck, 1853) [Hylaeus] (*Hylaeus flavidarsis* Schenck, 1853; *Halictus simillimus* Schenck, 1869; *H. porcus* Morawitz, 1872; *H. griseescens* Schenck, 1873; *H. longiceps* Saunders, 1879; *H. tinitinensis* Cockerell, 1938). Russia: EP (N, NW, C, E, NC), UR. – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran.

Lasioglossum (Evylaeus) puncticolle (Morawitz, 1872) [Halictus] (*Halictus puncticollis* Saunders, 1879; *H. saundersii* Dalla Torre et Friese, 1895). Russia: EP (C, S, NC, CR), UR. – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, Syria, Iran.

Lasioglossum (Evylaeus) pygmaeum patulum (Vachal, 1905) [Halictus] (*Halictus spretus* Pérez, 1911; *H. denislucus*: Blüthgen, 1925; *H. andinus* Warncke, 1982). Russia: EP (S, CR). – Europe (SE, EE), Caucasus, Turkey, Syria, Jordan, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan.

Lasioglossum (Evylaeus) quadrinotatum (Schenck, 1861) [Halictus] (*Halictus pallipes* Morawitz, 1865; *H. megacephalus* Schenck, 1869; *H. sexsignatus*

- Schenck, 1869; *H. pallidipes* Dalla Torre, 1896). Russia: **EP** (N, NW, C, E, S), **UR**, **ES** (KR, BR, YA, ZB). – Europe (WE, NE, SE, EE), Turkey, China (NC).
- Lasioglossum (Evylaeus) quadrisignatum** (Schenck, 1853) [*Hylaeus*] (*Hylaeus rufipes* Schenck, 1853; *H. atratulus* Schenck, 1853; *Halictus pleuralis* Morawitz, 1872; *H. nigerrimus* Schenck, 1875; *H. com mixtus* Dalla Torre et Friese, 1895). Russia: **EP** (C, S, E, CR). – Europe (WE, SE, EE), Caucasus, Turkey, Iran.
- Lasioglossum (Evylaeus) rufitarse** (Zetterstedt, 1838) [*Halictus*] (*Halictus parumpunctatus* Schenck, 1869; *H. lucidus* Schenck, 1869; *H. atricornis* Smith, 1870). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (TM, KM, AL), **ES** (IR, YA, ZB), **FE** (AM, PR, KA, SA, KU, MG). – Europe (WE, NE, SE, EE), Caucasus, Iran, Kazakhstan, Mongolia, China (NE, SE), Korean Peninsula
- Lasioglossum (Evylaeus) sabulosus** (Warncke, 1986) [*Halictus*]. Russia: **EP** (C, E). – Europe (WE, NE, SE, EE).
- Lasioglossum (Evylaeus) sakagamii** Ebmer, 1978. Russia: **FE** (KH, PR). – Mongolia, China (NE, NC, NW), Korean Peninsula, Japan (Hok, Hon).
- Lasioglossum (Evylaeus) samaricum** (Blüthgen, 1935) [*Halictus*] (*L. pendschakenticum*: Ebmer, 1974). Russia: **EP** (E, CR). – Europe (EE), Turkey, Turkmenistan.
- Lasioglossum (Evylaeus) scirpaceum** (Warncke, 1975) [*Halictus*]. Russia: **EP** (NC). – Turkey, Iran.
- Lasioglossum (Evylaeus) semilaeve** (Blüthgen, 1923) [*Halictus*] (*Halictus minutulus speculiferus* Cockerell, 1937; *H. minutulus speculigerus* Cockerell, 1938). Russia: **WS** (AL), **ES** (IR), **FE** (AM, KH, PR). – Mongolia, China (NE).
- Lasioglossum (Evylaeus) semilucens** (Alfken, 1914) [*Halictus*] (*Hylaeus pygmaeus* Schenck, 1861). Russia: **EP** (C, E, NC). – Europe (WE, NE, SE, EE), Azerbaijan, Afghanistan, Tajikistan, Kyrgyzstan.
- Lasioglossum (Evylaeus) setulellum** (Strand, 1909) [*Halictus*] (*Halictus foveolatus* Blüthgen, 1924; *H. cinclus* Warncke, 1984). Russia: **EP** (E, S, CR), **UR**. – Europe (WE, SE, EE), Armenia, Turkey, Syria, Iran.
- Lasioglossum (Evylaeus) sexstrigatum** (Schenck, 1869) [*Halictus*]. Russia: **EP** (C, E), **UR**. – Europe (WE, NE, SE, EE), Armenia, Turkey.
- Lasioglossum (Evylaeus) sibiriacum** (Blüthgen, 1923) [*Halictus*] (*Halictus solovieffi* Cockerell, 1925). Russia: FE (PR). – China (NE), Korean Peninsula, Japan (Hok, Hon, Shi, Izu, Tsu, Ryu).
- Lasioglossum (Evylaeus) simplicior** (Cockerell, 1931) [*Halictus*]. Russia: **FE** (KH, PR). – China (SE), Korean Peninsula, Japan (Izu)
- Lasioglossum (Evylaeus) speculinum** (Cockerell, 1925) [*Halictus*] (*Halictus pallilomus* Strand, 1914; *H. perplexans speculinus* Cockerell, 1925). Russia: **FE** (PR). – China (NE), Korean Peninsula.
- Lasioglossum (Evylaeus) subaenescens subaenescens** (Pérez, 1895) [*Halictus*] (*Lasioglossum illyricum* Ebmer, 1971). Russia: **EP** (CR). – Europe (WE, SE, EE).
- Lasioglossum (Evylaeus) subfulvicorne austriacum** Ebmer, 1974. Russia: **EP** (NC). – Europe (WE, SE, EE), Georgia, Turkey, Kyrgyzstan, Kazakhstan.
- Lasioglossum (Evylaeus) subfulvicorne subfulvicorne** (Blüthgen, 1934) [*Halictus*]. Russia: **WS** (KR, YA, ZB), **FE** (AM, PR, KA, SA, KU). – Mongolia, China (NC, NW), Korean Peninsula.
- Lasioglossum (Evylaeus) sulcatulum longifacies** Sakagami et Tadauchi, 1995. Russia: **FE** (KU). – Japan (Hok, Hon, Tsu).
- Lasioglossum (Evylaeus) sulcatulum sulcatulum** (Cockerell, 1925) [*Halictus*]. Russia: **FE** (AM, KH, PR). – Korean Peninsula.
- Lasioglossum (Evylaeus) tarsatum** (Schenck, 1869) [*Halictus*] (*Halictus berolinensis* Strand, 1909). Russia: **EP** (NW, C, E, S). – Europe (WE, NE, SE, EE), Afghanistan.
- Lasioglossum (Evylaeus) tauricum** Ebmer, 1972. Russia: **EP** (NC, CR).
- Lasioglossum (Evylaeus) transpositum** (Cockerell, 1925) [*Halictus*] (*Halictus tutihensis* Cockerell, 1925). Russia: **FE** (AM, KH, PR). – Korean Peninsula, Japan (Hok, Hon, Kyu).
- Lasioglossum (Evylaeus) trichorhinum** (Cockerell, 1925) [*Halictus*]. Russia: **FE** (PR). – Mongolia, China (NC).
- Lasioglossum (Evylaeus) tricinctum tricinctum** (Schenck, 1874) [*Halictus*] (*Halictus delmasi* Pérez, 1903). Russia: **EP** (E, S, CR). – Europe (WE, SE, EE).
- Lasioglossum (Evylaeus) trispine** (Vachal, 1903) [*Halictus*]. Russia: **FE** (PR). – Korean Peninsula, Japan (Hok, Hon, Shi, Ryu).
- Lasioglossum (Evylaeus) truncaticolle** (Morawitz, 1877) [*Halictus*] (*Halictus brevithorax* Pérez, 1903; *H. blidahensis* Strand, 1909; *H. sudaghensis* Strand, 1909). Russia: **EP** (S, CR). – Europe (WE, SE, EE), Caucasus, Turkey, Syria, Israel, Iran, Kazakhstan.

Lasioglossum (Evylaeus) villosulum (Kirby, 1802) [Melitta] (*Melitta punctulata* Kirby, 1802; *Halictus hirtellus* Schenck, 1869; *H. medinai* Vachal, 1895; *H. pauperatulellus* Strand, 1909; *H. trichopsis* Strand, 1909; *H. melanomitratus* Strand, 1914; *H. m. mitratolus* Strand, 1914; *H. villosulopsis* Blüthgen, 1926; *H. pahanganus* Blüthgen, 1928; *H. barkensis* Blüthgen, 1930; *H. villosulus perlautus* Cockerell, 1938; *H. rufotegularis* Cockerell, 1938; *H. villiersi* Benoist, 1941; *H. berberus* Benoist, 1941). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **ES** (IR), **FE** (AM, PR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Syria, Israel, Central Asia, Kazakhstan, India.

Lasioglossum (Evylaeus) virideglaucum Ebmer et Sakagami, 1994. Russia: **FE** (KH, PR). – China (NC, SW, SE), Korean Peninsula.

Lasioglossum (Evylaeus) viridellum (Cockerell, 1931) [Halictus]. Russia: **FE** (KH, PR). – China (NE, CC), Korean Peninsula.

Lasioglossum (Evylaeus) vulsum (Vachal, 1903) [Halictus] (*Halictus trispinis* Vachal, 1903). Russia: **ES** (KR), **FE** (AM, PR, KU). – Mongolia, Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu).

Lasioglossum (Evylaeus) yakuticum (Pesenko et Davydova, 2004) [Evylaeus]. Russia: **ES** (KR, YA, ZB), **FE** (AM).

SPHECODES Latreille, 1804. Type species *Nomada gibba* Fabricius, 1804 (= *Sphex gibba* Linnaeus, 1758). Species of this cosmopolitan bees are notable cleptoparasites in the nests of wide variety of short-tongued bees, particularly species of *Lasioglossum* Curtis and *Halictus* Latreille. Number of species: World – about 300 species, Palaearctic – about 70 species, Russia – 37.

Sphecodes albilabris (Fabricius, 1793) [Nomada] (*Diachroa fuscipennis* Germar, 1822; *Sphecodes latreillei* Wesmael, 1835; *S. rubripes* Spinola, 1838; *S. africanus* Lepeletier de Saint Fargeau, 1841; *S. nigripes* Lepeletier de Saint Fargeau, 1841; *S. rugosus* Smith, 1848; *S. nodicornis* Gistel, 1857; *S. fuscipennis basalis* Dalla Torre, 1877; *Sabulicola cirsii* Verhoeff, 1890; *Sphecodes grandis* Meyer, 1922; *S. rufipennis* Cockerell, 1931; *S. atrescens* Cockerell, 1931). Cleptoparasite of *Colletes cunicularius* (L.) (Colletidae), *Halictus quadricinctus* (F) (Halictidae), *Melitturga clavicornis* (Latr.) (Andrenidae), *Dasypoda hirtipes*

(F) (Melittidae). Russia: **EP** (NE, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, KM, AL), **ES** (KR, BR), **FE** (PR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Syria, Israel, Central Asia, Kazakhstan, India.

Sphecodes alternatus Smith, 1853 (*S. gibbus similis scariosus* Sichel, 1865; *S. punctiventris* Hagens, 1882; *S. gracilior* Morawitz, 1893; *S. antigae* Tournier, 1901; *S. reticulatus algeriensis* Alflen, 1914; *S. alternatus lindbergi* Pittioni, 1950). Cleptoparasite of *Halictus compressus* (Walckener), *H. langobardicus* Blüthgen, *H. patellatus* Mor. (Halictidae). Russia: **EP** (S, NC, CR), **UR**, **WS** (AL), **ES** (KR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Iran, Central Asia, Kazakhstan.

Sphecodes crassus Thomson, 1870 (*S. variegatus* Hagens, 1874; *S. divisus* Hagens, 1882; *S. valesianus* Frey-Gessner, 1903; *S. dyozankeanus* Tsuneki, 1983; *S. ohdeyamanus* Tsuneki, 1984). Cleptoparasite of *Lasioglossum pauxillum* (Schenck), *L. punctatissimum* (Schenck), *L. quadrinotatum* (Schenck), *L. nitidiusculum* (Kirby), *L. prasinum* (Smith) (Halictidae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KA). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Iran, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, Japan (Hok, Hon).

Sphecodes cristatus Hagens, 1882. Cleptoparasite of *Halictus confusus* Smith, *H. leucaheneus* Ebmer, *H. seladonius* (F), *H. subauratus* (Rossi), *Lasioglossum nigripes* (Lep.) (Halictidae). Russia: **EP** (C, E, S, NC), **UR**, **WS** (OM, KM, AL), **ES** (KS, TU, KR, BR, ZB), **FE** (AM, KH, PR). – Europe (WE, SE, EE), Caucasus, Turkey, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE).

Sphecodes croaticus Meyer, 1922. Cleptoparasite of *Lasioglossum interruptum* (Panz.) (Halictidae). Russia: **EP** (C, S, NC, CR). – Europe (WE, SE, EE), Caucasus, Turkey, Turkmenistan.

Sphecodes ephippius (Linnaeus, 1767) [Sphex] (*Apis minimus* Harris, 1776; *A. rufescens* Geoffroy in Fourcroy, 1785; *A. obscura* Geoffroy in Fourcroy, 1785; *A. rufescens* Gmelin, 1790; *A. labiata* Fabricius, 1793; *Melitta divisa* Kirby, 1802; *Andrena minuta* Fabricius, 1804; *Sphecodes similis* Wesmael, 1835; *S. zablockii* Blüthgen, 1924). Cleptoparasite of *Halictus tumulorum* (L.), *H. maculatus* Smith, *H. rubicundus* (Christ), *Lasioglossum laticeps* (Schenck), *L. leucozonium* (Schrank), *L. malachurum*

(Kirby), *L. pauxillum* Schenck, *L. quadrinotatum* (Schenck), *L. fratellum* (Pérez), *L. lativentre* (Schenck), *L. prasinum* (Smith) (Halictidae), *Andrena argentata* Smith, *A. barbilabris* (Kirby), *A. flavipes* Panzer, *A. chrysopyga* Schenck, *A. labialis* (Kirby), *A. minutula* (Kirby), *A. wilkella* (Kirby) (Andrenidae). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (OM, TK, NS, KM, AL), **ES** (KR, IR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Central Asia, Kazakhstan, Mongolia.

Sphecodes ferruginatus Hagens, 1882 (*S. rufescens alpestris* Frey-Gessner, 1903; *S. koikensis* Tsuneki, 1983; *S. hanedai* Tsuneki, 1983; *S. baratonis* Tsuneki, 1983; *S. sudai* Tsuneki, 1983; *S. chibaensis* Tsuneki, 1984; *S. izumindus* Tsuneki, 1986). Cleptoparasite of *Lasioglossum fulvicorne* (Kirby), *L. laticeps* (Schenck), *L. pauxillum* Schenck, *L. vulsum* (Vachal) (Halictidae). Russia: **EP** (N, NW, C, E, NC, CR), **UR**, **WS** (TK, NS, KM, AL), **ES** (IR), **FE** (AM, KH, PR, KU). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Kyrgyzstan, Kazakhstan, Japan (Hok, Hon, Shi, Kyu).

Sphecodes geoffrellus (Kirby, 1802) [Melitta] (*S. affinis* Hagens, 1882; *S. fasciatus* Hagens, 1882; *S. rimalis* Pérez, 1903; *S. impunctatus* Meyer, 1922; *S. silvicola* Tsuneki, 1983; *S. shirozui* Tsuneki, 1983; *S. kitamius* Tsuneki, 1983). Cleptoparasite of *Lasioglossum leucopus* (Kirby), *L. morio* (F), *L. nitidiusculum* (Kirby), *L. pauxillum* Schenck, *L. fratellum* (Pérez), *L. marginellum* (Schenck), *L. rufitarse* (Zetterstedt), *L. sexstrigatum* (Schenck) (Halictidae). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (TK, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, KU). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Near East, Central Asia, Kazakhstan, Japan (Hok, Hon, Shi, Kyu).

Sphecodes gibbus (Linnaeus, 1758) [Sphex] (*Apis glabra* Füessli, 1775; *Nomada gibba* Fabricius, 1775; *Andrena ferruginea* Olivier, 1789; *Apis gibbosa* Christ, 1791; *Melitta sphecoidea* Kirby, 1802; *M. picea* Kirby, 1802; *Sphecodes apicatus* Smith, 1853; *S. nigripennis* Morawitz, 1876; *S. sutor* Nurse, 1903; *S. gibbus rufispinosus* Meyer, 1919; *S. g. turkestanicus* Meyer, 1919; *S. castilianus* Blüthgen, 1924; *S. pergibbus* Blüthgen, 1937; *S. lustrans* Cockerell, 1931; *S. angarensis* Cockerell, 1937). Cleptoparasite of *Halictus maculatus* Smith, *H. quadricinctus* (F), *H. rubicundus* (Christ), *H. sexcinctus* (F), *H. simplex* Blüthgen, *Lasioglossum malachurum* (Kirby) (Halictidae), *Andrena vaga* Panz. (Andrenidae),

Colletes cunicularius (L.) (Colletidae). Russia: **EP** (N, NW, C, E, N, NC, CR), **UR**, **WS** (OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, YA). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NW), India.

Sphecodes hyalinatus Hagens, 1882. Cleptoparasite of *Lasioglossum fratellum* (Pérez), *L. fulvicorne* (Kirby) (Halictidae). Russia: **EP** (N, NW, C, E, NC, CR), **UR**, **WS** (KM, AL), **ES** (KS, TU, IR), **FE** (AM, KH, PR, KA, KU, MG). – Europe (WE, NE, SE, EE), Caucasus, Kazakhstan.

Sphecodes intermedius Blüthgen, 1923 (*S. lactipennis* Meyer, 1924). Parasite of *Halictus kessleri* Bramson (Halictidae). Russia: **EP** (C, S, CR), **UR**. – Europe (SE, EE), N Africa, Caucasus, Turkey, Central Asia, Kazakhstan.

Sphecodes laticaudatus Tsuneki, 1983. Russia: **FE** (KH, PR, SA, KU). – Japan (Hon).

Sphecodes longulus Hagens, 1882 (*S. longulus eupidus* Hagens, 1882; *S. nitidulus* Hagens, 1882; *S. amakusensis* Yasumatsu et Hirashima, 1951; *S. sabulosus* Tsuneki, 1983; *S. crassicornis* Tsuneki, 1983; *S. tsunekii* Haneda, 1994). Cleptoparasite of *Lasioglossum leucopus* (Kirby), *L. minutissimum* (Kirby), *L. punctatissimum* (Schenck), *L. semilucens* (Alfken), *L. zonulum* (Smith), *L. lucidulum* (Schenck), *L. sexstrigatum* (Schenck) (Halictidae). Russia: **EP** (NW, C, E, S, NC), **UR**, **WS** (KM), **ES** (KS, KR, IR), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Syria, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Japan.

Sphecodes majalis Pérez, 1903 (*S. gracilior* Pérez, 1903; *S. opacifrons* Pérez, 1903; *S. problematicus* Schulz, 1906). Cleptoparasite of *Lasioglossum pallens* (Brullé) (Halictidae). Russia: **EP** (C, NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Iran.

Sphecodes maruyamanus Tsuneki, 1983 (*S. convergens* Tsuneki, 1983). Russia: **FE** (PR). – Japan (Hok, Hon).

Sphecodes miniatus Hagens, 1882 (*S. dimidiatus* Hagens, 1882; *S. murithianus* Frey-Gessner, 1903; *S. pilicornis* Meyer, 1922). Cleptoparasite of *Lasioglossum nitidiusculum* (Kirby), *L. morio* (F), *L. pauxillum* Schenck, *L. politum* Schenck, *L. punctatissimum* (Schenck), *L. sexstrigatum* (Schenck) (Halictidae). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (TM, KM, AL), **ES** (KS, TU, BR), **FE**

(KH). – Europe (WE, NE, SE, EE), Caucasus, Kyrgyzstan, Kazakhstan, Mongolia.

Sphecodes monilicornis (Kirby, 1802) [Melitta]
S. maculatus Lepeletier de Saint Fargeau, 1841;
S. subquadratus Smith, 1845; *S. gibbus ephippium rufipes* Sichel, 1865; *S. g. e. dubious* Sichel, 1865;
S. g. e. incertus Sichel, 1865; *S. g. e. nigrescens* Sichel, 1865; *S. g. e. testaceipes* Sichel, 1865; *S. rufifcrus* Dalla Torre, 1896; *S. hanuman* Nurse, 1903;
S. monilicornis nigerrima Blüthgen, 1927; *S. cephalotes* Meyer, 1920; *S. smyrnensis* Meyer, 1920;
S. monilicornis quadratus Meyer, 1920; *S. m. berberus* Warncke, 1992; *S. quadratus cephalotiformis* Pittoni, 1950). Cleptoparasite of *Halictus maculatus* Smith, *H. rubicundus* (Christ), *H. tumulorum* (L.), *Lasioglossum albipes* (F), *L. calceatum* (Scop.), *L. leucozonium* (Schrink), *L. malachurum* (Kirby), *L. quadrinotatulum* (Schenck), *L. zonulum* (Smith), *L. laticeps* (Schenck), *L. pauxillum* Schenck, *L. prasinum* (Smith), *L. villosulum* (Kirby) (Halictidae), *Andrena flavipes* Panz. (Andrenidae). Russia: EP (NW, C, E, S, NC, CR), UR, WS (TM, OM, TK, NS, AL), ES (KS, TU, KR, IR, BR, YA), FE (AM, KH, PR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Pakistan, Central Asia, Kazakhstan, Mongolia.

Sphecodes murotai Tsuneki, 1983 (*S. akitanus* Tsuneki, 1983; *S. kaiensis* Tsuneki, 1983; *S. mutsuoides* Tsuneki, 1984; *S. maetai* Tsuneki, 1984). Cleptoparasite of *Lasioglossum baleicum* (Cockerell) (Halictidae). Russia: FE (PR). – Japan (Hon, Shi, Kyu).

Sphecodes niger Hagens, 1874. Cleptoparasite of *Lasioglossum morio* (F), *L. lucidulum* (Schenck) (Halictidae). Russia: EP (C, E, S, NC). – Europe (WE, NE, SE, EE), Turkey.

Sphecodes nippon Meyer, 1922 (*S. aino* Tsuneki, 1983; *S. awaensis* Tsuneki, 1983; *S. gibbus*: Warncke, 1992). Cleptoparasite of *Lasioglossum sakagamii* Ebmer (Halictidae). Russia: ES (BR, ZB), FE (AM, KH, PR, SA, KU). – Mongolia, Japan (Hok, Hon, Shi, Kyu).

Sphecodes nomioidis Pesenko, 1979. Russia: EP (C). – Europe (WE, SE, EE), Turkey, Jordan.

Sphecodes okuyetsu Tsuneki, 1984 (*S. duplipunctatus* Tsuneki, 1983; *S. hasshanus* Tsuneki, 1983; *S. etienensis* Tsuneki, 1983; *S. sapporoensis* Tsuneki, 1983; *S. brevicypeatus* Tsuneki, 1984). Cleptoparasite of *Lasioglossum allodatum* Ebmer et Sakagami (Halictidae). Russia: FE (KH, PR, SA, KU). – Japan (Hok, Hon, Shi, Kyu).

Sphecodes olivieri Lepeletier de Saint Fargeau, 1825
S. collaris Spinola, 1843; *S. hispanicus abyssinicus* Sichel, 1865; *S. ruficornis* Sichel, 1865; *S. punctulatus* Sichel, 1865; *S. subpunctulatus* Sichel, 1865; *S. rufithorax* Morawitz, 1876; *S. verticalis* Hagens, 1882; *S. desertus* Nurse, 1903; *S. chionospilus* Cockerell, 1911; *S. ch. sanguinatus* Cockerell, 1911; *S. tenuis* Meyer, 1919; *S. olivieri niveatus* Meyer, 1924). Cleptoparasite of *Lasioglossum aegyptiellum* (Strand), *L. vagans* (Smith) (Halictidae). Russia: EP (S, NC). – Europe (SE), N Africa, Caucasus, Turkey, Israel, UAE, Qatar, Iran, Pakistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, India.

Sphecodes orientalis Astafurova et Proshchalykin, 2014. Russia: FE (KH, PR, SA).

Sphecodes pellucidus Smith, 1845 (*S. pilifrons* Thomson, 1870; *S. brevicornis* Hagens, 1874; *S. volatilis* Smith, 1879; *S. pellucidus algirus* Alfken, 1914; *S. p. hybridus* Blüthgen, 1924; *S. p. niveipennis* Meyer, 1924). Cleptoparasite of *Lasioglossum leucozonium* (Schrink), *L. prasinum* (Smith) (Halictidae), *Andrena barbilabris* (Kirby), *A. nycthemera* Imhoff, *A. argentata* Smith, *A. bicolor* F, *A. humilis* Imhoff, *A. ventralis* Imhoff, *A. wilkella* (Kirby) (Andrenidae). Russia: EP (N, NW, C, E, S, NC, CR), UR, WS (TM, OM, TK, NS, AL), ES (KS, TU, KR, IR, BR, YA, ZB), FE (AM, KH, PR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Iran, Central Asia, Kazakhstan, Mongolia, China (NW, WP).

Sphecodes pinguiculus Pérez, 1903 (*S. sareptensis* Meyer, 1922; *S. excellens* Meyer, 1922; *S. punctatissimus* Meyer, 1922; *S. hungaricus* Blüthgen, 1923; *S. coelebs* Blüthgen, 1923; *S. consobrinus* Blüthgen, 1923; *S. persicus* Blüthgen, 1924; *S. capverdensis* Pauly et La Roche, 2002). Cleptoparasite of *Halictus lucidipennis* Smith, *H. smaragdulus* Vachal (Halictidae). Russia: EP (S, CR), ES (TU, BR). – Europe (SE, EE), N Africa, Caucasus, Turkey, Israel, UAE, Iran, Central Asia, Kazakhstan, Mongolia.

Sphecodes pseudofasciatus Blüthgen, 1924. Russia: EP (C, NC). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey.

Sphecodes puncticeps Thomson, 1870 (*S. opacifrons* Pérez, 1903; *S. bituberculatus* Pérez, 1903; *S. puncticeps cretanus* Strand, 1921). Cleptoparasite of *Lasioglossum villosulum* (Kirby), *L. brevicorne* (Schenck), *L. politum* Schenck, *L. sabulosum* (Warncke) (Halictidae). Russia: EP (NW, C, E, S, NC, CR), WS (AL), FE (PR). – Europe (WE, NE,

SE, EE), N Africa, Caucasus, Turkey, Central Asia, Kazakhstan, Mongolia.

Sphecodes reticulatus Thomson, 1870 (*S. distinguendus* Hagens, 1874). Cleptoparasite of *Andrena barbilabris* (Kirby), *A. argentata* Smith, *A. wilkella* (Kirby) (Andrenidae), *Lasioglossum leucozonium cedri* Ebmer, *L. prasinum* (Smith) (Halictidae). Russia: **EP** (NW, C, E, S, NC, CR), **WS** (OM, NS, KM), **ES** (KS, TU, KR, IR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Tajikistan, Kyrgyzstan, Kazakhstan.

Sphecodes rubicundus Hagens, 1875. Cleptoparasite of *Andrena labialis* (Kirby), *A. agilissima* (Scopoli), *A. nigroaenea* (Kirby) (Andrenidae). Russia: **EP** (E, S, NC), **UR**. – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey.

Sphecodes rufiventris (Panzer, 1798) [Tiphia] (*S. subovalis* Schenck, 1853; *S. brevis* Hagens, 1875; *S. singularis* Meyer, 1919; *S. combinatus* Blüthgen, 1927; *S. subovalis austrinus* Erlandsson, 1979; *S. s. austrinus balcanicus* Erlandsson, 1979; *S. rufiventris hethiticus* Warncke, 1992). Cleptoparasite of *Halictus maculatus* Smith (Halictidae). Russia: **EP** (C, E, S, NC), **UR**, **WS** (TK, MS, KM, AL), **ES** (KS). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Sphecodes scabricollis Wesmael, 1835 (*S. perversus* Ritsema, 1879; *S. japonicus* Cockerell, 1911; *S. sibiricus* Cockerell, 1924; *S. utinamius* Tsuneki, 1983; *S. asakura* Tsuneki, 1983; *S. hatogayuuus* Tsuneki, 1983; *S. taicho* Tsuneki, 1983; *S. daisi* Tsuneki, 1983). Cleptoparasite of *Lasioglossum zonulum* (Smith), *L. prasinum* (Smith), *Halictus compressus* (Walkener), *H. quadricinctus* (F.) (Halictidae). Russia: **EP** (NW, C, E, S, CR), **UR**, **WS** (NS, KM, AL), **FE** (AM, KH, PR, SA, KU). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Kazakhstan, Korean Peninsula, Japan (Hok, Hon, Shi, Kyu), India.

Sphecodes schenckii Hagens, 1882. Cleptoparasite of *Lasioglossum discum* (Smith), *Halictus simplex* Blüthgen (Halictidae). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), Caucasus, Turkey, Israel, Iran.

Sphecodes schwarzi Astafurova et Proshchalykin, 2014 Russia: **ES** (TU). – Kazakhstan, Mongolia.

Sphecodes spinulosus Hagens, 1875. Cleptoparasite of *Lasioglossum xanthopus* (Kirby) (Halictidae). Russia: **EP** (C, E, S, CR), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Central Asia, Kazakhstan.

Sphecodes simillimus Smith, 1873 (*S. esakii* Strand et Yasumatsu, 1938). Cleptoparasite of *Lasioglossum duplex* (Dalla Torre) (Halictidae). Russia: **FE** (KH, PR, SA, KU). – Japan (Hok, Hon, Shi, Kyu).

Sphecodes tanoi Tsuneki, 1983 (*S. coptis* Tsuneki, 1983; *S. mutsu* Tsuneki, 1983; *S. fukiensis* Tsuneki, 1983; *S. fudzi* Tsuneki, 1983; *S. chichibuus* Tsuneki, 1984; *S. kamafuse* Tsuneki, 1983; *S. ohtsukius* Tsuneki, 1984; *S. chichibuensis* Tsuneki, 1986). Russia: **FE** (PR, KU). – Japan (Hok, Hon, Kyu).

Subfamily ROPHITINAE

The overwhelming majority of rophitine species occur in the Holarctic region, only a few species also inhabit Afrotropical, Oriental, and Neotropical regions; no species are recorded from Australia. Many species of Rophitinae are montane, the majority of species are oligo- or monolectic. All rophitines are nesting species (non-parasitic), constructing their nests in soil; all behaviourally known species are solitary (not even subsocial).

DUFOUREA Lepeletier de Saint Fargeau, 1841. Type species *Dufourea minuta* Lepeletier de Saint Fargeau, 1841. Number of species: World – 130; Palaearctic – 82; Russia – 8.

Dufourea (Cephalictoides) paradoxa sibirica Pesenko, 1998. Russia: **WS** (AL), **ES** (TU, YA). – Mongolia.

Dufourea (Cyprizophites) coeruleocephala Morawitz, 1872 (*D. caeruleocephala* Dalla Torre, 1896). Russia: **EP** (C, S). – Europe (EE).

Dufourea (Dufourea) halictula (Nylander, 1852) [*Rhophites*] (*D. minuta* auct.). Russia: **EP** (NW, C, CR). – Europe (WE, NE, SE, EE).

Dufourea (Dufourea) minuta Lepeletier de Saint Fargeau, 1841 (*D. vulgaris* Schenck, 1861). Russia: **EP** (N, NW, C, E, NC), **UR**. – Europe (WE, NE, SE, EE), China (WP).

Dufourea (Halictoides) carinata (Popov, 1959) [Halictoides]. Russia: **ES** (BR, ZB), **FE** (AM, KH, PR). – China (NE, NC).

Dufourea (Halictoides) dentiventris (Nylander, 1841) [Halictoides] (*D. dejeanii* Lepeletier de Saint Fargeau, 1841; *Rophites bispinosa* Eversmann, 1852; *Dufourea putoniana* Dours, 1873; *D. odontogastra* Ebmer, 1978). Russia: **EP** (NW, C, E, NC), **UR**, **ES** (IR). – Europe (WE, NE, SE, EE), China (WP), Korean Peninsula.

Dufourea (Halictoides) graeca dubiosa (Warncke, 1979) [Rophites]. Russia: EP (NC). – Georgia, Armenia, Turkey.

Dufourea (Halictoides) inermis (Nylander, 1848) [Halictoides]. Russia: EP (NW, C, E, NC), UR, WS (TK, AL), ES (IR, YA), FE (AM, KH, PR). – Europe (WE, NE, SE, EE), Uzbekistan, China (NE, WP).

FLAVODUFOUREA Ebmer, 1984. Type species *Dufourea flavigornis* Friese, 1914. Number of species: World – 2; Palaearctic – 2; Russia – 1.

Flavodufourea flavigornis (Friese, 1914) [Dufourea]. Russia: WS (BR). – China (WP).

RHOPHITOIDES Schenck, 1861. Type species *Rhophitoides distinguendus* Schenck, 1861 (= *Rophites cana* Eversmann, 1852). Number of species: World – 4; Palaearctic – 4; Russia – 1.

Rhophitoides canus (Eversmann, 1852) [Rophites] (*Rhophitoides bifoveolatus* Sichel, 1854; *Rhophitoides distinguendus* Schenck, 1861). Russia: EP (C, E, S, NC, CR), UR, WS (TM, OM, TK, KM, AL), ES (KR, TU, BR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Central Asia, Kazakhstan, Mongolia, China (NW), Korean Peninsula.

ROPHITES Spinola, 1808. Type species *Rophites quinquespinosus* Spinola, 1808. Number of species: World – 16; Palaearctic – 16; Russia – 6.

Rophites algirus montanus Ebmer, 1978 (*R. algirus hethicus* Warncke, 1980). Russia: EP (NC). – Caucasus, Turkey, Iran.

Rophites algirus trispinosus Pérez, 1903. Russia: EP (C, E, S, NC, CR), UR. – Europe (WE, NE, SE, EE).

Rophites caucasicus Morawitz, 1876. Russia: EP (NC). – Caucasus, Turkey.

Rophites clypealis stepposus Astafurova, 2011. Russia: EP (S, CR). – Europe (EE), Kazakhstan.

Rophites gruenwaldti Ebmer, 1978. Russia: FE (AM). – Mongolia, China (NE, NC).

Rophites hartmanni Friese, 1902 (*R. bistrispinosus* Lebedev, 1931). Russia: EP (C, E, S, NC, CR). – Europe (WE, SE, EE), Turkey, Israel.

Rophites quinquespinosus Spinola, 1808 (*R. pilichi* Moczar, 1967; *R. moeschleri* Schwammberger, 1971; *R. bluethgeni* Benedek, 1973). Russia: EP (C, E, S, NC, CR), UR, WS (TK, NS, KM, AL), ES (KS, KR, IR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Uzbekistan, Kyrgyzstan, Kazakhstan, China (NW).

SYSTROPHA Illiger, 1806. Type species *Andrena spiralis* Olivier, 1789 (= *Eucera curvicornis* Scopoli, 1770). Number of species: World – 28; Palaearctic – 13; Russia – 2.

Systropha curvicornis (Scopoli, 1770) [Eucera] (*Tenthredo convolvuli* Pallas, 1773; *Andrena spiralis* Olivier, 1789; *A. labrosa* Eversmann, 1852). Russia: EP (C, E, S, NC, CR), UR, WS (NS, AL). – Europe (WE, NE, SE, EE), Armenia, Azerbaijan, Turkey, Iran, Uzbekistan, Kazakhstan, China (NW, WP).

Systropha planidens Giraud, 1861 (*S. chrysura* Pérez, 1905; ?*S. grandimargo* Pérez, 1905; *S. planidens anatolica* Warncke, 1977). Russia: EP (C, E, S, NC, CR), UR. – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Kazakhstan.

Subfamily NOMIINAE

The subfamily Nomiinae is nearly cosmopolitan group of bees (absent in South America). Many species of Nomiinae are plain, the majority of species are polyleptic. All species of the subfamily make nests; there are no cleptoparasitic forms. All species, well studied in relation their biology, are solitary (sometimes subsocial). They nest in the ground, usually on exposed warm dry areas. Females of many species form communal nests.

LIPOTRICHES Gerstaecker, 1858. Type species *Lipotriches abdominalis* Gerstaecker, 1858 (= *Sphecodes cribrosa* Spinola, 1843). Number of species: World – 270; Palaearctic – 9; Russia – 1.

Lipotriches (Austronomia) fruhstorferi (Pérez, 1905) [Nomia]. Russia: ES (ZB), FE (AM, KH, PR). – Mongolia, China (NE), Japan (Hok, Kyu, Tsu).

NOMIAPIS Cockerell, 1919. Type species *Nomia diversipes* Latreille, 1806. Number of species: World – 14; Palaearctic – 11; Russia – 6.

Nomiapis armata (Olivier, 1812) [Nomia] (*Nomia monstrosa* Costa, 1861; *N. rumelica* Friese, 1923). Russia: EP (CR). – Europe (SE, EE), Armenia, Azerbaijan, Turkey, Israel.

Nomiapis bispinosa (Brulle, 1832) [Nomia] (*Nomia rufiventris* Spinola, 1839; *N. ruficornis* Spinola, 1839; *N. perforata* Lucas, 1849; *N. albocincta* Lucas, 1849; *N. aureocincta* Costa, 1861; *N. polita* Costa, 1861; *N. basalis* Smith, 1875; *N. aureocincta turcomanica* Radoszkowski, 1893; *N. fletcheri* Cockerell,

1920; *N. basalicincta* Cockerell, 1922; *N. unidentata* auct.). Russia: **EP** (E, S, NC, CR). – Europe (SE, EE), N Africa, Caucasus, Turkey, Israel, Iraq, Afghanistan, Pakistan, Central Asia, Kazakhstan, China (NW), N India.

Nomiapis diversipes (Latreille, 1806) [Nomia] (*Andrena humeralis* Jurine, 1807; *Nomia hungarica* Förster, 1853). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (NS, AL). – Europe (WE, SE, EE), Caucasus, Turkey, Israel, Iran, Pakistan, Central Asia, Kazakhstan, Mongolia.

Nomiapis femoralis (Pallas, 1773) [Apis] (*Lasius difformis* Panzer, 1805; *Andrena brevitarsis* Eversmann, 1852). Russia: **EP** (C, E, S, NC), **UR**, **WS** (AL), **ES** (KS, TU, KR, IR, BR). – Europe (WE, SE, EE), Central Asia, Kazakhstan, Mongolia, China (NW).

Nomiapis fugax (Morawitz, 1877) [Nomia]. Russia: **EP** (S, NC), **ES** (IR). – N Africa, Armenia, Azerbaijan, Turkey, Iran, Pakistan, Central Asia, Kazakhstan, China (NW).

Nomiapis mandschurica (Hedicke, 1940) [Nomia]. Russia: **ES** (BR), **FE** (AM, PR). – Mongolia, China (NE, NC), Korean Peninsula, Japan (Hok, Hon, Kyu).

PSEUDAPIS Kirby, 1900. Type species *Pseudapis anomala* W.F. Kirby, 1900. Number of species: World – 41; Palaearctic – 24; Russia – 2.

Pseudapis anatolica (Warncke, 1976) [Nomia]. Russia: **EP** (NC). – Armenia, Turkey.

Pseudapis elegantissima (Popov, 1949) [Nomia]. Russia: **EP** (S, NC). – Azerbaijan, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Subfamily NOMIOIDINAE

The subfamily Nomioidinae is represented by small (body length at most 5 mm) and usually brightly colored forms and overwhelming majority of species occur in the Old Word. The most of species inhabit deserts and semi-desert and are polylectic. All nomioides are

nesting species (non-parasitic), constructing their nests in soil, preferring sandy and stony soils; all behaviourally known species are solitary or subsocial (not eusocial).

CEYLALICTUS Strand, 1913 (*Eunomiooides* Blüthgen, 1937). Type species *Halictus horni* Strand, 1913. Number of species: World – 28; Palaearctic – 1; Russia – 1.

Ceylalictus (Ceylalictus) variegatus (Olivier, 1789) [Apis] (*Andrena pulchella* Jurine, 1807; *A. flavopicta* Dours, 1873; *A. flavopicta* Dours, 1873; *Nomiooides jucunda* Morawitz, 1874; *N. fasciatus* Friese, 1898; *N. fasciatus intermedius* Alfken, 1924; *N. variegata simplex* Blüthgen, 1925; *N. variegata unifasciata* Blüthgen, 1925; *N. labiatarum* Cockerell, 1931; *N. variegata nigrita* Blüthgen, 1934; *N. variegata pseudocerea* Blüthgen, 1934; *N. variegata nigri-ventris* Blüthgen, 1934). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Arabian Peninsula, Israel, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NW), NW India.

NOMIOIDES Schenck, 1866. Type species *Apis minutissima* Rossi, 1790. Number of species: World – 62; Palaearctic – 39; Russia – 2.

Nomiooides (Nomiooides) minutissimus minutissimus (Rossi, 1790) [Apis] (?*Apis parvula* Fabricius, 1798; *Nomiooides minutissima obscurata* Blüthgen, 1925; *N. m. versicolor* Blüthgen, 1925; *N. m. violascens* Blüthgen, 1925; *N. m. schencki* Blüthgen, 1925; *N. m. tristis* Blüthgen, 1934; *N. m. fusca* Blüthgen, 1934). Russia: **EP** (E, S, NC, CR). – Europe (WE, SE, EE), Caucasus, Turkey, Near East, Iraq, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NW), NW India.

Nomiooides (Nomiooides) pulverosus Handlirsch, 1888 (*N. subgalerita* Blüthgen, 1933). Russia: **EP** (S, NC). – Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia.

76. FAMILY MELITTIDAE

M.YU. PROSHCHALYKIN AND YU.V. ASTAFUROVA

Melittid bees occur in temperate and xeric ecosystems of the Nearctic and the Old World. Ethiopian region shows the maximum of generic diversity but the maximum of species diversity is reached in the Palaearctic region. The African continent lumps clearly the maximum of both generic and specific diversity. Melittids are typically small to moderate-sized bees, which are well known for their specialist and oligoleptic foraging habits. Nests in burrows on the soil. Number of taxa: World – 15 genera, 198 species; Palaearctic – 5/85; Russia – 3/25.

R e f e r e n c e s. Warncke, 1973; Dathe, 1980b; Situdikov, 1986; Radchenko, Pesenko, 1989; Michez et al., 2004, 2012; Celary, 2005; Michez, Patiny, 2005; Michez, Eardley, 2007; Proshchalykin, 2007c, 2008b; 2010; 2012a; 2014a; Levchenko, 2009; Radchenko, 2009; 2016; Levchenko, Yuferev, 2013; Levchenko, Tomkovich, 2014.

Subfamily DASYPODAINAE

Tribe DASYPODAINI

DASYPODA Latreille, 1802 (*Podasys* Rafinesque, 1815; *Microdasypoda* Michez, 2004; *Megadasypoda* Michez, 2004; *Heterodasypoda* Michez, 2004). Type species *Andrena hirtipes* Fabricius, 1793. The genus is widespread in the Palaearctic and only 2 species known in Afrotropical region. Number of taxa: World – 39; Palaearctic – 37; Russia – 8.

Dasypoda argentina Panzer, 1809 (*D. villipes* Eversmann, 1852, nom. praeocc., nec Lepeletier de Saint Fargeau, 1841; *D. thoracica* Baer, 1853; *D. palleola* Baer, 1853; *D. decora* Baer, 1853; *D. nigrans* Baer, 1853; *D. italicica* Radoszkowski, 1876; *D. rhododaetyla* Dalla Torre, 1881; *D. canescens* De Stefani, 1887; *D. mlokoszewiczi* Radoszkowski, 1890; *D. wankowiczi* Radoszkowski, 1891; *D. carinata* Pérez, 1895; *D. argentata rufipes* Friese, 1901; *D. a. nigricans* Friese, 1901; *D. a. cinerascens* Friese, 1901; *D. a. griseithorax* Müller, 1923; *D. a. spadix* Müller, 1923; *D. thoracica*: Radchenko, Pesenko, 1989). Russia: **EP** (C, S, E, NC, CR), **UR**. – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Lebanon, Iran.

Dasypoda braccata Eversmann, 1852 (*D. tschertkovi* Baer, 1853; *D. melanopleura* Baer, 1853).

Russia: **EP** (E, NC), **UR**. – Europe (WE, SE, EE), Turkey, Kazakhstan.

Dasypoda hirtipes (Fabricius, 1793) [*Andrena*] (?*Apis altercator* Harris, 1776; *Andrena hirtipes* Fabricius, 1793; *Apis hirta* Fabricius, 1793, nom. praeocc., nec Schrank, 1781; *Andrena plumipes* Panzer, 1797; *Melitta swammerdamella* Kirby, 1802; *Dasypoda villosa* Lepeletier de Saint Fargeau, 1841; ?*D. germanica* Gistel, 1848; *D. palleola* Baer, 1853; *D. nemoralis* Baer, 1853; ?*D. rudis* Gistel, 1857; *D. plumipes nigrescens* Friese, 1901; *D. minor* Pérez, 1903, nom. praeocc., nec Morawitz, 1874; *D. illegalis* Schulz, 1906; *D. altercator* auct.). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (NS), **ES** (BR), **FE** (PR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Iran, Kazakhstan, Mongolia, China (NE, NC).

Dasypoda japonica Cockerell, 1911. Russia: **ES** (BR), **FE** (KH, PR). – China (NC), Korean Peninsula, Japan (Hon, Kyu).

Dasypoda morawitzi Radchenko, 2016 (?*D. hirtipes minor* Morawitz, 1874). Russia: **EP** (NW, E, S, NC, CR), **UR**. – Europe (EE), Turkey, Kazakhstan.

Dasypoda spinigera Kohl, 1905. Russia: **EP** (NC, CR). – Europe (WE, SE, EE), Armenia, Turkey.

Dasypoda suripes (Christ, 1791) [*Apis laniger* Christ, 1791; *Dasypoda speetabilis* Rudow, 1881; *D. aurata* Rudow, 1881; *D. leueoura* Rudow, 1882; *D. mixta* Radoszkowski, 1887; *D. thomsoni* Schletterer, 1890]. Russia: **EP** (C, E, NC, CR), **UR**. – Europe (WE, NE, SE, EE), Turkey.

Dasypoda toroki Michez, 2004. Russia: **EP** (CR). – Turkey, Syria, Israel.

Subfamily MELITTINAE

MACROPIS Panzer, 1809. Type species *Megilla labiata* Fabricius, 1805 (=*Megilla fulvipes* Fabricius, 1804). The Holarctic genus *Macropis* Panzer comprises 16 recognized species of bees (Michez, Patiny, 2005). All species of the genus are believed to be strictly oligoleptic and are specialized in collecting pollen exclusively from flowers of Lysimachia (Primulaceae). Number of taxa: World – 16; Palaearctic – 13; Russia – 4.

Macropis (Macropis) dimidiata amurensis Popov, 1958 (*M. fulvipes amurensis* Popov, 1958). Russia: **ES** (BR, ZB), **FE** (AM, KH, PR).

Macropis (Macropis) europaea Warncke, 1973 (*M. labiata* auct., nec Fabricius, 1804). Russia: **EP** (NW, C,

E, S, NC, CR), **UR**, **WS** (TM, AL). – Europe (WE, NE, SE, EE), Turkey.

Macropis (Macropis) fulvipes (Fabricius, 1804) [*Megilla*] (*Megilla labiata* Fabricius, 1805). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (TM, AL), **ES** (KS, KR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Kazakhstan, Mongolia.

Macropis (Paramacropis) ussuriana (Popov, 1936) [Ctenoplectra]. Russia: **FE** (KH, PR). – China (NE).

MELITTA Kirby, 1802 (*Cilissa* Leach, 1812; *Kirbya* Lepeletier de Saint Fargeau, 1841; *Pseudocilissa* Radoszkowski, 1891; *Brachycephalapis* Viereck, 1909). Type species *Melitta tricincta* Kirby, 1802. The most of species occur in the Palaearctic and a few species in southern and East Africa, and in the Nearctic; nest in the soil and many species are oligolectic. Number of taxa: World – 48; Palaearctic – 30; Russia – 13.

Melitta budashkini Radchenko et Ivanov, 2012. Russia: **EP** (CR).

Melitta budensis (Mocsáry, 1878) [*Cilissa*]. Russia: **EP** (S, CR). – Europe (SE, EE), Armenia, Turkey, Kazakhstan.

Melitta ezoana Yasumatsu et Hirashima, 1956 (*M. sinensis* Wu, 1978). Russia: **FE** (PR). – China (NE, NC), Korean Peninsula, Japan (Hok, Hon).

Melitta dimidiata Morawitz, 1876 (*Cilissa robusta* Radoszkowski, 1876; *C. dimidiata* var. *hungaria* Mocsáry, 1883). Russia: **EP** (S, NC, CR), **UR**. – Europe (WE, SE, EE), Caucasus, Turkey, Iran, Central Asia, Kazakhstan.

Melitta haemorrhoidalis (Fabricius, 1775) [*Andrena*] (*Apis dichroa* Gmelin, 1790; *Melitta chrysura* Kirby, 1802; *Cilissa nigra* Friese, 1885). Russia: **EP** (NW,

C, E, NC, CR), **UR**. – Europe (WE, NE, SE, EE), Armenia, Turkey.

Melitta japonica Yasumatsu et Hirashima, 1956 (*M. taishanensis* Wu, 1978). Russia: **WS** (NS), **ES** (ZB), **FE** (PR). – China (NC), Japan (Hok, Hon).

Melitta latronis Cockerell, 1924. Russia: **FE** (PR). This species even not mentioned since description and requies the revision.

Melitta leporina (Panzer, 1799) [*Apis*] (*Andrena fortipes* Imhoff, 1832; *Cilissa ruthenica* Radoszkowski, 1891; *Melitta centaureae* Torka, 1922; *M. leporina* var. *nigrinotum* Alfken, 1927; *M. sinkiangensis* Wu, 1978). Russia: **EP** (NW, C, E, NC, CR), **UR**, **WS** (AL). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Afghanistan, Central Asia, Kazakhstan, Mongolia, China (NW).

Melitta melanura (Nylander, 1852) [*Kirbya*] (*Cilissa wankowiczi* Radoszkowski, 1891). Russia: **EP** (C, S, NC, CR), **UR**. – Europe (WE, NE, EE), Armenia, Turkey, Mongolia.

Melitta nigricans Alfken, 1905. Russia: **EP** (C, S, CR), **UR**. – Europe (WE, SE, EE), N Africa, Turkey.

Melitta sibirica (Morawitz, 1888) [*Cilissa*] (*M. albofasciata* Friese, 1900; *Cilissa alticola* Hedicke, 1938; *Melitta borealis* Wu, 2000; *M. quinghaiensis* Wu, 2000). Russia: **EP** (E), **ES** (KR). – Tajikistan, Kyrgyzstan, Mongolia, China (NC), India.

Melitta tricincta Kirby, 1802 (*Andrena microstigma* Eversmann, 1852; *A. quadricincta* Eversmann, 1852; *Melitta meridionalis* Hedicke, 1933). Russia: **EP** (C, E, S, NC, CR), **UR**, **ES** (IR, BR, YA, ZB), **FE** (KH). – Europe (WE, NE, SE, EE), Kazakhstan, China (WP).

Melitta udmurtica Sitdikov, 1986. Russia: **EP** (E), **UR**. – Europe (EE).

77. FAMILY MEGACHILIDAE

M.YU. PROSHCHALYKIN AND A.V. FATERYGA

The family Megachilidae is one of the largest families of the bees. Megachilid bees are found in a wide diversity of habitats on all continents except Antarctica, ranging from lowland tropical rain forests to deserts or alpine environments. Bees of this family have high morphological diversity, broad range of trophic links, and extreme diversity of nesting behavior among all bee families. These solitary bees are both ecologically and economically relevant; they include many pollinators of natural, urban and agricultural vegetation. For example, *Megachile rotundata* has been introduced into many parts of the world as a pollinator of alfalfa (*Medicago sativa*, Fabaceae); several species of the genus *Osmia* is widely used for pollination of orchards (*Prunus* s. l., *Malus*, and *Pyrus*, Rosaceae) and some warmhouse plants (*Fragaria ananassa*, Rosaceae; *Cucumis sativus*, Cucurbitaceae). Megachilidae also contains a large number of cleptoparasitic bees (19 genera, including the entire tribe Dioxynini). Number of taxa: World – 76 genera, 4096 species; Palaearctic – 36/1100; Russia – 18/198.

R e f e r e n c e s . Popov, 1933b; Nikiforuk, 1957; Pesenko, 1971; Sedykh, 1974; Osytshnjuk et al., 1978; Dathe, 1980b; Warncke, 1980; Sittikov, 1986; Romankova, 1995a; Banaszak, Romasenko, 2001; Proshchalykin, Lelej, 2004; Humala, 2006; Proshchalykin, 2006, 2007d,f, 2009a, 2012a,b, 2013a,b, 2014b; Levchenko, 2007; Michener, 2007; Praz et al., 2008; Kuznetsova, 2009; Aguib et al., 2010; Humala, Polevoi, 2011; Gonzalez et al., 2012; Levchenko, Tomkovich, 2014; Kasparek, 2015; Paukkunen, Kozlov, 2015; Yakovleva, 2015; Litman et al., 2016; Trunz et al., 2016; Ascher, Pickering, 2017; Fateryga, 2017b; Fateryga, Popov, 2017; Johansson, Paukkunen, 2017; Kuhlmann et al., 2017; Müller, 2017; Praz, 2017; Rasmont et al., 2017; Proshchalykin et al., 2017a; Fateryga et al., 2018.

Subfamily MEGACHILINAE

Tribe LITHURGINI

LITHURGUS Berthold, 1827 (*Lithurge* Latreille, 1825; *Liturgus* (!): Ashmead, 1899). Type species *Andrena cornuta* Fabricius, 1787. The genus is widely distributed in the Palaearctic, Neotropical, Oriental, and Australian regions. Species of the subgenus *Lithurgus* s. str. are floral specialists,

some on Malvaceae, others on Asteraceae. Number of taxa: World – 26; Palaearctic – 15; Russia – 2.

Lithurgus (Lithurgus) chrysurus Fonscolombe, 1834 (*L. analis* Lepeletier de Saint-Fargeau, 1841; *L. haemorrhoidalis* Lepeletier de Saint-Fargeau, 1841; *L. sublaevis* Pérez, 1897; *L. cornutus* auct.). Russia: **EP** (C, CR), **UR**. – Europe (WE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Syria, Israel, Iran, ?Central Asia, N America (introduced).

Lithurgus (Lithurgus) cornutus (Fabricius, 1787) [*Andrena*] (*L. umbraculatus* Lepeletier de Saint-Fargeau, 1841; *L. fuseipennis* Lepeletier de Saint-Fargeau, 1841; *L. nasutus* Dufour, 1849; *Megachile monoceros* Eversmann, 1852; *M. dohrni* Radoszkowski, 1862; *Lithurgus maximus* Radoszkowski, 1872; *L. cornutus* var. *obscurus* Cockerell, 1931). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (KM, AL), **ES** (KS, KR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran, Uzbekistan, Kyrgyzstan, Kazakhstan, N America (introduced).

Tribe OSMIINI

CHELOSTOMA Latreille, 1809. Type species *Apis maxillosa* Linnaeus, 1767. The genus is widespread in the Holarctic region, ranging across North America and western and central Eurasia, but is unknown in Japan. It extends south to northern Mexico, northern Africa, and into the montane tropics in northeast India, Burma, and Thailand. Many and perhaps all species are oligoleptic on particular floral taxa. Number of species: World – 54; Palaearctic – 45; Russia – 6.

Chelostoma (Chelostoma) florisomne (Linnaeus, 1758) [*Apis*] (*Apis maxillosa* Linnaeus, 1767; *Chelostoma culmorum* Lepeletier de Saint-Fargeau, 1841; *Heriades parumcrinitus* Alfken, 1932). Russia: **EP** (NW, C, E, NC, CR), **UR**. – Europe (WE, NE, SE, EE), Georgia, Armenia, Turkey.

Chelostoma (Chelostoma) mocsaryi Schletterer, 1889. Russia: **EP** (CR). – Europe (WE, SE, EE), Caucasus, Turkey, Lebanon, Israel.

Chelostoma (Foveosmia) campanularum (Kirby, 1802) [*Apis*] (*Apis florisomnis minima* Christ, 1791, nom. praeocc., nec Poda, 1761; *Chelostoma florisomne* auct.). Russia: **EP** (C, CR), **UR**. – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, N America (introduced).

Chelostoma (Foveosmia) distinctum (Stoeckhert, 1929) [Eriades] (*Heriades cantabrica* Benoist, 1935). Russia: **EP** (C, S, CR), **UR**. – Europe (WE, SE, EE), Azerbaijan, Turkey, Israel, Iran.

Chelostoma (Foveosmia) foveolatum (Morawitz, 1868) [Heriades]. Russia: **EP** (S, NC, ?CR), **ES** (KR), **FE** (AM, MG). – Europe (WE, SE, EE), Georgia, Turkey, Iran.

Chelostoma (Gyrodromella) rapunculi (Lepeletier de Saint-Fargeau, 1841) [Heriades] (*Apis fuliginosa* Panzer, 1798, nom. praeocc., nec Scopoli, 1770, nec Christ, 1791; *Heriades nigricornis* Nylander, 1848; *Chelostoma inerme* Eversmann, 1852; *Heraclides casularum* Chevrier, 1872; *Chelostoma proximum* Schletterer, 1889; *Osmia archanensis* Cockerell, 1928; *O. platyodonta* Cockerell, 1928; *Heraclides confusa* Benoist, 1934). Russia: **EP** (N, C, E, S, NC, CR), **UR**, **WS** (TK, KM, AL), **ES** (KS, KR, IR, BR), **FE** (AM, KH, SA). – Europe (WE, NE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Syria, Jordan, Israel, Iran, Central Asia, Kazakhstan, Mongolia, China (NE), N America (introduced).

HERIADES Spinola, 1808 (*Trypetes* Schenck, 1861; *Eriades* Spinola, 1895; *Orientoheriades* Gupta, 1987). Type species *Apis truncorum* Linnaeus, 1758. The genus has a wide distribution, it occurs in the Palaearctic, Nearctic, Afro-tropical, and Oriental regions. The great majority of species nest in preexisting cavities, mainly in insect burrows in dead wood or in hollow stems. Number of species: World – 137; Palaearctic – 22; Russia – 3.

Heriades (Heriades) crenulata Nylander, 1856. Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Kazakhstan.

Heriades (Heriades) rubicola Pérez, 1890 (*H. copeatica* Morawitz, 1892; *H. crenulatus aurantifer* Cockerell, 1931; *H. hierosolomita* Benoist, 1935). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), N Africa, Turkey, Syria, Jordan, Israel, Turkmenistan, Kyrgyzstan, Kazakhstan.

Heriades (Heriades) truncorum (Linnaeus, 1758) [*Apis*] (*Apis xanthogastria* Schrank, 1802; *Heraclides sinuata* Spinola, 1808; *Osmia punctatissima* Lepeletier de Saint-Fargeau, 1841; *Heriades mordax* Schletterer, 1889). Russia: **EP** (N, C, E, S, NC, CR), **UR**, **FE** (AM, PR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Syria, Lebanon, Israel, Iran, Uzbekistan, Kyrgyzstan, Kazakhstan, N America (introduced).

HOPLITIS Klug, 1807 (*Ctenosmia* Thomson, 1872; *Bytinskia* Mavromoustakis, 1954; *Annosmia* Warncke, 1991; *Coloplitis* Griswold, 1998). Type species *Apis aduncta* Panzer, 1798. The genus is distributed in the Palaearctic, Nearctic, and Afro-tropical regions, a few species also occur in the Oriental region. Number of species: World – 376; Palaearctic – 300; Russia – 28.

Hoplitis (Alcidamea) acuticornis (Dufour et Perris, 1840) [*Osmia*] (*Osmia dentiventris* Morawitz, 1877). Russia: **EP** (E, NC, CR), **UR**, **WS** (KM), **ES** (KR). – Europe (WE, SE, EE), N Africa, Armenia, Turkey, Syria, Jordan, Israel, Iran, Turkmenistan, Tajikistan, Kyrgyzstan, Kazakhstan.

Hoplitis (Alcidamea) caularis (Morawitz, 1875) [*Osmia*] (*Osmia proxima* Morawitz, 1875; *O. turkestanica* Dalla Torre, 1896). Russia: **EP** (NC, CR), **UR**. – Europe (EE), Turkey, Syria, Central Asia, Kazakhstan, China.

Hoplitis (Alcidamea) claviventris (Thomson, 1872) [*Osmia*] (*Osmia interrupta* Schenck, 1861, nom. praeocc., nec Latreille, 1811; *O. foveolata* Schenck, 1873, nom. praeocc., nec Morawitz, 1868; *Hoplitis leucomelana* auct.). Russia: **EP** (C, CR), **WS** (KM). – Europe (WE, NE, SE, EE), Turkey, Kazakhstan, Mongolia, China.

Hoplitis (Alcidamea) fulva (Eversmann, 1852) [*Osmia*] (*Osmia grandis* Morawitz, 1872). Russia: **EP** (E, CR), **UR**, **ES** (TU, IR, ZB). – Europe (EE), Armenia, Azerbaijan, Turkey, Syria, Jordan, Kazakhstan, Mongolia, China (NE).

Hoplitis (Alcidamea) laboriosa (Smith, 1878) [*Osmia*] (*Osmia maculata* Morawitz, 1880; *O. magrettii* Ducke, 1899). Russia: **EP** (Müller, 2017). – Kazakhstan, Mongolia, China (WP).

Hoplitis (Alcidamea) leucomelana (Kirby, 1802) [*Osmia*] (*Apis caulincola* Schrank, 1802; *A. zonata* Gravenhorst, 1807; *Osmia parvula* Dufour et Perris, 1840; *Heriades bidenticulata* Costa, 1863; *Hoplitis leucomelaena* (!) Romankova, 1995a). Russia: **EP** (N, C, E, S, NC, CR), **UR**, **WS** (KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Iran, Tajikistan, Kyrgyzstan, Kazakhstan, Mongolia, China (SW).

Hoplitis (Alcidamea) mitis (Nylander, 1852) [*Osmia*] (*Osmia montivaga* Morawitz, 1872; *O. laevifrons* Pérez, 1879, nom. praeocc., nec Morawitz, 1872). Russia: **EP** (CR), **UR**, **WS** (KM), **ES** (KR). – Europe (WE, SE, EE), Armenia, Kazakhstan, Mongolia.

- Hoplitis (Alcidamea) praestans** (Morawitz, 1893) [Osmia] (*Osmia lineola* Pérez, 1895). Russia: **EP** (S, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Jordan, Israel, Iran, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan.
- Hoplitis (Alcidamea) princeps** (Morawitz, 1872) [Osmia]. Russia: **EP** (S, CR), **UR**, **ES** (BR). – Europe (EE), Kazakhstan, Mongolia, China (NW, WP).
- Hoplitis (Alcidamea) scita** (Eversmann, 1852) [Osmia] (*Osmia rubripes* Smith, 1879). Russia: **ES** (TU, IR, BR), **FE** (AM, KH, PR). – Kyrgyzstan, Mongolia, China (NW).
- Hoplitis (Alcidamea) tridentata** (Dufour et Perris, 1840) [Osmia] (*Osmia melanoleuca* Benoist, 1934; *Hoplitis tridentata asensioi* Tkalcú, 1975). Russia: **EP** (C, E, NC, CR), **UR**, **WS** (KM, AL). – Europe (WE, SE, EE), N Africa, Georgia, Armenia, Turkey, Syria, Israel, Iran, Uzbekistan, Kyrgyzstan, Kazakhstan.
- Hoplitis (Alcidamea) tuberculata** (Nylander, 1848) [Osmia] (*Osmia hamata* Eversmann, 1852; *Chelostoma signatum* Eversmann, 1852, nom. praeocc., nec Erichson, 1835; *Osmia macrodonta* Cockerell, 1924; *O. amaguensis* Cockerell, 1924). Russia: **EP** (N, C, E), **UR**, **WS** (TM, KM), **ES** (IR, YA, ZB), **FE** (MG, AM, KH). – Europe (WE, NE, SE, EE), Mongolia, China.
- Hoplitis (Anthocopa) caucasicola** Müller, 2012. Russia: **EP** (S). – Georgia, Turkey.
- Hoplitis (Anthocopa) daurica** (Radoszkowski, 1887) [Osmia]. Russia: **ES** (BR).
- Hoplitis (Anthocopa) jakovlevi** (Radoszkowski, 1874) [*Pseudosmia*] (*Osmia serrilabris* Morawitz, 1875; *O. quadridentata* Fonscombe, 1879). Russia: **EP** (S, CR). – Europe (WE, SE), N Africa, Armenia, Azerbaijan, Turkey, Iran, Turkmenistan, Tajikistan, Kyrgyzstan, Kazakhstan, China.
- Hoplitis (Anthocopa) mocsaryi** (Friese, 1895) [Osmia]. Russia: **EP** (NC, CR). – Europe (WE, SE, EE), Armenia, Turkey, Israel, Iran.
- Hoplitis (Anthocopa) papaveris** (Latreille, 1799) [*Apis*] (*Osmia hyalinipennis* Lepeletier de Saint-Fargeau, 1841). Russia: **EP** (C, NC, CR), **UR**, **ES** (BR). – Europe (WE, SE, EE), Caucasus, Turkey, Kazakhstan, China.
- Hoplitis (Anthocopa) saxialis** (van der Zanden, 1994) [Anthocopa]. Russia: **EP** (NC). – Europe (SE), Turkey, Iran.
- Hoplitis (Anthocopa) taurica** (Radoszkowski, 1874) [*Pseudosmia*]. Russia: **EP** (CR). This species was

even not mentioned since description and requires the revision.

- Hoplitis (Anthocopa) villosa** (Schenck, 1853) [Megachile] (*Osmia platycera* Gerstäcker, 1869). Russia: **EP** (Müller, 2017). – Europe (WE, SE, EE).
- Hoplitis (Formicapis) maritima** (Romankova, 1985) [Formicapis]. Russia: **FE** (PR). – Mongolia.
- Hoplitis (Formicapis) robusta** (Nylander, 1848) [Heriades] (*Osmia rhinoceros* Giraud, 1861; *Heriades trinacria* Morawitz, 1869; *Formicapis clypeata* Sladen, 1916). Russia: **ES** (IR, YA, ZB), **FE** (AM, MG, CH). – Europe (WE, NE, SE, EE), Mongolia, China (NE), N America.
- Hoplitis (Hoplitis) adunca** (Panzer, 1798) [Apis] (*Osmia spinolae* Lepeletier de Saint-Fargeau, 1841; *O. aduncoides* Strand, 1910; *O. adunca contrario* Tkalcú, 1974). Russia: **EP** (C, S, NC, ?CR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Turkmenistan, Kyrgyzstan.
- Hoplitis (Hoplitis) anthocopoides** (Schenck, 1853) [Osmia] (*Osmia caementaria* Gerstäcker, 1869; *O. hybrida* Pérez, 1879; *O. romana* Morice, 1901; *Fertonella algerica* Benoist, 1969). Russia: **EP** (S, NC, CR). – Europe (WE, NE, SE, EE), N Africa, Armenia, N America (introduced).
- Hoplitis (Hoplitis) manicata** Morice, 1901. Russia: **EP** (CR). – Europe (WE, SE, EE), ?N Africa, Caucasus, Turkey.
- Hoplitis (Hoplitis) ravouxi** (Pérez, 1902) [Osmia] (*Osmia rhinotropis* Pérez, 1902; *O. brachyceros* Blüthgen, 1920; *Hoplitis ebmeri* van der Zanden, 1989). Russia: **EP** (CR). – Europe (WE, SE, EE).
- Hoplitis (Pentadentosmia) laevifrons** (Morawitz, 1872) [Osmia]. Russia: ?**EP**, **UR**. – Europe (WE, SE, EE), Armenia, Turkey, Pakistan, India.
- Hoplitis (Pentadentosmia) nitidula** (Morawitz, 1877) [Osmia] (*Osmia transcaspica* Morawitz, 1888). Russia: **EP** (S). – Armenia, Iran, Pakistan, Kazakhstan.
- OSMIA** Panzer, 1806 (*Amblys* Klug, 1807; *Ceratosmia* Thomson, 1872; *Aceratosmia* Schmiedeknecht, 1885; *Pachyosmia* Ducke, 1900; *Monosmia* Tkalcú, 1974; *Orientosmia* Peters, 1978). Type species *Apis bicornis* Linnaeus, 1758. The genus is distributed in the Palaearctic and Nearctic regions, a few species also occur in the Oriental and Neotropical regions. Number of species: World – 339; Palaearctic – 204; Russia – 44.
- Osmia (Allosmia) rufohirta** Latreille, 1811 (*O. fulvohirta* Lepeletier de Saint-Fargeau, 1841;

- O. spineiventris* Giraud, 1857). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), N Africa, Georgia, Armenia, Turkey, Syria, Jordan, Israel, Iran, China.
- Osmia (Erythrosmia) andrenoides** Spinola, 1808. Russia: **EP** (NC, CR). – Europe (WE, SE, EE), Georgia, Turkey, Syria, Jordan, Israel, Iran.
- Osmia (Helicosmia) aurulenta** (Panzer, 1799) [Apis] (*Apis haematopoda* Panzer, 1801; *Osmia marginella* Lepeletier de Saint-Fargeau, 1841). Russia: **EP** (C, E, S, NC, CR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Lebanon, Iran.
- Osmia (Helicosmia) caerulescens** (Linnaeus, 1758) [Apis] (*Apis aenea* Linnaeus, 1761; *A. superbus* Harris, 1776; *A. muraria* Retzius, 1783, nom. praeocc., nec Geoffroy, 1785, nec Olivier, 1789; *A. cuprea* Geoffrey, 1785; *Andrena cupraria* Walckenaer, 1802; *Osmia rustica* Cresson, 1864, nom. praeocc., nec Geoffroy, 1785; *O. purpurea* Cresson, 1864). Russia: **EP** (C, E, S, NC, CR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Syria, Jordan, Israel, Iran, Central Asia, Kazakhstan, China, India, N America (introduced), New Zealand (introduced).
- Osmia (Helicosmia) cyanescens** Morawitz, 1875. Russia: **EP** (Müller, 2017). – Kyrgyzstan, Tajikistan, Kazakhstan.
- Osmia (Helicosmia) dimidiata** Morawitz, 1870. Russia: **EP** (NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Lebanon, Israel, Iran, Turkmenistan, Kyrgyzstan.
- Osmia (Helicosmia) leaiana** (Kirby, 1802) [Apis] (*Apis hirta* Geoffroy, 1785, nom. praeocc., nec Schrank, 1781; *Osmia atra* Schenck, 1853; *O. confusa* Morawitz, 1869; *O. solskyi* Morawitz, 1870; *O. truncatula* Thomson, 1872; *O. bidens* Pérez, 1879; *O. forsii* Alfken, 1924). Russia: **EP** (C, E, NC, CR), **UR**, **WS** (KM, AL), **ES** (KS). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Iran, Kazakhstan.
- Osmia (Helicosmia) melanogaster** Spinola, 1808 (*O. aterrima* Morawitz, 1872; *O. carniolica* Morawitz, 1872; *O. incerta* Radoszkowski, 1876). Russia: **EP** (NC, CR). – Europe (WE, NE, SE, EE), N Africa, Georgia, Armenia, Turkey, Syria, Jordan, Israel, Iran, ?China.
- Osmia (Helicosmia) niveata** (Fabricius, 1804) [Anthophora] (*Apis fulviventris* Panzer, 1798, nom. praeocc., nec Scopoli, 1763; *Osmia minuta* Bramson, 1879; *O. sieversi* Morawitz, 1886; *O. carneiventris* Dours, 1887; *O. fulviventris albiscopa* Alfken, 1914). Russia: **EP** (C, S, NC, CR). – Europe (WE, NE, SE, EE), N Africa, Georgia, Armenia, Turkey, Syria, Jordan, Israel, Iran, Turkmenistan, China.
- Osmia (Helicosmia) orientalis** Benoist, 1929. Russia: **FE** (AM, KH, PR, SA). – Japan (Hon, Shi, Kyu).
- Osmia (Helicosmia) signata** Erichson, 1835 (*O. jucunda* Smith, 1853; *O. vidua* Gerstäcker, 1869; *O. cincta* Dours, 1873; *O. laticincta* Pérez, 1879; *O. atriventris* Costa, 1883, nom. praeocc., nec Cresson, 1864; *O. costaniana* Dalla Torre et Friese, 1895). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), N Africa, Turkey, Syria, Jordan, Israel, Iran, Turkmenistan, China.
- Osmia (Hemiosmia) difficilis** Morawitz, 1875 (*O. falcatula* Morawitz, 1875; *O. alborufa* Alfken, 1935). Russia (Müller, 2017). – Turkey, Israel, Iran, Central Asia, Kazakhstan.
- Osmia (Hoplosmia) bidentata** Morawitz, 1876 (*O. affinis* Frivaldszky, 1877). Russia: **EP** (S, CR). – Europe (WE, SE, EE), N Africa, Armenia, Azerbaijan, Turkey, Syria, Jordan, Israel, Iran.
- Osmia (Hoplosmia) scutellaris** Morawitz, 1868 (*Heriades integra* Benoist, 1934). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Syria, Jordan, Israel, Iran.
- Osmia (Hoplosmia) spinulosa** (Kirby, 1802) [Apis] (*O. euchreiformis* Radoszkowski, 1882). Russia: **EP** (C, S, NC, CR), **UR**, **WS** (KM, AL), **ES** (KS). – Europe (WE, SE, EE), Armenia, Turkey, Kyrgyzstan, Kazakhstan.
- Osmia (Melanosmia) disjuncta** Tkalcú, 1995. Russia: **EP** (N). – Europe (NE), Mongolia.
- Osmia (Melanosmia) ephippiata** Smith, 1879. Russia: **ES** (IR). – Central Asia.
- Osmia (Melanosmia) inermis** (Zetterstedt, 1838) [*Anthophora*] (*O. globosa* Cresson, 1864; *O. vulpecula* Gerstäcker, 1869; *O. globosiformis* Cockerell, 1910). Russia: **EP** (N, C). – Europe (WE, NE, SE, EE), China, N America.
- Osmia (Melanosmia) laticeps** Thomson, 1872 (*O. hyperborea* Tkalcú, 1983). Russia: **EP** (N). – Europe (WE, NE, EE), N America.
- Osmia (Melanosmia) maritima** Friese, 1885. Russia: **EP** (N, C, E), **ES** (TU, KR, BR, YA, ZB), **FE** (KH, PR, SA, KA, CH). – Europe (WE, NE, EE), Mongolia, N America.
- Osmia (Melanosmia) nigriventris** (Zetterstedt, 1838) [*Anthophora*] (*O. baicalensis* Radoszkowski, 1867; *O. corticalis* Gerstäcker, 1869). Russia: **EP** (N, E, NC), **UR**, **WS** (TM), **ES** (BR, YA), **FE** (AM, KH, PR, SA, KA, MG, CH). – Europe (WE, NE, EE), Mongolia, China, N America.

- Osmia (Melanosmia) parietina** Curtis, 1828 (*Anthophora angustula* Zetterstedt, 1838; *Osmia vankovitzii* Radoszkowski, 1887). Russia: **EP (C), UR, WS (TM, KM), FE (MG)**. – Europe (WE, NE, SE, EE), Caucasus, Turkey.
- Osmia (Melanosmia) pilicornis** Smith, 1846. Russia: **EP (C, E), UR, WS (KM)**. – Europe (WE, NE, SE, EE), ?China (NE).
- Osmia (Melanosmia) uncinata** Gerstäcker, 1869. Russia: **EP (N, C, E), WS (TM, KM), ES (IR, YA, ZB), FE (AM, KH, PR, MG, CH)**. – Europe (WE, NE, SE, EE).
- Osmia (Melanosmia) xanthomelana** (Kirby, 1802) [*Apis*] (*O. fuciformis* Latreille, 1811; *O. chrysomelina* Panzer, 1812). Russia: **EP (N), UR**. – Europe (WE, NE, SE, EE).
- Osmia (Metallinella) brevicornis** (Fabricius, 1798) [*Eucera*] (*O. atrocaerulea* Schilling, 1849; *O. panzeri* Morawitz, 1869). Russia: **EP (S, NC, CR)**. – Europe (WE, NE, SE, EE), N Africa, Armenia, Azerbaijan, Turkey, Iran, Afghanistan, Central Asia, Kazakhstan.
- Osmia (Neosmia) bicolor** (Schrank, 1781) [*Apis*] (*Apis rustica* Geoffroy, 1785; *A. fusca* Christ, 1971; *A. hirundinaria* Christ, 1791; *Osmia pyrenaea* Lepeletier de Saint-Fargeau, 1841; *O. fusca* Gistel, 1857; *O. rufitarsis* Smith, 1879). Russia: **EP (C, E), WS (KM), ES (IR)**. – Europe (WE, NE, SE, EE), Central Asia.
- Osmia (Osmia) apicata** Smith, 1853 (*O. macroglossa* Gerstäcker, 1869). Russia (Müller, 2017). – Europe (WE, NE, EE), Georgia, Armenia, Turkey, Syria, Jordan, Lebanon, Israel, Iran.
- Osmia (Osmia) bicornis** (Linnaeus, 1758) [*Apis*] (*Apis rufa* Linnaeus, 1758). Russia: **EP (N, C, NC, CR), UR**. – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Syria, Israel, Iran, Turkmenistan, Kyrgyzstan, Kazakhstan.
- Osmia (Osmia) cerinthidis** Morawitz, 1876. Russia: **EP (S, NC, CR)**. – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, ?China.
- Osmia (Osmia) cornifrons** (Radoszkowski, 1887) [*Clalicodoma*]. Russia: **FE (KH, PR)**. – China, Korean Peninsula, Japan (Hok, Hon), N America (introduced).
- Osmia (Osmia) cornuta** (Latreille, 1805) [Megachile]. Russia: **EP (S, CR)**. – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Syria, Iran, Central Asia, Kazakhstan.

- Osmia (Osmia) melanocephala** Morawitz, 1875. Russia (Müller, 2017). – Central Asia, Kazakhstan, Mongolia, China (NW).
- Osmia (Osmia) mustelina** Gerstäcker, 1869. Russia: **EP (CR)**. – Europe (WE, SE, EE), N Africa, Georgia, Armenia, Turkey, Lebanon, Israel, Iran.
- Osmia (Osmia) opima** Romankova, 1985. Russia: **FE (KH, PR)**. – China (NE, NC), Korean Peninsula.
- Osmia (Osmia) pedicornis** Cockerell, 1920. Russia: **FE (KH, PR)**. – China (CE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Osmia (Osmia) taurus** Smith, 1873 (*O. matsumurae* Cockerell, 1911). Russia: **FE (KH, PR)**. – China (NE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Osmia (Pyrosma) cephalotes** Morawitz, 1870 (*O. longiceps* Morawitz, 1876; *O. bacillus* Pérez, 1879; *O. pulsata* Buysson, 1899). Russia: **EP (NC, CR)**. – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Syria, Jordan, Israel, Iran, Turkmenistan, Uzbekistan.
- Osmia (Pyrosma) gallarum** Spinola, 1808 (*O. ruborum* Dufour et Perris, 1840). Russia: **EP (CR)**. – Europe (WE, SE, EE), N Africa, Turkey.
- Osmia (Pyrosma) nana** Morawitz, 1874 (*O. tetrodonta* Benoist, 1934). Russia: **EP (S, NC)**. – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Syria, Jordan, Lebanon, Israel, Turkmenistan.
- Osmia (Pyrosma) submicans** Morawitz, 1870 (*O. giraudi* Schmiedeknecht, 1886). Russia: **EP (NC, CR)**. – Europe (WE, SE, EE), N Africa, Turkey, Syria, Jordan, Lebanon, Israel, Kazakhstan.
- Osmia (Pyrosma) versicolor** Latreille, 1811. Russia: **EP (CR)**. – Europe (WE, SE, EE), N Africa, Georgia, Turkey, Syria, Jordan, Lebanon, Israel.
- Osmia (Pyrosma) viridana** Morawitz, 1874 (*O. angulata* Pérez, 1896; *O. veridata mulleolus* van der Zanden, 1984). Russia: **EP (NC, CR)**. – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Syria, Jordan, Lebanon, Israel, Iran, Central Asia, Kazakhstan.
- Osmia (Tergosmia) tergestensis** Ducke, 1897 (*O. ononides* Ferton, 1897; *O. rondoui* Pérez, 1902; *O. wolhyonica* Noskiewicz, 1922; *O. atlantica* Benoist, 1934; *Anthocopa tergestensis remota* Tkalcú, 1979). Russia: **EP (CR)**. – Europe (WE, SE, EE), N Africa, Turkey, Kazakhstan.
- PROTOSMIA** Ducke, 1900 (*Rhodosmia* Michener, 1943). Type species *Heriades glutinosus* Giraud, 1871. The genus is mainly distributed in the

Palaearctic region with one species each occurring in the Nearctic and Oriental regions. Number of species: World – 27; Palaearctic – 25; Russia – 2.

Protosmia (Protosmia) tauricola Popov, 1961. Russia: EP (CR). – Europe (SE, EE), Turkey.

Protosmia (Protosmia) tiflensis (Morawitz, 1876) [Osmia] (*Osmia graeffei* Schmiedeknecht, 1890). Russia: EP (CR). – Europe (SE, EE), Georgia, Turkey, Jordan, Israel.

Tribe ANTHIDIINI

ANTHIDIELLUM Cockerell, 1904 (*Cerianthidium* Friese, 1923). Type species *Trachusa strigata* Panzer, 1805. The genus is widely distributed in the Palaearctic, Neotropical, Oriental, and Australian regions. Number of species: World – 61; Palaearctic – 7; Russia – 1.

Anthidiellum (Anthidiellum) strigatum (Panzer, 1805) [*Trachusa*] (*Anthidium minusculum* Nylander, 1852; *A. decoratum* Chevrier, 1972; *Dianthidium leucorhinum* Cockerell, 1924; *Anthidiellum pauperulum* Cockerell, 1928). Russia: EP (C, E, S, NC, CR), UR, WS (NS, KM, AL), ES (IR, BR, ZB), FE (AM, KH, PR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Syria, Lebanon, Israel, Iraq, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Korean Peninsula.

ANTHIDIUM Fabricius, 1804 (*Stenanthidium* Moure, 1947; *Tetranthidium* Moure, 1947; *Melanthidium* Cockerell, 1947; *Melanoanthidium* Tkalcú, 1967; *Echinanthidium* Pasteels, 1969; *Pontanthidium* Pasteels, 1969; *Ardenthidium* Pasteels, 1969; *Morphanthidium* Pasteels, 1969). Type species *Apis manicata* Linnaeus, 1758. The genus is distributed worldwide except Australia. Number of species: World – 186; Palaearctic – 90; Russia – 11.

Anthidium (Anthidium) amurensis Radoszkowski, 1876 (*A. comatum* Morawitz, 1896). Russia: ES (KH, KR, IR, BR, YA, ZB), FE (AM, KH, PR). – China (NW, NC).

Anthidium (Anthidium) cingulatum Latreille, 1809 (*A. oraniense* Lepeletier de Saint-Fargeau, 1841; *A. dissectum* Eversmann, 1852; *A. rufispinum* Costa, 1883; *A. manicatum* var. *undulatafasciatum* Friese, 1917). Russia: EP (C, S, CR), UR. – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Iran, Kyrgyzstan, Uzbekistan, Kazakhstan, China (NW), India.

Anthidium (Anthidium) diadema Latreille, 1809

(*A. albiventre* Lepeletier de Saint-Fargeau, 1841; *A. ornatum* Lepeletier de Saint-Fargeau, 1841; *A. radoszkowskyi* Mocsáry, 1887; *A. seraxense* Radoszkowski, 1893; *A. terminale* Morawitz, 1894; *A. diadema* var. *caucasicum* Friese, 1897; *A. diadema* var. *turcestanicum* Friese, 1897; *A. diadema* var. *obscurum* Friese, 1897). Russia: EP (NC, CR). – Europe (WE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Israel, Iran, Central Asia, Kazakhstan.

Anthidium (Anthidium) florentinum (Fabricius, 1775)

[*Apis*] (*A. subspinosum* Klug, 1832; *A. caucasicum* Radoszkowski, 1862; *A. florentinum* var. *hispanicum* Mocsáry, 1884; *A. florentinum* var. *ruiescente* Dusmet, 1908; *A. florentinum* var. *kissi* Alfken, 1935; *A. florentinum cypriacum* Mavromoustakis, 1949). Russia: EP (C, S, NC, CR), WS (NS, AL), ES (BR), FE (PR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Syria, Lebanon, Israel, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, China (NW).

Anthidium (Anthidium) loti Perris, 1852 (*Apis variegata* Fabricius, 1781, nom. praeocc., nec Linnaeus, 1758; *A. varia* Gmelin, 1790, nom. praeocc., nec Olivier, 1789; *Anthidium regulare* Eversmann, 1852; *A. mosaicum* Costa, 1863; *A. meridionale* Glraud, 1863; *A. quadriseriatum* Kriechbaumer, 1873; *A. quadriseriatum* var. *circumcinctum* Kriechbaumer, 1873; *A. variegatum* var. *incisum* Friese, 1917). Russia: EP (CR), UR. – Europe (WE, SE, EE), Azerbaijan, Turkey, Syria, Lebanon, Israel, Iran, Turkmenistan, Kazakhstan.

Anthidium (Anthidium) manicatum (Linnaeus, 1758) [*Apis pervigil* Harris, 1776; *A. maculata* Fabricius, 1781; *A. fulvipes* Vlllers, 1789, nec Fabricius, 1793; *A. modesta* Christ, 1791; *A. amoenita* Christ, 1791; *A. uncata* Schrank, 1802; *Anthidium marginatum* Latreille, 1809; *A. obtusatum* Lepeletier de Saint-Fargeau, 1841; *A. manicatum* var. *nigrithorax* Dalla Torre, 1877; *A. manicatum* var. *fasciatum* Schirmer, 1915; *A. manicatum* var. *nasicolle* Friese, 1917). Russia: EP (C, E, S, NC, CR), UR, WS (NS, KM, AL), ES (KS, TU, BR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Lebanon, Israel, Iran, Central Asia, Kazakhstan, China, N and S America (introduced), New Zealand (introduced).

Anthidium (Anthidium) montanum Morawitz, 1865

(*A. montanum* var. *flavomaculatum* Friese, 1897).

Russia: EP (CR). – Europe (WE, SE, EE), Tajikistan, Kyrgyzstan, China (NW, WP).

Anthidium (Anthidium) punctatum Latreille, 1809 (*A. minus* Nylander, 1848; *A. senile* Eversmann, 1852; *A. greyi* Radoszkowski, 1862; *A. albidulum* Chevrier, 1872; *A. nigrinum* Morawitz, 1893; *A. nitidulum* Morawitz, 1893; *A. mayeti* Pérez, 1895; *A. kohlii* Friese, 1897; *A. punctatum* var. *fulvipes* Friese, 1897; *A. punctatum* var. *albofasciatum* Friese, 1897; *A. punctatum* var. *bequaerti* Alfken, 1914; *A. amanuseNSE* Dusmet, 1915; *A. punctatum* var. *ariasi* Dusmet, 1915; *A. kohlii* var. *nigritulum* Friese, 1917; *A. punctatum* var. *sueicum* Alfken, 1927). Russia: EP (C, E, NC), UR, WS (KM, AL), ES (KS, TU, KR, IR, YA, ZB), FE (AM, KH, PR). – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey, Syria, Israel, Iran, Uzbekistan, Kyrgyzstan, Kazakhstan, China (NW, NC).

Anthidium (Anthidium) septemspinosum Lepeletier de Saint-Fargeau, 1841 (*A. nigripes* Eversmann, 1852; *A. nigriceps* Smith, 1854; *A. septemspinosum* var. *flavoguttatum* Friese, 1917; *A. dinurum* Cockerell, 1928). Russia: EP (C, E, S, CR), UR, WS (NS, KM, AL), ES (KS, TU, KR, IR, BR), FE (AM, KH, PR). – Europe (WE, SE, EE), Turkey, Afghanistan, Kazakhstan, Mongolia, China (CC, SW, SE), Korean Peninsula, Japan (Hon).

Anthidium (Proanthidium) fulviventre Friese, 1917. Russia: ES (IR). This species was even not mentioned since description and requires the revision.

Anthidium (Proanthidium) oblongatum (Illiger, 1806) [Anthophora] (*A. trochantericum* Morawitz, 1893; *A. oblongatum* var. *nigrum* Friese, 1897; *A. oblongatum xanthurum* Cockerell, 1925; *A. oblongatum australe* Alfken, 1937; *A. oblongatum* var. *flavens* Moczar, 1956; *Proanthidium tornense* Tkalcú, 1966). Russia: EP (C, NC, CR), UR, ES (TU). – Europe (WE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Iran, Turkmenistan, Tajikistan, Kyrgyzstan, Kazakhstan, N America (introduced).

BATHANTHIDIUM Mavromoustakis, 1953 (*Dianthidium*: Romankova, 1995a; *Lasanthidium* Romanova, 1995). Type species *Dianthidium bifoveolatum* Alfken, 1937. The genus is distributed in the Palaearctic and Oriental regions. Number of species: World – 13; Palaearctic – 11; Russia – 2.

Bathanthidium (Stenanthidiellum) malaisei (Popov, 1941) [Stelis]. Russia: FE (PR). – Korean Peninsula.

Bathanthidium (Stenanthidiellum) sibiricum (Eversmann, 1852) [Anthidium]. Russia: UR, WS (KM, AL), ES (BR, ZB), FE (AM, KH, PR). – Korean Peninsula, China (NE, NC, CC, SW).

EOANTHIDIUM Popov, 1950 (*Eoanthidiellum* Pasteels, 1969). Type species *Anthidium insulare* Morawitz, 1873. The genus is distributed in the Palaearctic, Oriental, and African regions. Number of species: World – 17; Palaearctic – 5; Russia – 1.

Eoanthidium (Eoanthidium) clypeare (Morawitz, 1874) [Anthidium] (*Anthidium elongatum* Friese, 1897; *Dianthidium hoplostomum* Mavromoustakis, 1945). Russia: EP (NC). – Europe (SE), Turkey, Syria, Jordan, Lebanon.

ICTERANTHIDIUM Michener, 1948. Type species *Anthidium limbiferum* Morawitz, 1875. The genus is distributed in the Palaearctic, Oriental, and African regions. Number of species: World – 24; Palaearctic – 20; Russia – 3.

Icteranthidium fedtschenkoi (Morawitz, 1875) [Anthidium] (*Anthidium transcaspicum* Lebedev, 1923). Russia: ES (TU). – Turkey, Israel, Iran, Pakistan, Central Asia, Kazakhstan.

Icteranthidium grohmanni (Spinola, 1838) [Anthidium] (*Anthidium latreillii* Lepeletier de Saint-Fargeau, 1841; *A. rubiginosum* Lepeletier de Saint-Fargeau, 1841; *A. provinciale* Lepeletier de Saint-Fargeau, 1841; *A. comptum* Lepeletier de Saint-Fargeau, 1841; *A. numida* Lepeletier de Saint-Fargeau, 1841; *A. lepeletieri* Fonscolombre, 1846; *A. klugi* Lucas, 1849; *A. coronatum* Smith, 1854; *A. latreillei* var. *obscurum* Dusmet, 1908; *Icteranthidium tergale* Pasteels, 1969). Russia: EP (NC, CR). – Europe (WE, SE, EE), N Africa, Armenia, Turkey, Syria, Lebanon, Israel, Iran, Kyrgyzstan.

Icteranthidium laterale (Latreille, 1809) [Anthidium] (*Anthidium quadrilobum* Lepeletier de Saint-Fargeau, 1841; *A. auripes* Eversmann, 1852; *A. annulare* Schenck, 1870; *A. sexmaculatum* Chevrier, 1872; *A. perrisi* Dours, 1873). Russia: EP (C, S, CR), UR, WS (NS, AL), ES (TU, KR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Iran, Turkmenistan, Kyrgyzstan, Kazakhstan, China (NW).

PSEUDOANTHIDIUM Friese, 1898 (*Paranthidiellum* Michener, 1948; *Paraanthidiellum* (!): Pasteels, 1969; *Carinellum* Pasteels, 1969; *Orientotrachusa* Gupta, 1993). Type species *Anthidium alpinum*

Morawitz, 1874. The genus is distributed in the Palaearctic, Oriental, and African regions. Number of species: World – 36; Palaearctic – 20; Russia – 8.

Pseudoanthidium (Pseudoanthidium) alpinum (Morawitz, 1874) [Anthidium] (*Anthidium alpinum* Morawitz, 1874; *A. barbatum* Mocsáry, 1884; *A. serraticeps* Friese, 1917; *Pseudoanthidium alpinum gregoriiense* Nobile, 1990). Russia: EP (NC). – Europe (SE, EE), Turkey.

Pseudoanthidium (Pseudoanthidium) eversmanni (Radoszkowski, 1886) [Anthidium]. Russia: UR. This species requires the revision; it was previously synonymized with "*Anthidium lituratum*" (Warncke, 1980) which is actually an unclear complex of different species (Aguib et al., 2010).

Pseudoanthidium (Pseudoanthidium) floripetum (Eversmann, 1852) [Anthidium]. Russia: UR. This species requires the revision; it was previously synonymized with "*Anthidium lituratum*" (Warncke, 1980) which is actually an unclear complex of different species (Aguib et al., 2010).

Pseudoanthidium (Pseudoanthidium) nanum (Mocsáry, 1881) [Anthidium] (*Apis liturata* Panzer, 1801, nom. praeocc., nec Gmelin, 1790; ?*Anthidium sinutatum* Lepeletier de Saint-Fargeau, 1841). Russia: EP (C, E, S, NC, CR), UR. – Europe (WE, SE, EE), Caucasus, Turkey, Israel, Iran, Pakistan, N America (introduced).

Pseudoanthidium (Pseudoanthidium) reptans (Eversmann, 1852) [Anthidium]. Russia: UR. This species requires the revision; it was previously synonymized with "*Anthidium lituratum*" (Warncke, 1980) which is actually an unclear complex of different species (Aguib et al., 2010).

Pseudoanthidium (Pseudoanthidium) tenellum (Mocsáry, 1881) [Anthidium]. Russia: EP (CR). – Europe (WE, EE), N Africa.

Pseudoanthidium (Royanthidium) melanurum (Klug, 1832) [Anthidium] (*Anthidium nigricolle* Morawitz, 1876; *A. dilobum* Pérez, 1895; *A. moricei* Friese, 1911; *A. royoii* Dusmet, 1915; *A. nigricolle* var. *flaviventre* Friese, 1917; *A. nigricolle* var. *flavicolle* Friese, 1917). Russia: EP (NC). – Europe (WE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Syria, Lebanon, Israel, Iran.

Pseudoanthidium (Royanthidium) reticulatum (Mocsáry, 1884) [Anthidium] (*Anthidium tegulare* Morawitz, 1886; *A. moscaryi* Friese, 1897; *A. reticulatum kindenmani* Mavromoustakis, 1954). Russia:

EP (CR). – Europe (WE, SE, EE), N Africa, Azerbaijan, Turkey, Lebanon, Israel, Iran.

STELIS Panzer, 1806 (*Gyrodroma* Klug, 1807; *Gymnus* Spinola, 1808; *Ceraplastes* Gistel, 1848; *Ghelynia* Provancher, 1888; *Melanostelis* Ashmead, 1898; *Stelidium* Robertson, 1902; *Microstelis* Robertson, 1903; *Pavostelis* Sladen, 1916; *Stelidina* Timberlake, 1941; *Stelidiella* (!): Timberlake, 1941; *Leucostelis* Noskiewicz, 1961). Type species *Apis aterrima* Panzer, 1798. Cleptoparasites of *Megachile*, *Osmia*, *Chelostoma*, *Heriades*, *Anthidium*, *Anthidiellum*, *Trachusa* (Megachilidae), *Ceratina*, *Anthophora* (*Clisodon*) (Apidae). The genus is distributed worldwide except Australia. Number of species: World – 118; Palaearctic – 30; Russia – 14.

Stelis (Heterostelis) annulata (Lepeletier de Saint-Fargeau, 1841) [Anthidium] (*S. freygessneri* Friese, 1885; *S. hispanica* Dusmet, 1921; ?*S. hungarica* Noskiewicz, 1962). Russia: EP (CR). – Europe (WE, SE, EE), N Africa, Turkey, Lebanon.

Stelis (Protostelis) signata (Latreille, 1809) [Anthidium] (*Anthidium parvulum* Lepeletier de Saint-Fargeau, 1841; *Stelis strigata* Kriechbaumer, 1874; *S. signata flavescentia* Friese, 1925). Russia: EP (C, E, S, NC, CR), UR. – Europe (WE, NE, SE, EE), N Africa, Armenia, Azerbaijan, Turkey, Syria, Lebanon, Israel, Iraq, Iran, Kazakstan.

Stelis (Stelidomorpha) nasuta (Latreille, 1809) [Anthidium] (*Anthidium nasale* Latreille, 1809). Russia: EP (S, CR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Syria, Lebanon, Israel, Iraq, Iran, Tajikistan, Uzbekistan, Kazakhstan.

Stelis (Stelis) aculeata Morawitz, 1880. Russia: EP (CR), ES (TU, BR). – Turkey, Central Asia, Kazakhstan, Mongolia, China (NE, NW).

Stelis (Stelis) breviuscula (Nylander, 1848) [Heriades] (*S. pygmaea* Schenck, 1853; *Heriades paxillorum* Chevrier, 1872). Russia: EP (NW, C, E, S, NC, CR), UR. – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Turkmenistan, China (NW, WP).

Stelis (Stelis) melanura Cockerell, 1924. Russia: ES (IR, BR), FE (KH, PR). – Mongolia, China (NC, NW, WP), Korean Peninsula.

Stelis (Stelis) odontopyga Noskiewicz, 1926. Russia: EP (CR). – Europe (WE, SE, EE), Azerbaijan, Turkey.

Stelis (Stelis) ornatula (Klug, 1807) [Gyrodroma] (*Trachusa sexpunctata* Stschegloff, 1826; *Stelis octomaculata* Smith, 1843; *S. sexsignata* Costa, 1858;

S. ornatula var. *immaculata* Noskiewicz, 1926). Russia: **EP** (NW, C, E, NC, CR), **UR**, **WS** (KM), **ES** (KR, IR, YA), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), N Africa, Turkey, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.

Stelis (Stelis) phaeoptera (Kirby, 1802) [Apis] (*Apis stigma* Christ, 1791; *A. aterrima* Christ, 1791, nom. praeocc., nec Panzer, 1798). Russia: **EP** (NW, C, E, S, NC, CR), **UR**. – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran, Uzbekistan, Kyrgyzstan, Kazakhstan.

Stelis (Stelis) punctulatissima (Kirby, 1802) [Apis] (*Apis aterrima* Panzer, 1798, nom. praeocc., nec Christ, 1791; *Stelis moravica* Tkalcu, 1971). Russia: **EP** (NW, C, E, NC, CR), **UR**, **WS** (NS, KM, AL). – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, Syria, Kazakhstan.

Stelis (Stelis) scutellaris Morawitz, 1894 (*S. scutellaris inamoena* Popov, 1933). Russia: **EP** (S, NC). – Europe (EE), Armenia, Azerbaijan, Turkey, Iran, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NE, NC, CC).

Stelis (Stelis) simillima Morawitz, 1876 (*S. cognata* Kohl, 1892; *S. genalis* Pasteels, 1969). Russia: **EP** (CR), **UR**, **WS** (TK, KM). – Europe (WE, SE, EE), N Africa, Armenia, Azerbaijan, Turkey, Syria, Israel, Iran, Turkmenistan.

Stelis (Pseudostelis) minima Schenck, 1861 (*S. minima* f. *unicolor* Alfken, 1944). Russia: **EP** (NW, C, NC), **UR**. – Europe (WE, NE, EE).

Stelis (Pseudostelis) minuta Lepeletier de Saint-Fargeau et Audinet-Serville, 1825 (*S. nana* Schenck, 1853). Russia: **EP** (N, NW, C, E, NC), **UR**. – Europe (WE, NE, SE, EE), N Africa, Armenia, Azerbaijan, Turkey, Lebanon.

TRACHUSA Panzer, 1804 (*Diphysis* Lepeletier de Saint-Fargeau, 1841; *Megachileoides* Radoszkowski, 1874; *Megachiloidea* (!): de Saussure, 1890). Type species *Trachusa serratulae* Panzer, 1804. The genus is distributed worldwide except Australia. Number of species: World – 48; Palaearctic – 12; Russia – 3.

Trachusa (Archianthidium) pubescens (Morawitz, 1872) [*Anthidium*] (*Anthidium pubescens* var. *maximum* Friese, 1931; *Archianthidium pubescens verhoeffi* Mavromoustakis, 1955). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), N Africa, Georgia, Turkey, Syria, Lebanon, Israel, Iran, Turkmenistan.

Trachusa (Paraanthidium) interrupta (Fabricius, 1781) [Apis] (*Apis rufipes* Fabricius, 1787, nom.

praeocc., nec Fabricius, 1781; *A. varia* Olivier, 1789, nom. praeocc., nec Gmelin, 1790; *A. erythropus* Gmelin, 1790; *A. fulvipes* Fabricius, 1793, nom. praeocc., nec Villers, 1789; *Anthidium flavilabre* Latreille, 1809; *A. luteipes* Lepeletier de Saint-Fargeau, 1841; *A. dufourii* Lepeletier de Saint-Fargeau, 1841; *A. integrum* Eversmann, 1852; *A. curvipes* Schmid, 1872; *A. melanostomum* Costa, 1884; *A. folivolutor* Ferton, 1921). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), N Africa, Azerbaijan, Turkey, Syria.

Trachusa (Trachusa) byssina (Panzer, 1798) [Apis] (*T. serratulae* Panzer, 1805; *Diphysis pyrenaica* Lepeletier de Saint-Fargeau, 1841; *Megachile resinana* Schilling, 1849; *M. rotundiventris* Perris, 1852; *Osmia bluethegeni* Maidl, 1922; *Trachusa byssina* var. *seitzi* Cockerell, 1925; *Megachile kychtacensis* Cockerell, 1928). Russia: **EP** (C, E, NC, ?CR), **UR**, **WS** (TK, KM), **ES** (KS, TU, KR, IR, ZB), **FE** (AM, KH). – Europe (WE, NE, SE, EE), Georgia, Armenia, Turkey, Mongolia.

Tribe DIOXYINI

AGLAOAPIS Cameron, 1901 (*Dioxoides* Popov, 1947). Type species *Aglaoopis brevipennis* Cameron, 1901. Cleptoparasites of *Hoplitis*, *Osmia*, and *Megachile* (Megachilidae). The distribution of the genus is disjunct: Europe and western Asia, western India, and South Africa. Number of species: World – 3; Palaearctic – 1; Russia – 1.

Aglaoopis tridentata (Nylander, 1848) [Coelioxys] (*Dioxys fasciata* Schenck, 1861; *D. kuntzei* Noskiewicz, 1940; *Dioxoides tridentata limassolica* Mavromoustakis, 1949). Russia: **EP** (C, E, S, CR), **UR**, **WS** (KM), **ES** (KS, TU, IR, BR, ZB), **FE** (KH, PR). – Europe (WE, NE, SE, EE), Azerbaijan, Turkey, Kyrgyzstan, Kazakhstan.

DIOXYS Lepeletier de Saint-Fargeau and Audinet-Serville, 1825 (*Hoplopasites* Ashmead, 1898; *Chrysopheon* Titus, 1901). Type species *Trachusa cincta* Jurine, 1807. Cleptoparasites of *Hoplitis*, *Osmia*, and *Megachile* (Megachilidae). The genus is distributed in the Holarctic region. Number of species: World – 18; Palaearctic – 12; Russia – 1.

Dioxys cinctus (Jurine, 1807) [Trachusa] (*D. pyrenaica* Lepeletier de Saint-Fargeau, 1841; *D. maura* Lepeletier de Saint-Fargeau, 1841; *D. cruenta* Gerstacker, 1869; *D. spinigera* Pérez, 1884; *D. cincta jucunda* Mocsáry, 1894; *D. cincta* ab. *friederikae* Mader,

1933). Russia: **EP** (CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Israel, ?Central Asia.

Tribe MEGACHILINI

COELIOXYS Latreille, 1809 (*Paracoelioxys* Gribodo, 1884). Type species *Apis conica* Linnaeus, 1758. Cleptoparasites of *Hoplitis*, *Osmia*, and *Megachile* (Megachilidae), *Anthophora* (Apidae). The genus is distributed worldwide except Australia. Number of species: World – 474; Palaearctic – 150; Russia – 23.

Coelioxys (Allococoelioxys) afer Lepeletier de Saint-Fargeau, 1841 (*C. coronata* Förster, 1853; *C. mandibularis* Chevreffir, 1872, nom. praeocc., nec Nylander, 1848; *C. afra* var. *tunensis* Gribodo, 1894). Russia: **EP** (S, NC, CR), **WS** (KM), **ES** (BR), **FE** (KH). – Europe (WE, NE, SE, EE), N and S Africa, Georgia, Armenia, Turkey, Israel, Saudi Arabia, Iran, Turkmenistan, Kyrgyzstan, Kazakhstan, China (NE, NC, NW, CC, SE), Indonesia.

Coelioxys (Allococoelioxys) brevis Eversmann, 1852 (*C. erythropyga* Förster, 1853; *C. erythropyga* var. *rubida* Gribodo, 1894; *C. inflatus* Alfken, 1933; *C. brevis armeniaca* Hedicke, 1938). Russia: **EP** (NC, CR), **UR**, **ES** (TU). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Israel, Tajikistan, Uzbekistan, Kazakhstan, China (NC, NW, CC, SW), Japan (Hon).

Coelioxys (Allococoelioxys) caudatus Spinola, 1838 (*C. foersteri* Morawitz, 1871). Russia: **EP** (S, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran, Central Asia, China (NW).

Coelioxys (Allococoelioxys) conspersus Morawitz, 1874. Russia: **EP** (NC).

Coelioxys (Allococoelioxys) echinatus Förster, 1853. Russia: **EP** (C, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Iraq, Iran.

Coelioxys (Allococoelioxys) elsei Schwarz, 2001. Russia: **EP** (CR). – Europe (SE, EE), Turkey, Jordan.

Coelioxys (Allococoelioxys) emarginatus Förster, 1853 (*C. minuta* Gistel, 1857; *C. robusta* Morawitz, 1875). Russia: **EP** (S, CR), **UR**, **WS** (NS, KM), **ES** (TU, BR), **FE** (AM, PR). – Europe (WE, SE, EE), N Africa, Armenia, Turkey, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan, China (NC, CC, SE), India.

Coelioxys (Allococoelioxys) formosicola Strand, 1913 (*C. taiwanensis* Cockerell, 1927; *C. manchurica*

Proshchalykin et Lelej, 2004). Russia: **FE** (PR). – China (SE), Japan (Hon, Kyu), Sri Lanka.

Coelioxys (Allococoelioxys) haemorrhoa Förster, 1853 (*C. pulchella* Morawitz, 1874). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Israel, Pakistan, Central Asia, China (NE), Korean Peninsula, India.

Coelioxys (Allococoelioxys) obtusus Pérez, 1884. Russia: **EP** (S, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Iraq, Iran, Turkmenistan.

Coelioxys (Allococoelioxys) polycentris Förster, 1853 (*C. conspersa* Morawitz, 1874; *C. orientalis* Friese, 1925). Russia: **EP** (CR), **ES** (BR). – Europe (WE, SE, EE), Turkey, Iran, Pakistan, China (NC).

Coelioxys (Boreocoelioxys) alatus Förster, 1853 (*C. montandoni* Gribodo, 1884). Russia: **EP** (C, E), **UR**, **WS** (KM), **ES** (IR, ZB), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), Turkey, China (NE), Japan (Hok, Hon).

Coelioxys (Boreocoelioxys) inermis (Kirby, 1802) [*Apis*] (*C. acuminata* Nylander, 1852, nom. praeocc., nec Gmelin, 1790; *C. divergens* Förster, 1853; *C. microdonta* Förster, 1853). Russia: **EP** (NW, N, C, E, S, CR), **UR**, **WS** (TM, OM, TK, KM), **ES** (BR, ZB), **FE** (AM, KH, PR, SA). – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, Kyrgyzstan, China (NE, NC), Japan (Hok, Hon).

Coelioxys (Boreocoelioxys) mandibularis Nylander, 1848 (*C. mandibularis olgae* Cockerell, 1924; *C. mandibularis jeholensis* Yasumatsu, 1935). Russia: **EP** (C, E, CR), **UR**, **WS** (KM, AL), **ES** (KS, TU, KR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU). – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, Kyrgyzstan, Mongolia, China (NE).

Coelioxys (Boreocoelioxys) rufescens Lepeletier de Saint-Fargeau et Audinet-Serville, 1825 (*C. umbrina* Smith, 1843; *C. hebescens* Nylander, 1848; *C. apiculata* Nylander, 1848; *C. trinacria* Förster, 1853; *C. diglypha* Förster, 1853; *C. carinata* Schenck, 1855, nom. praeocc., nec Smith, 1854; *C. longiuscula* Schenck, 1855; *C. obtusata* Schenck, 1855, nom. praeocc., nec Magretti, 1895; *C. parvula* Schenck, 1855; *C. fallax* Mocsáry, 1881; *C. rufescens* var. *agona* Alfken, 1912; *C. rufescens nigrescens* Cockerell, 1924; *C. rufescens* var. *turanica* Friese, 1925; *C. rufescens* var. *gigantea* Friese, 1925, nom. praeocc., nec Friese, 1922). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (KM), **ES** (KR, IR, YA, ZB), **FE** (AM, KH, PR, KU). – Europe (WE, NE, SE, EE), N Africa, Georgia, Armenia, Azerbaijan,

Turkey, Iran, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NW), Japan (Hon).

Coelioxys (Coelioxys) aurolimbatus Förster, 1853

(*C. apiculata* Förster, 1853, nom. praeocc., nec Nylander, 1848; *C. recurva* Schenck, 1853; *C. reflexa* Schenck, 1855; *C. ogivalis* Pérez, 1895; *C. aurolimbata* var. *algeriensis* Friese, 1895). Russia: EP (C, NC, CR), UR. – Europe (WE, NE, SE, EE), N Africa, Georgia, Armenia, Turkey, Lebanon, Iran, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NW).

Coelioxys (Coelioxys) conoideus (Illiger, 1806)

[Anthophora] (*C. vectis* Curtis, 1831; *C. decem-punctata* Spinola, 1838; *C. punctata* Lepeletier de Saint-Fargeau, 1841; *C. temporalis* Nylander, 1848; *C. ambigua* Schenck, 1855; *C. aegyptiaca* Radoszkowski, 1876; *C. spissicauda* Pasteels, 1968). Russia: EP (C, E, S, NC, CR), UR, WS (KM, AL), ES (TU, KR, BR), FE (AM, KH, PR). – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey.

Coelioxys (Coelioxys) elongatus Lepeletier de Saint-Fargeau, 1841 (*C. simplex* Nylander, 1852, nom. praeocc., nec Gmelin, 1790; *C. tricuspidata* Förster, 1853; *C. denticulata* Schenck, 1855; *C. stigmatica* Schenck, 1855; *C. distineta* Schenck, 1855; *C. sponsa* Smith, 1855; *C. tridenticulata* Schenck, 1861; *C. gracilis* Schenck, 1861; *C. obscura* Schenck, 1861; *C. claripennis* Schenck, 1870; *C. kudiana* Cockerell, 1924; *C. popovici* Friese, 1925). Russia: EP (C, E, CR), UR, WS (KM, AL), ES (BR, ZB), FE (AM, KH, PR, SA). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Iran.

Coelioxys (Coelioxys) lanceolatus Nylander, 1852 (*C. patula* Pérez, 1884). Russia: EP (N, C), ES (IR, BR, YA, ZB), WS (KM), FE (AM, KA). – Europe (WE, NE, SE, EE).

Coelioxys (Coelioxys) quadridentatus (Linnaeus, 1758) [Apis] (*Apis conica* Linnaeus, 1758; ?*A. agilis* Harris, 1776; *A. acuminata* Gmelin, 1790; *A. bidentata* Panzer, 1798; *Coelioxys acuta* Nylander, 1848; *C. fissidens* Förster, 1853; *C. fraterna* Förster, 1853; *C. convergens* Schenck, 1855; *C. truncata* Höppner, 1901). Russia: EP (C, E, CR), UR, WS (KM), ES (BR, YA, ZB), FE (AM, KH, PR, SA, KA, MG). – Europe (WE, NE, SE, EE), Turkey, Iran, China (NE, NW).

Coelioxys (uncertain) obtusispina Thomson, 1872. Russia: FE (AM, PR). – Europe (NE).

Coelioxys (uncertain) pielianus Friese, 1935. Russia: ES (BR), FE (AM, PR). – China (CC).

Coelioxys (uncertain) ruficinctus Cockerell, 1931 (*C. rufocincta* (!): Romankova, 1995a). Russia: FE (KH, PR). – China (CC).

MEGACHILE Latreille, 1802 (*Eumegachile* Friese, 1898; *Megalochila* Schulz, 1906; *Anthemois* Robertson, 1903; *Cyphopyga* Robertson, 1903; *Grosapis* Mitchell, 1980). Type species *Apis centuncularis* Linnaeus, 1758. The genus is distributed worldwide. Number of species: World – 1500; Palaearctic – 300; Russia – 45.

Megachile (Aethomegachile) remota Smith, 1879. Russia: FE (KH, PR). – China (NE, CC), Korean Peninsula, Japan (Hon, Kyu).

Megachile (Chalicodoma) desertorum Morawitz, 1875. Russia: ES (TU, BR). – Turkmenistan, Uzbekistan, Kazakhstan, Mongolia, China (NC).

Megachile (Chalicodoma) lefebvrei (Lepeletier de Saint-Fargeau, 1841) [Chalicodoma] (*Chalicodoma luctuosa* Dours, 1873; *Megachile lefebvrei* var. *albida* Prez, 1897; *M. lefebvrei* var. *albomaculata* Friese, 1898; *M. muraria* var. *variabilis* Friese, 1920). Russia: EP (NC, CR). – Europe (WE, SE, EE), N Africa, Georgia, Iran.

Megachile (Chalicodoma) parietina (Geoffroy, 1785) [Apis] (*Apis muraria* Olivier, 1789, nom. praeocc., nec Retzius, 1783; ?*A. picea* Christ, 1791; *A. varians* Rossi, 1792; *Megachile nestorea* Brüllé, 1832; *M. affinis* Brüllé, 1832; *M. aegyptica* Lepeletier de Saint-Fargeau, 1841; *Chalicodoma leueopogonata* Dours, 1873; *Megachile savignyi* Radoszkowski, 1874; *Chalicodoma valesina* Alfkén, 1931; *Megachile transitoria* Benoist, 1934; *Chalicodoma parietina carpathia* Tkalcú, 1992). Russia: EP (NC, CR). – Europe (WE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Syria, Jordan, Israel, Tajikistan, Uzbekistan, Kyrgyzstan, Kazakhstan.

Megachile (Creightonella) albisepta (Klug, 1817) [Anthophora] (*M. sericans* Fonscolombe, 1832; *M. caucasica* Lepeletier de Saint-Fargeau, 1841; *M. dufouri* Lepeletier de Saint-Fargeau, 1841; *M. odontura* Smith, 1849; *M. carinulata* Costa, 1882; *M. gerszabiensis* Radoszkowski, 1886; *M. germabensis* Radoszkowski, 1893). Russia: EP (NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Syria, Israel, Iran, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan.

Megachile (Eutricharaea) apicalis Spinola, 1808 (*M. mixta* Costa, 1863; *M. dimidiativentris* Dours, 1873; *M. massiliensis* Pérez, 1902). Russia: EP (C, S, NC, CR). – Europe (WE, NE, SE, EE), N Africa,

Georgia, Azerbaijan, Turkey, Israel, Iran, Pakistan, Tajikistan, Uzbekistan, Kazakhstan, China (NW), N America (introduced).

Megachile (Eutricharaea) deceptoria Pérez, 1890
(*M. mogadorensis* Benoist, 1934). Russia: **EP** (S, CR), **ES** (YA). – Europe (WE, SE, EE), N Africa, Azerbaijan, Turkey, Pakistan, Turkmenistan, Kazakhstan.

Megachile (Eutricharaea) giraudi Gerstäcker, 1869
(*M. vicina* Morawitz, 1894, nom. praeocc., nec Mocsáry, 1879; *M. intermixta* Gerstäcker, 1869; *M. bicoloriventris* Mocsáry, 1878; *M. giraudi erzurumensis* Tkalcú, 1980; *Chalicodoma atropyga* van der Zanden, 1995). Russia: **EP** (CR). – Europe (WE, SE, EE), Armenia, Turkey, Uzbekistan.

Megachile (Eutricharaea) leachella Curtis, 1828
(*M. dorsalis* Pérez, 1879; *M. argentata* var. *fossoria* Ferton, 1909; *M. leachella maadiensis* van der Zanden, 1986; *M. argentata* auct.). Russia: **EP** (S, NC, CR), **UR**, **WS** (KM), **ES** (BR, ZB), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Iran, China (NC, NW).

Megachile (Eutricharaea) leucomalla Gerstäcker, 1869 (*M. excellens* Morawitz, 1872; *M. laevifrons* Morawitz, 1878; *M. sedilloti* Pérez, 1895; *M. leucomalla* var. *panzeri* Friese, 1899). Russia: **EP** (CR). – Europe (WE, SE, EE), N Africa, Georgia, Turkey, Jordan, Kazakhstan.

Megachile (Eutricharaea) marginata Smith, 1853
(*M. picicornis* Morawitz, 1877; *M. albocineta* Pérez, 1879, nom. praeocc., nec Radoszkowski, 1874; *M. perezi* Mocsáry, 1887; *M. provincialis* Pérez, 1890; *Stelis megachiloides* Alfken, 1942). Russia: **EP** (CR). – Europe (WE, SE), N Africa, Azerbaijan, Turkey, Iraq, Iran, Afghanistan, Pakistan, Tajikistan, Kyrgyzstan.

Megachile (Eutricharaea) melanogaster Eversmann, 1852 (*M. sexmaculata* Alfken, 1942; *M. sexmaculata thracia* Tkalcú, 1979; *M. pilicrus flava* van der Zanden, 1998). Russia: **EP** (CR), **UR**. – Europe (WE, SE, EE), N Africa, Turkey, Kazakhstan.

Megachile (Eutricharaea) pilidens Alfken, 1924
(*M. argyrea* Cockerell, 1931). Russia: **EP** (C, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Georgia, Armenia, Turkey, Jordan, Kazakhstan.

Megachile (Eutricharaea) rotundata (Fabricius, 1787) [Apis] (*Apis pacifica* Panzer, 1798; *Megachile imbecilla* Gerstäcker, 1869; ?*M. nadia* Nurse, 1903). Russia: **EP** (C, S, NC, CR), **UR**, **WS** (KM), **ES** (KR, BR, YA, ZB), **FE** (AM, KH, PR). – Europe

(WE, NE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Iran, Pakistan, Turkmenistan, Kazakhstan, Mongolia, China (NE, NW), Japan (Hok), India, Africa, N and S America (introduced), Australia (introduced), New Zealand (introduced).

Megachile (Eutricharaea) rubrimana Morawitz, 1893. Russia: **FE** (PR). – Europe (SE, EE), Turkey, Tajikistan.

Megachile (Eutricharaea) semicircularis auct., nec van der Zanden, 1996. Russia: **EP** (CR). – Europe (SE), Turkey, Iran. The description of *M. semicircularis* by van der Zanden clearly points to this species, however the type series is composed of several taxa and the male holotype does not belong to this species but belongs to *M. apicalis* (Kuhlmann et al., 2017; C. Praz, in litt.).

Megachile (Megachile) alpicola Alfken, 1924
(*M. rubtzovi* Cockerell, 1928; *M. alpicola loeckenniana* Tkalcú, 1977). Russia: **EP** (NW, C, E), **WS** (NS, KM), **ES** (IR, BR, YA, ZB), **FE** (AM, KH, PR, SA). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Kazakhstan, Korean Peninsula.

Megachile (Megachile) bombycina Radoszkowski, 1874 (*M. maxillosa* Eversmann, 1852, nom. praeocc., nec Guerin, 1845). Russia: **EP** (C, E, S), **UR**, **WS** (TK, NS, KM, AL), **ES** (KS, BR, ZB), **FE** (AM, KH). – Europe (WE, NE, SE, EE), Georgia, China (NW).

Megachile (Megachile) centuncularis (Linnaeus, 1758) [Apis] (*M. parvula* Lepeletier de Saint-Fargeau, 1841; *M. appia* Nurse, 1903; *M. centuncularis theryi* Cockerell, 1931; *M. centuncularis nesiotica* Mavromoustakis, 1953). Russia: **EP** (N, C, E, S, NC, CR), **UR**, **WS** (KM), **ES** (BR, ZB), **FE** (AM, KH, PR, SA, MG). – Europe (WE, NE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Israel, Iran, Kazakhstan, China (NE, NC), Japan (Hok), India, Malasya, N America.

Megachile (Megachile) dacica Mocsáry, 1879. Russia: **ES** (BR). – Europe (EE). The status of this species remains in doubt, it may be a subspecies of *M. lapponica* (Praz, 2017).

Megachile (Megachile) genalis Morawitz, 1880
(*M. villosa* Henschel, 1888, nom. praeocc., nec Fabricius, 1775, nec Schenck, 1853; *M. buyssoni* Pérez, 1890; *M. henschelii* Dalla Torre, 1896; *M. genalis tortumensis* Tkalcú, 1980). Russia: **EP** (C, E, S, NC, CR), **WS** (KM), **FE** (AM, PR, SA). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Tajikistan, Kazakhstan, Mongolia, China (NW), Japan (Hok).

Megachile (Megachile) lapponica Thomson, 1872

(*M. melanopyga amaguella* Cockerell, 1924; *M. lapponica baicalica* Cockerell, 1928, nom. praeocc., nec *M. baicalica* Kokujev, 1927; *M. lapponica* var. *kurbati* Cockerell, 1928; *M. lapponica fuscifrons* Cockerell, 1930; *M. lapponica ishikawa* Hirashima, 1974). Russia: **EP** (N, C, E), **UR**, **WS** (TM, OM, NS, KM, AL), **ES** (IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KA, MG). – Europe (WE, NE, SE, EE), Kazakhstan, Korean Peninsula, Japan (Hok), N America.

Megachile (Megachile) ligniseca (Kirby, 1802) [Apis]

(*M. lignisecula* Cockerell, 1924; *M. ligniseca* var. *borealis* Niemelä, 1936). Russia: **EP** (N, C, E, NC, CR), **UR**, **WS** (KM), **ES** (IR, YA), **FE** (AM, KH, PR, SA, KU, KA). – Europe (WE, NE, SE, EE), Caucasus, Turkey, China (WP), Japan (Hok, Hon).

Megachile (Megachile) manipula Romankova, 1983.

Russia: **FE** (KH, PR).

Megachile (Megachile) melanopyga Costa, 1863

(*M. vestita* Giraud, 1863, nom. praeocc., nec Smith, 1853; *M. hymenaea* Gerstäcker, 1869; *M. melanopyga vulpecolor* Hedicke, 1938; *M. melanopyga zakakica* Mavromoustakis, 1957). Russia: **EP** (NC, CR), **WS** (KM), **ES** (BR). – Europe (WE, SE, EE), Azerbaijan, Turkey, Jordan, Israel, China, Korean Peninsula, Japan.

Megachile (Megachile) nipponica Cockerell, 1914.

Russia: **FE** (AM, KH, PR). – China (NE, NC), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Megachile (Megachile) octosignata Nylander, 1852.

Russia: **EP** (CR). – Europe (WE, SE, EE), Georgia, Turkey.

Megachile (Megachile) pilicrus Morawitz, 1877.

Russia: **EP** (NC, CR). – Europe (WE, SE, EE), Georgia, Turkey, Iran, Kyrgyzstan, China (NW).

Megachile (Megachile) versicolor Smith, 1844

(*M. rufiventris* Schenck, 1851; *M. distincta* Pérez, 1897; *M. semipleta* Cockerell, 1921; *M. versicolor* var. *hiberniae* Perkins, 1925; *M. pilicruriformis* Cockerell, 1928). Russia: **EP** (N, C, S, NC, CR), **UR**, **WS** (TM, KM), **ES** (YA, ZB), **FE** (AM, KH, PR, KA). – Europe (WE, NE, SE, EE), Azerbaijan, Turkey, Kazakhstan, Mongolia.

Megachile (Pseudomegachile) ericetorum Lepeletier de Saint-Fargeau, 1841

(*M. cristata* Dufour, 1841; *M. fasciata* Smith, 1844; *M. rufitarsis* Smith, 1844; *M. pyrina* Nylander, 1852, nom. praeocc., nec Lepeletier de Saint-Fargeau, 1841; *M. senex* Smith, 1853; *M. tsingtauensis* Strand, 1915; *M. ericetorum*

euroea Tkalcú, 1988; *M. ericetorum oraniensis* van der Zanden, 1989; *M. ericetorum melaleuca* van der Zanden, 1989). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, NE, SE, EE), N Africa, Armenia, Turkey, Syria, Kazakhstan, China (NW, CC, SE), N America (introduced).

Megachile (Pseudomegachile) saussurei Radoszkowski, 1874 (*M. multispinosa* Morawitz, 1875). Russia: **EP** (E, S). – Europe (SE), Turkey, Central Asia, Kazakhstan.

Megachile (Xanthosarus) analis Nylander, 1852

(*M. albicilla* Eversmann, 1852; *M. obscura* Eversmann, 1852; *M. kuennemanni* Alfken, 1897; *M. kuennemanni* var. *obscura* Alfken, 1897, nom. praeocc., nec *M. obscura* Eversmann, 1852; *M. analis* var. *albida* Friese, 1898; *M. analis chionura* Cockerell, 1924; *M. angarensis* Cockerell, 1928; *M. analis cana* Alfken, 1930; *M. analis* var. *holsatica* Alfken, 1936; *M. analis ulangomensis* Tkalcú, 1988; *M. analis batsumerensis* Tkalcú, 1988). Russia: **EP** (N, C, E, NC), **UR**, **WS** (KM), **ES** (IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KA, MG). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Turkmenistan, Tajikistan, Mongolia, China (NC, NW, CC, WP), Korean Peninsula.

Megachile (Xanthosarus) circumcincta (Kirby, 1802)

[Apis] (*M. circumcincta* var. *griseohirta* Alfken, 1929; *M. circumcincta* var. *fennica* Niemelä, 1936; *M. circumcincta* var. *insidiosa* Benoist, 1940; *M. circumcincta ozbeki* Tkalcú, 1977; *M. circumcincta numidica* Tkalcú, 1988; *M. circumcincta etnaensis* van der Zanden, 1989). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (KM), **ES** (IR, BR, YA, ZB), **FE** (KH, PR, SA, KU, KA, MG). – Europe (WE, NE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Iran, N America.

Megachile (Xanthosarus) fulvimana Eversmann, 1852 (*M. baicalica* Kokujev, 1927; *M. dybowskii* Cockerell, 1928). Russia: **ES** (KS, KR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KA). – Europe (EE), Kazakhstan, Mongolia.

Megachile (Xanthosarus) lagopoda (Linnaeus, 1761)

[Apis] (*M. pyrina* Lepeletier de Saint-Fargeau, 1841, nom. praeocc., nec Nylander, 1852; *M. lagopoda* var. *nigricans* Alfken, 1914; *M. baleina* Cockerell, 1928; *M. lagopoda albohirta* Alfken, 1930; *M. lagopoda pieli* Cockerell, 1931; *M. lagopoda* var. *fulvohirta* Alfken, 1935, nom. praeocc., nec Alfken, 1926; *M. furukawai* Yasumatsu, 1938; *M. lagopoda alkeni* van der Zanden, 1988). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (NS, KM, AL), **ES** (KS, KR,

BR, ZB), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey, Israel, Iran, Tajikistan, Kyrgyzstan, Kazakhstan, China (NW, CC, WP, SE), Korean Peninsula, Japan (Hok, Hon).

Megachile (Xanthosarus) maackii Radoszkowski, 1874. Russia: **EP** (N), **ES** (IR, BR, YA), **WS** (KM), **FE** (AM, KH, PR, MG). – Europe (WE, EE), Mongolia. The status of this species remains in doubt, it may be a subspecies of *M. nigriventris* (Praz, 2017).

Megachile (Xanthosarus) maritima (Kirby, 1802) [Apis] (*M. flaviventris* Schenck, 1853; *M. fuhescens* Smith, 1853, nom. praeocc., nec Walker, 1871; *Phyllotoma manicata* Dumeril, 1860, nom. praeocc., nec Giraud, 1861; *M. kashgarensis* Cockerell, 1913; *M. maritima continentalis* Hedicke, 1938). Russia: **EP** (C, S, NC, CR), **WS** (NS), **ES** (BR), **FE** (AM, KH, PR, CH). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Iran, Kyrgyzstan, Kazakhstan, China (CC), Korean Peninsula.

Megachile (Xanthosarus) nigriventris Schenck, 1870 (*M. curvicrus* Thomson, 1872; *M. hasticornis* Cockerell, 1924). Russia: **EP** (N), **UR**, **WS** (TU), **FE** (AM, MG). – Europe (WE, NE, SE, EE), Iran.

Megachile (Xanthosarus) willughbiella (Kirby, 1802) [Apis] (*M. atriventris* Schenck, 1853; *M. willughbiella kudiensis* Cockerell, 1924; *M. willughbiella provocata* Cockerell, 1924; *M. willughbiella* var. *hibernica*

Perkins, 1925; *M. centuncularis* var. *janssoni* Alfken, 1926; *M. korotnevi* Cockerell, 1928; *M. willughbiella munakatai* Hlrashima, 1974; *M. willughbiella zeraldae* Tkalcú, 1988). Russia: **EP** (N, C, E, S, NC, CR), **UR**, **WS** (TM, KM), **ES** (IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, MG). – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, Israel, Kyrgyzstan, China (NE, NC, NW), Japan (Hok, Hon).

Megachile (?) epovae Cockerell, 1928. Russia: **ES** (IR). This species was even not mentioned since description and requies the revision.

Megachile (?) ferritinata Cockerell, 1924. Russia: **FE** (PR). This species was even not mentioned since description and requies the revision.

Megachile (?) lucidula Mocsáry, 1901. Russia: **ES** (KR). This species was even not mentioned since description and requies the revision.

Megachile (?) scheviakovi Cockerell, 1928. Russia: **ES** (IR). This species was even not mentioned since description and requies the revision.

Megachile (?) sichotana Cockerell, 1924. Russia: **FE** (PR). This species was even not mentioned since description and requies the revision.

Megachile (?) polita Cockerell, 1924. Russia: **FE** (PR). This species was even not mentioned since description and requies the revision.

78. FAMILY APIDAE

**T.V. LEVCHENKO, A.M. BYVALTSEV
AND M.YU. PROSHCHALYKIN**

The most widespread family of bees, presenting by *Bombus* species even in Arctic island. Apidae includes many parasitic and nesting bees, both solitary (*Anthophora*, singly or in aggregations), communal (some *Eucera*) and social from primitive (*Ceratina*) to eusocial (*Apis*). Nests are builded mostly in ground, rare in stems, wood, empy holes or on open surface. Solitary species have one generation per year, cleptoparasites (*Nomada* and many other) have one or two generations depend on host life cycle. Larva of single nest parasites (*Melecta*) eats host egg and than host provision. Social parasites (*Psithyrus*) female drives out host queen and lives in her nest with wokers. The family is the most reach reperesented in New World, especially parasitic tribes. Many species are important orchard and crop pollinators. Beekeeping products of stingless bees (*Meliponini*) and honey bees (*Apini*) take significant role in human life and culture. Number of taxa: World – 176 genera, 5614 species; Palaearctic – 33/1358 species; Russia – 25/385 species.

References. Eversmann, 1852; Morawitz, 1874; Radoszkowski, 1888, 1893; Friese, 1895, 1896, 1897, 1901; Mocsáry, Szépligeti, 1901; Maidl, 1912; Keren-skij, 1919; Skorikov, 1922, 1931; Alfken, 1927, 1931; Lavrov, 1927; Wnukowski, 1927; Bischoff, 1930; Gur-vich, 1931; Popov, 1931, 1933a, 1947, 1948, 1950, 1951, 1954, 1956, 1958a, 1967a, b; Popov, Yasumatsu, 1935; Möschler, 1938; Stöckhert, 1944; Panfilov, 1951, 1956, 1957; 1981, 1982, 1984; Lith, 1956; Priesner, 1957; Iuga, 1958; Panfilov et al., 1961; Popov, Ponomareva, 1961; Schwarz, 1965, 1967a,b, 1980, 1981; Ponomareva, 1966, 1967; Elfving, 1968; Lieftinck, 1968, 1974, 1980; Løken, 1973; Pesenko, 1974; Tkalcú, 1961, 1974a, b; 1978, 1979, 1984a, b; Marikovskaya, 1979, 1982, 2001; Warncke, 1979, 1982b, 1983; Dathe, 1980b, 2009; Skhirtladze, 1981; Shumakova et al., 1982; Daly, 1983; Banaszak, 1984; Løken, 1984; Ito, 1985; Brooks, 1988; Pesenko, Sítdíkov, 1988; Sítdíkov, 1986, 1988; Sítdíkov, Pesenko, 1988; Kuznetsova, 1990; Williams, 1991, 1998; Banaszak, Ortiz-Sánchez, 1993; Alexander, Schwarz, 1994; Blagoveshchenskaya, Popova, 1994; Pagliano, 1994; Celary, 1995; Monsevičius, 1995; Rasmont, 1995; Rasmont et al., 1995; Terzo, Rasmont, 1997, 2004; Gogala, 1999; Kholodov, 1999; Risch, 1999, 2001; Konusova, Yanushkin, 2000; Lykov, 2000; Wu, 2000; Gusenleitner,

Schwarz, 2001; Schwarz, Gusenleitner, 2001, 2013, 2015, 2017; Davydova, Pesenko, 2002; Filatov, 2003, 2006, 2013, 2015; Iwata, 2003; Proshchalykin, 2004, 2007e, 2012a,b, 2013a; Sapaev, 2004; Chenikalova, 2005; Lopatin, Dobrynin, 2005; Proshchalykin, Kupianskaya, 2005, 2009; Ban-Calefariu, 2006; Danilov, 2006; Filatov et al., 2006; Amiet et al., 2007; Filatov, Ivanov, 2007; Mi-chener, 2007; Ortiz-Sánchez, Roberts, 2007; Standfuss, Schwarz, 2007; Straka et al., 2007; Byvaltsev, 2008, 2013; Engel, 2008, 2009; Mitai, Tadauchi, 2008; Right-myer, 2008; Schaefer, Renner, 2008; Scheuchl, 2008; Sidorov, Eremeeva, 2008; Williams et al., 2008, 2009, 2011, 2012, 2013, 2015, 2016; Ivanov et al., 2009, 2011; Quest, 2009; Sung et al., 2009; Aliev, Qamarli, 2010; Eardley, Urban, 2010; Grace, 2010; Konovalova, 2010; Proshchalykin, Lelej, 2010, 2013, 2014; Aliev, 2011; An et al., 2011; Ebmer, 2011; Lecocq et al., 2011, 2013, 2014, 2015; Levchenko, 2011, 2013; Szentgyörgyi et al., 2011; Guerschon, Ioneschu-Hirsch, 2012; Long et al., 2012; Shibaev, Polumordvinov, 2012; Straka, Rozen, 2012; Byvaltsev et al., 2013, 2016; Levchenko, Yuferev, 2013; Danilov, Byvaltsev, 2014; Kupianskaya et al., 2014; Levchenko, Tomkovich, 2014; Maharramov, Bayramov, 2014; Proshchalykin et al., 2014, 2017b; Tadauchi, Murao, 2014; Polumordvinov, Shibaev, 2015; Rasmont et al., 2015, 2017; Proshchalykin, Schwarz, 2017; Lev-chenko et al., 2017.

Subfamily XYLOCOPINAE

Tribe XYLOCOPINI

XYLOCOPA Latreille, 1802 (*Xilocopa* Latreille, 1802). Type species *Apis violacea* Linnaeus, 1758. Polylectic bees, borrowing nests in ground or wood, mostly tropical species. Number of taxa: World – 469; Palaearctic – 49; Russia – 6.

Xylocopa (Alloxylocopa) appendiculata Smith, 1852 (*Coptorthsoma appendiculata*: Popov, Ponomareva, 1961). Russia: **FE** (PR). – China (NE, NC, CC, SW, WP, SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Xylocopa (Copoxyla) iris (Christ, 1791) [*Apis*] (?*Apis nasuta* Christ, 1791; *Xilocopa cyanescens* Brülle, 1832; *X. minuta* Lepeletier de Saint-Fargeau, 1841; *X. canuta* Rondani, 1874). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Syria, Lebanon, Israel, Jordan, Iran, Afghanistan, Turkmenistan, Tajikistan, Kazakhstan.

Xylocopa (Proxylocopa) olivieri Lepeletier de Saint-Fargeau, 1841 (*X. binominata* Gribodo, 1894; *X. fusca* Smith, 1854; *X. hellenica* Spinola, 1843; *X. lana-ta* Smith, 1854; *X. lunata* Klug, 1807). Russia: **EP** (S, NC). – Europe (SE, EE), N Africa, Armenia, Azerbaijan, Turkey, Cyprus, Syria, Iraq, Iran, Turkmenistan.

Xylocopa (Proxylocopa) altaica Popov, 1947 (*X. nitidiventris altaica* Popov, 1947). Russia: **WS** (AL). – China (NC, NW).

Xylocopa (Xylocopa) valga Gerstäcker, 1872 (*X. ramulorum* Rondani, 1874; *X. convexa* Smith, 1878). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (KM, AL), **ES** (KS, TU, KR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Iraq, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NW, WP).

Xylocopa (Xylocopa) violacea (Linnaeus, 1758) [Apis] (*Apis insubrica* Müller, 1766; *Xylocopa femorata* Fabricius, 1804; *X. violacea* ab. *heteropennis* Vici domini, 2003). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Lebanon, Syria, Iraq, Iran, Turkmenistan.

Tribe CERATININI

CERATINA Latreille, 1802 (*Clavicera* Latreille, 1802; *Ceratinidia* Cockerell et Porter, 1899). Type species *Hylaeus albilabris* Fabricius, 1793. Polylectic bees, nesting in pithy dead stems or twigs, mostly tropic and subtropic species. Number of taxa: World – 199; Palaearctic – 67; Russia – 14.

Ceratina (Ceratina) cucurbitina (Rossi, 1792) [Apis] (*Hylaeus albilabris* Fabricius, 1793; *Ceratina decolorans* Brullé, 1832). Russia: **EP** (C, NC, CR). – Europe (WE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Israel, Jordan, China (CC).

Ceratina (Ceratina) satoi Yasumatsu, 1936. Russia: **ES** (ZB), **FE** (AM, PR). – China (NC), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu, Ryu).

Ceratina (Ceratinidia) flavipes Smith, 1879 (*C. flavovaria* Gussakovskij, 1932; *Ceratinidia flavipes*: Romankova, 1995b). Russia: **FE** (KH, AM, PR, SA). – China (NE, NC, CC, SW), Japan (Hok, Hon, Shi, Kyu).

Ceratina (Euceratina) acuta Friese, 1896. Russia: **EP** (S, CR), **UR**. – Europe (WE, SE, EE), Georgia, Azerbaijan, Turkey, Israel, Iran, Kazakhstan.

Ceratina (Euceratina) cyanea (Kirby, 1802) [Apis] (*C. nitidula* Spinola, 1806; *C. coerulea* Chevrier,

1872, nom. praeocc., nec Villers, 1789; *C. chevrieri* Tournier, 1876; *C. cyanea* var. *imitatrix* Markowsky, 1938). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (NS, KM), **ES** (KR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Cyprus, Iran, Turkmenistan, Kazakhstan, China (NW).

Ceratina (Euceratina) chalcites Germar, 1839 (*C. aenea* Brülle, 1832, nom. praeocc., nec Fabricius, 1798; *C. egregia* Gerstäcker, 1869). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), Caucasus, Turkey, Iran, Turkmenistan.

Ceratina (Euceratina) chalybea Chevrier, 1872 (*C. hungarica* Mocsáry, 1881; *C. callosa algeriensis* Friese, 1896; *C. c. cephalica* Cockerell, 1931; *C. callosa* auct., nec Fabricius, 1794). Russia: **EP** (C, E, S, NC, CR). – Europe (WE, SE, EE), N Africa, Abkhasia, Georgia, Azerbaijan, Turkey, Jordan, Kazakhstan.

Ceratina (Euceratina) chrysomalla Gerstäcker, 1869. Russia: **EP** (S). – Europe (SE), Azerbaijan, Turkey, Cyprus, Iran.

Ceratina (Euceratina) dallatorreana Friese, 1896. Russia: **EP** (CR). – Europe (WE, SE, EE), N Africa, Abkhasia, Turkey, Syria, Palestine, Israel, Central Asia, introduced into N America (California).

Ceratina (Euceratina) gravidula Gerstäcker, 1869. Russia: **EP** (CR). – Europe (WE, SE, EE), Georgia, Azerbaijan.

Ceratina (Euceratina) laevifrons Morawitz, 1894. Russia: **UR**. – Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan.

Ceratina (Euceratina) nigroaenea Gerstäcker, 1869. Russia: **EP** (S, NC, CR). – Europe (SE), Azerbaijan, Turkey, Israel.

Ceratina (Euceratina) nigrolabiata Friese, 1896. Russia: **EP** (CR). – Europe (WE, SE, EE), Azerbaijan, Turkey, Syria, Lebanon, Israel.

Ceratina (Euceratina) zwakhalsi Terzo et Rasmont, 1997. Russia: **EP** (NC). – Azerbaijan, Turkey, Iran, Turkmenistan, Kazakhstan.

Subfamily NOMADINAE

Tribe NOMADINI

NOMADA Scopoli, 1770 (*Hypochrotaenia* Holmberg, 1886; *Nomadita* Mocsáry, 1894; *Lamproapis* Cameron, 1902; *Centrias* Robertson, 1903; *Cephen* Robertson, 1903; *Gnathias* Robertson, 1903; *Holonomada* Robertson, 1903; *Xanthidium* Robertson,

1903; *Phor* Robertson, 1903; *Nomadosoma* Rohwer, 1911; *Polybiapis* Cockerell, 1916; *Acanthonomada* Schwarz, 1966). Type species *Apis ruficornis* Linnaeus, 1758. Cleptoparasites of *Andrena*, rarely *Panurgus* (Andrenidae), *Halictus*, *Lasioglossum* (Halictidae), *Eucera* (Apidae), *Melitta* (Melittidae) and *Colletes* (Colletidae). The only genus of Nomadini with the greatest diversity presented in Holarctic. Number of taxa: World – more than 800, Palaearctic – 280; Russia – 116.

Nomada albidemaculata Łoziński, 1922. Russia: **ES** (IR, BR, ZB).

Nomada alboguttata Herrich-Schäffer, 1839 (*N. ochrostroma* Zetterstedt, 1838, nom. praeocc., nec Kirby, 1802; *N. baccata* Smith, 1844; *N. quinquespinosa* Thomson, 1870; *N. laeta* Thomson, 1870; *N. alboguttata* var. *liburnica* Friese, 1921; *N. alboguttata* var. *suerinensis* Friese, 1921). Cleptoparasite of *Andrena argentata* Sm., *A. barbilabris* (Kby.) and probably of *A. ventralis* Imh. (Andrenidae). Russia: **EP** (N, NW, C, E, S, CR), **UR**, **ES** (KR), **FE** (KA). – Europe (WE, NE, SE, EE), Georgia, Azerbaijan, Japan (Hok, Hon, Kyu).

Nomada amurensis Radoszkowski, 1876 (*N. esakii* Yasumatsu et Hirashima, 1953; *N. sudai* Tsuneki, 1976). Russia: **WS** (AL), **ES** (KS, IR), **FE** (PR). – Kazakhstan, Japan (Hok, Hon, Shi, Kyu).

Nomada argentata Herrich-Schäffer, 1839 (*N. atrata* Smith, 1846; *N. brevicornis* Schmiedeknecht, 1882). Cleptoparasite of *Andrena argentata* (H.-Sch.) (Andrenidae). Russia: **EP** (NW, C). – Europe (WE, NE, SE, EE).

Nomada armata Herrich-Schäffer, 1839 (*N. rostrata* Lepeletier de Saint-Fargeau, 1841, nom. praeocc., nec Herrich-Schäffer, 1839; *N. lanceolata* Lepeletier de Saint-Fargeau, 1841; *N. compta* Lepeletier de Saint-Fargeau, 1841; *N. kirbyella* Stephens, 1846; *N. cineticornis* Nylander, 1848). Cleptoparasite of *Andrena hattofiana* (F.) (Andrenidae). Russia: **EP** (N, NW, C, E, CR), **UR**. – Europe (WE, NE, SE, EE), Georgia, Azerbaijan.

Nomada arrogans Schmiedeknecht, 1882. Russia: **EP** (CR). – Europe (EE, SE), Turkey, Israel, Tajikistan.

Nomada atroscutellaris Strand, 1921. Cleptoparasite of *Andrena viridescens* Vier. (Andrenidae). Russia: **EP** (C, CR). – Europe (WE, SE, EE), Turkey.

Nomada basalis Herrich-Schäffer, 1839 (*N. rubra* Smith, 1849; *N. flavomaculata* Lucas, 1849; *N. bimaculata* Eversmann, 1852, nom. praeocc., nec

Schilling, 1849; *N. regalis* Morawitz, 1870; *N. tripunctata* Morawitz, 1872; *N. andalusica* Schmiedeknecht, 1882; *N. astarte* Pérez, 1895; *N. emendata* Schulz, 1906; *N. flavomaculata* var. *carnica* Friese, 1921; *N. grossa* Friese, 1921; *N. regalis* var. *indereskiana* Łoziński, 1922). Cleptoparasite of *Andrena truncatilabris* Mor. (Andrenidae). Russia: **EP** (S, CR), **UR**, **WS** (NS). – Europe (WE, SE, EE), N Africa, Armenia, Turkey, Syria, Palestine, Israel, Iraq, Iran, Turkmenistan, Kazakhstan.

Nomada bifasciata Olivier, 1811 (*N. zonata* Panzer in Lepeletier de Saint-Fargeau, 1841; *N. lepeletieri* Pérez, 1884; *N. navasi* Dusmet, 1913; *N. pusilla* auct., nec Lepeletier de Saint-Fargeau, 1841). Cleptoparasite of *Andrena gravida* Imh. (Andrenidae). Russia: **EP** (CR). – Europe (WE, SE, EE), N Africa, Turkey.

Nomada bispinosa Mocsáry, 1883 (*N. excisa* Pérez, 1890). Probably cleptoparasite of *Andrena hypopolia* Schmied. (Andrenidae). Russia: **EP** (CR). – Europe (WE, SE, EE), Azerbaijan, Turkey.

Nomada blepharipes Schmiedeknecht, 1882. Russia: **UR**, **ES** (IR). – Europe (WE, SE, EE), Turkey, Kazakhstan.

Nomada bluethgeni Stoeckhert, 1944. Cleptoparasite of *Lasioglossum marginellum* (Schck.) (Halictidae). Russia: **EP** (C, S, NC, CR). – Europe (WE, SE, EE), Azerbaijan.

Nomada braunsiana Schmiedeknecht, 1882. Cleptoparasite of *Andrena pandellei* (Pér.) and probably of *A. curvungula* Thoms. (Andrenidae). Russia: **EP** (C, S, NC, CR). – Europe (WE, SE, EE), Turkey, Jordan.

Nomada calimorpha Schmiedeknecht, 1882 (*N. copeatica* Morawitz, 1894). Russia: **EP** (E, S, CR). – Europe (WE, SE, EE), Turkey.

Nomada castellana Dusmet, 1913 (*N. baeri* E. Stoeckhert, 1930). Probably cleptoparasite of *Andrena semilaevis* Pér. (Andrenidae). Russia: **EP** (C, E, S), **ES** (IR). – Europe (WE, NE, SE, EE), Azerbaijan, Kazakhstan.

Nomada comparata Cockerell, 1911 (*N. maculifrons* var. *comparata* Cockerell, 1911). Probably cleptoparasite of *Andrena mikado* Strand et Yasumatsu and *A. richardsi* Hirashima (Andrenidae). Russia: **FE** (PR). – Japan (Hok, Hon, Shi, Kyu).

Nomada conjungens Herrich-Schäffer, 1839 (*N. dailaturreana* Schmiedeknecht, 1882; *N. bipunctis* Pérez, 1902). Cleptoparasite of *Andrena proxima* (Kby.) (Andrenidae). Russia: **EP** (NW, C, E, S). – Europe (WE, SE, EE), N Africa, Turkey, Kazakhstan.

- Nomada corcyraea** Schmiedeknecht, 1882 (*N. transitoria* Schmiedeknecht, 1882). Russia: **EP** (S). – Europe (SE, EE), Azerbaijan, Turkey.
- Nomada coxalis** Morawitz, 1877. Russia: **EP** (CR), **UR**. – Europe (SE), Azerbaijan, Turkey, Israel, Iran, Turkmenistan.
- Nomada cruenta** Schmiedeknecht, 1882 (*N. sexspinosa* Friese, 1921; *N. chrysopyga* var. *rubescens* Lozinski, 1922). Cleptoparasite of *Andrena scita* Ev. (Andrenidae). Russia: **EP** (S, CR). – Europe (WE, SE, EE), Azerbaijan, Turkey.
- Nomada diacantha** Schwarz, 1981. Russia: **EP** (S, CR). – Europe (SE, EE), Turkey, Syria.
- Nomada distinguenda** Morawitz, 1874 (*N. rugithorax* Pérez, 1902). Cleptoparasite of *Lasioglossum villosum* (Kby.) and probably of *L. parvulum* (Schck.) (Halictidae). Russia: **EP** (C, S, NC, CR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Syria, Afghanistan, Turkmenistan.
- Nomada dybovskij** Radoszkowski, 1876. Russia: **ES** (IR).
- Nomada ecarinata** Morawitz, 1888 (*N. munakatai* Tsuneki, 1973). Russia: **EP** (C), **UR**, **WS** (TK, AL), **ES** (KS, KR, IR, BR, ZB), **FE** (PR). – Mongolia, China (NE), Japan (Hok, Hon).
- Nomada emarginata** Morawitz, 1877. Cleptoparasite of *Melitta haemorrhoidalis* (F.) (Melittidae). Russia: **EP** (C, S, NC), **UR**. – Europe (WE, NE, SE, EE), Georgia, Azerbaijan.
- Nomada errans** Lepeletier de Saint-Fargeau, 1841 (*N. errans* var. *korlevicina* Friese, 1921; *N. errans* var. *sibirica* Friese, 1921; *N. transylvatica* Zilahi-Kiss, 1927). Cleptoparasite of *Andrena nitidiuscula* Schck. and *A. pallitarsis* Pér. (Andrenidae). Russia: **EP** (C, E), **UR**, **ES** (IR). – Europe (WE, SE, EE), Kazakhstan.
- Nomada erythrocephala** Morawitz, 1871. Russia: **EP** (S, NC). – Europe (WE, SE, EE), Armenia, Turkey, Cyprus.
- Nomada esana** Tsuneki, 1973. Russia: **FE** (AM, KH, PR). – Korean Peninsula, Japan (Hok, Hon).
- Nomada fabriciana** (Linnaeus, 1767) [Apis] (*N. germanica* Panzer, 1799, nom. praeocc., nec Fabricius, 1804; *Apis fabriciella* Kirby, 1802; *A. quadri-notata* Kirby, 1802; *Nomada nigricornis* Olivier, 1811; *N. nigrita* Schenck, 1861, nom. praeocc., nec Fabricius, 1775, nec Panzer, 1800, nec Pérez, 1895; *N. fabriciana* var. *aestivalis* Friese, 1921; *N. fabriciana* var. *ruficrus* E. Stoeckhert, 1930). Cleptoparasite of *Andrena bicolor* F., *A. chrysosceles* (Kby.), *A. angustior* (Kby.) and probably of *A. nigroaenea* (Kby.) (Andrenidae). Russia: **EP** (NW, C, E, CR), **UR**, **WS** (TK), **ES** (KS, IR). – Europe (WE, NE, SE, EE), Azerbaijan.
- Nomada facilis** Schwarz, 1967. Probably cleptoparasite of *Andrena humilis* Imh. (Andrenidae). Russia: **EP** (C). – Europe (WE, NE, SE, EE), Azerbaijan, Kazakhstan.
- Nomada femoralis** Morawitz, 1869 (*N. dentipes* Rudow, 1879; *N. femoralis* var. *flavopunctata* Friese, 1921). Cleptoparasite of *Andrena humilis* Imh. (Andrenidae). Russia: **EP** (C, CR), **UR**. – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey, Syria, Palestine, Iran.
- Nomada ferruginata** (Linnaeus, 1767) [Apis] (*Apis xanthosticta* Kirby, 1802; *Nomada lateralis* Panzer, 1804; *N. bridgmaniana* Smith, 1876; *N. lateralis* var. *blancoburgensis* Schmiedeknecht, 1882; *N. lateralis* var. *megapolitana* Schmiedeknecht, 1882). Cleptoparasite of *Andrena praecox* (Scop.) (Andrenidae). Russia: **EP** (NW, C, E, S), **UR**. – Europe (WE, NE, SE, EE), Caucasus, Uzbekistan.
- Nomada flava** Panzer, 1798 (*N. intermedia* Imhoff, 1834; *N. proteus* Lepeletier de Saint-Fargeau, 1841; *N. hirsuta* Verhoeff, 1890). Cleptoparasite of *Andrena carantonica* (Pér.), *A. nitida* (Müll.), *A. fulva* (Müll.) and probably of *A. nigroaenea* Kby. (Andrenidae). Russia: **EP** (C, E, S), **UR**. – Europe (WE, NE, SE, EE), Caucasus, Kazakhstan.
- Nomada flavoguttata** (Kirby, 1802) [Apis] (*Apis rufocincta* Kirby, 1802; *Nomada minuta* Fabricius, 1804, nom. praeocc., nec Swenk, 1913; *N. nana* Schenck, 1874; *N. pygmaea* Schenck, 1874, nom. praeocc., nec Cresson, 1863; *N. flavoguttata* var. *serotina* Schmiedeknecht, 1882; *N. flavoguttata* var. *hoeppneri* Alfken, 1898; *N. aljkeni* Cockerell, 1907, nom. praeocc., nec Gaunitz, 1935, nec Noskiewicz, 1939; *N. kurilensis* Yasumatsu, 1939; *N. flavoguttata japonensis* Tsuneki, 1973; *N. tridentata* Tsuneki, 1986). Cleptoparasite of *Andrena minutula* (Kirby), *A. minutuloides* Perkins, *A. falsifica* Perkins, *A. subopaca* Nylander and *A. semilaevigata* Pérez (Andrenidae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, TK, AL), **ES** (KS, TU, IR, YA), **FE** (PR, KU). – Europe (WE, NE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Iran, Turkmenistan, Tajikistan, Kazakhstan, China (SE), Korean Peninsula, Japan (Hok, Hon, Shi, Kyu), India.
- Nomada flavopicta** (Kirby, 1802) [Apis] (*N. jacobaeae* var. *miranda* Schmiedeknecht, 1882; *N. jacobaeae*

var. haematodes Schmiedeknecht, 1882; *N. jacobaeae* var. *konowiana* Friese, 1921; *N. jacobaeae* var. *parva* Friese, 1921, nom. praeocc., nec Robertson, 1900; *N. jasnitskii* Cockerell, 1928. Cleptoparasite of *Melitta leporina* (Pz.), *M. haemorrhoidalis* (F.), *M. tricincta* Kby. and probably of *M. nigricans* Alfk. (Melittidae). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (OM, TK, AL), **ES** (KS, TU, KR, IR, BR, ZB). – Europe (WE, NE, SE, EE), Georgia, Azerbaijan, Tajikistan, Uzbekistan, Kazakhstan.

Nomada fucata Panzer, 1798 (*N. varia* Panzer, 1798; *N. fucata* var. *taeniata* Gribodo, 1894; *N. f.* var. *iberica* Dalla Torre et Friese, 1894; *N. f.* var. *pretiosa* Friese, 1921; *N. f.* var. *flavescens* Friese, 1921, nom. praeocc., nec Friese, 1916; *N. f.* var. *caucasica* Lozinski, 1922). Cleptoparasite of *Andrena flavi-pes* Pz. (Andrenidae). Russia: **EP** (NW, C, S, NC, CR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Iraq, Afghanistan, Pakistan, Turkmenistan, Uzbekistan, Kyrgyzstan, Kazakhstan.

Nomada fulvicornis Fabricius, 1793 (*N. lineola* Panzer, 1798; *Apis sexcincta* Kirby, 1802, nom. praeocc., nec Fabricius, 1775; *A. capreae* Kirby, 1802; *A. cornigera* Kirby, 1802, nom. praeocc., nec Rossi, 1790; *A. subcornuta* Kirby, 1802; *Nomada ventralis* Imhoff, 1834; *N. rubra* Eversmann, 1852, nom. praeocc., nec Geofrey, 1785, nom. praeocc., nec Provancher, 1888, nec Smith, 1849; *N. robusta* Morawitz, 1871; *N. aurantiaca* Morawitz, 1875; *N. schmiedeknechti* Schmiedeknecht, 1882; *N. lineola* var. *aurigena* Schmiedeknecht, 1882; *N. l.* var. *melanothorax* Schmiedeknecht, 1882; *N. l.* var. *meridionalis* Schmiedeknecht, 1882; *N. l.* var. *flavolimbata* Schmiedeknecht, 1882; *N. l.* var. *latistriga* Schmiedeknecht, 1882; *N. l.* var. *rossica* Schmiedeknecht, 1882; *N. l.* var. *sagemeqli* Schmiedeknecht, 1882; *N. pallidenotata* Schmiedeknecht, 1882; *N. l.* var. *diluta* Gribodo, 1893, nom. praeocc., nec Pérez, 1884; *N. l.* var. *melanocera* Gribodo, 1893; *N. l.* var. *sagemeqli* Dallatorre, 1896; *N. lineola* var. *sibirica* Mocsáry, 1901; *N. l.* var. *bremensis* Alfken, 1913; *N. l.* var. *erubescens* Friese, 1921; *N. rakosiensis* Friese, 1921; *N. lineola* var. *nigrina* Lozinski, 1922; *N. l.* var. *aurolimbata* Lozinski, 1922; *N. l.* var. *brunneofasciata* Lozinski, 1922; *N. l.* var. *rufomaculata* Lozinski, 1922; *N. l.* var. *quadrivittata* Lozinski, 1922; *N. l.* var. *eversmanni* Lozinski, 1922; *N. belikovi* Cockerell, 1928; *N. lineola* var. *decorata* Markowsky, 1939, nom. praeocc., nec Smith, 1854). Cleptoparasite of *Andrena tibialis* (Kby.), *A. agilissima* (Scop.), *A. pilipes* F.,

A. bimaculata (Kby.) and *A. thoracica* (F.) (Andrenidae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (OM, TK, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (PR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, UAE, Pakistan, Turkmenistan, Uzbekistan, Japan (Hon, Hok).

Nomada furva Panzer, 1798. Cleptoparasite of *Lasioglossum leucopus* (Kby.), *L. morio* (F.), *L. punctatissimum* (Schck.) (Halictidae). Russia: **EP** (C, E, S, NC), **UR**, **WS** (TK), **FE** (PR). – Europe (WE, SE, EE), N Africa, Georgia, Turkey, Turkmenistan, Uzbekistan, Kazakhstan.

Nomada furvoides Stoechert, 1944. Cleptoparasite of *Lasioglossum politum* (Schck.) and probably of *L. semilucens* (Alfk.) (Halictidae). Russia: **EP** (S). – Europe (WE, SE, EE), Turkey.

Nomada fusca Schwarz, 1986. Russia: **EP** (NW, C), **WS** (AL), **ES** (KS, IR, YA), **FE** (KH, PR). – Europe (NE), Japan (Hok, Hon).

Nomada fuscicornis Nylander, 1848 (*N. megacephala* Schenck, 1874). Cleptoparasite of *Panurgus calceatus* (Scop.) (Andrenidae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TK, AL). – Europe (WE, NE, SE, EE).

Nomada ginran Tsuneki, 1973 (*N. yasha* Tsuneki, 1986). Russia: **FE** (PR). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Nomada glaberrima Schmiedeknecht, 1882 (*N. graja* Mocsáry, 1883). Russia: **EP** (C, S, CR). – Europe (SE), Turkey, Lebanon, Syria.

Nomada glaucopis Pérez, 1890. Russia: **UR**. – Europe (WE, SE, EE), N Africa, Israel, Kazakhstan.

Nomada goodeniana (Kirby, 1802) [*Apis alternata* Kirby, 1802; *Nomada Integra* Imhoff, 1834, nom. praeocc., nec Brüllé, 1832, nec Robertson, 1893; *N. cineta* Herrich-Schäffer, 1839, nom. praeocc., nec Rossi, 1792, nec Lepeletier de Saint-Fargeau, 1841; *Nomada batava* Vollenhoven, 1858; *N. succineta* var. *lineolata* Friese, 1921; *N. scheviakovi* Cockerell, 1928; *N. goodeniana danuvia* Pittioni, 1951]. Cleptoparasite of *Andrena nitida* (Müll.), *A. nigroaenea* (Kby.), *A. cineraria* (L.), *A. tibialis* (Kby.) and *A. thoracica* (F.) (Andrenidae). Russia: **EP** (N, NW, C, E, S, CR), **UR**, **WS** (TM, AL), **ES** (IR). – Europe (WE, NE, SE, EE), Azerbaijan, Turkey, Kyrgyzstan, Kazakhstan.

Nomada guttulata Schenck, 1861 (*N. rufilabris* Thomson, 1870; *N. mishimana* Tsuneki, 1976). Cleptoparasite of *Andrena labiata* F. and probably of *A. potentillae* Pz. (Andrenidae). Russia: **EP** (C, E, S), **UR**,

- ES (KS), FE (PR).** – Europe (WE, NE, SE, EE), Kazakhstan, Japan (Hon, Kyu).
- Nomada hakusana** Tsuneki, 1973. Russia: **FE (KH, SA, KU).** – Japan (Hok, Hon).
- Nomada hammarstroemi** Morawitz, 1888. Russia: **WS (AL), ES (KR, IR, BR).**
- Nomada hera** Schwarz, 1965. Russia: **EP (NC).** – Europe (SE), Turkey.
- Nomada hungarica** Dalla Torre et Friese, 1894 (*N. scita* Schmiedeknecht, 1882, nom. praeocc., nec Cresson, 1878; *N. lagreciae* Nobile, 1990). Russia: **EP (S), UR.** – Europe (SE, EE), Turkey.
- Nomada immaculata** Morawitz, 1874. Russia: **EP (S, NC).** – Europe (SE, EE), Armenia, Cyprus, Afghanistan, Pakistan.
- Nomada imperialis** Schmiedeknecht, 1882. Probably cleptoparasite of *Eucera pannonica* Mocs. (Apidae). Russia: **EP (NC, CR).** – Europe (SE), Turkey.
- Nomada incisa** Schmiedeknecht, 1882. Russia: **EP (S).** – Europe (WE, SE, EE), Cyprus.
- Nomada integra** Brüllé, 1832 (*N. germanica* Fabricius, 1804, nom. praeocc., nec Panzer, 1799; *N. ferruginata* var. *cinctiventris* Friese, 1921; *N. stigma* auct., nec Fabricius, 1805; *N. ferruginata* auct., nec Linnaeus, 1767). Cleptoparasite of *Andrena humilis* Imh. and *A. taraxaci* Gir. (Andrenidae). Russia: **EP (N, NW, C, E, S, NC), UR, ES (KR).** – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Cyprus, Israel, Kazakhstan.
- Nomada irkutskiensis** Proshchalykin et Schwarz, 2017. Russia: **ES (IR, BR).**
- Nomada italica** Dalla Torre et Friese, 1894. Cleptoparasite of *Andrena assimilis* Rad. (Andrenidae). Russia: **EP (C), UR, WS (AL), ES (KS, KR, BR).** – Europe (WE, SE, EE), Azerbaijan, Turkey, Kazakhstan.
- Nomada issikii** Yasumatsu, 1939. Russia: **FE (AM, KH, PR, SA, KU).** – Japan (Hok, Hon, Kyu, Shi).
- Nomada japonica** Smith, 1873. Cleptoparasite of *Eucera spurcatipes* Pér., *E. nipponensis* Pér., *E. longicornis* (L.) (Apidae). Russia: **FE (PR).** – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Nomada kohli** Schmiedeknecht, 1882 (*N. hippomensis* Saunders, 1908). Cleptoparasite of *LasioGLOSSUM puncticolle* (Mor.) (Halictidae). Russia: **EP (C, S, NC, CR).** – Europe (WE, SE, EE) N Africa.
- Nomada koikensis** Tsuneki, 1973 (*N. buyoo* Tsuneki, 1976). Russia: **FE (PR).** – Japan (Hok, Hon).
- Nomada lathburiana** (Kirby, 1802) [Apis] (*Apis rufiventris* Kirby, 1802). Cleptoparasite of *Andrena vaga* Pz. and *A. cineraria* (L.) (Andrenidae). Russia: **EP (N, NW, C, E, S, NC), UR, WS (TM, TK, AL), ES (KS, TU, KR, IR, BR, YA, ZB).** – Europe (WE, NE, SE, EE), Caucasus, Kazakhstan.
- Nomada laticrus** Mocsáry, 1883 (*N. erythrosoma* Mocsáry, 1883). Russia: **EP (S).** – Turkey.
- Nomada leucophthalma** (Kirby, 1802) [Apis] (*N. borealis* Zetterstedt, 1838; *N. inquilina* Smith, 1844; *N. kuro* Tsuneki, 1986; *N. muinensis* Tsuneki, 1986). Cleptoparasite of *Andrena nycthemera* Imh., *A. apicata* Sm., *A. clarkella* (Kby.) (Andrenidae). Russia: **EP (N, NW, C, E, S), WS (TK), ES (YA), FE (PR, KA).** – Europe (WE, NE, SE, EE), Kazakhstan, Mongolia, Japan (Hok, Hon, Kyu).
- Nomada lutea** Eversmann, 1852 (*N. rubricosa* Eversmann, 1982; *N. fasciculata* Morawitz, 1888). Russia: **EP (CR), UR.** – Kazakhstan.
- Nomada maculifrons** Smith, 1869. Russia: **FE (SA, KU).** – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Nomada margelanica** Schwarz, 1987. Russia: **ES (KS, BR, YA).** – Uzbekistan, Kazakhstan.
- Nomada marshamella** (Kirby, 1802) [Apis] (*N. marshamella* var. *dubia* Schmiedeknecht, 1882; *N. m.* var. *modesta* Schmiedeknecht, 1882, nom. praeocc., nec Cresson, 1863, nec Herrich-Schäffer, 1839). Cleptoparasite of *Andrena carantonica* Pér., *A. nigroaenea* (Kby.), *A. trimmerana* (Kby.), *A. rosae* Pz., *A. ferox* Sm. or probably *A. nitida* (Müll.) (Andrenidae). Russia: **EP (N, NW, C, E, CR), UR.** – Europe (WE, NE, SE, EE), Azerbaijan, Kazakhstan.
- Nomada mauritanica** Lepeletier de Saint-Fargeau, 1841 (*N. chrysopyga* Morawitz, 1872; *N. mephisto* Schmiedeknecht, 1882; *N. speciosissima* Schmiedeknecht, 1882; *N. superba* Pérez, 1902, nom. praeocc., nec Cresson, 1863; *N. sericultorum* Schulz, 1906; *N. syriaca* Friese, 1921). Cleptoparasite of *Andrena fuscosa* Erichson and probably of *A. pilipes* F. Russia: **EP (S, CR), ES (TU, BR).** – Europe (WE, SE, EE), N Africa, Azerbaijan, Turkey, Cyprus, Syria, Israel, Iraq, UAE.
- Nomada mitaii** Proshchalykin, 2010. Russia: **ES (KS, IR, ZB), FE (AM, KH, PR).** – Mongolia.
- Nomada mocsaryi** Schmiedeknecht, 1882. Russia: **EP (CR).** – Europe (SE, EE), Turkey, Iran.
- Nomada moeschleri** Alfken, 1913. Russia: **EP (NW, C, E), UR, ES (KR, IR, ZB).** – Europe (WE, NE, EE), Azerbaijan, Uzbekistan, Kazakhstan.
- Nomada mutabilis** Morawitz, 1871 (*N. mutabilis* var. *lucifera* Schmiedeknecht, 1882; *N. antigana*

- Pérez, 1895). Cleptoparasite of *Andrena chrysopyga* Schck. and probably of *A. labialis* (Kby.) (Andrenidae). Russia: **EP** (C, E, S, CR), **UR**, **ES** (KS, IR). – Europe (WE, NE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Iran, Turkmenistan, Kyrgyzstan, Kazakhstan, Nepal, India.
- Nomada mutica** Morawitz, 1872. Cleptoparasite of *Andrena ferox* Sm. (Andrenidae). Russia: **EP** (NC). – Europe (WE, SE, EE), Turkey, Azerbaijan.
- Nomada nobilis** Herrich-Schäffer, 1839 (*N. aber-rans* Eversmann, 1852; *N. calabra* Morawitz, 1872; *N. nobilis* var. *magrettiana* Schmiedeknecht, 1882). Cleptoparasite of *Andrena nasuta* Gir. (Andrenidae). Russia: **EP** (C, S, CR), **UR**. – Europe (WE, SE, EE), N Africa, Armenia, Azerbaijan, Turkey, Israel.
- Nomada noskiewiczi** Schwarz, 1966 (*N. aljkeni* Noskiewicz, 1939, nom. praeocc., nec Cockerell, 1907, nec Gaunitz, 1935). Russia: **EP** (E, NC). – Europe (WE, EE).
- Nomada obscura** Zetterstedt, 1838 (*N. obscura* var. *bitineta* Alfken, 1924; *N. o.* var. *aljkeni* Gaunitz, 1935, nom. praeocc., nec Cockerell, 1907, nec Noskiewicz, 1939). Cleptoparasite of *Andrena ruficrus* Nyl. (Andrenidae). Russia: **EP** (N, NW, C, E), **UR**, **WS** (TK), **ES** (IR, BR, ZB), **FE** (PR). – Europe (WE, NE, EE).
- Nomada obscuriceps** Schwarz et Levchenko, 2017. Russia: **EP** (E), **ES** (IR).
- Nomada obtusifrons** Nylander, 1848 (*N. mistura* Smith, 1851; *N. robertjeotiana* var. *alpina* Morawitz, 1867). Cleptoparasite of *Andrena coitana* (Kby.) and probably of *A. tarsata* Nyl. (Andrenidae). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (AL), **ES** (IR). – Europe (WE, NE, SE, EE), Kazakhstan.
- Nomada oculata** Friese, 1821. Russia: **EP** (NC, CR). – Europe (SE), Azerbaijan, Turkey, Syria, Israel.
- Nomada okamotonis** Matsumura, 1912 (*N. okamotonis kaiensis* Tsuneki, 1976). Cleptoparasite of *Andrena valeriana* Hirashima (Andrenidae). Russia: **FE** (AM, PR). – Japan (Hok, Hon, Kyu).
- Nomada opaca** Alfken, 1913. Cleptoparasite of *Andrena fulvida* Schck. (Andrenidae). Russia: **EP** (N, NW), **ES** (IR), **FE** (PR). – Europe (WE, NE, EE).
- Nomada pacifica** Tsuneki, 1973. Russia: **FE** (PR). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Nomada pallispinosa** Schwarz, 1967. Russia: **EP** (NC). – Europe (SE), Turkey, Cyprus, Israel.
- Nomada palmeni** Morawitz, 1888. Russia: **ES** (KR, IR), **FE** (KH).
- Nomada panzeri** Lepeletier de Saint-Fargeau, 1841 (*N. glabella* Thomson, 1870; *N. ruficornis* var. *phoenicia* Schmiedeknecht, 1882; *N. ruficornis* var. *incompleta* Schmiedeknecht, 1882; *N. ruficornis* var. *hybrida* Schmiedeknecht, 1882; *N. diluta* Pérez, 1884, nom. praeocc., nec Gribodo, 1893; *N. ruficornis* var. *fasciata* Friese, 1921). Cleptoparasite of *Andrena helvola* (L.), *A. synadelpha* Perk., *A. lapponica* Zett., *A. varians* (Kby.), *A. fucata* Sm. and *A. lapponica* Zett. (Andrenidae). Russia: **EP** (N, NW, C, E, S), **UR**, **WS** (TM, TK, AL), **ES** (TU, IR, BR, YA, ZB), **FE** (AM, PR, SA, KU, KA, MG). – Europe (WE, NE, SE, EE), N Africa, Japan (Hok, Hon, Shi, Kyu), Korean Peninsula.
- Nomada pastoralis** Eversmann, 1852. Russia: **UR**. – Europe (SE), Turkey.
- Nomada pectoralis** Morawitz, 1877 (*N. difficilis* Friese, 1921; *N. pectoralis* var. *atroflava* Friese, 1921). Russia: **EP** (C, S, NC), **UR**. – Azerbaijan, Turkmenistan.
- Nomada piliventris** Morawitz, 1877. Russia: **EP** (NC). – Azerbaijan, Turkey.
- Nomada pilosa** Schwarz et Gusenleitner, 2017. Russia: **EP** (S, CR). – Turkey.
- Nomada piccioliana** Magretti, 1883 (*N. piccioliana* Magretti, 1883; *N. novioregensis* Pérez, 1902; *N. piccioliana jurassica* E. Stoeckhert, 1941). Cleptoparasite of *Andrena combinata* Christ (Andrenidae). Russia: **EP** (C, CR). – Europe (WE, SE, EE).
- Nomada posthuma** Blüthgen, 1949 (*N. mixtella* Noskiewicz, 959). Cleptoparasite of *Lasioglossum glabriusculum* (Mor.) (Halictidae). Russia: **EP** (C), **ES** (IR), **FE** (PR). – Europe (WE, EE), Nepal.
- Nomada pulchra** Arnold, 1888 (*N. olhae* Cockerell, 1928). Probably cleptoparasite of *Melitta melanura* Nyl. (Melittidae). Russia: **EP** (C), **UR**, **WS** (AL), **ES** (TU, IR, ZB). – Europe (WE, NE, EE), Kazakhstan.
- Nomada pygidialis** Schwarz, 1981. Russia: **EP** (S). – Europe (SE, EE), Georgia, Turkey.
- Nomada radoszkowskii** Łoziński, 1922. Russia: **EP** (NC), **ES** (BR). – Europe (SE), Turkey, Armenia, Azerbaijan.
- Nomada rhenana** Morawitz, 1872 (*N. rufipes* Schenck, 1870, nom. praeocc., nec Fabricius, 1793). Cleptoparasite of *Andrena ovatula* (Kby.) (Andrenidae). Russia: **EP** (C, S), **UR**. – Europe (WE, NE, SE, EE), N Africa.
- Nomada roberjeotiana** Panzer, 1799 (*N. panzeriana* Walckenaer, 1802; *N. neglecta* Herrich-Schäffer, 1839; *Nomadita montana* Mocsáry, 1894, nom. praeocc., nec Scopoli, 1763; *Nomada roberjeotiana tormentillae* Alfken, 1901). Cleptoparasite of *Andrena*

tarsata Nyl., and probably of *A. fuscipes* (Kby.) and *A. coitana* (Kby.) (Andrenidae). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (TK, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR). – Europe (WE, NE, EE), Azerbaijan, Kyrgyzstan, Mongolia, Japan (Hok, Hon).

Nomada rubricollis Schwarz, 1966. Russia: **EP** (NC). – Europe (SE), Turkey, Syria, Jordan, Israel.

Nomada rubricosa Eversmann, 1852. Russia: **EP** (S), **UR**. – Kazakhstan.

Nomada ruficornis (Linnaeus, 1758) [Apis] (*N. bifida* Thomson, 1872; *N. bifida orenburgensis* Łoziński, 1922). Cleptoparasite of *Andrena haemorrhoa* F. (Andrenidae). Russia: **EP** (N, NW, C, E, S, CR), **UR**, **WS** (TK, AL), **ES** (KR, IR, BR, YA), **FE** (AM, SA, KU). – Europe (WE, NE, SE, EE), Azerbaijan, Turkey, Uzbekistan, Kazakhstan.

Nomada rufipes Fabricius, 1793 (*N. vaga* Panzer, 1798; *N. solidaginis* Panzer, 1799; *Apis picta* Kirby, 1802; *A. rufopicta* Kirby, 1802; *Nomada solidaginis* var. *punctulifera* Friese, 1921; *N. s.* var. *minutula* Friese, 1921; *N. fennica* Alfken, 1924; *N. rufipes nigriventris* Alfken, 1926). Cleptoparasite of *Andrena denticulata* (Kby.), *A. fuscipes* (Kby.), probably of *A. nigriceps* (Kby.) (Andrenidae) and *Colletes cunicularius* (L.) (Colletidae). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (AL), **ES** (KS, YA), **FE** (AM). – Europe (WE, NE, SE, EE), N Africa, Kazakhstan.

Nomada setteri Proshchalykin, 2010. Russia: **ES** (IR), **FE** (PR).

Nomada sexfasciata Panzer, 1799 (*Apis connexa* Kirby, 1802; *A. schaefferella* Kirby, 1802). Cleptoparasite of *Eucera longicornis* (L.) and *E. nigrescens* Pér. (Apidae). Russia: **EP** (C, E, NC, CR), **UR**, **WS** (TK, AL), **ES** (KS, IR, BR, ZB), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Israel, Iran, Turkmenistan, Kazakhstan, Korean Peninsula.

Nomada sheppardana (Kirby, 1802) [Apis] (*N. dalii* Curtis, 1832; *N. minuscula* Noskiewicz, 1930; *N. microsticta* Cockerell, 1931). Cleptoparasite of *Lasiosiglossum nitidiusculum* (Kby.) and *L. sexstrigatum* (Schck.) (Halictidae). Russia: **EP** (C, E, S). – Europe (WE, NE, SE, EE), N Africa.

Nomada silvicola Tsuneki, 1973 (*N. hangani* Tsuneki, 1973; *N. diervillae* Tsuneki, 1973). Russia: **FE** (PR). – Japan (Hok, Hon).

Nomada stigma Fabricius, 1804 (*N. stigma* Fabricius, 1804; *N. cinnaberina* Morawitz, 1871; *N. cinnabarina*

Morawitz, 1872; *N. austriaca* Schmiedeknecht, 1882; *N. stigma villipes* Stoeckhert, 1930; *N. cinnabarina* var. *slovakica* Mavromoustakis, 1955). Cleptoparasite of *Andrena labialis* (Kby.), *A. schencki* Mor. and *A. decipiens* Schck. (Andrenidae). Russia: **EP** (N, C, E, S, NC, CR), **UR**, **WS** (TK, AL), **ES** (KR, IR, YA). – Europe (WE, NE, SE, EE), N Africa, Armenia, Azerbaijan, Turkey, Cyprus, Uzbekistan, Kazakhstan.

Nomada striata Fabricius, 1793 (*Apis hillana* Kirby, 1802; *A. ochrostoma* Kirby, 1802, nom. praecocc., nec Zetterstedt, 1838; *Nomada vidua* Smith, 1844; *N. punetiscuta* Thomson, 1870; *N. simillima* Pérez, 1913; *N. dzieduszyckii* Noskiewicz, 1924). Cleptoparasite of *Andrena wilkella* (Kby.), *A. aberrans* Ev., *A. similis* Sm., *A. gelriae* van der Vecht and probably of *A. pandellei* Pér., *A. intermedia* Thoms. (Andrenidae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TK, AL), **ES** (BR, YA, ZB), **FE** (PR). – Europe (WE, NE, SE, EE), Cyprus, Kazakhstan.

Nomada stoeckherti Pittioni, 1951. Russia: **EP** (S, CR). – Europe (WE, EE), Afghanistan.

Nomada succincta Panzer, 1798 (*N. succineta* var. *mixta* Dalla Torre, 1877; *N. succineta* var. *rufula* Friese, 1921; *N. fidvicornis* auct., nec Fabricius, 1793). Cleptoparasite of *Andrena nigroaenea* (Kby.), *A. nitida* (Müll.), *A. cineraria* (L.), *A. tibialis* (Kby.), *A. thoracica* (F) or probably *A. carantonica* Pér. (Andrenidae). Russia: **FE** (PR). – Europe (WE, NE, SE, EE), N Africa, Azerbaijan, Turkey, Jordan.

Nomada sybarita Schmiedeknecht, 1882. Russia: **UR**, **WS** (TK). – Europe (WE, SE, EE), Azerbaijan.

Nomada tenella Mocsáry, 1883. Russia: **EP** (E, S, NC, CR), **UR**, **ES** (BR, YA). – Azerbaijan, Iran, Turkmenistan, Kazakhstan.

Nomada thersites Schmiedeknecht, 1882. Russia: **EP** (S). – Europe (EE), Cyprus, Israel, Kyrgyzstan, Kazakhstan.

Nomada trapeziformis Schmiedeknecht, 1882 (*N. ruficornis* var. *auctumnalis* Ducke, 1900). Probably cleptoparasite of *Andrena nitida limata* Sm. (Andrenidae). Russia: **EP** (C, E), **WS** (TK), **ES** (KS, KR, IR, BR). – Europe (WE, NE, EE).

Nomada tridentirostris Dours, 1873 (*N. cyphognatha* Pérez, 1884; *N. amblystoma* Pérez, 1884). Russia: **EP** (NC). – Europe (WE, SE, EE), N Africa, Turkey, Israel.

Nomada trispinosa Schmiedeknecht, 1882. Cleptoparasite of *Andrena taraxaci* Gir. (Andrenidae).

Russia: **EP** (CR). – Europe (WE, SE, EE), Cyprus, Azerbaijan.

Nomada yarrowi Schwarz, 1981. Russia: **EP** (S). – Turkey.

Nomada zonata Panzer, 1798 (*N. bofillana* Pérez, 1913; *N. banatica* Zilahi-Kiss, 1915; *N. zonata sarmatica* Stoeckhert, 1941). Cleptoparasite of *Andrena congreuens* Schmied. and *A. dorsata* (Kby.) (Andrenidae). Russia: **EP** (C, E, S), **UR**, **ES** (KS, IR). – Europe (WE, NE, SE, EE), N Africa, Armenia, Azerbaijan, Turkey, Iran, Turkmenistan, Uzbekistan, Kazakhstan.

Tribe EPEOLINI

EPEOLUS Latreille, 1802 (*Trophoclepria* Holmberg, 1886; *Pyrrhonelecta* Ashmead, 1899; *Argyrosoleenis* Robertson, 1903; *Oxybiastes* Mavromoustakis, 1854). Type species *Apis variegata* Linnaeus, 1758. Species with known biology are nest parasites of *Colletes* (Colletidae). Number of taxa: World – 109, absent in Australasia; Palaearctic – 35; Russia – 12.

Epeolus alpinus Friese, 1893 (*E. glacialis* Alfken, 1913; *E. montanus* Bischoff, 1930; *E. pilosus* Bischoff, 1930; *E. alpinus* Bischoff, 1930). Cleptoparasite of *Colletes floralis* Ev. and *C. impunctatus* Nyl. (Colletidae). Russia: **EP** (N, NW, C), **ES** (IR). – Europe (WE, NE, SE).

Epeolus coreanus Yasumatsu, 1933. Russia: **FE** (PR, SA). – Korean Peninsula, Japan (Hon, Kyu).

Epeolus cruciger (Panzer, 1799) [Nomada] (*E. rufipes* Thomson, 1870; *E. similis* Höppner, 1899; *E. cruciger* var. *elegans* Müller, 1921; *E. c.* var. *rufiventris* Müller, 1921, nom. praeocc., nec Friese, 1908; *E. laevifrons* Bischoff, 1930; *E. marginatus* Bischoff, 1930). Cleptoparasite of *Colletes marginatus* Sm. and *C. succinctus* (L.) (Colletidae). Russia: **EP** (NW, C, E, S), **WS** (NS, TK, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (SA). – Europe (WE, NE, SE, EE), Iran.

Epeolus fasciatus Friese, 1895. Cleptoparasite of *Colletes punctatus* Mocs. (Colletidae). Russia: **EP** (S, CR). – Europe (WE, SE, EE), Caucasus, Turkey.

Epeolus julliani Pérez, 1884. Russia: **EP** (S). – Europe (WE, SE), Georgia.

Epeolus laticauda Bischoff, 1930. Probably cleptoparasite of *Colletes anceps* Rad. (Colletidae). Russia: **ES** (BR). – Turkmenistan, Uzbekistan, Tajikistan.

Epeolus melectiformis Yasumatsu, 1938. Cleptoparasite of *Colletes esakii* Hirashima (Colletidae). Russia: **ES** (BR), **FE** (PR). – Japan (Hok, Hon, Shi, Kyu, Ryu).

Epeolus productulus Bischoff, 1930. Russia: **EP** (S). – Europe (WE, SE), Turkey, Syria.

Epeolus schummeli Schilling, 1849 (*E. ruthenicus* Radoszkowski, 1891). Cleptoparasite of *Colletes nasutus* Sm. (Colletidae). Russia: **EP** (S). – Europe (WE, SE, EE), Turkey.

Epeolus tarsalis Morawitz, 1874 (*E. praestans* Pérez, 1884; *E. rozenburgensis* van Lith, 1949; *E. tarsalis tirolensis* van Lith, 1956). Cleptoparasite of *Colletes collaris* Dours and probably of *C. halophilus* Verhoeff (Colletidae). Russia: **EP** (S, NC), **ES** (ZB, BR), **FE** (AM, KH, PR). – Europe (WE, EE), Mongolia, Korean Peninsula, Japan (Hon, Shi).

Epeolus transitorius Eversmann, 1852 (*E. transitorius* var. *seraxensis* Radoszkowski, 1893). Russia: **EP** (S), **UR**. – Europe (SE), Georgia, Iran, Turkmenistan, Uzbekistan, Kazakhstan.

Epeolus variegatus (Linnaeus, 1758) [Apis] (*Apis muscaria* Christ, 1791; *Epeolus pictus* Nylander, 1848; *E. productus* Thomson, 1870; *Apis muraria* Christ, 1791 in Friese, 1895; *Epeolus variegatus* var. *rossicus* Friese, 1925). Cleptoparasite of *Colletes daviesanus* Sm., *C. fodiens* (Geoffr.), *C. similis* Schck. and *C. halophilus* Verhoeff (Colletidae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TK, KM), **ES** (IR, YA). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Central Asia.

TRIEPEOLUS Robertson, 1901 (*Doeringiella* Holmberg, 1886). Type species *Epeolus concavus* Cresson, 1878. American cleptoparasitic genus with only 2 Old World species. Hosts – various Apiformes, especially Eucerini (Apidae). Number of taxa: World – 103; Palaearctic – 2; Russia – 2.

Triepeolus tristis (Smith, 1854) [Epeolus] (*Epeolus luctuosus* Eversmann, 1852, nom. praeocc., nec Spinola, 1851; *E. speciosus* Gerstäcker, 1869). Probably cleptoparasite of *Tetralonia malvae* (Rossi) and *Tetraloniella nana* (Mor.) (Apidae). Russia: **EP** (C, E, S), **UR**, **WS** (AL). – Europe (WE, SE, EE), Turkey, Kazakhstan.

Triepeolus ventralis (Meade-Waldo, 1913) [Epeolus] (*T. signatus* Hedicke, 1940). Cleptoparasite of *Tetraloniella mitsukurii* (Cock.) (Apidae). Russia: **WS** (AL), **ES** (BR), **FE** (AM, KH, PR). – China (NE, NC, CC, SE), Japan (Hon, Kyu, Shi).

Tribe AMMOBATOIDINI

AMMOBATOIDES Radoszkowski, 1867 (*Phiarus Gerstäcker, 1869; Euglages* Gerstäcker, 1869; *Paidia* Radoszkowski, 1872; *Paedia* Dalla Torre, 1891). Type species *Phileremus abdominalis* Eversmann, 1852. Cleptoparasits in nests of *Melitturga* and *Melitturgula* (Andrenidae). Number of taxa: World – 8, includes 1 in S Africa; Palaearctic – 7; Russia – 2.

Ammobatoides abdominalis (Eversmann, 1852) [Phileremus] (*Phileremus hirsutus* Eversmann, 1852; *Ammobates extraneus* Förster, 1855; *A. rufitarsis* Smith, 1879; *Phiarus abdominalis* var. *sanguineus* Friese, 1911; *Ph. a. var. rufus* Friese, 1911; *Ph. angarensis* Cockerell, 1928; *Ammobatoides abdominalis* ssp. *marchicus* Bischoff, 1952; *A. lebanensis* Mavromoustakis, 1959). Cleptoparasite of *Melitturga clavicornis* (Latr.), *M. praestans* Gir. and *M. pictipes* Mor. (Andrenidae). Russia: **EP** (C, E, S, CR), **UR**, **WS** (OM, NS, TK, KM, AL), **ES** (KS, TU, KR, IR). – Europe (WE, SE, EE), Caucasus, Turkey, Syria, Lebanon, Iran, Central Asia, Kazakhstan, China (NW).

Ammobatoides radoszkowskii Proshchalykin et Lelej, 2014. Probably cleptoparasite of *Melitturga mongolica* Alfken (Andrenidae). Russia: **ES** (BR), **FE** (AM, PR). – China (NC).

Tribe BIASTINI

BIASTES Panzer, 1806 (*Rhineta* Illiger, 1807; *Melitoxena* Morawitz, 1873; *Biastoides* Schenck, 1874). Type species *Tiphia brevicornis* Panzer, 1798. Parasites in nests of *Dufourea*, *Rophites* and *Systropha* (Halictidae). Number of taxa: World – 5 Palaearctic species, including 1 Eastern Palaearctic and 1 Transpalaearctic species; Russia – 4.

Biastes brevicornis (Panzer, 1798) [Tiphia] (*Nomada atrata* Fabricius, 1804; *N. schottii* Fabricius, 1804; *Pasites unicolor* Jurine, 1807; *P. ater* Spinola, 1808; *Stelis aberrans* Eversmann, 1852; *Pasites fasciatus* Eversmann, 1852). Cleptoparasite of *Systropha curvifrons* (Scop.) and *S. planidens* Gir. (Halictidae). Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, SE, EE), Armenia, Azerbaijan, Turkey, Syria, Iran, Kazakhstan.

Biastes emarginatus (Schenck, 1853) [*Phileremus*] (*Phileremus nasutus* Gerstäcker, 1869; *Pasites*

punctatus Schenck, 1870). Cleptoparasite of *Rophites algirus* Pér. and *R. quinquespinosus* (Halictidae). Russia: **EP** (C, E, S, CR), **UR**, **WS** (KM), **ES** (KR, IR). – Europe (WE, NE, SE, EE), Armenia, Azerbaijan, Turkey, Kazakhstan.

Biastes popovi Proshchalykin et Lelej, 2004. Probably cleptoparasite of *Dufourea carinata* Popov and *D. inermis* (Nyl.) (Halictidae). Russia: **FE** (AM, KH).

Biastes truncatus (Nylander, 1848) [Nomada] (*Phileremus punetatus* Gerstäcker, 1869; *Pasites rostratus* Zilahi-Kiss, 1927). Cleptoparasite of *Dufourea inermis* (Nyl.) and *D. dentiventris* (Nyl.) (Halictidae). Russia: **EP** (NW, C), **WS** (NS, AL), **ES** (KR, ZB), **FE** (PR). – Europe (WE, NE, SE, EE), Turkey.

Tribe AMMOBATINI

AMMOBATES Latreille, 1809 (*Phileremus* Latreille, 1809; *Ammobatoides* Schenck, 1869, nom. praeocc., nec Radoszkowski, 1867). Type species *Ammobates rufiventris* Latreille, 1809. Cleptoparasites of *Tetraloniella*, *Anthophora* and, with hight probability, *Ancyla*. Number of taxa: World – 50, including 1 Afrotropic and 1 Oriental species; Palaearctic – 48; Russia – 4.

Ammobates (Ammobates) opacus Popov, 1951 (*A. arménicus*: Popov, 1951). Russia: **EP** (S). – Europe (EE), Azerbaijan, Kazakhstan.

Ammobates (Ammobates) punctatus (Fabricius, 1804) [Epeolus] (*Epeolus kirbienus* Latreille, 1805; *Phileremus dufouri* Latreille, 1809; *Ammobates bicolor* Lepeletier de Saint-Fargeau, 1825; *A. apunctatus* var. *atratus* Mocsáry, 1894). Cleptoparasite of *Anthophora bimaculata* (Pz.) (Apidae). Russia: **EP** (C), **UR**. – Europe (WE, SE, EE), N Africa.

Ammobates (Ammobates) vinctus Gerstäcker, 1869 (*A. setosus* Morawitz, 1870). Cleptoparasite of *Tetraloniella dentata* (Germ.) and *T. ruficornis* (F.) (Apidae). Russia: **EP** (C, S, CR), **UR**. – Europe (WE, SE, EE), N Africa, Turkey, Kazakhstan.

Ammobates (Euphilermus) oraniensis (Lepeletier de Saint-Fargeau, 1841) [*Phileremus*] (*Phileremus melectoides* Smith, 1854; *Ammobates oraniensis* var. *manni* Friese, 1895; *A. o. tenuicornis* Popov, 1951; *Pasites oraniensis anatolicus* Warncke, 1983). Cleptoparasite of *Eucera armeniaca* (Mor.) (Apidae). Russia: **EP** (CR). – Europe (SE, EE), N Africa, Armenia, Turkey, Jordan, Israel, Turkmenistan.

PARAMMOBATODES Popov, 1951. Type species *Phiarus minutus* Mocsáry, 1878 (= *Ammobates rufiventris* Latreille, 1809). Cleptoparasites of *Camtopoeum* (Andrenidae) and probably of *Nomioides* (Halictidae). Number of taxa: World – 6, including 1 Oriental species; Palaearctic – 5 Westpalaearctic species; Russia – 1.

Parammobatodes minutus (Mocsáry, 1878) [Phiarus] (*Parammobatodes schmidti* Alfken, 1936). Cleptoparasite of *Camtopoeum friesei* Mocs. and *C. frontale* (F.) (Andrenidae). Russia: **EP** (NC). – Europe (WE, SE, EE), Turkey.

PASITES Jurine, 1807 (*Morgania* Smith, 1854; *Omachthes* Gerstäcker, 1869; *Homachthes* Dalla Torre, 1896; *Pasitomachthes* Bischoff, 1923). Type species *Pasites maculata* Jurine, 1807. Species with known biology are cleptoparasites of *Pseudapis* and *Nomiapis* (Halictidae). Number of taxa: World – 29 in Old World, including 23 Afrotropic species; Palaearctic – 2; Russia – 2.

Pasites esakii Popov et Yasumatsu, 1935 (*P. maculatus esakii* Popov, Yasumatsu, 1935). Cleptoparasite of *Nomiapis mandschurica* (Hedicke) (Halictidae). Russia: **FE** (PR). – China (NE), Korean Peninsula, Japan (Kyu).

Pasites maculatus Jurine, 1807 (*Phileremus kirbyanus* Latreille, 1809; *Nomada albomaculatus* Lucas, 1849; *Pasites schottii* Eversmann, 1852; *Ammobates variegatus* Smith, 1854; *Phileremus rufiventris* Förster, 1855, nom. praeocc., nec Latreille, 1809; *Phiarus maculatus* var. *pusillus* Radoszkowski, 1872; *Pasites maculatus* var. *aschabadensis* Radoszkowski, 1893; *P. maculatus* var. *brunneus* Friese, 1895; *Pasites comptus* Alfken, 1929). Cleptoparasite of *Pseudapis bispinosa* (Brull.), *P. femoralis* (Pall.) and *P. diversipes* Latr. (Halictidae). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (AL), **ES** (KS, IR, BR, ZB), **FE** (AM). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Cyprus, Israel, Iraq, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NE, NC, NW).

Subfamily APINAE

Tribe OSIRINI

EPEOLOIDES Giraud, 1863 (*Viereckella* Swenk, 1907). Type species *Epeoloides ambiguus* Giraud, 1863

= *Apis coecutiens* Fabricius, 1775. Cleptoparasites of *Macropis* (Melittidae). Number of taxa: World – 2 Holarctic species; Palaearctic – 1; Russia – 1.

Epeoloides coecutiens (Fabricius, 1775) [*Apis*] (*Saropoda fulva* Eversmann, 1852; *Epeoloides ambiguus* Giraud, 1863; *E. fulviventris* Schenck, 1866). Cleptoparasite of *Macropis europaea* Warncke and *M. fulvipes* (F.) (Melittidae). Russia: **EP** (NW, C, E, CR), **UR**, **WS** (KM), **ES** (KR). – Europe (WE, NE, SE, EE), Kazakhstan.

Tribe ANCYLINI

ANCYLA Lepeletier de Saint-Fargeau, 1841 (*Plistotrichia* Morawitz, 1873). Type species *Ancyla oraniensis* Lepeletier de Saint-Fargeau, 1841. Ground nesting summer oligolects, possible hosts of *Ammobates* (Apidae). Number of taxa: World – 8 West Palaearctic species; Russia – 1.

Ancyla asiatica Friese, 1922 (*Plistotrichia flavilabris*: Morawitz, 1874). Oligolect on *Daucus* (Apiaceae). Russia: **EP** (NC). – Turkey, Lebanon.

Tribe CTENOPLECTRINI

CTENOPLECTRA Kirby, 1826. Type species *Ctenoplectra chalybea* Smith, 1857. Wood holes nesting and oil-collecting oligolects on Cucurbitaceae. Number of taxa: World – 24 in the Old World, mostly Afrotropic and Oriental species; Palaearctic – 2 East Palaearctic; Russia – 1.

Ctenoplectra davidi Vachal, 1903 (*Scrapter simpliciceps* Strand, 1913). Russia: **FE** (PR). – China (NE, NC, SW, SE).

Tribe EUCERINI

CUBITALIA Friese, 1911. Type species *Eucera breviceps* Friese, 1911. Summer or spring-summer oligo- and polylect species. Nesting unknown. Number of taxa: World – 8 West Palaearctic species; Russia – 3.

Cubitalia (Cubitalia) morio (Friese, 1911) [Eucera]. Oligolect on *Onosma* (Boraginaceae). Russia: **EP** (NC, CR). – Europe (SE), Turkey.

Cubitalia (Cubitalia) tristis (Morawitz, 1876) [Eucera]. Russia: **EP** (NC, CR). – Georgia, Turkey.

Cubitalia (Pseudoeucera) parvicornis (Mocsáry, 1878) [Eucera]. Polylect, prefer Boraginaceae. Russia: **EP** (NC). – Europe (WE, SE, EE).

EUCERA Scopoli, 1770 (*Tetralonia* Spinola, 1839, part.). Type species *Apis longicornis* Linnaeus, 1758. Summer or spring-summer oligo- and polylect ground nesting species. Number of taxa: World – 223, overwhelming majority in Holarctic; Palaearctic – 173; Russia – 34.

Eucera (Eucera) chrysopyga Pérez, 1879 (*E. pollinosa* Smith, 1854, nom. praeocc., nec Lepeletier de Saint-Fargeau, 1841; *E. favosa* Mocsáry, 1879; *E. polonica* Ruszkowski, 1994). Oligolect on Fabaceae. Russia: **EP** (S, CR), **UR**. – Europe (WE, SE, EE), N Africa, Caucasus, Iran, Kazakhstan, China (NC).

Eucera (Eucera) dalmatica Lepeletier de Saint-Fargeau, 1841 (*E. robusta* Baer, 1850; *E. echii* Mocsáry, 1878; *E. flabellifera* Mocsáry, 1879). Polylect, prefer Boraginaceae. Russia: **EP** (NC, CR). – Europe (WE, SE, EE), Azerbaijan, Armenia, Turkey, Cyprus, Iran.

Eucera (Eucera) interrupta Baer, 1850 (*E. semistrigosa* Dours, 1873; *E. confusa* Gribodo, 1881). Oligolect on Fabaceae. Russia: **EP** (C, E, S, CR), **UR**, **WS** (AL). – Europe (WE, SE, EE), Caucasus, Turkey, China (NC, NW).

Eucera (Eucera) kullenbergi Tkalcú, 1984. Polylect, prefer Boraginaceae. Russia: **EP** (CR). – Europe (SE, EE), Azerbaijan, Turkey, Iran.

Eucera (Eucera) longicornis (Linnaeus, 1758) [*Apis*] (*Apis linguaria* Fabricius, 1775; *A. tuberculata* Fabricius, 1793; *Eucera difficilis* Pérez in Friese, 1896; *E. atricollis* Friese, 1922; *E. pillichi* Alfken, 1932). Oligolect on Fabaceae. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (OM, TK, KM), **ES** (IR, BR, ZB), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Kazakhstan, China (NE, NC, SW).

Eucera (Eucera) nigrescens Pérez, 1879 (*E. tuberculata* auct., nec Fabricius, 1793). Oligolect on Fabaceae. Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), Caucasus, Turkey, Jordan, Iran, Turkmenistan.

Eucera (Eucera) nigrilabris Lepeletier de Saint-Fargeau, 1841 (*E. melanostoma* Morawitz, 1875; *E. terminalis* Smith, 1879; *Macroceras grisea*: Eversmann, 1852; *Eucera nigrilabris rufitarsis* Tkalcú, 1984; *E. n. orientalis* Tkalcú, 1984). Russia: **EP** (NC), **UR**. – Europe (WE, SE, EE), N Africa, Azerbaijan, Turkey, Israel, Palestine, Iran, Turkmenistan, Uzbekistan, Tajikistan, Kazakhstan.

Eucera (Eucera) proxima Morawitz, 1875 (*E. gracca* Radoszkowski, 1876; *E. nitidiventris* Mocsáry, 1879; *E. bipartita* Pérez, 1911). Russia: **EP** (E, S, NC, CR), **UR**. – Europe (WE, SE, EE), Azerbaijan, Turkey, Syria, Iran, Tajikistan.

Eucera (Eucera) rufipes Smith, 1879 (*E. platyrhina* Pérez, 1902). Oligolect on Fabaceae. Russia: **EP** (C, S, CR), **UR**, **WS** (KM, AL), **ES** (KR, YA). – Europe (EE), Kazakhstan, Mongolia, China (NC).

Eucera (Eucera) sociabilis Smith, 1873 (*E. andreae* Friese, 1910). Oligolect on Fabaceae. Russia: **FE** (PR). – China (NC, SW, SE), Japan (Hon, Hok).

Eucera (Hetereucera) atriceps Morawitz, 1878. Russia: **EP** (CR). – Armenia, Kazakhstan.

Eucera (Hetereucera) caspica Morawitz, 1874 (*E. albopunctata* Dours, 1873; *E. frivaldszkyi* Mocsáry, 1877; *E. perézi* Mocsáry, 1878; *E. amplitarsis* Mocsáry, 1878; *E. caspica* var. *pallida* Kerenskij, 1919). Russia: **EP** (S, NC, CR). – Europe (WE, SE, EE), Azerbaijan, Turkey.

Eucera (Hetereucera) cineraria Eversmann, 1852 (?*E. cinerea* Lepeletier de Saint-Fargeau, 1841 (patrice inconnue); ?*E. cana* Baer, 1850; ?*E. cantatrix* Baer, 1850; ?*E. crinita* Baer, 1850, nom. praeocc., nec Klug, 1845; *E. concinna* Gribodo, 1873; ?*E. consimilis* Dours, 1873; *E. nigritrons* Smith, 1879; *Tetralonia tarsata* Mocsáry, 1879). Polylect, prefers Fabaceae. Russia: **EP** (S, NC, CR), **UR**. – Europe (WE, SE, EE), Caucasus, Turkey, Iran, Afghanistan, China (NE).

Eucera (Hetereucera) clypeata Erichson, 1835 (*E. clypeata* Erichson, 1835; *E. similis* Lepeletier de Saint-Fargeau, 1841; *E. punctilabris* Lepeletier de Saint-Fargeau, 1841; *E. coarcata* Eversmann, 1852; *E. medusa* Nurse, 1904). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (AL). – Europe (WE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Syria, Israel, Palestine, Jordan, Iraq, Iran, Afghanistan, Pakistan, Turkmenistan, Uzbekistan, Tajikistan, Kazakhstan.

Eucera (Hetereucera) curvitarsis Mocsáry, 1879. Russia: **EP** (S, CR). – Europe (SE, EE), Turkey.

Eucera (Hetereucera) excisa Mocsáry, 1879. Russia: **EP** (E, NC). – Europe (SE), Azerbaijan, Turkey.

Eucera (Hetereucera) ferghanica Morawitz, 1875 (*E. tomentosa* Morawitz, 1875). Russia: **EP** (S). – Iran, Uzbekistan.

Eucera (Hetereucera) nigrita Friese, 1895 (?*E. nigra* Lepeletier de Saint-Fargeau, 1841, nom. praeocc., nec Lepeletier de Saint-Fargeau, 1841; *E. caspica* var. *nigrita* Friese, 1895; *E. albofasciata* Friese, 1895).

- Russia: EP (S, NC, CR). – Europe (SE), N Africa, Azerbaijan, Turkey, Syria, Lebanon, Jordan, Israel.
- Eucera (Hetereucera) pannonica** Mocsáry, 1878.
Russia: EP (CR). – Europe (EE).
- Eucera (Hetereucera) paraclypeata** Sitzikov, 1988.
Russia: EP (NC, CR). – Europe (SE), Turkey, Syria, Israel, Jordan.
- Eucera (Hetereucera) taurica** Morawitz, 1870
(*E. pulveracea* Dours, 1873; *E. spectabilis* Mocsáry, 1881; *E. asiatica* Alfken, 1936; *E. hispana* auct., nec Lepeletier de Saint-Fargeau, 1841). Russia: EP (S, NC, CR), UR. – Europe (WE, SE, EE), N Africa, Iran, China (NW).
- Eucera (Hetereucera) seminuda** Brülle, 1832 (*E. trivittata* Brülle, 1832; *E. paradoxa* Mocsáry, 1878). Russia: EP (C, S, CR). – Europe (WE, SE, EE), N Africa, Azerbaijan, Turkey.
- Eucera (Hetereucera) sogdiana** Morawitz, 1875
(*E. kouchkiana* Radoszkowski, 1886; *E. melanostoma* Morawitz, 1875). Russia: EP (S). – Europe (SE), Turkey, Turkmenistan, Uzbekistan, Tajikistan, Afghanistan.
- Eucera (Hetereucera) vittulata** Noskiewicz, 1936
(*E. parvula* Friese, 1895; *E. cineraria* Eversmann, 1852). Russia: EP (C, S, NC). – Europe (WE, SE, EE), Azerbaijan, Armenia, Turkey.
- Eucera (Pteneucera) eucnemidea** Dours, 1873. Russia: EP (S, NC). – Europe (SE), N Africa, Armenia, Turkey, Syria, Kazakhstan.
- Eucera (Pteneucera) nigrifacies** Lepeletier de Saint-Fargeau, 1841 (*E. caucasica* Morawitz, 1874; *E. sedula* Mocsáry, 1879; *E. cornuta* Stefani, 1882). Russia: EP (S, NC, CR), WS (AL). – Europe (WE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Syria, Israel, Jordan, Iran, Uzbekistan, Tajikistan, Kazakhstan.
- Eucera (Synhalonia) alborufa** (Radoszkowski, 1871) [Tetralonia] (*Tetralonia amoena* Walker, 1871; *T. rufescens* Morawitz, 1875; *T. radoszkowskii* Morawitz, 1875; *Macrocerca clypeata* Radoszkowski, 1893; *Eucera sarachensis* Dalla Torre, 1896; *E. turcestanica* Dalla Torre, 1895). Russia: EP (E, S). – N Africa, Azerbaijan, Iran, Turkmenistan, Uzbekistan, Kazakhstan.
- Eucera (Synhalonia) alternans** Brullé, 1832 [Macrocerca]. Russia: EP (S, CR). – Europe (WE, SE, EE), N Africa, Azerbaijan, Turkey, Iran, Kazakhstan.
- Eucera (Synhalonia) armeniaca** (Morawitz, 1877) [Tetralonia]. Russia: EP (C, S, CR). – Europe (EE), Armenia, Turkmenistan.

- Eucera (Synhalonia) chinensis** (Smith, 1854) [Tetralonia]. Russia: FE (PR). – China (NE, NC, CC, SW, SE), Japan (Ryu), N Vietnam.
- Eucera (Synhalonia) hungarica** Friese, 1895. Polylect, prefer Fabaceae. Russia: EP (C, E, S, CR), UR. – Europe (WE, SE, EE), N Africa, Armenia, Kazakhstan.
- Eucera (Synhalonia) ruficollis** (Brullé, 1832) [Macrocerca] (*E. atricornis*: Eversmann, 1852). Russia: EP (S, NC), UR. – Europe (SE, EE), N Africa, Caucasus, Turkey, Syria, Iran, Turkmenistan, Uzbekistan, Tajikistan, Kazakhstan.
- Eucera (Synhalonia) tricincta** Erichson, 1835. Russia: EP (S, NC, CR). – Europe (WE, SE, EE), N Africa, Azerbaijan, Israel, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan.
- Eucera (Synhalonia) velutina** (Morawitz, 1874) [Tetralonia]. Visit Rindera tetraspis (Boraginaceae). Russia: EP (S, NC), UR. – Turkey, Israel.
- TETRALONIA** Spinola, 1839 (*Macrocerca* Latreille, 1810, nom. praeocc., nec Meigen, 1803). Type species *Apis malvae* Rossi, 1790. Oligolect ground-nesting bees. Number of taxa: World – 18, mostly Afrotropic species; Palaearctic – 1; Russia – 1.
- Tetralonia malvae** (Rossi, 1790) [Eucera] (*Eucera antennata* Fabricius, 1793; *E. macroglossa* Illiger, 1806). Oligolect on Malvaceae. Russia: EP (C, E, S, NC, CR), UR, WS (AL). – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, Syria, Iran, Uzbekistan, Tajikistan, Kazakhstan, China (NW).
- TETRALONIELLA** Ashmead, 1899 (*Xenoglossodes* Ashmead, 1899; *Melissina* Cockerell, 1911; *Tetralonia* part.: Proshchalykin, 2007e). Type species *Macrocerca graja* Eversmann, 1852. Oligolect or polylect ground-nesting bees. Number of taxa: World – 137, mostly in Old World; Palaearctic – 50; Russia – 14 species of subgenus *Tetraloniella* s. str.
- Tetraloniella alticincta** (Lepeletier de Saint-Fargeau, 1841) [Macrocerca]. Russia: EP (CR). – Europe (WE, SE, EE).
- Tetraloniella dentata** (Germar, 1839) [Eucera] (*Tetralonia dentata* Klug, 1835; *Macrocerca tricincta* Lepeletier de Saint-Fargeau, 1841; *M. nigricornis* Lepeletier de Saint-Fargeau, 1841). Oligolect on Asteraceae. Russia: EP (NW, C, E, S, NC), UR, WS (OM). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Syria, Iran, Uzbekistan, Kazakhstan, China (NW, NC).

Tetraloniella fulvescens (Giraud, 1863) [Tetralonia] (*Tetralonia acutangula* Morawitz, 1876; *Macroceria dufouri* Pérez, 1879). Oligolect on Inula (Asteraceae). Russia: **EP** (C, S, NC), **UR**, **WS** (KM). – Europe (WE, SE, EE), N Africa, Turkey, Azerbaijan, Kazakhstan, China (NE, NC, NW).

Tetraloniella graja (Eversmann, 1852) [Macroceria] (*Macroceria ruficornis* Lepeletier de Saint-Fargeau, 1841, nom. praeocc., nec Fabricius, 1804). Polylect, prefer Asteraceae. Russia: **EP** (C, S, NC, CR), **UR**. – Europe (WE, SE, EE), Azerbaijan, Turkey, Syria, Iran, Kazakhstan.

Tetraloniella inulae (Tkalcú, 1979) [Tetralonia]. Oligolect on Inula (Asteraceae). Russia: **EP** (S, NC, CR), **UR**. – Europe (WE, SE, EE), Cyprus, Iran.

Tetraloniella julliani (Pérez, 1879) [Tetralonia] (*Tetralonia julliani biroj* Mocsáry, 1879). Russia: **EP** (S, CR). – Europe (WE, SE, EE), Turkey, Iran, Turkmenistan.

Tetraloniella lyncea Mocsáry, 1879 (*Tetralonia lyncea*: Chenikalova, 2005). Russia: **EP** (S, NC, CR). – Europe (SE, EE).

Tetraloniella mitsukurii (Cockerell, 1911) [Tetralonia]. Oligolect on Fabaceae. Russia: **FE** (AM, KH, PR). – China (NE, NC, CC), Korean Peninsula, Japan (Hon, Shi, Kyu.).

Tetraloniella nana (Morawitz, 1874) [Tetralonia] (*Macroceria griseola* Pérez, 1879; *Tetralonia tenella* Mocsáry, 1879). Russia: **EP** (C, S, NC), **UR**. – Europe (WE, SE, EE), Azerbaijan, Turkmenistan, Kazakhstan.

Tetraloniella pollinosa (Lepeletier de Saint-Fargeau, 1841) [Macroceria] (*Macroceria mediocris* Eversmann, 1853; *Eucera canescens* Dours, 1873; *Tetralonia fossulata* Morawitz, 1874; *T. adusta* Mocsáry, 1877). Russia: **EP** (C, S, NC), **UR**. – Europe (WE, SE, EE), N Africa, Georgia, Armenia.

Tetraloniella salicariae (Lepeletier de Saint-Fargeau, 1841) [Macroceria] (*Macroceria meridiana* Dufour, 1841; *Tetralonia lythri* Schenck, 1870; *T. basalis* Morawitz, 1871; *T. salicariae* var. *albiclypeata* Dalla Torre, 1877; *T. s.* var. *flavicypeata* Dalla Torre, 1877). Collect pollen exclusively on Lythrum (Lythraceae) and Melilotus (Fabaceae). Russia: **EP** (C, E, S, NC), **UR**. – Europe (WE, SE, EE), N Africa, Caucasus, Uzbekistan, Kazakhstan.

Tetraloniella scabiosae (Mocsáry, 1881) [Eucera]. Possible oligolect on Dipsacaceae. Russia: **EP** (S). – Europe (WE, SE, EE).

Tetraloniella strigata (Lepeletier de Saint-Fargeau, 1841) [Macroceria]. Russia: **EP** (S). – Europe (WE, SE).

Tetraloniella vicina (Morawitz, 1876) [Tetralonia]. Russia: **EP** (S). – Europe (WE, EE), Georgia.

Tribe ANTHOPHORINI

AMEGILLA Friese, 1897 (*Alfkenella* Börner, 1919; *Asaropoda* Cockerell, 1926). Type species *Apis quadrifasciata* Villers, 1789. Mostly polylects, fly at summer in temperate zone. Number of taxa: World – 253 Old World species, mostly Afrotropic and Oriental; Palaearctic – 47; Russia – 9.

Amegilla (Amegilla) garrula (Rossi, 1790) [*Apis*]. Russia: **EP** (NC). – Europe (WE, SE, EE), Abkhasia, Georgia, Iran, Turkmenistan, Kazakhstan.

Amegilla (Amegilla) ochroleuca (Pérez, 1879) [*Anthophora*]. Russia: **EP** (S, CR). – Europe (WE, SE, EE), N Africa.

Amegilla (Amegilla) quadrifasciata (de Villers, 1789) [*Apis*] (*Apis nidulans* Fabricius, 1793; *Anthophora quadrifasciata* var. *albescens* Dours, 1869; *A. q.* var. *nana* Radoszkowski, 1869; *A. maderae* Sichel, 1868; *A. mervensis* Radoszkowski, 1893; *A. mediterranea* Alfken, 1927; *A. quadrifasciata* var. *teneriffensis* Cockerell, 1930; *A. klugi* Priesner, 1957; *A. litorana* Priesner, 1957). Russia: **EP** (C, E, S, CR), **UR**, **FE** (PR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Cyprus, Syria, Lebanon, Iran, Central Asia, Kazakhstan, Mongolia, China (NC, NW), Japan (Hon, Shi, Kyu), India.

Amegilla (Glossamegilla) florea (Smith, 1879) [*Anthophora*] (*Anthophora tsushimensis* Cockerell, 1926). Russia: **FE** (PR). – China (NC, CC, SE), Korean Peninsula, Japan (Hon, Shi, Kyu).

Amegilla (Micramegilla) nigricornis (Morawitz, 1872) [*Anthophora*] (*Anthophora picicornis* Fedtschenko, 1875). Probably polylect, prefer Alhagi (Fabaceae). Russia: **EP** (S, NC). – N Africa, Azerbaijan, Iran, Turkmenistan, Uzbekistan, Tajikistan, Mongolia, China (NC, NW).

Amegilla (Micramegilla) velocissima (Fedtschenko, 1875) [*Anthophora*]. Russia: **EP** (S). – Europe (EE), Armenia, Azerbaijan, Iran, Pakistan, Turkmenistan, Uzbekistan, Tajikistan, Kazakhstan, Mongolia, China (NC, NW, WP).

Amegilla (Zebramegilla) albigena (Lepeletier de Saint-Fargeau, 1841) [*Anthophora*] (*Anthophora*

binotata Lepeletier de Saint-Fargeau, 1841; *A. albigena* var. *albida* Dours, 1869; *A. quadrifasciata* var. *nana* Radoszkowski, 1870; *Podalirius albigena* var. *nigrithorax* Dalla Torre, 1877; *P. pyramidalis* Kirby, 1900; *Anthophora albigena quadrata* Cockerell, 1911; *A. a. afra* Priesner, 1957). Polylect, prefer Lamiaceae and Fabaceae. Russia: **EP** (C, S, NC, CR). – Europe (WE, SE, EE), N Africa, Georgia, Armenia, Turkey, Cyprus, Syria, Jordan, Iran, Central Asia, Kazakhstan, China (NC, NW), India.

Amegilla (Zebramegilla) magnilabris (Fedtschenko, 1875) [Anthophora]. Russia: **EP** (S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Turkey, Syria, Turkmenistan, Uzbekistan.

Amegilla (Zebramegilla) salviae (Morawitz, 1876) [Anthophora] (*Anthophora pipiens* Mocsáry, 1879). Russia: **EP** (NC, CR). – Europe (SE, EE), N Africa, Armenia, Cyprus, Turkey, Iran, China (NC, NW).

ANTHOPHORA Latreille, 1803 (*Podalirius* Latreille, 1802; *Lasius* Panzer, 1804; *Megilla* Fabricius, 1804). Type species *Apis pilipes* Fabricius, 1775. Polylects or olicolects, fly at spring or summer in temperate zone. Nests in ground or rotten wood (*Clisodon*). Number of taxa: World – 348, mostly Holarctic species; Palaearctic – 203; Russia – 42.

Anthophora (Anthomegilla) arctica Morawitz, 1883. Russia: **ES** (YA, ZB), **FE** (AM). – Tajikistan, China (NW).

Anthophora (Anthophora) fulvitarsis Brullé, 1832 (*Megilla personata* Erichson, 1835; *Anthophora nasuta* Lepeletier de Saint-Fargeau, 1841; *A. personata* var. *euris* Dours, 1870; *A. arietina* Dours, 1870). Russia: **EP** (S, NC), **UR**, **WS** (TK, KM). – Europe (WE, SE, EE), N Africa, Georgia, Turkey, Israel, Kazakhstan, China (NC, WP).

Anthophora (Anthophora) plumipes (Pallas, 1772) [Apis] (*Apis pilipes* Fabricius, 1775; *Andrena hirsuta* Fabricius, 1787; *Apis rufipes* Christ, 1791, nom. praeocc., nec Fabricius, 1781, nec Christ, 1791; *A. cochlearipes* Christ, 1791; *A. palmipes* Rossi, 1792; *Anthophora pennata* Lepeletier de Saint-Fargeau, 1841; *A. nigrofulva* Lepeletier de Saint-Fargeau, 1841; *A. sicula* Smith, 1854; *A. villosula* Smith, 1854; *A. personata* var. *squalens* Dours, 1869; *Podalirius acervorum* var. *niger* Friese, 1896; *P. a. var. nigripes* Friese, 1896; *P. acervorum* var. *albipes* Friese, 1896; *Anthophora soror* Pérez, 1905; *A. acervorum* var. *dimidiata* Alfken, 1913; *A. pingshiangensis* Strand, 1913; *A. acervorum* var. *intermixta* Alfken, 1913;

A. a. var. varians Friese, 1922; *A. a. var. lisbonensis* Cockerell, 1922; *A. a. palestinensis* Hedicke, 1936; *A. a. cypriaca* Mavromoustakis, 1957; *Apis acerorum* auct., nec Linnaeus, 1758). Russia: **EP** (N, NW, C, E, S, NC, CR), **FE** (PR). – Europe (WE, NE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Israel, Iran, Kazakhstan, China (NE, NC, NW, CC, SW, WP, SE), Japan (Hon, Shi, Kyu).

Anthophora (Anthophora) salviae (Panzer, 1804) [*Lasius*] (*A. senescens* Lepeletier de Saint-Fargeau, 1841; *A. ephippium* Lepeletier de Saint-Fargeau, 1841; *A. crinipes* Smith, 1854). Russia: **EP** (NC). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran, China (CC).

Anthophora (Caranthophora) dufourii Lepeletier de Saint-Fargeau, 1841. Russia: **EP** (S, CR). – Europe (SE), Georgia, Turkey, Syria, Israel, Iran, Turkmenistan.

Anthophora (Caranthophora) pubescens (Fabricius, 1781) [Apis] (*A. flabellifera* Lepeletier de Saint-Fargeau, 1841). Russia: **EP** (S). – Europe (WE, SE, EE), N Africa, Israel, China (NW).

Anthophora (Clisodon) furcata (Panzer, 1798) [Apis] (*Apis dumetorum* Panzer, 1798; *Megilla furcata* var. *norvegica* Nylander, 1852; *Podalirius furcata* var. *caucasicus* Friese, 1897, nom. praeocc., nec Radoszkowski, 1874; *P. furcatus montislinguarum* Schulz, 1906; *Anthophora furcata* f. *atrrata* Hedicke, 1929; *A. f. caucasicola* Hedicke, 1929; *A. f. var. obscuricolla* Alfken, 1926). Oligolect on Lamiaceae. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TK, KM, AL), **ES** (TU, IR, BR, YA). – Europe (WE, NE, SE, EE), Abkhasia, Georgia, Turkey, Iran, Kazakhstan, China (NE, NC, CC, SW, WP).

Anthophora (Clisodon) terminalis Cresson, 1869. Prefer Lamiaceae. Russia: **UR**, **WS** (KM), **ES** (TU, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA). – Mongolia, China (NE, NC, CC, SW, WP), N America.

Anthophora (Dasymegilla) quadrimaculata (Panzer, 1798) [Apis] (*Apis vulpina* Panzer, 1798, nom. praeocc., nec Christ, 1791; *A. subglobosa* Kirby, 1802; *Anthophora vara* Lepeletier de Saint-Fargeau, 1841; *A. mixta* Lepeletier de Saint-Fargeau, 1841; *A. seguina* Gribodo, 1873; *A. vulpina paehypoda* Cockerell, 1924; *A. v. alticola* Hedicke, 1931). Oligolect on Lamiaceae. Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (OM). – Europe (WE, NE, SE, EE), Georgia, Turkey, Tajikistan, Kazakhstan, China (NC, CC).

Anthophora (Heliophila) bimaculata (Panzer, 1798) [Apis] (*Apis rotundata* Panzer, 1798, nom. praeocc.,

nec Fabricius, 1787; *Anthophora saropoda* Lamarck, 1817; *A. squalida* Lepeletier de Saint-Fargeau, 1841; *A. albifrons* Eversmann, 1852). Polylect, prefer Asteraceae. Russia: **EP** (NW, C, E, S). – Europe (WE, NE, SE, EE), Georgia, Turkey, Iran, Uzbekistan, Tajikistan, Kazakhstan, China (NW).

Anthophora (Lophanthophora) affinis Brullé, 1832 (*A. biciiliata* Lepeletier de Saint-Fargeau, 1841; *A. caucasica* Radoszkowski, 1874; *A. morawitzi* Alfken et Blüthgen, 1937). Polylect, prefer Fabaceae. Russia: **EP** (CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Israel, Iran, Uzbekistan, Kazakhstan, China (NW).

Anthophora (Lophanthophora) atricilla Eversmann, 1846 (*A. atricilla aegyptorum* Priesner, 1957). Russia: **EP** (E, S, NC, CR). – N Africa, Caucasus, Turkey, Pakistan, Turkmenistan, Tajikistan, Kazakhstan, China (NW).

Anthophora (Lophanthophora) robusta (Klug, 1845) [*Megilla*] (*Megilla caliginosa* Klug, 1845; *Anthophora nigromaculata* Lucas, 1846; *A. oxygona* Dours, 1870; *A. atroferruginea* Dours, 1870; *A. robusta atra* Friese, 1919; *A. r. atratula* Friese, 1919; *A. r. atroscopacea* Friese, 1919; *A. fuliginosa pamirica* Hedicke, 1931). Russia: **EP** (NC, CR). – Europe (WE, SE, EE), Georgia, Azerbaijan, Turkey, Syria, Israel, Iran, Turkmenistan.

Anthophora (Lophanthophora) rutilans Dour, 1870 (*A. moderna* Morawitz, 1877). Russia: **EP** (NC). – Europe (SE), Armenia, Turkey, Cyprus, Israel, Palestine.

Anthophora (Melea) plagiata (Illiger, 1806) [*Megilla*] (*Apis parietina* Fabricius, 1793, nom. praeocc., nec Fourcroy, 1785; *Anthophora villosa* Herrich-Schffer, 1840; *A. parietina* var. *fulvocinerea* Dours, 1870; *A. turanica* Fedtschenko, 1875; *A. parietina* var. *schenkii* Dalla Torre, 1877; *A. simplicipes* Morawitz, 1880; *A. mlokosewitzii* Radoszkowski, 1884; *A. nigripes* Morawitz, 1887, nom. praeocc., nec Pérez, 1879; *Podalirius simplicipes semiater* Friese, 1896; *Anthophora pulcherrima* Bingham, 1897; *Podalirius parietinus nigrescens* Friese, 1897; *Anthophora filchnerae* Friese, 1908; *A. khambana* Cockerell, 1910; *A. khambana* var. *atramentata* Cockerell, 1911; *A. pilosella* Friese, 1919; *A. semenovi* Kuznetsov-Ugamsky, 1927; *A. parietina pamiricola* Hedicke, 1931; *A. khambana* f. *chodjana* Hedicke, 1938; *A. parietina baltistanica* Hedicke, 1940; *A. parietina ladakhiana* Hedicke, 1940; *A. pulcherrima himalayaensis* Wu, 1982). Polylect, prefer Lamiaceae. Russia: **EP** (NC),

UR. – Europe (WE, NE, SE, EE), N Africa, Georgia, Turkey, Iran, Tajikistan, Kyrgyzstan, Kazakhstan, Mongolia, China (NE, NC, NW, CC, SW, WP).

Anthophora (Mystacanthophora) borealis Morawitz, 1865. Russia: **EP** (NW, C, E), **UR**, **WS** (OM, KM, AL), **ES** (IR, BU, YA, ZB), **FE** (AM, KH). – Europe (WE, SE, EE), Armenia, Turkey, Tajikistan, Kazakhstan, Mongolia, China (NE).

Anthophora (Paramegilla) aeneiventris Hedicke, 1931. Prefer Fabaceae. Russia: **FE** (PR). – Kazakhstan, Mongolia, China (NW, NC).

Anthophora (Paramegilla) astragali Morawitz, 1878. Russia: **EP** (S). – Georgia, Armenia.

Anthophora (Paramegilla) balassogloei (Radoszkowski, 1877) [*Habropoda*] (*A. christofi* Morawitz, 1880). Russia: **EP** (E, NC), **UR**. – Armenia, Turkmenistan.

Anthophora (Paramegilla) deserticola Morawitz, 1872. Russia: **EP** (E, S), **UR**, **WS** (TK), **ES** (KS, KR, ZB, BR). – Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China (NW, NC).

Anthophora (Paramegilla) dubia Eversmann, 1852 (*A. albomaculata* Radoszkowski, 1874; *A. saussurei* Fedtschenko, 1875; *A. semperi* Fedtschenko, 1875; *A. carbonaria* Morawitz, 1876; *A. faddei* Radoszkowski, 1882; *A. semperi cerberus* Friese, 1919). Probably polylect, prefer Fabaceae. Russia: **EP** (NC), **UR**. – Armenia, Azerbaijan, Turkey, Iran, Central Asia, Kazakhstan, Mongolia, China (NE, NC, NW, WP).

Anthophora (Paramegilla) fulvipes Eversmann, 1846 (*Podalirius larvata* Giraud, 1863; *Anthophora croceipes* Morawitz, 1876). Russia: **EP** (C, E, S), **UR**. – Europe (SE, EE), Armenia, Turkey, China (NW, NC).

Anthophora (Paramegilla) gracilipes Morawitz, 1872. Russia: **EP** (S, NC). – Caucasus, Iran, Turkmenistan, Uzbekistan, Tajikistan, Kazakhstan, Mongolia, China (NW).

Anthophora (Paramegilla) ireos (Pallas, 1773) [*Apis*] (*A. lepida* Pallas, 1773). Oligolect on Phlomis tuberosa (Lamiaceae). Russia: **EP** (S), **UR**. – Iran, Kazakhstan.

Anthophora (Paramegilla) podagra Lepeletier de Saint-Fargeau, 1841 (*A. rufa* Lepeletier de Saint-Fargeau, 1841; *A. segnis* Eversmann, 1852; *A. cinerea* Eversmann, 1852; *A. tomentosa* Mocsáry, 1878; *A. taurica* Friese, 1922). Polylect, prefer Asteraceae. Russia: **EP** (C, E, S, CR), **UR**. – Europe (WE, SE,

- EE), Armenia, Turkey, Iran, Turkmenistan, Tajikistan, Kazakhstan, China (NC, NW).
- Anthophora (Paramegilla) ponomarevae** Brooks, 1988 (*A. morawitzi* Ponomareva, 1966, nom. praecocc., nec Alfken et Blüthgen, 1937). Russia: **EP** (S, CR). – Israel, Turkmenistan.
- Anthophora (Paramegilla) prshewalskii** Morawitz, 1880 (*A. pilosa* Morawitz, 1880; *A. przewalskii*: Osytshnjuk et al., 1978). Russia: **EP** (CR). – Turkey, Turkmenistan, Kazakhstan, China (NW, SW).
- Anthophora (Paramegilla) raddei** Morawitz, 1876. Russia: **EP** (NC). – Europe (EE), Armenia, Iran.
- Anthophora (Petalosternon) crassipes** Lepeletier de Saint-Fargeau, 1841 (*A. quadristrigata* Dours, 1870; *A. denticrus* Morawitz, 1872; *A. perplexa* Radoszkowski, 1884). Russia: **UR**. – Europe (WE, SE, EE), N Africa, Turkey, Israel.
- Anthophora (Petalosternon) hansenii** Morawitz, 1883. Russia: **ES** (IR).
- Anthophora (Petalosternon) radoszkowskyi** Fedtschenko, 1875. Russia: **EP** (E, S, CR). – Caucasus, Tajikistan, Kyrgyzstan, Kazakhstan.
- Anthophora (Pyganthophora) aestivalis** (Panzer, 1801) [Apis] (*A. intermedia* Lepeletier de Saint-Fargeau, 1841). Russia: **EP** (C, E), **UR**, **ES** (ZB). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Israel, Jordan, Iran.
- Anthophora (Pyganthophora) altaica** Radoszkowski, 1882 (*Megilla tersa* Erichson, 1849). Russia: **UR**, **WS** (AL). – Mongolia, China (NC).
- Anthophora (Pyganthophora) cinereus** (Friese, 1896) [Podalirius] (*Podalirius cinereus* Friese, 1897). Russia: **EP** (S, NC), **UR**. – Caucasus, Kazakhstan, Turkmenistan.
- Anthophora (Pyganthophora) erschowi** Fedtschenko, 1875. Russia: **EP** (S, NC, CR), **UR**. – Europe (EE), N Africa, Georgia, Azerbaijan, Turkey, Israel, Iran, Turkmenistan, Uzbekistan, Tajikistan, Kazakhstan, China (NC, NW).
- Anthophora (Pyganthophora) nigriceps** Morawitz, 1886. Russia: **EP** (NC). – Azerbaijan, Syria, Lebanon, Israel.
- Anthophora (Pyganthophora) orientalis** Morawitz, 1878. Russia: **EP** (S, NC). – Europe (SE), N Africa, Caucasus, Turkey, Syria, Israel.
- Anthophora (Pyganthophora) pedata** Eversmann, 1852. Polylect, prefer Astragalus (Fabaceae). Russia: **EP** (S, NC), **UR**. – Azerbaijan, Turkey, Uzbekistan, Kazakhstan, China (NW).
- Anthophora (Pyganthophora) retusa** (Linnaeus, 1758) [Apis] (*Apis haworthana* Kirby, 1802; *A. penipes* Kirby, 1802; *Anthophora intermedia* Lepeletier de Saint-Fargeau, 1841, part.; *Megilla monacha* Erichson, 1849; *Anthophora hispanica*: Eversmann, 1852; *A. ruthenica* Morawitz, 1870; *A. retusa* var. *meridionalis* Pérez, 1879; *Podalirius retusus sareptanus* Friese, 1896; *Anthophora retusiformis* Cockrell, 1911; *A. retusa fasciata* Alfken, 1913; *A. aestivalis baicalensis* Hedicke, 1929; *A. retusa semi-nigra* Benoist, 1930; *A. aestivalis alaica* Hedicke, 1931; *A. monacha tschelcarica* Ponomareva, 1967; *A. rudolphae* Romankova, 2003). Polylect, prefer Lamiaceae. Russia: **EP** (C, E, S), **UR**, **WS** (TK, AL), **ES** (KR, IR, BR, YA), **FE** (AM, PR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Central Asia, Kazakhstan, Mongolia, China (NC, NW).
- Anthophora (Pyganthophora) sichelii** Radoszkowski, 1869. Russia: **EP** (S). – Europe (WE, SE), N Africa, Caucasus, Turkmenistan, Uzbekistan.
- Anthophora (Pyganthophora) testaceipes** Morawitz, 1888. Oligolect on Fabaceae. Russia: **UR**, **ES** (BR). – Uzbekistan, Kazakhstan.
- Anthophora (Pyganthophora) vernalis** Morawitz, 1877 (*A. atricilla*: Eversmann, 1852). Russia: **EP** (NC), **UR**. – Israel, Uzbekistan, Kazakhstan.
- HABROPODA** Smith, 1854 (*Habrophora* Smith, 1854, nom. praecocc., nec Erichson, 1846; *Emphoropsis* Ashmead, 1899; *Meliturgopsis* Ashmead, 1899). Type species *Habropoda ezonata* Smith, 1854 = *Tetralonia tarsata* Spinola, 1838. Early summer, summer or autumn species, probably polylects, nests in borrows in soil. Number of taxa: World – 50; Palaearctic – 17; Russia – 1.
- Habropoda zonatula** Smith, 1854. Russia: **EP** (NC, CR). – Europe (WE, SE, EE), Azerbaijan, Turkey, Syria.

Tribe MELECTINI

- MELECTA** Latreille, 1802 (*Symmorpha* Klug, 1807; *Bombomelecta* Patton, 1879). Type species *Apis punctata* Fabricius, 1775. Cleptoparasites of *Anthophora* and *Habropoda* (Apidae). Number of taxa: World – 59 in Holarctic; Palaearctic – 54; Russia – 11.

- Melecta (Melecta) albifrons** (Förster, 1771) [Apis] (*Apis punctata* Fabricius, 1775; *Andrena armata*

Panzer, 1799; *Melecta nigra* Spinola, 1806; *M. fasciculata* Spinola, 1806; *Crocisa atra* Jurine, 1807; *Melecta tisiphone* Newman, 1835; *M. alecto* Newman, 1835; *M. clotho* Newman, 1835; *M. megaera* Newman, 1835; *M. albovaria* Erichson, 1841; *M. bipunctata* Lepeletier de Saint-Fargeau, 1841; *M. atcrrima* Lepeletier de Saint-Fargeau, 1841; *M. calabrina* Radoszkowski, 1876; *M. pseudoarmata* Radoszkowski, 1893; *M. armata* var. *mediterranea* Gribodo, 1893; *M. novellai* Dusmet, 1915; *M. armata* var. *gigantea* Friese, 1925; *M. luctuosa* var. *minima* Friese, 1925). Cleptoparasite of *Anthophora plumipes* (Pall.), *A. plagiata* (Ill.) and *A. fulvitarsis* Brull. (Apidae). Russia: **EP** (NW, C, E, CR), **UR**. – Europe (NE, SE, EE), N Africa, Caucasus, Turkey, Cyprus, Syria, Lebanon, Palestine, Israel, Iran.

Melecta (Melecta) amanda Lieftinck, 1980. Russia: **EP** (S). – Kazakhstan.

Melecta (Melecta) baerii (Radoszkowski, 1865) [Pseudomelecta]. Russia: **UR**, **WS** (AL). – Caucasus, Kazakhstan.

Melecta (Melecta) diacantha Eversmann, 1852. Russia: **EP** (C), **UR**.

Melecta (Melecta) duodecimmaculata (Rossi, 1790) [*Nomada*] (*M. plurinotata* Brullé, 1832; *M. quatuordecimpunctata* Fischer de Waldheim, 1843; *M. soederbomi* Alfken, 1936; *M. jakovlewii* Radoszkowski, 1877). Russia: **EP** (S, CR), **UR**, **ES** (KR). – Europe (WE, SE, EE), N Africa, Turkey, Iraq, Iran, Central Asia, Kazakhstan, China (NE, NC, NW).

Melecta (Melecta) eversmanni Radoszkowski, 1893. Russia: **EP** (S), **UR**. – Uzbekistan.

Melecta (Melecta) festiva Lieftinck, 1980. Russia: **EP** (NC, CR). – Europe (WE, SE, EE), Georgia, Armenia, Turkey, Palestine, Israel, Turkmenistan, Uzbekistan.

Melecta (Melecta) luctuosa (Scopoli, 1770) [*Apis*] (*M. lachesis* Newman, 1835; *M. atropos* Newman, 1835; *M. fasciculata* Fischer de Waldheim, 1843, nom. praeocc., nec Spinola, 1806; *M. eversmanni* Radoszkowski, 1893; *M. eczmiadzini* Radoszkowski, 1893). Cleptoparasite of *Anthophora aestivalis* (Pz.), *A. retusa* (L.), *A. plagiata* Ill. and *A. crinipes* Sm. (Apidae). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (OM, TK, KM, AL), **ES** (TU, IR, BR, YA, ZB), **FE** (AM). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Cyprus, Lebanon, Israel, Jordan, Iraq, Iran, Turkmenistan, Uzbekistan, Tajikistan, Kazakhstan.

Melecta (Melecta) mundula Lieftinck, 1983 (*M. megaera* sensu Lieftinck, 1980, non Newman, 1835; *M. cypriaca* Schwarz, 1999). Russia: **EP** (S). – Europe (SE), Turkey, Palestine, Israel, Iran.

Melecta (Melecta) sibirica Radoszkowski, 1891. Russia: **ES** (IR). – Afghanistan, Turkmenistan, Tajikistan, Kyrgyzstan.

Melecta (Melecta) tuberculata Lieftinck, 1980. Russia: **EP** (NC). – Europe (WE, SE, EE), Turkey, Cyprus, Lebanon, Israel.

THYREOMELECTA Rightmyer et Engel, 2003. Type species *Thyreomelecta kirghisia* Rightmyer et Engel, 2003. Nest parasites of *Anthophora* (Apidae). Number of taxa: World – 7 Palaearctic species; Russia – 2.

Thyreomelecta propinqua (Lieftinck, 1968) [Thyreus]. Russia: **ES** (TU, BR), **FE** (PR). – China (NE), Korean Peninsula.

Thyreomelecta sibirica (Radoszkowski, 1893) [Crocisa] (*Thyreus sibirica*: Romankova, 1995b). Russia: **ES** (IR, KR, BR, YA, ZB), **FE** (AM, MG). – Armenia, China (NE).

THYREUS Panzer, 1806 (*Crocisa* Jurine, 1807). Type species *Nomada scutellaris* Fabricius, 1781. Cleptoparasites of *Amegilla*, some species of *Anthophora* and *Eucera* subgen. *Synchalonia* (Apidae). Number of taxa: World – 123 species in Old World, mostly Indo-Australian; Palaearctic – 40; Russia – 10.

Thyreus aberrans (Morawitz, 1875) [Crocisa]. Russia: **EP** (S). – Uzbekistan, Tajikistan.

Thyreus affinis (Morawitz, 1874) [Crocisa] (*Crocisa portchinski* Radoszkowski, 1893; *C. transcaspica* Radoszkowski, 1893; *C. tunensis* Pérez, 1895; *C. affinis* var. *villosa* Meyer, 1921). Cleptoparasite of *Amegilla salvia* (Mor.) (Apidae). Russia: **EP** (C, S, NC, CR), **UR**. – Europe (WE, SE, EE), N Africa, Azerbaijan, Armenia, Turkey, Cyprus, Pakistan, Iran, Uzbekistan, Kyrgyzstan, Kazakhstan.

Thyreus altaicus (Radoszkowski, 1893) [Crocisa]. Russia: **WS** (AL), **ES** (KS, KR, BR, ZB), **FE** (PR). – Kazakhstan, Mongolia, China (NE, NC).

Thyreus decorus (Smith, 1852) [Crocisa] (*Crocisa japonica* Friese, 1905; *C. kanshireana* Cockerell, 1911; *C. pallescens* Cockerell, 1927). Russia: **FE** (PR). – China (SE), Korean Peninsula, Japan (Hon, Shi, Kyu), Thailand, Malaysia.

Thyreus hirtus (de Beaumont, 1940) [Crocisa]. Russia: EP (S). – Europe (WE, SE), N Africa, Turkey.

Thyreus histrionicus (Illiger, 1806) [Melecta] (*Crocisa major* Morawitz, 1875; *C. divisa* Pérez, 1905; *C. histrionica* var. *alboscutellata* Meyer, 1921; *C. rimosiscutum* Alfken, 1927). Cleptoparasite of *Amegilla quadrifasciata* (de Villers) (Apidae). Russia: EP (C, E, S, CR), UR, WS (KM). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Cyprus, Yemen, Iran, Central Asia, Kazakhstan, China (NW).

Thyreus orbatus (Lepeletier de Saint-Fargeau, 1841) [Crocisa] (*Crocisa scutellaris* auct., nec Fabricius, 1781). Cleptoparasite of *Anthophora parietina* Mor. and *A. quadrimaculata* (Pz.) (Apidae). Russia: EP (S). – Europe (WE, SE, EE), N Africa, Turkey.

Thyreus ramosus (Lepeletier de Saint-Fargeau, 1841) [Crocisa] (*Crocisa rufa* Radoszkowski, 1886; *C. ashabadiensis* Radoszkowski, 1893; *C. caucasica* Radoszkowski, 1893; *C. ramosa* var. *albociliata* Meyer, 1921; *C. affinis* var. *minor* Friese, 1925; *C. circulata* Alfken, 1927). Cleptoparasite of *Amegilla albigena* (Lep.) and possible of *Anthophora pubescens* (F.) (Apidae). Russia: EP (C, E, S, CR). – Europe (WE, SE, EE), N Africa, Caucasus, Turkey, Cyprus, Syria, Palestine, Israel, Iraq, Saudi Arabia, Yemen, Iran, Afghanistan, Turkmenistan, Uzbekistan, Kazakhstan, China (NE, NC, WP).

Thyreus scutellaris (Fabricius, 1781) [Nomada] (*Crocisa crassicornis* Morawitz, 1890). Cleptoparasite of *Anthophora borealis* Mor. (Apidae). Russia: EP (C, S, NC), UR, WS (TK, KM), ES (ZB), FE (AM, PR). – Europe (SE, EE), N Africa, Armenia, Turkey, Syria, Iran, Afghanistan, Turkmenistan, Kazakhstan, China (NC).

Thyreus truncatus (Pérez, 1883) [Crocisa] (*Crocisa ramosa* var. *mucorea* Friese, 1925; *C. nadigi* Alfken, 1933; *C. curviscutum* Alfken, 1934). Cleptoparasite of *Amegilla garrula* (Rossi) (Apidae). Russia: EP (C, CR), UR. – Europe (WE, SE, EE), N Africa, Georgia, Azerbaijan, Turkey, Palestine, Kuwait, Iran, Tajikistan.

Tribe BOMBINI

BOMBUS Latreille, 1802 (*Psithyrus* Lepeletier de Saint-Fargeau, 1833). Type species *Apis terristris* Linnaeus, 1758. Almost worldwide distributed genus, absent in Afrotropics and Oceania and most reachly represented in highland Palaearctic. Some species were introduced into New Zealand

and Australia. Most species are social insects with annual colonies, some species are cleptoparasitic, including subgenus *Psithyrus*. Number of taxa: World – 264 species, Palaearctic – 121 species, Russia – 90.

Bombus (Alpigenobombus) wurlenii Radoszkowski, 1859. Russia: EP (NC), UR. – Europe (WE, NE, SE, EE), Caucasus, Turkey.

Bombus (Alpinobombus) alpinus (Linnaeus, 1758) [Apis]. Russia: EP (N). – Europe (WE, NE, SE, EE).

Bombus (Alpinobonbus) balteatus Dahlbom, 1832 (*B. balteatus lysholmi* Friese, 1905; *B. kirbyellus friesei* Skorikov, 1908; *Alpinobombus kirbyellus gmelini* Skorikov, 1914; *A. k. subbalteatus* Skorikov, 1914; *A. k. subcollaris* Skorikov, 1914; *A. k. appropinquans* Skorikov, 1914). Russia: EP (N), WS (TM), ES (KR, YA), FE (KH, KU, KA, MG, CH). – Europe (NE), Mongolia.

Bombus (Alpinobonbus) hyperboreus Schönherr, 1809. Facultative inquiline in nests *Bombus pyrrhopygus* Friese and *B. jonellus* (Kirby) (Apidae). Russia: EP (N), WS (TM), ES (KR, YA), FE (KA, MG, CH). – Europe (NE).

Bombus (Alpinobombus) pyrrhopygus Friese, 1902 (*B. polaris* auct., nec Curtis, 1835). Russia: EP (N), WS (TM), ES (KR, YA), FE (KA, CH). – Europe (NE).

Bombus (Bombias) confusus confusus Schenck, 1859. Russia: EP (NW, C, E, S), UR. – Europe (WE, NE, SE, EE).

Bombus (Bombias) confusus paradoxus Dalla Torre, 1882. Russia: EP (C, E, S), UR, WS (OM, TK, NS, KM, AL), ES (KS, KR). – Europe (WE, NE, SE, EE), Kazakhstan.

Bombus (Bombus) cryptarum (Fabricius, 1775) [Apis] (*B. lucorum albocinctus* Smith, 1854; *B. florilegus* Panfilov, 1956). Russia: EP (N, NW, C, E, NC), UR, WS (TM, NS, AL), ES (KS, KR, IR, BR, ZB), FE (AM, KH, PR, SA, KU, KA, MG, CH). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Kyrgyzstan, Kazakhstan, Mongolia, China (NW, NC), Japan (Hok), India, Nepal, N America.

Bombus (Bombus) hypocrita Pérez, 1905 (*B. sapporoensis* Cockerell, 1911; *B. jesoenensis* Matsumura, 1911; *B. ikonnikovi* Skorikov, 1914; *B. i. mariae* Skorikov, 1914; *B. i. catagraphus* Skorikov, 1914; *B. i. invitabilis* Skorikov, 1914; *B. i. vanus* Skorikov, 1914). Russia: FE (KH, PR, SA, KU). – Korean Peninsula, Japan (Hok, Hon, Shi).

Bombus (Bombus) ignitus Smith, 1869 (*B. ignitus subcollaris* Skorikov, 1914). Russia: **FE** (PR, SA). – China (NE, NC, NW, SW, WP), Korean Peninsula, Japan (Hon, Kyu).

Bombus (Bombus) lucorum (Linnaeus, 1761) [Apis]. Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (NS, AL), **ES** (KS, YA, ZB). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Kyrgyzstan, Mongolia, China (NE, NW, NC).

Bombus (Bombus) magnus Vogt, 1911. Russia: **EP** (N, NW, C). – Europe (WE, NE, SE, EE).

Bombus (Bombus) patagiatus Nylander, 1848. Russia: **EP** (N, NW, C, E), **UR**, **WS** (TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, MG). – Europe (NE), Mongolia, China (NE, NC, CC, WP), Korean Peninsula.

Bombus (Bombus) sporadicus Nylander, 1848 (*B. terrestris czerskianus* Vogt, 1911; *Terrestribombus terrestris malaisei* Bischoff, 1930; *Bombus sporadicus sachalinensis* Krüger, 1956). Russia: **EP** (N, NW, C), **UR**, **WS** (TM, TK, NS, KM, AL), **ES** (TU, KR, IR, BR, YA, ZB), **FE** (KH, PR, SA, KA, MG). – Europe (NE), Mongolia, China (NC, NW, WP), Korean Peninsula.

Bombus (Bombus) terrestris (Linnaeus, 1758) [Apis]. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (OM, NS, AL). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Syria, Jordan, Lebanon, Israel, Iraq, Iran, Central Asia, Kazakhstan, China (NW), Japan (introduced), N and S America (introduced), Australia (introduced), New Zealand (introduced).

Bombus (Cullumanobombus) cullumanus apollineus Skorikov, 1910. Russia: **EP** (NC). – Caucasus, Turkey, Iran.

Bombus (Cullumanobombus) cullumanus serrisquamis Morawitz, 1888 (*B. silantjewi* Morawitz, 1891; *B. praemarinus* Panfilov, 1951). Russia: **EP** (C, E, S, NC), **UR**, **WS** (TM, OM, NS, KM, AL), **ES** (KS, KR, IR), **FE** (PR). – Europe (WE, SE, EE), Caucasus, Turkey, Iran, Afghanistan, Central Asia, Kazakhstan, China (NW).

Bombus (Cullumanobombus) semenoviellus Skorikov, 1910. Russia: **EP** (N, NW, C, E), **UR**, **WS** (TM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB). – Europe (WE, NE, SE, EE), Kazakhstan.

Bombus (Cullumanobombus) unicus Morawitz, 1883 (*B. jankowskii* Radoszkowski, 1888; *B. controversus* Skorikov, 1910). Russia: **FE** (AM, KH, PR).

Bombus (Kallobombus) soroeensis (Fabricius, 1777) [Apis] (*B. proteus* Gerstcker, 1869). Russia: **EP** (N,

NW, C, E, S, NC), **UR**, **WS** (OM, TK, NS, KM, AL), **ES** (KS, TU, KR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Syria, Iraq, Iran, Central Asia, Kazakhstan, Mongolia.

Bombus (Megabombus) argillaceus (Scopoli, 1763) [Apis]. Russia: **EP** (C, E, S, NC, CR). – Europe (WE, SE, EE), Turkey, Lebanon, Iraq, Iran, Turkmenistan.

Bombus (Megabombus) consobrinus Dahlbom, 1832 (*Hortobombus consobrinus wittenburgi* Vogt, 1911, *H. consobrinus subvinogradovi* Vogt, 1911; *H. c. nigroventralis* Vogt, 1911; *Bombus solowjowfkae* Matsu-mura, 1911; *Hortobombus consobrinus ochroleucus* Skorikov, 1914; *H. c. submonochromos* Skorikov, 1914; *H. c. derzhavini* Skorikov, 1914; *H. c. nigrociliatus* Skorikov, 1914; *H. c. albociliatus* Skorikov, 1914; *H. c. bianchii* Skorikov, 1914; *H. c. dianae* Skorikov, 1914; *H. c. ferrugineociliatus* Skorikov, 1914; *Bombus yezoensis* Matsumura, 1932). Russia: **EP** (N, NW, C, E), **UR**, **WS** (TM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KA, MG). – Europe (NE), Kazakhstan, Mongolia, China (NW, WP, N), Korean Peninsula, Japan (Hon).

Bombus (Megabombus) czerskii Skorikov, 1910 (*B. czerskii suberiophoroides* Skorikov, 1915). Russia: **ES** (BR), **FE** (PR). – Mongolia, China (NC, NW), Korean Peninsula.

Bombus (Megabombus) diversus Smith, 1869 (*B. ussurensis atrocaudatus* Vogt, 1911). Russia: **FE** (PR, SA, KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).

Bombus (Megabombus) gerstaeckeri Morawitz, 1881. Russia: **EP** (NC). – Europe (WE, SE, EE), Georgia.

Bombus (Megabombus) hortorum (Linnaeus, 1761) [Apis]. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA), **FE** (PR). – Europe (WE, NE, SE, EE), Turkey, Iran, Central Asia, Kazakhstan, Mongolia, New Zeland (introduced).

Bombus (Megabombus) portchinskyi Radoszkowski, 1883. Russia: **EP** (NC). – Caucasus, Turkey, Iran.

Bombus (Megabombus) ruderatus (Scopoli, 1763) [Apis]. Russia: **EP** (NW, C, E, S, CR), **UR**. – Europe (WE, NE, SE, EE), N Africa, S America (introduced).

Bombus (Megabombus) saltuarius (Skorikov, 1931) [*Hortobombus*]. Russia: **EP** (N), **WS** (TM, NS), **ES** (KS, KR, TU), **FE** (MG).

Bombus (Megabombus) tichenkoi (Skorikov, 1926) [*Hortobombus*] (*Hortobombus tichenkoi*

- pseudoligisticus* Skorikov, 1926; *Bombus tersatus kurilensis* Sakagami, 1954). Russia: **FE** (KU, KA).
- Bombus (Megabombus) ussurensis** Radoszkowski, 1877 (*Diversobombus ussuriensis postzonatus* Skorikov, 1915; *D. u. olivaceus* Skorikov, 1915). Russia: **FE** (AM, KH, PR, SA). – China (NE, NC), Korean Peninsula, Japan (Hon).
- Bombus (Melanobombus) eriophorus** Klug, 1807 (*B. caucasicus* Radoszkowski, 1860). Russia: **EP** (NC). – Caucasus, Turkey.
- Bombus (Melanobombus) keriensis** Morawitz, 1887 (*B. alagesianus* Reinig, 1930). Russia: **EP** (NC), **WS** (AL), **ES** (TU). – Caucasus, Turkey, Iran, Afghanistan, Pakistan, Central Asia, Kazakhstan, Mongolia, China (NC, NW, SW, WP), India.
- Bombus (Melanobombus) lapidarius** (Linnaeus, 1758) [Apis]. Russia: **EP** (N, NW, C, E, S, CR), **UR**. – Europe (WE, NE, SE, EE), N Africa, Turkey, New Zealand (introduced).
- Bombus (Melanobombus) sichelii** Radoszkowski, 1860 (*Lapidariobombus sicheli czerskiae* Vogt, 1911). Russia: **EP** (N, C, E, NC), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KA, MG). – Europe (WE, SE, EE), Caucasus, Turkey, Kazakhstan, Mongolia, China (NE, NC, NW, WP), Korean Peninsula.
- Bombus (Mendacibombus) margreiteri** Skorikov, 1910 (*B. altaicus* Skorikov, 1910). Russia: **WS** (AL), **ES** (TU, IR, BR, ZB), **FE** (KA). – Kyrgyzstan, Kazakhstan, Mongolia, China (NW, WP).
- Bombus (Mendacibombus) handlirschianus** Vogt, 1909 (*B. shaposhnikovi* Skorikov, 1910). Russia: **EP** (NC). – Caucasus, Turkey, Iraq, Iran.
- Bombus (Psithyrus) barbutellus** (Kirby, 1802) [Apis] (*B. maxillosus* Klug, 1817). Parasite in nests of *Bombus hortorum* (L.) and *B. hypnorum* (L.) (Apidae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, BR, YA, ZB), **FE** (PR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Kyrgyzstan, Kazakhstan, Mongolia, China (NC).
- Bombus (Psithyrus) bohemicus** Seidl, 1837 (*Psithyrus distinctus* Pérez, 1884; *P. vestalis* auct., nec Fourcroy, 1785; *Bombus naiptchianus* Matsumura, 1911). Parasite in nests of *Bombus lucorum* (L.) (Apidae). Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KA, CH). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran,
- Central Asia, Kazakhstan, Mongolia, China (NW, NC, NE, SE), India, N America.
- Bombus (Psithyrus) branickii** (Radoszkowski, 1893) [*Psithyrus*] (*Apathon chloronotus* Morawitz, 1893; *Psithyrus rupestris eriophoroides* Reinig, 1930; *P. r. elisabethae* Reinig, 1940). Host is unknown. Russia: **WS** (AL), **ES** (ZB). – Afghanistan, Pakistan, Central Asia, Mongolia, China (NC, NW, SW, WP), Korean Peninsula, India.
- Bombus (Psithyrus) campestris** (Panzer, 1801) [Apis]. Parasite in nests of *Bombus pascuorum* (Scop.), *B. humilis* Illiger, *B. pomorum* (Panz.) and *B. pratorum* (L.) (Apidae). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA), **FE** (AM, PR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Uzbekistan, Kazakhstan, Mongolia, China (NW, NC).
- Bombus (Psithyrus) flavidus** Eversmann, 1852 (*Psithyrus flavidus frisoni* Popov, 1931). Parasite in nests of *Bombus jonellus* (Kirby) (Apidae). Russia: **EP** (N, NW), **WS** (TM), **ES** (KS, KR, BR, YA), **FE** (SA, KU, KA, MG). – Europe (WE, NE, SE, EE), N America.
- Bombus (Psithyrus) norvegicus** (Sparre-Schneider, 1918) [Psithyrus]. Parasite in nests of *Bombus hypnorum* (L.) (Apidae). Russia: **EP** (N, NW, C, E), **UR**, **WS** (TM, TK, NS, KM, AL), **ES** (KS, KR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KA). – Europe (WE, NE, SE, EE), Kazakhstan, Mongolia, China (NC, NE, SE), Japan (Hok, Hon), N America.
- Bombus (Psithyrus) quadricolor** (Lepeletier de Saint-Fargeau, 1832) [Psithyrus]. Parasite in nests of *Bombus soroeensis* (F.) (Apidae). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (TM, TK, NS, KM, AL), **ES** (KR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran.
- Bombus (Psithyrus) rupestris** (Fabricius, 1793) [Apis] (*Psithyrus rupestris buyssoni* Vogt, 1911). Parasite in nests of *Bombus lapidarius* (L.), *B. sichelii* Rad., *B. sylvarum* (L.) and *B. pascuorum* (Scop.) (Apidae). Russia: **EP** (N, NW, C, E, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, MG). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Mongolia, China (NW, NC, NE, SE).
- Bombus (Psithyrus) sylvestris** (Lepeletier de Saint-Fargeau, 1832) [Psithyrus]. Parasite in nests of *Bombus pratorum* (L.), *B. jonellus* (Kirby) and *B. schrencki* Mor. (Apidae). Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (TM, TK, NS, KM, AL), **ES** (KS,

- KR, YA), **FE** (AM, KH, PR, SA, MG). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Kazakhstan, Mongolia, China (NC, NE), Korean Peninsula.
- Bombus (Psithyrus) vestalis** (Fourcroy, 1785) [Apis]. Parasite in nests of *Bombus terrestris* (L.) (Apidae). Russia: **EP** (N, C, E, NC, CR). – Europe (WE, NE, SE, EE), N Africa, Caucasus, Turkey, Iran.
- Bombus (Pyrobombus) ardens** Smith, 1879 (*B. ardens sakagamii* Tkalcú, 1962). Russia: **FE** (SA, KU). – Korean Peninsula, Japan (Hok, Hon, Shi, Kyu).
- Bombus (Pyrobombus) beaticola** (Tkalcú, 1968) [*Pyrobombus*] (*B. beaticola moskhareppus* Sakagami et Ishikawa, 1969; *B. b. shikotanensis* Ito et Sakagami, 1980). Russia: **FE** (SA, KU). – Japan (Hok, Hon).
- Bombus (Pyrobombus) biroi** Vogt, 1911. Russia: **WS** (AL). – Afghanistan, Central Asia, Kazakhstan, Mongolia, India.
- Bombus (Pyrobombus) brodmannicus** Vogt, 1909. Russia: **EP** (NC). – Europe (WE, SE), Caucasus, Turkey.
- Bombus (Pyrobombus) cingulatus** Wahlberg, 1854 (*B. hypnorum tilingi* Morawitz, 1881; *B. cingulatus pseudocalidus* Reinig, 1936). Russia: **EP** (N), **UR**, **WS** (TM), **ES** (KS, KR, YA), **FE** (KH, PR, SA, KU, KA, CH, MG). – Europe (NE).
- Bombus (Pyrobombus) glacialis** Friese, 1902 (*B. glacialis* Sparre-Schneider, 1909). Russia: **EP** (N), **FE** (KA).
- Bombus (Pyrobombus) haematurus** Kriechbaumer, 1870. Russia: **EP** (NC, CR). – Europe (SE, EE), Azerbaijan, Turkey, Iran, Turkmenistan.
- Bombus (Pyrobombus) hypnorum** (Linnaeus, 1758) [Apis] (*B. calidus* Erichson, 1851; *Pratobombus hypnorum klutschianis* Bischoff, 1930; *Bombus koropokrus* Sakagami and Ishikawa, 1972). Russia: **EP** (N, NW, C, E, NC, CR), **UR**, **WS** (TM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (KH, PR, SA, KU, KA, MG, CH). – Europe (WE, NE, SE, EE), Georgia, Kazakhstan, Mongolia, China (NE, NC, NW, CC, SW, WP, SE), Korean Peninsula, Japan (Hok), Nepal, Myanmar, India.
- Bombus (Pyrobombus) jonellus** (Kirby, 1802) [Apis]. Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (TM, TK, NS, AL), **ES** (TU, KR, IR, BR, YA, ZB), **FE** (KH, SA, KA, MG, CH). – Europe (WE, NE, SE, EE), Georgia, Mongolia, N America.
- Bombus (Pyrobombus) lapponicus** (Fabricius, 1793) [Apis] (*B. lapponicus kamtshaticus* Skorikov, 1912; *B. l. obscurus* Skorikov, 1912; *B. l. occultodistinctus* Skorikov, 1912; *B. l. rarior* Skorikov, 1912; *B. l. kaganinus* Skorikov, 1912; *B. l. commutabilis* Skorikov, 1912; *B. l. virgatus* Skorikov, 1912; *B. l. korjak* Skorikov, 1912; *B. l. ceciliae* Skorikov, 1912; *B. l. pallido-caudatus* Skorikov, 1912; *B. l. cecilioides* Skorikov, 1912; *B. l. simius* Skorikov, 1912). Russia: **EP** (N), **WS** (TM, AL), **ES** (KR, BR, YA), **FE** (KH, KA, MG, CH). – Europe (NE), Kyrgyzstan.
- Bombus (Pyrobombus) modestus** Eversmann, 1852 (*B. nymphae* Skorikov, 1910; *Pratobombus modestus krylovi* Skorikov, 1915). Russia: **EP** (N, C, E), **UR**, **WS** (TM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KA, MG). – Kazakhstan, Mongolia, China (NC), Korean Peninsula.
- Bombus (Pyrobombus) monticola** Smith, 1849. Russia: **EP** (N). – Europe (WE, SE, NE).
- Bombus (Pyrobombus) oceanicus** Friese, 1909. Russia: **FE** (KU).
- Bombus (Pyrobombus) pratorum** (Linnaeus, 1761) [Apis]. Russia: **EP** (N, NW, C, E, NC, CR), **UR**, **WS** (TM, TK, KM, AL), **ES** (KR, YA). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Australia (introduced).
- Bombus (Sibiricobombus) asiaticus** Morawitz, 1875. Russia: **WS** (AL), **ES** (TU). – Afghanistan, Pakistan, Central Asia, Mongolia, China (NW, NC, WP), India, Nepal.
- Bombus (Sibiricobombus) niveatus** Kriechbaumer, 1870. Russia: **EP** (NC). – Europe (SE, EE), Turkey, Syria, Lebanon, Iran, Turkmenistan.
- Bombus (Sibiricobombus) sibiricus** (Fabricius, 1781) [Apis]. Russia: **ES** (KS, TU, KR, IR, BR, ZB). – Kazakhstan, Mongolia, China (NE, NC, NW, WP), Korean Peninsula.
- Bombus (Subterraneobombus) amurensis** Radozowski, 1862 (*B. fragrans mongol* Skorikov, 1931). Russia: **ES** (KS, KR, BR, ZB). – Kazakhstan, Mongolia, China (NC, NW).
- Bombus (Subterraneobombus) distinguendus** Morawitz, 1869. Russia: **EP** (N, NW, C, E, S, NC), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (KH, PR, SA, KA, MG). – Europe (WE, NE, SE, EE), Uzbekistan, Kazakhstan, Mongolia, N America.
- Bombus (Subterraneobombus) fragrans** (Pallas, 1771) [Apis]. Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (OM, NS, AL), **ES** (KS, TU, KR, BR). – Europe (WE, SE, EE), Caucasus, Turkey, Iran, Kyrgyzstan, Kazakhstan, China (NW).

Bombus (Subterraneobombus) melanurus Lepeletier de Saint-Fargeau, 1836. Russia: **WS** (AL), **ES** (TU). – Caucasus, Turkey, Lebanon, Iran, Afghanistan, Central Asia, Mongolia, China (NC, NW, WP), India, Nepal.

Bombus (Subterraneobombus) subterraneus (Linnaeus, 1758) [Apis]. Russia: **EP** (N, NW, C, E, NC, CR), **UR**, **WS** (OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, ZB). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Kyrgyzstan, Kazakhstan, Mongolia, China (NC, NW), New Zealand (introduced).

Bombus (Subterraneobombus) tshitscherini Radoszkowski, 1862. Russia: **WS** (AL), **ES** (TU, BR, ZB). – Mongolia, China (NC, NW).

Bombus (Thoracobombus) anachoreta (Skorikov, 1914) [Agrobombus] (*Agrobombus anachoreta czerwinski* Skorikov, 1914). Russia: **FE** (PR). – China (NE, NC), Korean Peninsula.

Bombus (Thoracobombus) armeniacus Radozhkowski, 1877 (*B. dumoucheli* auct., nec Radoszkowski, 1884; *Fervidobombus scythes* Skorikov, 1925). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (OM, NS, KM, AL), **ES** (KS, TU, KR). – Europe (SE, EE), Caucasus, Turkey, Iraq, Iran, Central Asia, Kazakhstan, China (WP).

Bombus (Thoracobombus) deuteronymus Schulz, 1906 (*B. subbaicalensis* auct., nec. Vogt, 1911; *B. humilis* Illiger, 1806; *Agrobombus superequester* Skorikov, 1926). Russia: **EP** (NW, C, E), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KR, IR, BR, YA, ZB), **FE** (AM, PR, SA). – Europe (SE, EE), Kazakhstan, Mongolia, China (NE, NC, WP), Korean Peninsula, Japan (Hok).

Bombus (Thoracobombus) exil Skorikov, 1922 (*B. nymphae* Skorikov, 1910). Russia: **ES** (TU, IR, YA, ZB). – Mongolia, China (NC).

Bombus (Thoracobombus) filchnerae Vogt, 1908 (*Agrobombus adventor* Skorikov, 1914; *A. a. ermolaevi* Skorikov, 1914; *A. a. napalkovi* Skorikov, 1914). Russia: **WS** (AL), **ES** (KS, TU, KR). – Mongolia, China (N).

Bombus (Thoracobombus) humilis Illiger, 1806 (*B. solstitialis* Panzer, 1806; *B. helferanus* Seidl, 1838; *B. tristis* Seidl, 1838; *B. variabilis* Schmiedeknecht, 1878; *B. helferanus subbaicalensis* Vogt, 1911). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, ZB), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Kazakhstan, Mongolia, China (NW, NC, NE, SE), Korean Peninsula.

Bombus (Thoracobombus) laesus Morawitz, 1875 (*B. laesus ferrugifer* Skorikov, 1909). Russia: **EP** (C, E, S, NC, CR), **UR**, **WS** (OM, NS, AL), **ES** (KR). – Europe (WE, SE, EE), Caucasus, Turkey, Iran, Central Asia, Kazakhstan.

Bombus (Thoracobombus) mesomelas Gerstäcker, 1869. Russia: **EP** (NC, CR). – Europe (WE, SE, EE), Caucasus, Turkey, Iran.

Bombus (Thoracobombus) mlokosievitzii Radoszkowski, 1877. Russia: **EP** (NC). – Europe (SE), Caucasus, Turkey, Iran.

Bombus (Thoracobombus) mocsaryi Kriechbaumer, 1877 (*B. mocsaryi sidemii* Radoszkowski, 1888; *Agrobombus maculidorsis* Skorikov, 1922; *Bombus maculidorsis angarensis* Panfilov, 1956). Russia: **EP** (N, NW, C, E), **UR**, **WS** (TM, OM, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, ZB), **FE** (PR). – Europe (WE, NE, SE, EE), N Africa Kazakhstan, Mongolia, China (NE, NC, SE), Korean Peninsula.

Bombus (Thoracobombus) muscorum (Linnaeus, 1758) [Apis]. Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Kyrgyzstan, Kazakhstan, Mongolia, China (NE, NC).

Bombus (Thoracobombus) pascuorum (Scopoli, 1763) [Apis] (*Apis agrorum* Fabricius, 1787; *Bombus agrorum flavobarbatus* sensu Skorikov, 1915). Russia: **EP** (N, NW, C, E, S, NC, CR), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA), **FE** (AM, KH, PR, SA, KA, MG). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Syria, Iran, Kazakhstan, Mongolia, China (NC, NW), Korean Peninsula.

Bombus (Thoracobombus) persicus Radoszkowski, 1881. Russia: **EP** (NC). – Caucasus, Turkey, Iran.

Bombus (Thoracobombus) pomorum (Panzer, 1805) [Bremus]. Russia: **EP** (C, E, S, NC, CR), **UR**. – Europe (WE, NE, SE, EE), Caucasus, Turkey.

Bombus (Thoracobombus) pseudobaicalensis Vogt, 1911 (*Agrobombus gilvus* Skorikov, 1925). Russia: **WS** (TM, TK, NS, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU). – Mongolia, China (N), Korean Peninsula, Japan (Hok, Hon).

Bombus (Thoracobombus) ruderarius (Müller, 1776) [Apis]. Russia: **EP** (N, NW, C, E, NC), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR, BR, YA, ZB). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Iran, Central Asia, Kazakhstan.

Bombus (Thoracobombus) schrencki Morawitz, 1881 (*Agrobombus schrencki mironowianus* Vogt, 1911;

A. schrencki albidopleuralis Skorikov, 1915; *Bombus konakovi* Panfilov, 1956; *B. schrencki kuwayamai* Sakagami et Ishikawa, 1969). Russia: **EP** (N, NW, C, E), **UR**, **WS** (TM, OM, TK, NS, KM, AL), **ES** (KS, TU, KR, BR, YA, ZB), **FE** (AM, KH, PR, SA, KU, KA, MG). – Europe (NE, EE), Kazakhstan, China (NC), Korean Peninsula, Japan (Hok).

Bombus (Thoracobombus) sylvarum (Linnaeus, 1761) [*Apis*] (*B. daghestanicus* Radoszkowski, 1877). Russia: **EP** (NW, C, E, S, NC, CR), **UR**, **WS** (OM, NS, AL). – Europe (WE, NE, SE, EE), Caucasus, Turkey, Syria, Iran, Kazakhstan.

Bombus (Thoracobombus) tricornis Radoszkowski, 1888 (*B. tricornis biciliatus* Skorikov, 1915; *B. t. triciliatus* Skorikov, 1915). Russia: **FE** (AM, KH, PR). – China (NE, NC), Korean Peninsula.

Bombus (Thoracobombus) velox (Skorikov, 1914) [Agrobombus]. Russia: **EP** (NC). – Turkey.

Bombus (Thoracobombus) veteranus (Fabricius, 1793) [*Apis*] (*B. equestris* auct., nec Fabricius, 1793). Russia: **EP** (N, NW, C, E), **UR**, **WS** (OM, TK, NS, KM, AL), **ES** (KS, TU, KR, IR). – Europe (WE, NE, SE, EE), Central Asia, Kazakhstan, Mongolia.

Bombus (Thoracobombus) zonatus Smith, 1854. Russia: **EP** (S, NC, CR). – Europe (SE, EE), Caucasus, Turkey, Iran.

Tribe APINI

APIS Linnaeus, 1758 (*Apicula* Rafineque, 1814; *Apia-*
rus Rafineque, 1815; *Megapis* Ashmead, 1904; *Hauf-*
fapis Armbruster, 1938). Type species *Apis mellifera* Linnaeus, 1761. Live in huge family in wax nest with honey reserve. Number of taxa: World – 11 originally Old World and mostly Oriental species; Palaearctic – 3; Russia – 2.

Apis cerana Fabricius, 1793 (*A. indica* Fabricius, 1798; *A. sinensis* Smith, 1865; *A. mellifera* var. *japonica* Radoszkowski, 1887; *A. johni* Skorikov, 1929; *A. indica sinensis ussuriensis* Goetze, 1964). Russia: **FE** (KH, AM, PR). – Afghanistan, Pakistan, India, Sri-Lanka, Nepal, China, Korean Peninsula, Japan, N Vietnam, Thailand, Philippines, Mailasya, Indonesia.

Apis mellifera Linnaeus, 1758. Cosmopolitan all-world cultivated species.

REFERENCES

- Abasheev R.Yu.** 2012. *Social wasps (Hymenoptera, Vespidae: Vespinae, Polistinae) of South-West Transbaikalia*. Buryat State University Publishing Department, Ulan-Ude, 106 p. [In Russian].
- Abasheev R.Yu.** 2016. New data on the fauna and distribution of vespid wasps (Hymenoptera, Vespidae) of the Western Transbaikalia. In: Ubugunov L.L., Badmaev N.B., Baldanova D.R., Moroldoev I.V. (Eds). *Biodiversity: global and regional processes*. BSC SD RAS, Ulan-Ude: 46–47. [In Russian].
- Abasheev R.Yu. and Buyanjargal B.** 2015. To the fauna of solitary wasp of genera *Eumenes* Latreille and *Katamenes* Meado-Waldo (Hymenoptera: Vespidae, Eumeninae) of Mongolia. *A.I. Kurentsov's Annual Memorial Meetings*, 26: 277–283. [In Russian].
- Abasheev R.Yu. and Dabaev N.-B.M.** 2014. To vespid wasp fauna (Hymenoptera, Vespidae) of National Park "Alkhanai" (South-Eastern Transbaikalia). *Vestnik Buryatskogo gosudarstvennogo universiteta*, 2014(4/1): 55–58. [In Russian].
- Abe M.** 1988. A biosystematic study of the genus *Athalia* Leach of Japan (Hymenoptera: Tenthredinidae). *Esakia*, 26: 91–131.
- Achterberg C. van and Aarsten B. van.** 1986. The European Pamphiliidae (Hymenoptera: Symphyta), with special reference to the Netherlands. *Zoologische Verhandelingen*, 234: 3–98.
- Achterberg C. van and Kats R.J.M. van.** 2000. Revision of the Palaearctic Embolemidae (Hymenoptera). *Zoologische Mededelingen*, 74: 251–269.
- Agnoli G.L.** 2005. The genus *Methocha* in Europe: a discussion on taxonomy, distribution and likely origin of its known species and subspecies (Hymenoptera Tiphiidae Methochinae). *Bulletin of Insectology*, 58(1): 35–47.
- Aguiar A.P., Deans A.R., Engel M.S., Forshage M., Huber J.T., Jennings J.T., Johnson N.F., Lelej A.S., Longino J.T., Lohrmann V., Mikó I., Ohl M., Rasmussen C., Taeger A. and Yu D.S.K.** 2013. Order Hymenoptera. In: Zhang Z.-Q. (Ed.). *Animal Biodiversity: An Outline of Higher-level Classification and Survey of Taxonomic Richness (Addenda 2013)*. Zootaxa, 3703: 51–62.
- Aguib S., Louadi K. and Schwarz M.** 2010. Les Anthidiini (Megachilidae, Megachilinae) d'Algérie avec trois espèces nouvelles pour ce pays: *Anthidium (Anthidium) florentinum* (Fabricius, 1775), *Anthidium (Proanthidium) amabile* Alfken, 1932 et *Pseudaanthidium (Exanthidium) enslini* (Alfken, 1928). *Entomofauna*, 31(12): 121–152.
- Ahn N.H., Jeon M.J., Seo H.Y., Kim K.G., Park S.J. and Han S.H.** 2013. *National list of species of Korea [Insect] (Insects of North Korea)*. NIBR, Incheon, xvi + 386 p.
- Akre R.D.** 1982. Social wasps. In: Hermann H.R. (Ed.). *Social insects*. Academic Press, New York, 4: 1–105.
- Akulov E.N. and Proshchalykin M.Yu.** 2013. Contribution to the digger wasp fauna (Hymenoptera, Sphecidae, Crabronidae) of Krasnoyarsk Territory. *A.I. Kurentsov's Annual Memorial Meetings*, 24: 107–121. [In Russian].
- Alexander B.A.** 1990. A preliminary phylogenetic analysis of sphecid wasps and bees. *Sphecos*, 20: 7–16.
- Alexander B.A.** 1992. An exploratory analysis of cladistic relationships within the superfamily Apoidea, with special reference to sphecid wasps (Hymenoptera). *Journal of Hymenoptera Research*, 1: 25–61.
- Alexander B.A. and Schwarz M.** 1994. A Catalog of the species of *Nomada* (Hymenoptera: Apoidea) of the World. *The University of Kansas Science Bulletin*, 55(7): 239–270.
- Alfken J.D.** 1915. Verzeichnis der Faltenwespen (Vespiden) Nordwestdeutschlands. *Abhandlungen herausgeben vom Naturwissenschaftlichen Verein zu Bremen*, 23: 296–304.
- Alfken J.D.** 1927. Über einige Arten der *Anthophora quadri-fasciata*-Gruppe (Hym.). *Entomologische Mitteilungen*, 16: 120–122.
- Alfken J.D.** 1931. Eine neue paläarktische *Tetralonia*-Art. (Apid. Hym.). *Mitteilungen der Deutschen Entomologischen Gesellschaft*, 2: 111–112.
- Aliev Kh.A.** 2011. The Aculeate Hymenoptera in the collection of the institute of Zoologi of the Natural Academy of Sciences of Azerbaijan, Bacu. Part. 3. Bees of the genus *Nomada* Scopoli, 1770 (Hymenoptera: Anthophoridae). *Caucasian Entomological Bulletin*, 7(2): 219–225.
- Aliev Kh.A. and Qamarli V.P.** 2010. Bees of genera *Ceratina* Latreille, 1802 and *Xylocopa* Latreille, 1802 (Hymenoptera: Apoidea: Anthophoridae) of Azerbaijan. *Caucasian Entomological Bulletin*, 6(1): 109–111.
- Aliyev Kh.A., Proshchalykin M.Yu., Maharramov M.M. and Huseinzade G.A.** 2017. To the knowledge of the genus *Andrena* Fabricius, 1775 (Hymenoptera: Apoidea: Andrenidae) of Azerbaijan. *Caucasian Entomological Bulletin*, 13(1): 99–109. [In Russian].
- Aliyeva M.G.** 2010. Flight dynamics of the wasps from the Eumenidae family (Hymenoptera, Vespoidea) in the Nakhchivan AR. *Baki universitetinin xəbərləri. Təbiət elmləri seriyası*, 2010(2): 55–60. [In Russian].
- Altenhofer E. and Pschorn-Walcher H.** 2003. Biologische Notizen über die Blattwespen-Gattungen *Metallus* Forbes, *Monostegia* A. Costa und *Phymatocera* Dahlbom (Hymenoptera: Tenthredinidae). *Linzer Biologische Beiträge*, 35(1): 405–417.
- Altenhofer E. and Zombori L.** 1987. The species of *Heterarthrus* Stephens, 1835 feeding on maple (Hymenoptera, Tenthredinidae). *Annales Historico-Naturales Musei Nationalis Hungarici*, 79: 185–197.
- Amiet F., Herrmann M., Müller A. and Neumeyer R.** 2007. Apidae 5: *Ammobates*, *Ammobatoides*, *Anthophora*, *Biastes*, *Ceratina*, *Dasyopoda*, *Epeoloides*, *Epeolus*, *Eucera*, *Macropis*, *Melecta*, *Melitta*, *Nomada*, *Pasites*, *Tetralonia*, *Thyreus*, *Xylocopa*. *Fauna Helvetica*, 20: 1–356.

- Amini A., Lelej A.S., Sadeghi H. and Karimi J.** 2014. First record of the velvet ants (Hymenoptera: Mutillidae) reared from puparia of the ber fruit fly *Carpomya vesuviana* Costa (Diptera: Tephritidae) in Iran. *Zootaxa*, **3861**(6): 585–590.
- Amolin A.V.** 2009. *Ecologically-faunistic review of the wasps of the subfamily Eumeninae (Hymenoptera: Vespidae) of the Southeastern Ukraine*. DonNU, Donetsk, 123 p. [In Russian].
- Amolin A.V. and Artokhin K.S.** 2014. To the study of potter wasps (Hymenoptera: Vespidae: Eumeninae) fauna in Rostov-Don Province. *Eversmannia*, **37**: 10–18. [In Russian].
- An J.-D., Huang J.-X., Shao Y.-Q., Zhang S.-W., Wang B., Liu X.-Y., Wu J. and Williams P.H.** 2014. The bumblebees of North China (Apidae, *Bombus* Latreille). *Zootaxa*, **3830**: 1–89.
- An J.-D., Williams P.H., Zhou B.-F., Miao Z.-Y. and Qi W.-Z.** 2011. The bumblebees of Gansu, Northwest China (Hymenoptera, Apidae). *Zootaxa*, **2865**: 1–36.
- Anan'eva S.I. and Kochetkov D.N.** 1999. The aculeate fauna of the Ryazan' Oblast'. In: Anan'eva S.I. (Ed.). *Fauna i ekologiya zhivotnykh. Sbornik nauchnykh dokladov zoologicheskogo obshchestva RGPU*. Ryazanskiy Oblastnoy Institut Razvitiya Obrazovaniya, Ryazan': 4–12. [In Russian].
- André E.** 1881. *Species des Hyménoptères d'Europe & d'Algérie*. Beaune (Côte-d'Or), **1**[1879–1882](11): 565–596.
- André E.** 1881–1886. *Species des Hyménoptères d'Europe et d'Algérie*. E. André, Beaune, 2, 919 + 48 p.
- André E.** 1902 (1903). Fam. Mutillidae. In: Wytsman P., *Genera Insectorum, Fasc. 11*. Bruxelles, 77 p. + 3 pls.
- Andreeva E.M.** 1998. Some results of faunal studies of sawflies (Hymenoptera, Symphyta, Tenthredinidae) in the Urals. In: Mikhailova I.N., Golovachov I.B. (Eds). *Sovremennye problemy populatsionnoy, istoricheskoy i prikladnoy ekologii. Materialy konferentsii*. Ekaterinburg: 138–141. [In Russian].
- Antropov A.V.** 1986a. New and little known sphecid wasps of the Soviet Far East. In: Lehr P.A., Belokobylskij S.A., Storozheva N.A. (Eds). *Hymenopterous Insects of Eastern Siberia and Far East*. BPI DVNTS AN SSSR: 81–91. [In Russian].
- Antropov A.V.** 1986b. To knowledge of digger wasps from the genus *Trypoxyylon* (Hymenoptera, Sphecidae) in the Palearctic fauna. *Zoologicheskiy Zhurnal*, **65**: 624–628. [In Russian].
- Antropov A.V.** 1987. A contribution to the sphecid wasps genus *Trypoxyylon* of the eastern Palearctic. In: Lehr P.A., Storozheva N.A. (Eds). *Novye dannye po sistematike nasekomykh Dal'nego Vostoka*. BPI DVNTS AN SSSR, Vladivostok: 57–61. [In Russian].
- Antropov A.V.** 1988. A contribution to the fauna of Trypoxylini (Hymenoptera, Sphecidae) of the Soviet Far East. In: Kapustina O.G. (Ed). *Taksonomiya nasekomykh Sibirii i Dal'nego Vostoka SSR*. BPI DVNTS AN SSSR, Vladivostok: 85–88. [In Russian].
- Antropov A.V.** 1989. To the knowledge of the digger wasps of the tribe Trypoxylini (Hymenoptera, Sphecidae, Laricinae) of the Palearctic fauna. *Bulletin of the Moscow Society of Naturalists, Section of Biology*, **94**: 55–58. [In Russian].
- Antropov A.V.** 1991. On taxonomic rank of *Trypoxyylon attenuatum* Smith and adjacent species (Hymenoptera, Sphecidae). *Entomologicheskoe Obozrenie*, **70**: 672–685. [In Russian].
- Antropov A.V.** 1993. Notes on three little known Palearctic species of digger wasps (Hymenoptera, Sphecidae). *Zoologicheskiy Zhurnal*, **72**(10): 156–158. [In Russian].
- Antropov A.V.** 1994. A review of the *agile* species group of *Pison* (Hymenoptera: Sphecidae: Trypoxylini). *Journal of Hymenoptera Research*, **3**: 119–132.
- Antropov A.V. and Mokrousov M.V.** 2016. New data on distribution of four species of the genus *Trypoxyylon* (Hymenoptera: Crabronidae: Trypoxylini) in European Russia. *Russian Entomological Journal*, **25**(3): 265–269.
- Antropov A.V. and Proshchalykin M.Yu.** 2014. To the fauna of social wasps (Hymenoptera: Vespidae: Vespinae, Polistinae) of Southern Siberia. *A.I. Kurentsov's Annual Memorial Meetings*, **25**: 127–133. [In Russian].
- AntWeb.** 2017. AntWeb: *Ants of Palearctic*. Available from: <https://www.antweb.org/bioregion.do?name=Palearctic> (accessed 22 March 2017).
- Arakelian G.R.** 1994. *Fauna of the Republic of Armenia. Hymenopterous insects. Ants (Formicidae)*. Gitutyum, Erevan, 153 p. [In Russian].
- Archer M.E.** 1980. A new species of *Dolichovespula* and subspecies of *D. pacifica* (Hymenoptera: Vespidae) from China. *Entomon*, **5**: 341–344.
- Archer M.E.** 1981a. The Euro-Asian species of the *Vespula rufa* Group (Hymenoptera, Vespidae), with descriptions of two new species and one new subspecies. *Kontyû*, **49**(1): 54–64.
- Archer M.E.** 1981b. Taxonomy of the *sylvestris* group (Hymenoptera: Vespula, *Dolichovespula*) with the introduction of a new name and notes on distribution. *Entomologica Scandinavica*, **12**: 187–193.
- Archer M.E.** 1982. A revision of the subgenus *Rugovespula* nov. of the genus *Vespula* (Hymenoptera, Vespidae). *Kontyû*, **50**: 261–269.
- Archer M.E.** 1989a. A key to the world species of Vespinae (Hymenoptera): Part 1, keys, checklist, and distribution. *Research Monograph of the College of Ripon and York, St. John*, **2**: 1–47.
- Archer M.E.** 1989b. A key to the world species of Vespinae (Hymenoptera): Part 2, figures. *Research Monograph of the College of Ripon and York, St. John*, **2**: 1–34.

- Archer M.E.** 1992. The taxonomy of *Vespa crabro* L. and *V. dybowskii* André (Hym., Vespinae). *The Entomologist's Monthly Magazine*, **128**: 157–163.
- Archer M.E.** 1995. Taxonomy, distribution and nesting biology of the *Vespa mandarinia* group (Hym., Vespinae). *The Entomologist's Monthly Magazine*, **131**: 47–53.
- Archer M.E.** 1997. Taxonomy, distribution and nesting biology of the species of the Euro-Asian *Vespula rufa* group (Hym., Vespinae). *The Entomologist's Monthly Magazine*, **133**: 107–114.
- Archer M.E.** 1998a. Taxonomy, distribution and nesting biology of *Vespa orientalis* L. (Hym., Vespidae). *The Entomologist's Monthly Magazine*, **134**: 45–51.
- Archer M.E.** 1998b. The world distribution of the Eurasian species of *Paravespula* (Hym., Vespinae). *The Entomologist's Monthly Magazine*, **134**: 279–284.
- Archer M.E.** 1999a. Taxonomy, distribution and nesting biology of *Vespa binghami*, *V. basalis*, *V. variabilis*, *V. fervida*, *V. luctuosa*, *V. multimaculata* and *V. bellicosa* (Hym., Vespinae). *The Entomologist's Monthly Magazine*, **135**: 43–50.
- Archer M.E.** 1999b. Taxonomy and world distribution of the Euro-Asian species of *Dolichovespula* (Hym., Vespinae). *The Entomologist's Monthly Magazine*, **135**: 153–160.
- Archer M.E.** 2006. Taxonomy, distribution and nesting biology of species of the genus *Dolichovespula* (Hymenoptera, Vespidae). *Entomological Science*, **9**(3): 281–293.
- Archer M.E.** 2008. Taxonomy, distribution and nesting biology of species of the genera *Provespa* Ashmead and *Vespa* Linnaeus (Hymenoptera, Vespidae). *The Entomologist's Monthly Magazine*, **144**: 69–101.
- Archer M.E. and Penney D.** 2012. *Vespine wasps of the world: behaviour, ecology & taxonomy of the Vespinae*. Siri Scientific Press, Monograph series, Manchester, UK, 352 p.
- Arens W.** 2010. Revision der *Hedychridium roseum*-Gruppe in Kleinasien (Hymenoptera: Chrysididae), mit Neubewertung zahlreicher europäischer Taxa und Beschreibung zweier neuer Arten. *Linzer Biologische Beiträge*, **42**: 401–458.
- Arens W.** 2014. Die Goldwespen der Peloponnes (Hymenoptera: Chrysididae) 1. Teil: Die Gattungen *Cleptes*, *Omalus*, *Holopyga*, *Hedychrum*, *Hedychridium* und *Euchroeus*; mit Beschreibung einer neuen *Cleptes*-Art. *Linzer Biologische Beiträge*, **46**(1): 553–621.
- Argaman Q.** 1990. Generic synopsis of Sierolomorphidae (Hymenoptera). *Israel Journal of Entomology*, **24**: 29–33.
- Argaman Q.** 1992. Reclassification of Tiphiidae (Hymenoptera, Aculeata) with description of a new subfamily from Turkey. *Türkiye Entomoloji Dergisi*, **16**(1): 3–12.
- Argaman Q.** 1994a. Generic synopsis of Apterogyninae (Hymenoptera: Apterogynidae). *Folia Entomologica Hungarica*, **55**: 41–58.
- Argaman Q.** 1994b. Generic synopsis of Myzinidae (Hymenoptera: Scolioidea). *Annales Musei Historico-Naturalis Hungarici*, **86**: 85–104.
- Argaman Q.** 1996. Generic synopsis of Scoliidae (Hymenoptera, Scolioidea). *Annales Musei Historico-Naturalis Hungarici*, **88**: 171–222.
- Argaman Q.** 2003. Generic synopsis of Mesitinae Kieffer, 1914 (Hymenoptera: Bethylidae). *Entomofauna*, **24**: 61–96.
- Artokhin K.S., Ignatova P.K. and Terskov E.N.** 2012. New insects including invasive species for the fauna of Rostov Region (Russia). *Caucasian Entomological Bulletin*, **8**: 199–202. [In Russian].
- Arutyunyan R.G., Arutyunyan G.A. and Manukyan D.V.** 2009. Biological features of some leaf-miner insects under conditions of Armenia. *Zoocenosis – 2009. Biodiversity and Role of Animals in Ecosystems The International Conference*. DNU, Dnipropetrovsk: 194–195.
- Ascher J.S. and Pickering J.** 2017. Discover Life bee species guide and world checklist (Hymenoptera: Apoidea: Anthophila). http://www.discoverlife.org/mp/20q?guide=Apoidea_species (accessed 18 July 2017).
- Assmuss E.P.** 1859. Symbola ad faunam hymenopterologicam Mosquensem. Enumeratio Hymenopterorum Spheciiformium gubernii Mosquensis. *Bulletin de la Société Impériale des Naturalistes de Moscou. Première Partie*, **32**: 604–620.
- Assmuss E.P.** 1862. Erste Fortsetzung zu meinem "Symbola ad Faunam Hymenopterorum Mosquensem". Enumeratio Hymenopterorum Chrysidiiformium gubernii Mosquensis. *Bulletin de la Société imperiale des naturalistes de Moscou*, **35**: 264–270.
- Astafurova Yu.V.** 2014. *Bees of the subfamilies Rophitinae and Nomiinae (Hymenoptera, Halictidae) of the Russia and adjacent territories*. KMK Scientific Press Ltd., Moscow, 383 p. [In Russian].
- Astafurova Yu.V. and Proshchalykin M.Yu.** 2014. The bees of the genus *Sphecodes* Latreille 1804 of the Russian Far East, with keys to species (Hymenoptera: Apoidea: Halictidae). *Zootaxa*, **3887**(5): 501–528.
- Astafurova Yu.V. and Proshchalykin M.Yu.** 2015. Bees of the genus *Sphecodes* Latreille 1804 of Siberia, with a key to species (Hymenoptera: Apoidea: Halictidae). *Zootaxa*, **4052**(1): 65–95.
- Astafurova Yu.V. and Proshchalykin M.Yu.** 2016. The bees of the genus *Sphecodes* Latreille (Hymenoptera: Halictidae) of the European part of Russia. *Far Eastern Entomologist*, **321**: 1–21.
- Astafurova Yu.V. and Proshchalykin M.Yu.** 2017a. To the knowledge of the *Sphecodes hyalinatus* Hagens species-group (Hymenoptera, Halictidae). *Entomologcheskoe Obozrenie*, **96**(2): 356–365.
- Astafurova Yu.V. and Proshchalykin M.Yu.** 2017b. The genus *Sphecodes* Latreille 1804 (Hymenoptera: Apoidea: Halictidae) in Central Asia. *Zootaxa*, **4324**(2): 249–284.

- Astafurova Yu.V. and Proshchalykin M.Yu. 2017c.** Additional data on the short-tongued bee fauna (Hymenoptera, Apoidea: Andrenidae, Halictidae, Melittidae) of Russia. *Proceedings of the Russian Entomological Society*, **88**(2). [In press].
- Astafurova Yu.V., Proshchalykin M.Yu. and Maharramov M.M. 2017.** Contribution to the knowledge of the *Seladonia* Robertson and *Vestitohalictus* Blüthgen (Hymenoptera: Halictidae: *Halictus* Latreille) of European part of Russia. *Linzer Biologische Beiträge*, **49**(1): 377–386.
- Astafurova Yu.V. and Romankova T.G. 2012.** Additional data on three bee species of the genus *Panurginus* from *P. clavatus* species group (Hymenoptera: Andrenidae: Panurginae). *Zoosystematica Rossica*, **21**(1): 145–157.
- Astapenko S.A., Baranchikov Yu.N. and Palnikova E.N. 2006.** Ecological niche peculiarities in mass species of sawflies (Symphyta: Hymenoptera) at spruce forests of Krasnoyarsk Kray. *Vestnik Krasnoyarskogo gosudarstvennogo universiteta, Seria "Estestvennye nauki"*, **5**: 73–77. [In Russian].
- Astapenko S.A. and Palnikova E.N. 2011.** Factors of spruce sawfly (Symphyta, Hymenoptera) population dynamics in fir man-made forests in south of Krasnoyarsk Region. *Khvoynye boreal'noi zony*, **28**(1–2): 118–125. [In Russian].
- Averenskiy A.I., Bagachanova A.K., Burnasheva A.P., Vinokurov N.N., Ermakova Yu.V., Kajmuk E.L., Novikov D.A., Nogovitsyna S.N., Popov A.A. and Potapova N.K. 2006.** The composition of the arthropod fauna of Lensky District. In: Volpert Ya.L. (Ed.). *Pochvy, rastitel'nyi i zhivotnyi mir Yugo-Zapadnoy Yakutii*. Nauka, Novosibirsk: 103–155. [In Russian].
- Baer I. 1848.** De mutillis nonnullis rossicis. *Bulletin de la Société Impériale des Naturalistes de Moscou*, **21**(1): 228–232, pl. 2, figs 1–8.
- Baghirov R.T.-o. 2010.** New records of digger wasps (Hymenoptera: Sphecidae, Crabronidae) from Western Siberia. *Euroasian Entomological Journal*, **9**: 677–680. [In Russian].
- Baghirov R.T.-o. 2011a.** Fauna of the digger wasps (Hymenoptera: Sphecidae, Crabronidae) of the Altai Territory and Republic of Altai. *Entomologicheskoe Obozrenie*, **90**: 138–158. [In Russian].
- Baghirov R.T.-o. 2011b.** First record of the digger wasp *Chalybion turanicum* (Gussakovskij, 1935) (Hymenoptera: Sphecidae) from Russia. *Far Eastern Entomologist*, **222**: 24.
- Baghirov R.T.-o. 2014.** New data on the spider wasps (Hymenoptera, Pompilidae) from the Western Siberia. *Far Eastern Entomologist*, **279**: 1–10.
- Balthasar V. 1953(1951).** Monographie des chrysidiides de Palestine et des pays limitrophes. *Acta Entomologica Musei Nationalis Prague*, **27**(Suppl. 2): 1–317.
- Balthasar V. 1954.** Zlatěnky – Chryridoidea (Řád: Blanokřidli – Hymenoptera). *Fauna ČSR*, **3**: 1–272.
- Banaszak J. 1980.** Złotolitki Chrysidiidae. *Katalog Fauny Polski*, **26**: 1–52.
- Banaszak J. 1984.** Anthophorini from Mongolia (Hymenoptera: Apoidea, Anthophoridae). *Folia Entomologica Hungarica (S.N.)*, **45**(1): 15–18.
- Banaszak J. and Ortiz-Sánchez F.J. 1993.** Nuevas aportaciones al conocimiento de la tribu Eucerini en el sureste de España (Hymenoptera: Anthophoridae). *Boletín de la Asociación Española de Entomología*, **17**(2): 263–274.
- Banaszak J. and Romasenko L. 2001.** Megachilid bees of Europe (Hymenoptera, Apoidea, Megachilidae). Second edition. Bydgoszcz University of Kazimierz Wielki, Bydgoszcz, 239 p.
- Ban-Calefariu C. 2006.** The systematics and distribution of genus *Nomada* (Hymenoptera: Anthophoridae) in Romania. *Complexul muzeal de Științele naturii "Ion Borcea" Baucău, Studii și comunicări*, **21**: 360 – 368.
- Barovskij V.V. 1929.** To the knowledge of the entomofauna of the Lake district. II. *Russkoe Entomologicheskoe Obozrenie*, **23**: 143–151. [In Russian].
- Becker A. 1865.** Mittheilungen einer botanischen und entomologischen Reise. *Bulletin de la Société Impériale des Naturalistes de Moscou*, **38**(2): 562–582.
- Becker A. 1880.** Beiträge zu meinem Verzeichnissen der um Sarepta und am Bogdo verkommenen Pflanzen und Insekten, und Beschreibung einer Mylabris-Larve. *Bulletin de la Société Impériale des Naturalistes de Moscou*, **55**: 145–156.
- Belke G. 1859.** Esquisse de l'histoire naturelle de Kamienietz-Podolski, précédée d'un coup-d'œil sur les travaux des Naturalistes des provinces occidentales de la Russie et du Royaume de Pologne au XIX siècle. *Bulletin de la Société Impériale des Naturalistes de Moscou*, **32** (1): 24–106.
- Belke G. 1866.** Notice sur l'histoire naturelle du district de Radomysl (Gouvernement de Kief). *Bulletin de la Société Impériale des Naturalistes de Moscou*, **39**: 214–251, 491–526.
- Belo kobylskij S.A. 1990.** Embolemidae is a new to the Soviet Far East fauna family of bethyloids wasp (Hymenoptera: Bethyloidea). In: Lelej A.S. (Ed.). *News of the insect systematic of Soviet Far East*. DVO AN SSSR, Vladivostok: 64–70. [In Russian].
- Belov D.A. 2011.** Features of the complex leaf-mining insects in Moscow. *Forestry Bulletin*, **7**: 103–108. [In Russian].
- Beneš K. 1972a.** A new species of the *histrio*-group of *Pamphilius* Latr. from East Siberia with notes on *P. brevicornis* Hellén (Hymenoptera, Pamphiliidae). *Acta Entomologica Bohemoslovaca*, **69**(1): 46–53.
- Beneš K. 1972b.** Generic classification of the tribe Pamphiliini (Hymenoptera, Pamphiliidae). *Entomologica Bohemoslovaca*, **69**(6): 378–395.

- Beneš K.** 1974. The Siberian species of *Pamphilius* Latr. related to *P. histrio* Latr. (Hymenoptera, Pamphiliidae). *Entomologica Bohemoslovaca*, **71**(5): 298–314.
- Beneš K.** 1976a. Revision of the European species of *Cephalcia* Panzer, 1805 (Hymenoptera, Pamphiliidae). *Studie CSAV*, **3**: 1–67.
- Beneš K.** 1976b. The Siberian species of the genus *Pamphilius* related to *Pamphilius wafer* (L.) (Hymenoptera, Pamphiliidae). *Entomologica Bohemoslovaca*, **73**(3): 159–173.
- Beneš K.** 2014a. Additions to the Czech list of *Euura* Newman, 1837 (Hymenoptera, Tenthredinidae). *Acta Musei Moraviae, Scientiae Biologicae (Brno)*, **99**(1): 69–75.
- Beneš K.** 2014b. Sawflies (Hymenoptera, Symphyta) of the Bohemian Forest and its foothills. *Silva Gabreta*, **20**(3): 131–147.
- Beneš K.** 2014c. A new species of *Pristiphora* (*Lygaeoneurus*) Konow, 1890 on spruce from East Siberia (Hymenoptera, Tenthredinidae). *Beiträge zur Entomologie*, **64**(1): 39–46.
- Beneš K.** 2015. Czech species of the gall-making sawflies of the genera *Phyllocolpa*, *Tubpontania* and *Pontania* (Hymenoptera, Nematinae). *Acta Musei Moraviae, Scientiae biologicae*, **100**(1): 137–156.
- Beneš K. and Holuša J.** 2015. Sawflies (Hymenoptera: Symphyta) in the northeast of the Czech Republic with special regard to spruce forests. *Journal of Forest Science*, **61**(3): 112–130.
- Benson R.B.** 1962. A revision of the Athaliini (Hymenoptera: Symphyta). *Bulletin of the British Museum (Natural History). Entomology series*, **11**: 333–382.
- Benson R.B.** 1965a. Some sawflies new to the Himalayan region (Hymenoptera, Symphyta). *The Annals & Magazine of Natural History (incorporating "The Journal of Botany")*, *Zoology, Botany, and Geology; 13th Series*, **8**: 141–144.
- Benson R.B.** 1965b. The classification of *Rhogogaster* Konow (Hymenoptera: Tenthredinidae). *Proceedings of the Royal Entomological Society of London. Series B: Taxonomy*, **34**: 105–112.
- Bequaert J.** 1930. On the generic and subgeneric divisions of the Vespinae (Hymenoptera). *Bulletin of the Brooklyn Entomological Society*, **25**: 59–70.
- Białyńicki-Birula A.** 1926. Ein Beitrag zur Kenntnis der paläarktischen *Hoplomerus*-Arten (Hymenoptera – Diptera). *Bulletin de l'Académie des Sciences de l'URSS*, **20**(10–11): 885–906.
- Birula A.** 1925. Sur la distribution géographique du frelon commun (*Vespa crabro* L.) et sur ses races climatiques. *Comptes Rendus de l'Académie des Sciences de Russie*, **1925**: 53–56.
- Birula A.** 1927. Über die russischen Wespen und ihre geographische Verbreitung. (Zweiter Beitrag). *Annuaire du Musée Zoologique de l'Academie des Sciences de l'URSS*, **28**(1): 72–82.
- Birula A.** 1930a. Über die russischen Wespen und ihre geographische Verbreitung. (Dritter Beitrag). *Annuaire du Musée Zoologique de l'Academie des Sciences de l'URSS*, **31**: 291–339.
- Birula A.** 1930b. Über die russischen Wespen und ihre geographische Verbreitung (Vierter Beitrag). *Zoologischer Anzeiger*, **87**: 127–143.
- Biryukova O.B. and Novgorodova T.A.** 2008. Trophobiotic interactions between representatives of the order Hymenoptera: ants (Formicidae) and sawfly larvae (Blasticotomidae). *Eurasian Entomological Journal*, **7**(3): 227–233. [In Russian].
- Bischoff H.** 1910. Die Chrysididen des Königlichen Zoologischen Museums zu Berlin. *Mitteilungen aus dem Zoologischen Museum in Berlin*, **4**: 426–493.
- Bischoff H.** 1913. Hymenoptera. Fam. Chrysidae. In: Wytsman P. (Ed.). *Genera insectorum. Fascicule 151*. L. Desmet-Verteneuil, Bruxelles, 86 p. + 5 pls.
- Bischoff H.** 1930. Beitrag zur Kenntnis paläarktischer Arterender Gattung *Epeolus*. (Hym. Apid.). *Deutsche Entomologische Zeitschrift*, **1**: 1–15.
- Bischoff H. and Bradley J.C.** 1929. The Scoliidae described by Pallas (Hym.). *Deutsche Entomologische Zeitschrift*, **5**: 301–302.
- Bitsch J., Dollfuss H., Bouček Z., Schmidt K., Schmid-Egger Ch., Gayubo S.F., Antropov A.V. and Barbier Y.** 2001. *Faune de France. France et régions limitrophes. 86. Hyménoptères Sphecidae d'Europe occidentale. Volume 3*. Fédération Française des Sociétés de Sciences Naturelles, Paris, 459 p.
- Bitsch J. and Leclercq J.** 1993. *Faune de France. France et régions limitrophes. 79. Hyménoptères Sphecidae d'Europe occidentale. Volume 1. Généralités - Crabroninae*. Fédération Française des Sociétés de Sciences Naturelles, Paris, 325 p.
- Blagoveshchenskaya N.N.** 1970. Results of the study of solitary bees in the Middle Volga basin. In: Popov V.A. (Ed.). *The resulting scientific conference of zoologists of the Volgo-Kama region. Proceedings*. Ministry of the Higher and Middle Special Education of the USSR, Kazan: 212–217. [In Russian].
- Blagoveshchenskaya N.N.** 1994. Catalog of aculeate Hymenoptera of Ul'yanovsk Oblast'. In: Shustov M.V. (Ed.). *Priroda Ul'yanovskoy Oblasti. Vypusk 5. Nasekomye Ul'yanovskoy Oblasti*. Rossiyskaya Akademiya Nauk, Institut Ekologii Volzhskogo Basseina, Ul'yanovskiy Filial MGU im. M.V. Lomonosova, Ul'yanovskiy Oblastnoy Komitet po Okhrane Prirody, Ul'yanovsk: 82–93. [In Russian].
- Blagoveshchenskaya N.N. and Popova L.M.** 1994. A catalogue of the fauna of bees (Hymenoptera, Apoidea) in the Ulyanovsk Province. In: Isaev A.Yu. (Ed.). *Nature of the Ulyanovsk Province. No. 5. Insects of the Ulyanovsk Province*. Ulyanovsk Branch of the Moscow State University, Ulyanovsk: 94–116. [In Russian].

- Blank S.M. 1993.** Eine neue Blattwespenart für die Fauna Österreichs: *Dolerus vernalis* Ermolenko, 1964 (Insecta: Hymenoptera, Tenthredinidae). *Berichte des Naturwissenschaftlich-Medizinischen Vereins in Innsbruck*, **80**: 363–371.
- Blank S.M. 1996.** Revision of the sawflies described by Lothar Zirngiebl (Preliminary studies for a Catalogue of Symphyta, part 2) (Insecta, Hymenoptera, Symphyta). *Spixiana*, **19**(2): 195–219.
- Blank S.M. 2002a.** Taxonomic notes on strongylogasterini (Hymenoptera: Tenthredinidae). *Proceedings of the Entomological Society of Washington*, **104**: 692–701.
- Blank S.M. 2002b.** The Western Palaearctic Xyelidae (Hymenoptera). In: Viitasaari M. (Ed.). *Sawflies (Hymenoptera, Symphyta) I. A review of the suborder, the Western Palaearctic taxa of Xyeloidea and Pamphilioidea*. Tremex, Helsinki, **1**: 197–233.
- Blank S.M. 2011.** Case 3538. Coryninae Benson, 1938 (Insecta, Hymenoptera, Cimbicidae): proposed emendation of spelling to Corynidinae to remove homonymy with Corynidae Johnston, 1836 (Cnidaria, Anthoathecata). *Bulletin of Zoological Nomenclature*, **68**(2): 113–116.
- Blank S.M., Gninenko Y.I. and Telegina O.S. 2011.** New invasive sawfly *Arge captiva* in Kazakhstan. *Zertzeushi – Issledovatel*, **1**–**2**(57–58): 131–134. [In Russian].
- Blank S.M., Köhler T., Pfannenstill T., Neuenfeldt N., Zimmer B., Jansen E., Taeger A. and Liston A.D. 2014.** Zig-zagging across Central Europe: recent range extension, dispersal speed and larval hosts of *Aproceros leucopoda* (Hymenoptera, Argidae) in Germany. *Journal of Hymenoptera Research*, **41**: 57–74.
- Blank S.M., Kramp K., Smith D.R., Sundukov Yu.N., Wei M. and Shinohara A. 2017.** Big and beautiful: The *Megaxyela* species (Hymenoptera, Xyelidae) of East Asia and North America. *European Journal of Taxonomy*, **348**: 1–46
- Blank S.M., Kraus M. and Taeger A. 2006.** *Orussus smithi* sp. n. and notes on other West Palaearctic Orussidae (Hymenoptera). In: Blank S.M., Schmidt S., Taeger A. (Eds). *Recent Sawfly Research – Synthesis and Prospects*. Goecke & Evers, Keltern: 265–278.
- Blank S.M. and Ritzau C. 1998.** Die Tenthredopsini Deutschlands (Hymenoptera: Tenthredinidae). In: Taeger A., Blank S.M. (Eds). *Pflanzenwespen Deutschlands (Hymenoptera, Symphyta)*. Goecke & Evers, Keltern: 227–246.
- Blank S.M., Shinohara A. and Altenhofer E. 2013.** The Eurasian species of *Xyela* (Hymenoptera: Xyelidae): taxonomy, host plants and distribution. *Zootaxa*, **3629**(1): 1–106.
- Blank S.M., Shinohara A. and Byun B.-K. 2005.** The East Asian *Xyela* species (Hymenoptera: Xyelidae) associated with Japanese Red Pine (*Pinus densiflora*; Pinaceae) and their distribution history. *Insect Systematics & Evolution*, **36**: 259–278.
- Blank S.M. and Taeger A. 1998a.** Comments on the taxonomy of Symphyta (Hymenoptera) (Preliminary studies for a catalogue of Symphyta, part 4). In: Taeger A., Blank S.M. (Eds). *Pflanzenwespen Deutschlands (Hymenoptera, Symphyta). Kommentierte Bestandsaufnahme*. Goecke & Evers, Keltern: 141–174.
- Blank S.M. and Taeger A. 1998b.** Revision of the sawfly genera *Asiarge* Gussakovskij, 1935 and *Kokujewia* Konow, 1902 (Hymenoptera: Argidae). *Beiträge zur Entomologie*, **48**: 505–515.
- Blank S.M. and Taeger A. 2006.** Taxonomy and evolution of *Tenthredo* (*Elinora*) species similar to *T. dahlii* and *T. koehleri* (Hymenoptera: Tenthredinidae). In: Blank S.M., Schmidt S., Taeger A. (Eds). *Recent Sawfly Research – Synthesis and Prospects*. Goecke & Evers, Keltern: 199–227.
- Blank S.M. and Taeger A. 2011.** *Trachelus* Jurine, 1807 (Hymenoptera, Cephidae): proposed precedence over *Astatas* Panzer, [1801]. *The Bulletin of Zoological Nomenclature*, **68**(1): 65–70.
- Blank S.M., Taeger A., Liston A.D., Smith D.R., Rasnitsyn A.P., Shinohara A., Heidemaa M. and Viitasaari M. 2009.** Studies toward a world catalog of Symphyta (Hymenoptera). *Zootaxa*, **2254**: 1–96.
- Blüthgen P. 1923.** Beiträge zur Kenntnis der Bienengattung *Halictus* Latr. *Archiv Naturgeschihte* (A), **89**(5): 232–332.
- Blüthgen P. 1924.** Notas sobre “*Halictus*” paleárticos (Hymenopt. Apidae). *Boletín de la Real Sociedad Española de Historia Natural*, **24**(5): 262–266.
- Blüthgen P. 1925.** Beiträge zur Kenntnis der Bienengattung *Halictus* Latr. II. *Archiv Naturgeschihte* (A), (1924), **90**(10): 86–136.
- Blüthgen P. 1931.** Beiträge zur Kenntnis der Bienengattung *Halictus* Latr. III. *Mitteilungen aus dem Zoologischen Museum in Berlin*, **17**(3): 319–398.
- Blüthgen P. 1935.** Neue paläarktische *Halictus*-Arten (Hym., Apidae). [II]. *Deutsche Entomologische Zeitschrift*, **1935**(1/2): 111–120.
- Blüthgen P. 1955.** The Halictinae (Hymen., Apoidea) of Israel. I. Genus *Halictus* (subgenera *Halictus* s. str. and *Thrincohalictus*). *Bulletin of the Research Council of Israel* (B), **5**(1): 5–23.
- Blüthgen P. 1961.** Die Faltenwespen Mitteleuropas (Hymenoptera, Dipteroptera). *Abhandlungen der Deutschen Akademie der Wissenschaften zu Berlin. Klasse für Chemie, Geologie und Biologie*, **1961**(2): 1–252.
- Bogacheva I.A. 1977.** Sawflies (Hymenoptera, Tenthredinoidea) and their role in ecosystems of the northern Ob basin. In: Danilov N.N. (Ed.). *The biocoenotical significance of animals in the forest tundra of Yamal. Trudy Instituta Ekologii Rastenii i Zhivotnykh, Sverdlovsk*, **106**: 85–103. [In Russian].
- Bogusch P. 2006.** The velvet ants (Hymenoptera: Mutilidae) of the Czech Republic and Slovakia: an identification key and annotated checklist. *Acta Musei Moraviae, Scientiae biologicae*, **91**: 103–148.

- Bogusch P. and Straka J.** 2012. Review and identification of the cuckoo bees of Central Europe (Hymenoptera: Halictidae: Sphecodes). *Zootaxa*, **3311**: 1–41.
- Bohart R.M.** 2000. A review of Gorytini in the Neotropical Region (Hymenoptera: Sphecidae: Bembicinae). *Contributions on Entomology, International*, **4**: 111–259.
- Bohart R.M. and Menke A.S.** 1976. *Sphecid Wasps of the World. A generic revision*. University of California Press, Berkeley, Los Angeles, London, ix + 695 p.
- Bolton B.** 2016. *An online catalog of the ants of the world*. Available from <http://antcat.org> (accessed 28 February 2017).
- Boni Bartalucci M.** 2010. A review of the genus *Pseudotiphia* Ashmead 1903 (Hymenoptera, Tiphidae). *Linzer Biologische Beiträge*, **42**(2): 1183–1236.
- Boni Bartalucci M.** 2016. Meriini from Western Palaearctic and northern Afrotropical Regions (Hymenoptera: Tiphidae: Myzininae): new taxa and records. *Onychium*, **12**: 47–81.
- Borowiec L.** 2014. Catalogue of ants of Europe, the Mediterranean basin and adjacent regions (Hymenoptera: Formicidae). *Genus*, **25**(1–2): 1–340.
- Borowski J. and Marczak D.** 2015. *Gilpinia fennica* (Forsius, 1911) new to Poland (Hymenoptera, Symphyta: Diprionidae), and a review of the species. *Entomologica Fennica*, **26**: 177–180.
- Bouček Z.** 2001. Palaearctic species of *Ammoplanus* (Hymenoptera: Sphecidae). *Journal of Natural History*, **35**: 849–929.
- Bradley J.C.** 1972. Scoliid types in the Museum für Naturkunde of the Humboldt-Universität zu Berlin. *Mitteilungen aus dem Zoologischen Museum in Berlin*, **48**(1): 3–19.
- Brischke C.G.A.** 1862. Die Hymenopteren der Provinz Preussen (Erste Fortsetzung). *Schriften der Königlichen Physikalisch-Ökonomischen Gesellschaft zu Königsberg*, **2**: 97–118.
- Brischke C.G.A.** 1883. Beobachtungen über die Arten der Blatt- und Holzwespen von C.G.A. Brischke, Hauptlehrer a. D. in Langfuhr und Dr. Gustav Zaddach weiland Professor in Königsberg. Zweite Abtheilung. *Schriften der Naturforschenden Gesellschaft in Danzig, N. S.*, **5**(4): 201–328.
- Brischke C.G.A.** 1884. Beobachtungen über die Arten der Blatt- und Holzwespen von C.G.A. Brischke, Hauptlehrer a. D. in Langfuhr und Dr. Gustav Zaddach, Professor in Königsberg, mitgetheilt von Brischke aus Zaddach's Manuscripten. (Schluss). *Schriften der physikalisch-ökonomischen Gesellschaft zu Königsberg*, **24**: 121–173.
- Brooks R.W.** 1988. Systematics and phylogeny of the Anthophorine bees (Hymenoptera Anthophoridae; Anthophorini). *The University of Kansas Science Bulletin*, **53**(9): 436–575.
- Brothers D.J.** 1975. Phylogeny and classification of the aculeate Hymenoptera, with special reference to Mutilidae. *The University of Kansas Science Bulletin*, **50**(11): 483–648.
- Brothers D.J.** 1989. Alternative life-history styles of mutiliid wasps (Insecta, Hymenoptera). In: Bruton M.N. (Ed.). *Alternative life-history styles of animals*. Kluwer Academic Publishers, Dordrecht: 279–291.
- Brothers D.J.** 1999. Phylogeny and evolution of wasps, ants and bees (Hymenoptera, Chrysidoidea, Vespoidea and Apoidea). *Zoologica Scripta*, **28**(1–2): 233–249.
- Brothers D.J. and Carpenter J.M.** 1993. Phylogeny of Aculeata: Chrysidoidea and Vespoidea. *Journal of Hymenoptera Research*, **2**: 227–302.
- Brothers D.J. and Finnimore A.T.** 1993. Superfamily Vespoidea. In: Goulet H., Huber J.T. (Eds). *Hymenoptera of the World. An identification Guide to Families*. Agriculture Canada, Ottawa: 161–278.
- Brothers D.J., Tschuch G. and Burger F.** 2000. Associations of mutiliid wasps (Hymenoptera, Mutilidae) with eusocial insects. *Insectes Sociaux*, **47**: 201–211.
- Budrys E.R.** 1985. New and little known species of the genus *Mimesa* Shuck. (Hymenoptera, Sphecidae) of Middle Asia and Kazakhstan and a key to species of the USSR fauna. *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, **142**: 58–76. [In Russian].
- Budrys E.R.** 1986. Digger wasps of the genus *Psen* (Hymenoptera, Sphecidae) of the Far East of USSR. In: Lehr P.A., Belokobylskij S.A., Storoževa N.A. (Eds). *Hymenopterous Insects of Eastern Siberia and Far East*. BPI DVNTS AN SSSR, Vladivostok: 138–147. [In Russian].
- Budrys E.R.** 1987. Digger wasps of the genus *Stigmus* Panzer and *Carinostigmus* Tsuneki (Hymenoptera, Sphecidae) of the Far East of USSR. In: Lehr P.A., Storoževa N.A. (Eds). *Novye dannye po sistematike nasekomykh Dalnego Vostoka*. BPI DO AN SSSR, Vladivostok: 49–56. [In Russian].
- Budrys E.R.** 1990. Digger-wasps of the genus *Mimumesa* Malloch of the fauna of the USSR. *Entomologicheskoe Obozrenie*, **69**(4): 941–947. [In Russian].
- Buysson R. du 1891–1896.** Les Chrysides. In: André E. (Ed.). *Species des Hyménoptères d'Europe & d'Algérie. Tome Sixième*. Vve Dubosclard, Paris, i–xii + 13–758 + 64 unnumbered pages + pls. 1–32 (1891) 1–88, (1892) 89–208, (1893) 209–272, (1894) 273–400, (1895) 401–624, (1896) 625–756 + 1*–22*, (1891–1896) 64 unnumbered pages + pls. 1–32.
- Buyanjargal B., Dorzhiev Ts.Z., Abashev R.Yu. and Bataa D.** 2016. Geographical range of vespid wasps (Hymenoptera, Vespidae) of northern Mongolia. *Mongolian Journal of Biological Sciences*, **14**(1–2): 21–31.
- Bytinski-Salz H. and Ebmer A.W.** 1974. The Halictidae of Israel (Hymenoptera, Apoidea). II. Genus *Lasioglossum*. *Israel Journal of Entomology*, **9**: 175–217.

- Byvaltsev A.M.** 2008. Bumblebee (Hymenoptera: Apidae, Bombini) fauna of the forest-steppe and steppe zones of the West Siberian Plain. *Euroasian Entomological Journal*, 7(2): 141–147. [In Russian].
- Byvaltsev A.M.** 2013. The communities of bumblebees (Hymenoptera, Apidae, Bombini) of the Kulunda Plain of West Siberia, Russia. *Euroasian Entomological Journal*, 12(6): 575–586. [In Russian].
- Byvaltsev A.M., Belova K.A., Kupianskaya A.N. and Proshchalykin M.Yu.** 2015. The diversity and abundance of bumblebees (Hymenoptera: Apidae, *Bombus*) in the steppes of Khakassia. *A.I. Kurentsov's Annual Memorial Meetings*, 26: 264–276. [In Russian].
- Byvaltsev A.M., Belova K.A. and Proskuryakova A.A.** 2013. Populations and ecology of bumblebees (Hymenoptera: Apidae, *Bombus*) of the Irtysh steppe in early summer. *Vestnik Novosibirskogo Gosudarstvennogo Universiteta. Seriya: Biologiya, Klinicheskaya meditsina*, 11(1): 40–46. [In Russian].
- Byvaltsev A.M., Proshchalykin M.Yu., Levchenko T.V., Kupianskaya A.N. and Akulov E.N.** 2016. Bumble bee fauna (Hymenoptera, Apidae: *Bombus* Latreille) of Krasnoyarsk Territory. *A.I. Kurentsov's Annual Memorial Meetings*, 27: 137–154. [In Russian].
- Carpenter J.M. and Kojima J.** 2002. Correction to the nomenclature in the genus *Dolichovespula* (Hymenoptera: Vespidae). *Entomological Science*, 5(3): 389–390.
- Carpenter J.M.** 1986. A synonymic generic checklist of the Eumeninae (Hymenoptera: Vespidae). *Psyche*, 93(1–2): 61–90.
- Carpenter J.M.** 1996. Distribution checklist of species of the genus *Polistes* (Hymenoptera: Vespidae: Polistinae: Polistini). *American Museum Novitates*, 3188: 1–36.
- Carpenter J.M.** 2001. Checklist of species of the subfamily Masariniae (Hymenoptera: Vespidae). *American Museum Novitates*, 3325: 1–39.
- Carpenter J.M., Dvořák L. and Pickett K.M.** 2011. *Dolichovespula albida* (Sladen), a valid species, not a synonym of *D. norwegica* (Fabricius). *Entomologica Americana*, 117: 113–116.
- Carpenter J.M. and Glare T.R.** 2010. Misidentification of *Vespula alascensis* as *V. vulgaris* in North America (Hymenoptera: Vespidae; Vespinae). *American Museum Novitates*, 3690: 1–7.
- Carpenter J.M. and Kojima J.** 1996. Checklist of the subfamily Stenogastrinae (Hymenoptera: Vespidae). *Journal of the New York Entomological Society*, 104: 21–36.
- Carpenter J.M. and Kojima J.** 1997. Checklist of the species in the subfamily Vespinae (Insecta: Hymenoptera: Vespidae). *Natural History Bulletin of Ibaraki University*, 1997(1): 51–92.
- Carpenter J.M. and Nguyen L.P.T.** 2003. Keys to the genera of social wasps of South-East Asia (Hymenoptera: Vespidae). *Entomological Science*, 6: 183–192.
- Castro L.** 2003. Notas sobre véspidos solitarios (Hymenoptera: Vespoidea: Eumeninae). *Boletín de la Sociedad Entomológica Aragonesa*, 32: 181–183.
- Castro L. and Dvořák L.** 2009. New and noteworthy records of vespid wasps (Hymenoptera: Vespidae) from the Palaearctic region (II). *Boletín de la Sociedad Entomológica Aragonesa*, 44: 295–304.
- Castro L. and Dvořák L.** 2010. New and noteworthy records of vespid wasps (Hymenoptera: Vespidae) from the Palaearctic region (III). *Acta Musei Moraviae, Scientiae Biologicae (Brno)*, 95(2): 37–53.
- Castro L., Sanza F., Aguirre-Segura A. and Ortiz-Sánchez F.J.** 2008. Vespidae y Scoliidae (Hymenoptera) del Parque Natural Cabo de Gata-Níjar (Almería, España): lista preliminar. *Boletín de la Sociedad Entomológica Aragonesa*, 43: 319–324.
- Cavro E.** 1950. Catalogue des Hyménoptères de département du Nord et régions limitrophes. *Bulletin de la Société entomologique du Nord de la France*, 52 (Suppl. 30): 1–84.
- Celary W.** 1995. Nomadini (Hymenoptera, Apoidea, Anthophoridae) of Poland. *Monografie Fauny Polski*, 20: 1–281.
- Celary W.** 2005. *Melittidae of Poland (Hymenoptera: Apoidea: Anthophila): Their Biodiversity and Biology*. Polish Academia of Sciences, Kraków, 175 p.
- Cervo R.** 2006. *Polistes* wasps and their social parasites: an overview. *Annales Entomologici Fennici*, 43: 531–549.
- Ćetković A., Mokrousov M.V., Plečaš M., Bogush P., Antic D., Đorović-Jovanović L., Krpo-Ćetković J. and Karaman M.** 2011. Status of the potentially invasive Asian species *Sceliphron deforme* in Europe, and an update on the distribution of *S. curvatum* (Hymenoptera: Sphecidae). *Acta Entomologica Serbica*, 16: 91–114.
- Chen C.-W.** 1957. A revision of the velvety ants or Mutilidae of China. *Quarterly Journal of the Taiwan Museum*, 10(3–4): 135–224.
- Chen S., Hoelmer K.A., Chen H., Liu A. and Shanower T.G.** 2004. A review of wheat stem sawfly (Hymenoptera: Cephidae) research in China. *Journal of Agricultural and Urban Entomology*, 21(4): 249–256.
- Chenikalova E.V.** 2005. Wild bees of the Stavropol Kray, their efficiency and reservation in agrolandscapes. Argus, Stavropol, 112 p. [In Russian].
- Chinin A.A.** 1991. Contribution to the sphecid fauna of Samara Oblast'. *Samarskaya Luka*, 1: 110–112. [In Russian].
- Choi J.-K., Lee S.-B., Wei M. and Lee J.-W.** 2015. Study on two genera, *Yasumatsua* Togashi and *Aproceros* Malaise (Hymenoptera: Argidae: Sterictiphorinae), from South Korea with description of a new species. *Entomological Research*, 45(6): 323–327. [In Russian].
- Choi J.-K., Lee S.-B., Wei M. and Lee J.-W.** 2016. Three new species of the genus *Arge* (Hymenoptera: Symphyta: Argidae) from South Korea with key to species of the subfamily Arginae. *Journal of Asia-Pacific Biodiversity*, 9: 183–193.

- Choi J.-K., Wei M., Vilhelmsen L. and Lee J.-W. 2014.** A new *Orussus* species from South Korea, and a key to the East Asian Orussidae (Hymenoptera). *Zootaxa*, **3873**(3): 250–258.
- Clerc V. and Clerc M. 1907.** Materials for the Entomology of the Urals. List of insects collected in the district of Ekaterinburg by V. and M. Clerc in 1896–1897, and determined by Dr. Frey-Gessner, assistant to the Museum of Natural History of Geneva. *Zapiski Sostoyashchego pod Avgusteyshim Pokrovitel'stvo Ego Imperatorskogo Vysochestva Velikago Knyazya Mikhaila Nikolaevicha Ural'skago Obshchestva Lyubiteley Estestvoznaniya*, **26**: 75–80. [In Russian].
- Conde O. 1935.** Oryssodea et Tenthredinoidea collecta in Ussuri et Sachalin ab N. Delle. *Notulae Entomologicae*, **14**: 67–87.
- Csösz S., Heinze J. and Mikó I. 2015.** Taxonomic Synopsis of the Ponto-Mediterranean ants of *Temnothorax nylanderi* species-group. *PLoS ONE*, **10**(11): 1–62.
- Csösz S., Radchenko A. and Schulz A. 2007.** Taxonomic revision of the Palaearctic *Tetramorium chefketi* species-complex (Hymenoptera: Formicidae). *Zootaxa*, **1405**: 1–38.
- Csösz S. and Schulz A. 2010.** Taxonomic review of the Palaearctic *Tetramorium ferox* species-complex (Hymenoptera, Formicidae). *Zootaxa*, **2401**: 1–29.
- Cumming J.M. 1989.** Classification and evolution of the eumenine wasp genus *Symmorphus* Wesmael (Hymenoptera: Vespidae). *Memoirs of the Entomological Society of Canada*, **148**: 5–168.
- Czechowski W., Radchenko A. and Czechowska W. 2002.** *The ants (Hymenoptera, Formicidae) of Poland*. Museum and Institute of Zoology, Polish Academy of Sciences, Warsaw, 200 p.
- Czechowski W., Radchenko A., Czechowska W. and Vejsäläinen K. 2012.** *The ants of Poland with reference to the myrmecofauna of Europe*. Fauna Poloniae, Vol. 4. Natura Optima Dux Foundation, Warsaw, 496 p.
- Dalla Torre C.G. de. 1892.** *Catalogus hymenopterorum 6. Chrysidae (Tubulifera)*. Wilhelm Engelmann, Lipsiae, ix + 118 p.
- Dalla Torre C.G. de. 1894.** *Catalogus Hymenopterorum hucusque descriptorum systematicus et synonymicus. Vol. 1: Tenthredinidae incl. Uroceridae (Phyllophaga et Xylophaga)*. Sumptibus Guilelmi Engelmann, Lipsiae. [6] + VIII + 459 p.
- Dalla Torre C.G. de. 1897.** *Catalogus Hymenopterorum hucusque descriptorum systematicus et synonymicus. Volumen VIII. Fossores (Shegidae)*. Guilelmi Engelmann, Lipsiae, i–viii + 1–749.
- Daly H.V. 1983.** Taxonomy and ecology of Ceratinini of North Africa and the Iberian Peninsula (Hymenoptera: Apoidea). *Systematic Entomology*, **8**(1): 29–62.
- Danilov Yu.N. 2006.** Preliminary data on fauna of wasps (Hymenoptera, Vespoformes) and bees (Hymenoptera, Apoidea) from the environs of Barnaul (Altai Territory). In: *Entomological investigations in North Asia. Materials of VII interregional conference of entomologists of Siberia and Far East in the framework of the Siberian zoological conference*. Institute of Systematic and Ecology of Animals Siberian Branch of RAS, Novosibirsk: 55–56. [In Russian].
- Danilov Yu.N. 2008.** The digger wasps (Hymenoptera: Sphecidae, Crabronidae) of the Altai forest-steppe near Barnaul. *Euroasian Entomological Journal*, **7**: 345–352. [In Russian].
- Danilov Yu.N. 2009.** The fauna of digger wasps (Hymenoptera: Sphecidae) of the Kulundinskaya Steppe and adjacent territories. *A.I. Kurentsov's Annual Memorial Meetings*, **20**: 52–62. [In Russian].
- Danilov Yu.N. 2011a.** Digger wasps of the family Sphecidae (Hymenoptera: Apoidea) of Daurskii Reserve. *A.I. Kurentsov's Annual Memorial Meetings*, **22**: 199–206. [In Russian].
- Danilov Yu.N. 2011b.** New records of digger wasps of the family Sphecidae (Hymenoptera, Apoidea) from the Asian part of Russia. *Euroasian Entomological Journal*, **10**: 188–190. [In Russian].
- Danilov Yu.N. 2012a.** To the knowledge of the digger wasps of the tribe Prionychini Bohart and Menke, 1963 (Hymenoptera, Sphecidae) of Russia and adjacent countries. *Proceedings of the Russian Entomological Society*, **83**: 159–165. [In Russian].
- Danilov Yu.N. 2012b.** The digger wasps of the genus *Prionyx* Vander Linden (Hymenoptera: Sphecidae) of Russia and Central Asia, with a key to species, new synonymies, and lectotype designations. *Zootaxa*, **3526**: 59–71.
- Danilov Yu.N. 2013.** New records of the digger wasp *Palmodes orientalis* (Mocsáry) (Apoidea, Sphecidae) in Eurasia. *Euroasian Entomological Journal*, **12**: 46–50. [In Russian].
- Danilov Yu.N. 2014a.** Review of Sphecidae wasps (Hymenoptera: Apoidea) of Siberia. Part I. List of species. *Euroasian Entomological Journal*, **13**: 422–429. [In Russian].
- Danilov Yu.N. 2014b.** Review of Sphecidae wasps (Hymenoptera: Apoidea) of Siberia. Part 2. Key to genera and species. *Euroasian Entomological Journal*, **13**: 511–521. [In Russian].
- Danilov Yu.N. 2015.** New and little known digger wasps of the genus *Ammophila* W. Kirby, 1798 (Hymenoptera: Apoidea: Sphecidae) from Central Asia. *Zootaxa*, **3964**: 537–545.
- Danilov Yu.N. 2017.** Taxonomic notes on Palearctic *Podalonia* Fernald, 1927 (Hymenoptera, Sphecidae), with descriptions of four new species. *Zootaxa*, **4320**(3): 554–570.

- Danilov Yu.N. and Byvaltsev A.M. 2014.** Carpenter bee *Xylocopa valga* Gerstäcker, 1872. In: Savchenko A.P. (Ed.). *Red book of Khakasia Republic*, 2 ed. SFU, Krasnoyarsk-Abakan: 31. [In Russian].
- Danilov Yu.N. and Mokrousov M.V. 2017a.** New data on distribution and taxonomy of some Palaearctic species of Sphecidae (Hymenoptera: Apoidea). *Euroasian Entomological Journal*, **16**(2): 107–113.
- Danilov Yu.N. and Mokrousov M.V. 2017b.** New records of sphecid digger wasps (Hymenoptera: Apoidea: Sphecidae) in Russia. *Proceedings of the Russian Entomological Society*, **88**(2). [In press].
- Danilov Yu.N. and Odintseva A.A. 2013.** To our knowledge of digger wasp fauna (Hymenoptera: Sphecidae, Crabronidae) of south taiga of Irtysch River, West Siberia, Russia. *Euroasian Entomological Journal*, **12**: 331–335. [In Russian].
- Danilov Yu.N. and Tshernyshev S.E. 2008.** Digger wasps (Hymenoptera: Sphecidae, Crabronidae) in the Kulundinskaya forest-steppe of West Siberia. *Euroasian Entomological Journal*, **7**: 40–46. [In Russian].
- Dathe H.H. 1980a.** Die arten der gattung *Hylaeus* F. in Europa (Hymenoptera: Apoidea, Colletidae). *Mitteilungen aus dem Zoologischen Museum in Berlin*, **56**: 207–294.
- Dathe H.H. 1980b.** Zur Hymenopterenfauna des Naturschutzgebietes Teberda im Westkaukasus. *Milu*, **5**(1/2): 194–217.
- Dathe H.H. 2009.** Order Hymenoptera, superfamily Apoidea. In: van Harten A. (Ed.). *Arthropod fauna of the UAE*. Vol. 1. Dar Al Ummah Printing, Abu Dhabi: 335–432.
- Dathe H.H. and Proshchalykin M.Yu. 2016.** The genus *Hylaeus* Fabricius in Mongolia, an updated species inventory (Hymenoptera: Apoidea: Colletidae). *Zootaxa*, **4121**(4): 351–382.
- Dathe H.H. and Proshchalykin M.Yu. 2017.** Type revision of Asiatic bees of the genus *Hylaeus* F. described by Ferdinand Morawitz (Hymenoptera: Apoidea, Colletidae). *Zootaxa*, **4227**(1): 1–48.
- Davydova N.G. and Pesenko Yu.A. 2002.** Bee fauna (Hymenoptera, Apoidea) of Yakutia. I. *Entomologicheskoe Obozrenie*, **81**(3): 382–399. [In Russian].
- Day M.C. 1979.** The species of Hymenoptera described by Linnaeus in the genera *Sphex*, *Chrysis*, *Vespa*, *Apis* and *Mutilla*. *Biological Journal of the Linnean Society*, **12**: 45–84.
- Day M.C. 1981.** A revision of *Pompilus* Fabricius (Hymenoptera: Pompilidae), with further nomenclatural and biological considerations. *Bulletin of the British Museum (Natural History), Entomology*, **42**(1): 1–42.
- Day M.C. 1988.** *Spider wasps Hymenoptera: Pompilidae*. Royal Entomological Society of London, Dorchester, 60 p.
- de Andrade N.F. 1954.** Palaearctic *Miscophus* of the *gallicus* group (Hymenoptera, Sphecidae). *Memórias e Estudos do Museu Zoológico da Universidade de Coimbra*, **226**: 1–87.
- de Andrade N.F. 1956a.** Note of the palaearctic *Miscophus* of the *soikai* group (Hymenoptera, Sphecidae). *Memórias e Estudos do Museu Zoológico da Universidade de Coimbra*, **238**: 1–2.
- de Andrade N.F. 1956b.** Western palaearctic *Miscophus* of the *handlirschi* group (Hymenoptera, Sphecidae). *Memórias e Estudos do Museu Zoológico da Universidade de Coimbra*, **239**: 1–40.
- de Andrade N.F. 1960.** Palaearctic *Miscophus: bicolor* group and isolated species (Hymenoptera, Sphecidae). *Memórias e Estudos do Museu Zoológico da Universidade de Coimbra*, **262**: 1–136.
- de Beaumont J. 1944.** Les guêpes (*Vespa* L. s.l.) de la Suisse. *Bulletin de la Société Vaudoise des Sciences Naturelles*, **57**(261): 329–362.
- de Beaumont J. 1947.** Contribution à l'étude du genre *Tachysphex* (Hym. Sphecid.). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, **20**: 661–677.
- de Beaumont J. 1949.** Contribution à l'étude du genre *Palarus* Latr. (Hym. Sphecid.). *Revue Suisse de Zoologie*, **56**: 627–673.
- de Beaumont J. 1952.** Les *Hoplisoides* et les *Psammaecius* de la région paléarctique (Hym. Sphecid.). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, **25**: 211–238.
- de Beaumont J. 1953a.** Le genre *Olgia* Radoszk. (Hym. Sphecid.). *Revue Suisse de Zoologie*, **60**: 205–223.
- de Beaumont J. 1953b.** Les *Gorytes* s.s. (= *Hoplisus*) de la région paléarctique (Hym. Sphecid.). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, **26**: 161–200.
- de Beaumont J. 1954a.** Les *Bembecinus* de la région paléarctique (Hym. Sphecid.). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, **27**: 241–276.
- de Beaumont J. 1954b.** Notes sur le genre *Prosopigastra* (Hym. Sphecid.). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, **27**: 153–156.
- de Beaumont J. 1956.** Notes sur les *Lindenius* paléarctiques (Hym. Sphecid.). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, **29**: 145–185.
- de Stefani T. 1888.** Note sulle Crisididi di Sicilia. *Il Naturalista Siciliano*, **7**: 88–95, 114–125, 139–145, 156–161, 177–182, 215–224, 273–291.
- DLussky G.M., Soyunov O.S. and Zabelin S.I. 1990.** *Ants of Turkmenistan*. Ylym Press, Ashkabad, 273 p. [In Russian].
- Dollfuss H. 1986.** Eine Revision der Gattung *Spilomena* Shuckard der westlichen und zentralen palaärktischen Region (Hymenoptera, Sphecidae). *Annalen des Naturhistorischen Museums in Wien*, **88–89B**: 481–510.
- Dollfuss H. 1991.** Bestimmungsschlüssel der Grabwespen Nord- und Zentraleuropas (Hymenoptera, Sphecidae) mit speziellen Angaben zur Grabwespenfauna Österreichs. *Stapfia*, **24**: 1–247.

- Dollfuss H.** 1995. A worldwide revision of *Pemphredon* Latreille 1796 (Hymenoptera, Sphecidae). *Linzer Biologische Beiträge*, **27**: 905–1019.
- Dollfuss H.** 2004. The Crabroninae wasps of “Biologiezentrum Linz” – collection in Linz, Austria (Hymenoptera, Apoidea, Crabronidae), Part 1. *Linzer Biologische Beiträge*, **36**: 761–784.
- Dollfuss H.** 2006. The Crabroninae wasps of “Biologiezentrum Linz” – collection in Linz, Austria, (Hymenoptera, Apoidea, Crabronidae), Part 2. *Linzer Biologische Beiträge*, **38**: 505–532.
- Dollfuss H.** 2008. The Crabronid wasps of the genus *Oxybelus* Latreille 1796 and *Brimocelus* Arnold 1927 of “Biologiezentrum Linz” collection in Linz, Austria, (Hymenoptera, Apoidea, Crabronidae). *Linzer Biologische Beiträge*, **40**: 463–505.
- Dollfuss H.** 2010. A Key to Wasps of the Genus *Podalonia* Fernald 1927 (Hymenoptera: Apoidea: Sphecidae) of the Old World. *Linzer Biologische Beiträge*, **42**: 1241–1291.
- Dollfuss H.** 2013. Revision of the wasp genus *Ammophila* Kirby 1798 (Hymenoptera: Apoidea: Sphecidae) of the Palaearctic Region and India. *Linzer Biologische Beiträge*, **45**: 383–564.
- Douwes P., Abenius J., Cederberg B. and Wahlstedt U.** 2012. Nationalnyckeln till Sveriges flora och fauna. Steklar: Myror-getingar. Hymenoptera: Formicidae-Vespidae. ArtDatabanken, SLU, Uppsala: 1–382.
- Dovnar-Zapolskij D.P.** 1925. Sawflies (Chalastogastra) of the steppes of the Pre-Caucasus. *Izvestiya Stavropol'skogo Entomologicheskogo Obshchestva*, **1**: 33–38. [In Russian].
- Dovnar-Zapolskij D.P.** 1926a. On stem sawflies (Cephidae) of the North Caucasus area. *Izvestiya Severo-Kavkazkoy Kraevoy Stantsii zashchity rasteniy*, **1**: 127–132. [In Russian].
- Dovnar-Zapolskij D.P.** 1926b. Remarks on sawflies. *Izvestiya Stavropol'skogo Entomologicheskogo Obshchestva*, **2**: 17–19. [In Russian].
- Dovnar-Zapolskij D.P.** 1928. Ueber Halmwespen aus der Gattung *Cephus* Latr. *Izvestiya Severo-Kavkazkoy Kraevoy Stantsii zashchity rasteniy*, **4**: 249–254. [In Russian].
- Dovnar-Zapolskij D.P.** 1929. Einige neue oder wenig bekannte Arten der Gattung *Empria* Lep. (Hymenoptera), mit einer Bestimmungstabelle der paläarktischen Arten. *Russkoe Entomologicheskoe Obozrenie*, **23**(1–2): 37–47.
- Dovnar-Zapolskij D.P.** 1930. Neue oder wenig bekannte Chalastogastren. *Russkoe Entomologicheskoe Obozrenie*, **24**(1–2): 86–94.
- Dovnar-Zapolskij D.P.** 1931a. Cephiden Studien (Hymenoptera, Chalastogastra) (I. Beitrag). *Ezhegodnik Zoologicheskogo Muzeya Akademii nauk SSSR*, **32**: 37–49.
- Dovnar-Zapolskij D.P.** 1931b. Eine Uebersicht über die Blattwespen (Chalastogastra) des nord kaukasischen Gebietes. *Izvestiya Severo-Kavkazkoy Kraevoy Stantsii zashchity rasteniy*, **6–7**: 33–62. [In Russian].
- Dovnar-Zapolskij D.P.** 1940. Contribution to the knowledge of the Hymenoptera of Kursk Oblast', Russia. *Trudy Tsentral'no-Chernozemnogo Gosudarstvennogo Zapovednika im. Prof. Alekhina*, **1**: 302–306. [In Russian].
- Dubatolov V.V.** 1998. Social wasps (Hymenoptera, Vespidae: Polistinae, Vespinae) of Siberia in the collection of Siberian Zoological Museum. *Far Eastern Entomologist*, **57**: 1–11.
- Dubatolov V.V.** 1999. Social wasps (Insecta, Hymenoptera, Vespidae: Vespinae, Polistinae) of the Daurian State Nature Reserve (South-Eastern Siberia). In: Insects of Dauria and adjacent territories. Vol. II. Proceedings of the Daurian State Biosphere Nature Reserve. Novosibirsk: 67–69. [In Russian].
- Dubatolov V.V.** 2011. Social wasps (Hymenoptera, Vespidae: Polistinae, Vespinae) of the Bastak nature reserve (Jewish autonomous province, Amur basin). *Amurian Zoological Journal*, **3**(1): 60–63.
- Dubatolov V.V. and Dolgikh A.M.** 2009. Social wasps (Hymenoptera, Vespidae: Polistinae, Vespinae) of the Bolshekhekhtsirsky nature reserve (the Khabarovsk suburbs), with notes on their distribution in the Lower Amur. *Amurian Zoological Journal*, **1**(1): 76–82.
- Dubatolov V.V. and Novomodnyi E.V.** 2005. New data on distribution of social wasps (Hymenoptera, Vespidae: Polistinae, Vespinae) in the Russian Far East. In: Streletsov A.N. (Ed.). *Zhivotnyi mir Dal'nego Vostoka: sbornik nauchnykh trudov*. Blagoveshchensk, **5**: 157–160. [In Russian].
- Dubatolov V.V., Streletzov A.N. and Malikova E.I.** 2002. New data on distribution of social wasps (Hymenoptera, Vespidae: Polistinae, Vespinae) in the Asian Russia and North China. In: Streletsov A.N. (Ed.). *Zhivotnyi mir Dal'nego Vostoka: sbornik nauchnykh trudov*. Blagoveshchensk, **4**: 117–122.
- Dubovikoff D.A.** 2002. New species of the ant genus *Bothriomyrmex* Emery, 1869 (Hymenoptera, Formicidae) from the Caucasus. *Entomological Review*, **81**: 918–922.
- Dubovikoff D.A.** 2005. A new species of the genus *Proformica* Ruzsky, 1902 (Hymenoptera, Formicidae) from North Ossetia and key to identification of *Proformica* species of the Caucasian Isthmus. *Caucasian Entomological Bulletin*, **1**: 189–191. [In Russian].
- Dubovikoff D.A. and Yusupov Z.M.** 2017. New and interesting finds of ants (Hymenoptera, Formicidae) in fauna of the Russia. *Caucasian Entomological Bulletin*, **13**(2). [In press].
- Dvořák L. and Castro L.** 2007. New and noteworthy records of vespid wasps (Hymenoptera: Vespidae) from the Palaearctic region. *Acta Entomologica Musei Nationalis Pragae*, **47**: 229–236.
- Dvořák L. and Roberts S.P.M.** 2006. Key to the paper and social wasps of Central Europe (Hymenoptera: Vespidae). *Acta Entomologica Musei Nationalis Pragae*, **46**: 221–244.

- Eaerdley C. and Urban R. 2010.** Catalogue of Afrotropical bees (Hymenoptera: Apoidea: Apiformes). *Zootaxa*, **2455**: 1–548.
- Ebmer A.W. 1972.** Neue westpaläarktische Halictidae (Halictinae, Apoidea). *Mitteilungen aus dem zoologischen Museum in Berlin*, **48**(2): 225–263.
- Ebmer A.W. 1974.** Die Bienen des Genus *Halictus* Latr. s. 1. im Großraum von Linz (Hymenoptera Apoidea). Nachtrag und zweiter Anhang. *Naturkundliches Jahrbuch der Stadt Linz*, **1973**: 123–158.
- Ebmer A.W. 1975.** Neue westpaläarktische Halictidae (Halictinae, Apoidea), Teil II. – Die Gruppe des *Halictus (Vestitohalictus) mucoreus* (Ev.). *Mitteilungen aus dem zoologischen Museum in Berlin*, **51**(2): 161–177.
- Ebmer A.W. 1978.** *Halictus*, *Lasioglossum*, *Rophites* und *Systropha* aus dem Iran (Halictidae, Apoidea) sowie neue Arten aus der Paläarktis. *Linzer Biologische Beiträge*, **10**(1): 1–109.
- Ebmer A.W. 1980.** Asiatische Halictidae (Apoidea, Hymenoptera). *Linzer biologische Beiträge*, **12**(2): 469–506.
- Ebmer A.W. 1986.** Die Artgruppe des *Lasioglossum strictifrons* (Vachal 1895) mit einer Bestimmungstabelle der Weibchen (Hymenoptera, Apoidea, Halictidae). *Linzer Biologische Beiträge*, **18**(2): 417–443.
- Ebmer A.W. 1988a.** Die europäischen Arten der Gattungen *Halictus* Latreille 1804 und *Lasioglossum* Curtis 1833 mit illustrierten Bestimmungstabellen (Insecta: Hymenoptera: Apoidea: Halictidae: Halictinae). 2 – Die Untergattung *Seladonia* Robertson 1918. *Senckenbergiana Biologica*, **68**(4/6): 323–375.
- Ebmer A.W. 1988b.** Kritische Liste der nicht-parasitischen Halictidae Österreichs mit Berücksichtigung aller mitteleuropäischen Arten (Insecta: Hymenoptera: Apoidea: Halictidae). *Linzer Biologische Beiträge*, **20**(2): 527–711.
- Ebmer A.W. 1995.** Asiatische Halictidae, 3. Die Artengruppe der *Lasioglossum carinata-Evylaeus* (Insecta: Hymenoptera: Apoidea: Halictidae: Halictinae). *Linzer Biologische Beiträge*, **27**(2): 525–652.
- Ebmer A.W. 1998.** Asiatische Halictidae - 7. Neue *Lasioglossum*-Arten mit einer Übersicht der *Lasioglossum* s. str.-Arten der nepalischen und yunnanischen Subregion, sowie des nördlichen Zentral-China (Insecta: Hymenoptera: Apoidea: Halictidae: Halictinae). *Linzer Biologische Beiträge*, **30**(1): 365–430.
- Ebmer A.W. 2000.** Asiatische Halictidae - 9. Die Artengruppe des *Lasioglossum pauperatum* (Insecta: Hymenoptera: Apoidea: Halictidae: Halictinae). *Linzer Biologische Beiträge*, **32**(2): 399–453.
- Ebmer A.W. 2004.** Zur Bienenfauna Nepals: Arten der Gattungen *Halictus*, *Lasioglossum* und *Dufourea* (Insecta: Hymenoptera: Apoidea: Halictidae). *Veröffentlichungen Naturkundemuseum Erfurt*, **23**: 123–150.
- Ebmer A.W. 2005.** Zur Bienenfauna der Mongolei. Die Arten der Gattungen *Halictus* Latr. und *Lasioglossum* Curt. (Insecta: Hymenoptera: Apoidea: Halictidae: Halictinae). *Linzer Biologische Beiträge*, **37**(1): 343–392.
- Ebmer A.W. 2011.** Holarktische Bienenarten – autochthon, eingeführt, eingeschleppt. *Linzer Biologische Beiträge*, **43**(1): 5–83.
- Ebmer A.W. and Sakagami S.F. 1990.** *Lasioglossum (Evylaeus) algirum pseudannulipes* (Blüthgen), erstmals in Japan gefunden, mit Notizen über die *L. (E.) leucopus*-Gruppe (Hymenoptera, Halictidae). *Japanese Journal of Entomology*, **58**: 835–838.
- Ebrahimi E. and Carpenter J.M. 2008.** Catalog of the vespid wasps of Iran (Hymenoptera, Vespidae). *Zootaxa*, **1785**: 1–42.
- Elfving R. 1968.** Die Bienen Finnlands. *Fauna Flora Fennica*, **21**: 1–69.
- Elliott M.G. 2007.** Annotated catalogue of the Pompilidae (Hymenoptera) of Australia. *Zootaxa*, **1428**: 1–83.
- Engel M.S. 2008.** New species and records of Ammobatine bees from Pakistan, Kyrgyzstan and Sri Lanka (Hymenoptera: Apoidea). *Acta Entomologica Slovenica*, **16**(1): 19–36.
- Engel M.S. 2009.** Two new species of *Ammobates* from the Arabian Peninsula and Egypt (Hymenoptera: Apidae). *Transactions of the Kansas Academy of Science*, **112**(3/4): 191–197.
- Engel M.S. and Grimaldi D.A. 2006.** The first cretaceous spider wasp (Hymenoptera: Pompilidae). *Journal of the Kansas Entomological Society*, **79**(4): 259–368.
- Enslin E. 1915.** Die Tenthredinoidea Mitteleuropas IV. *Deutsche Entomologische Zeitschrift*, **1915**(4): 311–412.
- Enslin E. 1919.** Tenthredinidae. Résultats scientifiques de l'Expédition des frères Kuznecov (Kouznetsov) à l'Oural Arctique en 1909, sous la direction de H. Backlund. *Zapiski Rossiyskoy Akademii Nauk po otdelejnyu fizikomatematiceskikh nauk, Ser. 8*, **28**(14): 1–10.
- Enslin E. 1920.** Die Blattwespengattung *Tenthredo* L. (*Tenthredella* Rohwer). *Abhandlungen der Zoologisch-Botanischen Gesellschaft in Österreich*, **11**(1): 1–96.
- Enslin E. 1927.** Die Tenthrediniden (Hymenoptera) der Kamtschatka-Expedition, 1908–1909. *Ezhegodnik Zoolicheskogo Muzeya Akademii Nauk SSSR*, **27**[1926]: 363–381.
- Erichson W.F. 1849.** *Scolia garrula* Erichs. In: Ménétriés E. *Catalogue des insectes recueillis par feu M. Lehmann. Mémoires de l'Académie Impériale des Sciences de Saint-Petersbourg*, **8**(6): 304–305.
- Erichson W.F. 1851.** The new species of Hymenoptera, Diptera, Neuroptera. In: Ménétriés E. Die Insekten (außer Parasiten). In: Middendorff A.T. von (1851–1853). *Reise in den äussersten Norden und Osten Sibiriens während der Jahre 1843 u. 1844 auf Veranstaltung der Kaiserl. Akademie der Wissenschaften zu St. Petersburg*, Vol. 2(1). Imperatorskaya Akademiya nauk, St. Petersburg: 60–69. [In Russian].
- Ermolenko V.M. 1955.** Materials to the fauna of sawflies (Hymenoptera, Tenthredinidae) of Kuril Islands. *Soobshcheniya Dal'nevostochnogo filiala AN SSSR*, **8**: 65–67. [In Russian].

- Ermolenko V.M.** 1959. To study of the fauna of horntails and sawflies (Hymenoptera, Symphyta) Crimea. IV'sezd Vsesoyuznogo entomologicheskogo obshchestva: Tezisy dokladov. AN SSSR, Moscow–Leningrad, 1: 48–50. [In Russian].
- Ermolenko V.M.** 1968. New desert genus and species of the family Tenthredinidae (Hymenoptera) from the Mongolian People's Republic. *Vestnik Zoologii*, 6: 32–37. [In Russian].
- Ermolenko V.M.** 1969. New species *Acantholyda angarica* sp. nov. (Hymenoptera, Pamphiliidae) from central Siberia. *Zbirnyk prac' Zoolohichnogo muzeyu AN URSR*, 33: 64–68. [In Russian].
- Ermolenko V.M.** 1971a. New species and genus of sawflies (Hymenoptera, Tenthredinidae) from the Sakhalin island. Communication I. *Vestnik Zoologii*, 5: 18–24. [In Russian].
- Ermolenko V.M.** 1971b. A new species of stem sawflies (Hymenoptera, Cephidae) from the Mountain Crimea. *Zoologicheskiy Zhurnal*, 50(4): 593–595. [In Russian].
- Ermolenko V.M.** 1972. The sawflies of Tenthredinoidea. Cimbicidae. Blasticotomidae. In: Ermolenko V.M. (Ed.). *Fauna Ukrayny. Rogokhvosty ta pil'shchyky*. Naukova Dumka, Kiev, 10(2): 1–204. [In Ukrainian].
- Ermolenko V.M.** 1973. New species of sawflies – *Pamphilus alnicola* sp. nov. and *Trichiosoma ushinskii* sp. nov. (Hymenoptera: Symphyta) from the Kuril Islands. Communication I. *Zbirnyk prac' Zoolohichnogo muzeyu AN URSR*, 35: 24–29. [In Russian].
- Ermolenko V.M.** 1975. The sawflies Tenthredinoidea. Argidae. Diprionidae. Tenthredinidae (Selanderiini, Dolerini). In: Ermolenko V.M. (Ed.). *Fauna Ukrayny. Rogokhvosty ta pil'shchyky*. Naukova Dumka, Kiev, 10(3): 1–378. [In Ukrainian].
- Ermolenko V.M.** 1976. An account of the fauna of sawflies (Hymenoptera, Symphyta) of Mongolia, I. Familes Pamphiliidae, Cimbicidae and Argidae. In: Kerzhner I.M., Kozlov M.A. (Eds). *Insects of Mongolia*. Vol. 4. Nauka, Leningrad: 259–267. [In Russian].
- Ermolenko V.M.** 1979. A new species of Xiphydriidae horntails (Hymenoptera, Xiphydriidae) from the Sikhote-Alin. *Trudy Vsesoyuznogo Entomologicheskogo Obshchestva*, 61: 128–131. [In Russian].
- Ermolenko V.M.** 1981. On the Symphyta (Hymenoptera) fauna from the northern Kuril Islands. In: Ler P.A., Kuipanskaya A.N., Lelej A.S., Fedikova V.S. (Eds). *Hymenopterous Insects of Eastern Siberia and Far East*. DVNTS AN SSSR, Vladivostok: 3–10. [In Russian].
- Ermolenko V.M.** 1984. On the fauna of horntails and sawflies (Hymenoptera, Symphyta) of Crimea. In: Dolin V.G., Savchenko E.N. (Eds). *Taksonomiya i zoogeografiya nasekomykh. Sbornik nauchnykh trudov*, Naukova Dumka, Kiev: 47–56. [In Russian].
- Ermolenko V.M.** 1992. An account of the fauna of the horntails, the stem sawflies and the sawflies (Hymenoptera, Symphyta) of Daurian Reserve. In: Amirkhanov A.M. (Ed.). *Nasekomye Daurii i sopredel'iykh territoriy*, CNIL, Moscow, 1: 71–76. [In Russian].
- Ermolenko V.M. and Zemkova R.I.** 1979. On finding of *Gilpinia fennica* (Hymenoptera, Diprionidae) in the Tuva Autonomous SSR. *Vestnik Zoologii*, 13(1): 70–71. [In Russian].
- Evans H.E.** 1964. A synopsis of the American Bethylidae (Hymenoptera, Aculeata). *Bulletin of the Museum of Comparative Zoology*, 132(1): 1–222.
- Eversmann E.** 1846. Hymenopterorum Rossicorum species novae vel parum cognitae, descriptae et ex parte depictae. *Bulletin de la Société Impériale des Naturalistes de Moscou*, 19: 436–443.
- Eversmann E.** 1847. Fauna hymenopterologica volgoruralensis exhibens Hymenopterorum species quas in provinciis Volgam fluvium inter et montes Uralenses sitis observavit et nunc descripsit. *Bulletin de la Société Impériale des Naturalistes de Moscou*, 20(1): 3–68.
- Eversmann E.** 1848. Hymenopterorum Rossicorum species novae vel parum cognitae, descriptae et ex parte depictae. Decas prima. *Bulletin de la Société Impériale des Naturalistes de Moscou*, 19(2): 436–443.
- Eversmann E.** 1849. Fauna Hymenopterologica Volgo-Uralensis. Fam. III. Sphegidae Latr. *Bulletin de la Société Impériale des Naturalistes de Moscou*, 22(4): 359–436.
- Eversmann E.** 1852. Fauna Hymenopterologica Volgo-Uralensis. *Bulletin de la Imperiale Society d'Naturalistes de Moscou*, 25(2): 3–137.
- Eversmann E.** 1858. Fauna Hymenopterologica Volgo-Uralensis. Continuatio. Familia Chrysidarum. *Bulletin de la Société Imperiale des Naturalistes de Moscou*, 30: 544–567.
- Fadeev K.I.** 2017. New data on the distribution of bethylid wasps (Hymenoptera: Bethylidae) in Russia. *Proceedings of the Russian Entomological Society*, 88(2). [In press].
- Fan J.** 1991. A study on Chinese subgenera *Vestitolahalictus* and *Seladonia* with description of new species (Apoidea: Halictidae). *Acta Entomologica Sinica*, 34(1): 478–482.
- Fateryga A.V.** 2015. *Dasypoda toroki* Michez, 2004. In: Ivanov S.P., Fateryga A.V. (Eds). *Red book of the Republic of Crimea. Animals*. PP "ARIAL" LLC, Simferopol: 229. [In Russian].
- Fateryga A.V.** 2017a. New records of solitary vespid wasps (Hymenoptera: Vespidae: Eumeninae, Masarinae) from Russia and adjacent countries. *Far Eastern Entomologist*, 334: 1–16.
- Fateryga A.V.** 2017b. New data on megachilid bees (Hymenoptera: Megachilidae) of the European part of Russia. *Proceedings of the Russian Entomological Society*, 88(2). [In press].
- Fateryga A.V.** 2018. Wasps of the family Vespidae (Hymenoptera) of the Crimean Peninsula. *Entomofauna*, 39. [In press].

- Fateryga A.V., Ivanov S.P. and Filatov M.A.** 2018. Megachilid-bees (Hymenoptera: Megachilidae) of the Crimean Peninsula. *Entomofauna*, **39**. [In press].
- Fateryga A.V., Mokrousov M.V. and Danilov Yu.N.** 2017. New and noteworthy records of solitary vespid wasps (Hymenoptera: Vespidae: Masarinae, Eumeninae) in Russia. *Proceedings of the Russian Entomological Society*, **88**(2). [In press].
- Fateryga A.V. and Popov I.B.** 2017. New records of Vespidae and Megachilidae (Hymenoptera) in Russia. *Ekosistemy*, **9**. [In press].
- Fateryga A.V., Protsenko Yu.V. and Zhidkov V.Yu.** 2014. *Isodontia mexicana* (Hymenoptera, Sphecidae), a new invasive wasp species in the fauna of Ukraine reared from trap-nests in the Crimea. *Vestnik Zooligii*, **48**: 185–188.
- Filatov M.A.** 2003. List of solitary bees of (Hymenoptera, Apoidea) in the Karadag natural reserve. In: *Chronical of nature*. Tom XVIII, 2001. SONAT, Simferopol: 82–86. [In Russian].
- Filatov M.A.** 2006. Contribution to the fauna of bees (Hymenoptera, Apoidea) of the Opuk nature reserve. *Trudy Nikitskogo Botanicheskogo Sada*, **126**: 110–117. [In Russian].
- Filatov M.A.** 2013. Fauna of solitary bees of the family Apidae (Hymenoptera, Apoidea) of Karadag, Opuk and Kazantip nature reserves. In: *The Nature reservations of Crimea. Biodiversity and Conservation in Sea of Azov and Black Sea Region. Materials of the 7th International Scientific-Practical Conference*. Simferopol: 409–411. [In Russian].
- Filatov M.A.** 2015. Dark cubitalia – *Cubitalia morio* Friese, 1911. In: Ivanov S.P., Fateryga A.V. (Eds). *Red book of the Republic of Crimea. Animals*. PP “ARIAL” LLC, Simferopol: 246. [In Russian].
- Filatov M.A., Ivanov S.P. and Budashkin Yu.I.** 2006. Bees (Hymenoptera, Apoidea) of the Kazantip nature reserve. *Trudy Nikitskogo Botanicheskogo Sada*, **126**: 258–262. [In Russian].
- Filatov M.A. and Ivanov S.P.** 2007. On the question of biodiversity of bees of the family Apidae (Hymenoptera, Apoidea) in natural reserves of Eastern Crimea. In: Artov A.M., Bokov V.A., Goldin P.E., Dulitskiy A.I., Ena A.V., Prokopov G.A., Rudyk A.N. (Eds). *Materials of IV International scientific and practical conference “Crimea natural reservations-2007”. Part. 2. Zoology*. Simferopol: 165–168. [In Russian].
- Finnamore A.T.** 1995. Revision of the world genera of the subtribe Stigmina (Hymenoptera: Apoidea: Sphecidae: Pempredoninae), Part 1. *Journal of Hymenoptera Research*, **4**: 204–284.
- Fischer von Waldheim G.** 1806. Monsieur le Directeur G. Fischer a communiquée à la même société (le 20. Janv. 1806) une notice d'un nouveau genre d'hyménoptères, qu'il appelle Nyctéridium *Mémoires de la Société des Naturalistes de l'Université Impériale de Moscou*, **1**: 287.
- Forsius R.** 1928. Über die von Wuorentaus in Kamtschatka gesammelten Tenthredinoiden. *Notulae Entomologicae*, **8**: 43–50.
- Forsius R.** 1929. Entomologische Ergebnisse der schwedischen Kamtschatka-Expedition 1920–1922. *Arkiv för Zoologi*, **20**(4): 1–4.
- Friese H.** 1895. *Die Bienen Europa's (Apidae europaea)*. Theil I. Schmarotzerbienen. R. Friedländer & Sohn, Berlin, 218 s.
- Friese H.** 1896. *Die Bienen Europa's (Apidae europaea)*. Theil II. Solitäre Apiden. Genus *Eucera*. R. Friedländer & Sohn, Berlin, 216 s.
- Friese H.** 1897. *Die Bienen Europa's (Apidae europaea)*. Theil III. Solitäre Apiden. Genus *Podalirius*. R. Friedländer & Sohn, Berlin, 316 s.
- Friese H.** 1901. *Die Bienen Europa's (Apidae europaea)*. Theil VI. Solitäre Apiden: Subfam. *Panurginae*, *Melittinae*, *Xylocopinae*. C. Lampe, Innsbruck, 284 s.
- Gauld I. and Bolton B.** 1988. *The Hymenoptera*. British Museum (Natural History) and Oxford University Press, London, Oxford, 332 p.
- Gayubo S.F. and Felton J.C.** 2000. The European species of the genus *Nitela* Latreille, 1809 (Hymenoptera: Sphecidae). *Annales de la Société Entomologique de France (N.S.)*, **36**: 291–313.
- Gayubo S.F. and Schmid-Egger Ch.** 2014. Order Hymenoptera, family Crabronidae. Genus *Nitela* Latreille, 1809. *Arthropod fauna of the United Arab Emirates*, **5**: 640–648.
- Gereys B.** 2008. Clé d'identification des *Microdynerus* de France métropolitaine et liste commentée des espèces (Hymenoptera: Vespidae: Eumeninae). *Bulletin Mensuel de la Société Linnéenne de Lyon*, **77**(5–6): 73–85.
- Gess S.K.** 1996. *The pollen wasps: ecology and natural history of the Masarinae*. Harvard University Press, Cambridge, Massachusetts, 340 p.
- Giordani Soika A.** 1958. Notulae vespidiologicae V. Biogeografia e sistematica del sottogenere *Katamenes* (M.W.). *Bollettino del Museo Civico di Storia Naturale di Venezia*, **11**: 57–67.
- Giordani Soika A.** 1976. Vespidi ed Eumenidi raccolti in Corea (Hym.). *Annales Historico-Naturales Musei Nationalis Hungarici*, **68**: 287–293.
- Gninenko Ju.I. and Seryj G.A.** 2009. Complexes of needle browsing sawflies in the pine stands of Low Volga Region. *Lisivnytstvo i Agrolisomelioratsiya*, **116**: 49–55. [In Russian].
- Gogala A.** 1999. Bee Fauna of Slovenia: Checklist of Species (Hymenoptera: Apoidea). *Scoparia*, **42**: 1–79.
- Gonzalez V.H., Griswold T., Praz Ch.J. and Danforth B.** 2012. Phylogeny of the bee family Megachilidae (Hymenoptera: Apoidea) based on adult morphology. *Systematic Entomology*, **37**: 261–286.
- Gorbatovsky V.V.** 1995. Family Bethylidae – Bethylids. In: Lehr P.A. (Ed.). *Key to the insects of the Russian Far East*, Vol. IV, pt 1. Nauka, St Petersburg: 179–190. [In Russian].

- Gorbatovsky V.V.** 1998. Family Bethylidae – Bethylids (addition). In: Lehr P.A. (Ed.). *Key to the insects of the Russian Far East*. Vol. IV, pt 3. Dalnauka, Vladivostok: 680–682. [In Russian].
- Gorbatovsky V.V. and Lelej A.S.** 1998. Family Sierolomorphidae. In: Lelej A.S. (Ed.). *Key to the insects of Russian Far East*. Vol. IV, pt. 3. Dalnauka, Vladivostok: 683–684. [In Russian].
- Gordh G. and Moczar L.** 1990. A catalog of the world Bethylidae (Hymenoptera, Aculeata). *Memoirs of the American Entomological Institute*, **46**: 1–364.
- Goulet H.** 1986. The genera and species of the Nearctic Dolerini (Symphyta: Tenthredinidae: Selandriinae): classification and phylogeny. *Memoirs of the Entomological Society of Canada*, **135**: 1–208.
- Goulet H., Boudreault C. and Schiff N.M.** 2015. Revision of the World species of *Xeris* Costa (Hymenoptera: Siricidae). *Canadian Journal of Arthropod Identification*, **28**: 1–127.
- Grace A.** 2010. Introductory Biogeography to Bees of the Eastern Mediterranean and Near East. Bexhill Museum, Sussex, 284 p.
- Grebennikov K.A. and Zryanyin V.A.** 2017. Interesting records of the ants (Hymenoptera, Formicidae) from Volga region and Dagestan. *Proceedings of the Russian Entomological Society*, **88**(2). [In press].
- Guénard B. and Dunn R.R.** 2012. A checklist of the ants of China. *Zootaxa*, **3358**: 1–77.
- Guershon M. and Ionescu-Hirsch A.** 2012. A review of the *Xylocopa* species (Hymenoptera: Apidae) of Israel. *Israel Journal of Entomology*, **41**: 145–163.
- Guglielmino A., Olmi M. and Buckle C.** 2013. An updated host-parasite catalogue of world Dryinidae (Hymenoptera: Chrysidoidea). *Zootaxa*, **3740**(1): 1–113.
- Guglielmino A. and Buckle C.** 2013. Description of the mature larva of *Ampulicomorpha schajovskoyi* De Santis & Vidal Sarmiento (Hymenoptera: Chrysidoidea: Embolemididae). *Zootaxa*, **3637**: 385–393.
- Guichard K.M.** 1991. Old World species of *Belomicrus* A. Costa, 1871. *Entomofauna*, **12**: 353–369.
- Guichard K.M.** 1993. European *Oxybelus* with a note on *Oxybelus dusmeti* Perez, 1966 (Hymenoptera, Sphecidae). *Entomofauna*, **15**: 529–536.
- Guiglia D.** 1948. Le Vespa d'Italia. *Memorie della Società Entomologica Italiana*, **17**: 1–84.
- Guiglia D.** 1969. The history of the peculiar genus *Fedtschenkia* Saussure (1880) (Hymenoptera: Fedtschenkiidae). *Israel Journal of Entomology*, **4**: 339–342.
- Guiglia D.** 1971. De la distribution géographique du genre *Fedtschenkia* (Saussure, 1880) (Hymenoptera: Fedtschenkiidae). *Proceedings of the XIII International Congress of Entomology, Moscow, August 2–9, 1968*, 1, Nauka, Leningrad: 138–140.
- Gupta S.K., Gayubo S.F. and Pulawski W.J.** 2008. On two Asian species of the genus *Mellinus* Fabricius, 1790 (Hymenoptera: Crabronidae). *Journal of Hymenoptera Research*, **17**: 210–25.
- Gurov A.V. and Battisti A.** 2009. Web-spinning sawflies of the genus *Cephalcia* Panzer (Hymenoptera: Pampiliidae) in Siberia. *Izvestiya Irkutskogo gosudarstvennogo universiteta. Seriya "Biologiya, ekologiya"*, **2**(1): 26–30.
- Gurvich Yu.M.** 1931. Contributions to the study of the bees in northern Caucasus. 1. Bees collected in the Aksay Biological Station of the Northern Caucasian State University. *Izvestiya Severo-Kavkazskogo Gosudarstvennogo Universiteta*, **21**(4): 92–187.
- Gusenleitner F. and Schwarz M.** 2002. Weltweite Checkliste der Bienengattung *Andrena* mit bemerkungen und ergänzungen zu paläarktischen arten (Hymenoptera, Apidae, Andreninae, *Andrena*). *Entomofauna*, Suppl. **12**: 1–1280.
- Gusenleitner F., Schwarz M., Ascher J.S. and Scheuchl E.** 2005. Korrekturen und Nachträge zu Gusenleitner & Schwarz (2002): "Weltweite Checkliste der Bienengattung *Andrena* mit Bemerkungen und Ergänzungen zu paläarktischen Arten (Hymenoptera, Apidae, Andreninae, *Andrena*)". *Entomofauna*, **26**(26): 437–472.
- Gusenleitner J.** 1966a. Vespidae, Eumenidae und Masaridae aus der Turkey, Tail I. *Polskie Pismo Entomologiczne*, **36**: 343–363.
- Gusenleitner J.** 1966b. Vespidae, Eumenidae und Masaridae aus der Turkey, Tail II. *Polskie Pismo Entomologiczne*, **37**: 663–667.
- Gusenleitner J.** 1972a. Übersicht über die derzeit bekannten westpaläarktischen Arten der Gattung *Eumenes* Latr. (Hymenoptera: Vespoidea). *Bollettino del Museo Civico di Storia Naturale di Venezia*, **22/23**: 67–117.
- Gusenleitner J.** 1972b. Beiträge zur Kenntnis der Fauna Afghanistans. Vespoidea, Hymenoptera. *Acta Musei Moraviae*, **56/57**: 315–334.
- Gusenleitner J.** 1977. Über Faltenwespen aus Marokko (Hym. Diptoptera). *Linzer Biologische Beiträge*, **9**(1): 131–161.
- Gusenleitner J.** 1981. Revision der paläarktischen *Stenodynerus*-Arten (Hymenoptera, Eumenidae). *Polskie Pismo Entomologiczne*, **51**: 209–305.
- Gusenleitner J.** 1991. Über Vespoidea (Hymenoptera) aus der Mongolei und der Sowjetunion. *Linzer Biologische Beiträge*, **23**(2): 631–641.
- Gusenleitner J.** 2004. Über Faltenwespen aus dem Oman (Hymenoptera: Vespidae, Eumenidae, Masaridae). *Linzer Biologische Beiträge*, **36**(1): 151–158.
- Gusenleitner J.** 2007a. Eine neue *Celonites*-Art aus Sibirien (Hymenoptera: Vespidae, Masarinae). *Linzer Biologische Beiträge*, **39**(1): 133–135.

- Gusenleitner J.** 2007b. Über Vespidae aus Pakistan (Hymenoptera: Vespidae). *Linzer Biologische Beiträge*, **39**(2): 969–972.
- Gusenleitner J.** 2013a. Die Gattungen der Eumeninae im Nahen Osten, in Nordafrika und in Arabien (Hymenoptera: Vespidae: Eumeninae). *Linzer Biologische Beiträge*, **45**(1): 5–107.
- Gusenleitner J.** 2013b. Fauna Europaea: Family Vespidae. Fauna Europaea version 2.6.2. http://www.faunaeur.org/full_results.php?id=11363 (accessed 30 January 2017).
- Gusenleitner J.** 2013c. Fauna Europaea: Family Sapygidae. Fauna Europaea version 2.6.2. https://fauna-eu.org/cdm_dataportal/taxon/ (accessed 30 January 2017)
- Gussakovskij V.V.** 1927. Palaearctic species of the genus *Astatus* Latr. (Hymenoptera, Sphecidae). *Annals of the Zoological Museum of the Academy of Sciences of USSR*, **28**: 265–296. [In Russian].
- Gussakovskij V.V.** 1928. Generis *Solierella* Spin. species palaearcticae (Hymenoptera, Sphecidae). *Russkoe Entomologicheskoe Obozrenie*, **22**: 78–84. [In Russian].
- Gussakovskij V.V.** 1930a. Review of Palaearctic species of the genus *Priocnemis* Schdte (Hymenoptera, Psamnochidae). *Annals of the Zoological Museum of the Academy of Sciences of USSR*, **31**(2): 227–290.
- Gussakovskij V.V.** 1930b. Eine neue Sphecidengattung von Transcaspien. *Eos. Revista Española de Entomología*, **6**: 275–286.
- Gussakovskij V.V.** 1931. Revision der Gattung *Ammoplanus* Giraud und einigen verwandten Sphegidengattungen. *Boletín de la Real Sociedad Española de Historia Natural*, **31**: 437–465.
- Gussakovskij V.V.** 1932. Verzeichnis der von Herrn D:r R. Malaise im Ussuri und Kamtschatka gesammelten aculeaten Hymenopteren. *Arkiv för Zoologi*, **24A**(10): 1–66.
- Gussakovskij V.V.** 1933. Revisio goneris *Prosopigastra* Costa (s. lat.) (Hymenoptera, Sphecidae). *Entomologicheskoe Obozrenie*, **25**: 154–173. [In Russian].
- Gussakovskij V.V.** 1935. Insectes Hyménoptères, Chalastrogastera 1. *Fauna SSSR*, **2**(1): 1–453. [In Russian].
- Gussakovskij V.V.** 1936. Les espèces paléarctiques du genre *Trypoxylon* Latr. (Hymenoptera, Sphecidae). *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, **3**: 639–667. [In Russian].
- Gussakovskij V.V.** 1937. A review of Palaearctic species of the genera *Didineis* Wesm., *Pison* Latr. and *Psen* Latr. (Hymenoptera, Sphecoidea). *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, **4**: 599–698. [In Russian].
- Gussakovskij V.V.** 1947. Insectes Hyménoptères, Chalastrogastera 2. *Fauna SSSR*, **2**(2): 1–235. [In Russian].
- Hammer K.** 1950. On the insect fauna of Cyprus. Results of the expedition of 1939 by Harald, Håkan and P.H. Lindberg. VI. Hymenoptera aculeata II. Chrysidae, Scoliidae und Mutillidae der Insel Cypern. *Commentationes Biologicae*, **10**(13): 1–7.
- Hamon J.** 1993. Observations sur *Scolia (Scolia) galbula* (Pallas, 1771), *Scolia (Scolia) fallax* Eversmann, 1849 et *Scolia (Discolia) hirta* (Schrank, 1781). *Nouvelle Revue d'Entomologie (N.S.)*, **10**(1): 87–96.
- Handlirsch A.** 1887. Monographie der mit *Nysson* und *Bembex* verwandten Grabwespen. *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe. Abtheilung I*, **95**: 246–421, pl. 1–5.
- Handlirsch A.** 1888a. Monographie der mit *Nysson* und *Bembex* verwandten Grabwespen. II. *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe. Abtheilung I*, **96**: 219–311, pl. 1–2.
- Handlirsch A.** 1888b. Monographie der mit *Nysson* und *Bembex* verwandten Grabwespen. III. *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe. Abtheilung I*, **97**: 316–565, pl. 1–3.
- Handlirsch A.** 1889. Monographie der mit *Nysson* und *Bembex* verwandten Grabwespen. IV. *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe. Abtheilung I*, **98**: 440–517, pls. 1–2.
- Handlirsch A.** 1892. Monographie der mit *Nysson* und *Bembex* verwandten Grabwespen. VI. *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe. Abtheilung I*, **101**: 25–205, pl. 1–3.
- Handlirsch A.** 1893. Monographider mit *Nysson* und *Bembex* verwandten Grabwespen. VII. (Schluss). *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe. Abtheilung I*, **102**: 657–942, pl. 1–7.
- Handlirsch A.** 1895. Nachträge und Schlusswort zur Monographie der mit *Nysson* und *Bembex* verwandten Grabwespen. *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe. Abtheilung I*, **104**: 801–1079.
- Hara H.** 2010. Sawflies (Hymenoptera, Symphyta) injurious to trees and shrubs in Hokkaido I: Xyelidae, Pampiliidae, Argidae and Cimbicidae. *Bulletin of the Hokkaido Forestry Research Institute*, **4**: 51–68.
- Hara H., Kojima H. and Shinohara A.** 2007. Arge solo-wiyofka (Hymenoptera, Argidae) feeding on *Betula ermanii*, newly recorded from Japan. *Japanese Journal of Systematic Entomology*, **13**(1): 85–89.
- Hara H. and Shinohara A.** 2000. A systematic study on the sawfly genus *Cimbex* of East Asia (Hymenoptera, Cimbicidae). *Japanese Journal of Systematic Entomology*, **6**(2): 199–224.
- Hara H. and Shinohara A.** 2006. The sawfly genus *Spinarge* (Hymenoptera, Argidae). *Bulletin of the National Science Museum, Series A, Zoology*, **32**(2): 61–94.

- Hara H. and Shinohara A.** 2008. The species-group of *Arge aenea* (Insecta, Hymenoptera, Argidae). *Bulletin of the National Science Museum, Series A, Zoology*, **34**(2): 77–94.
- Hara H. and Shinohara A.** 2012. *Arge aruncus* n. sp. (Hymenoptera, Argidae) feeding on *Aruncus dioicus* in Japan. *Bulletin of the National Science Museum, Series A, Zoology*, **38**(2): 75–87.
- Hara H. and Shinohara A.** 2014. The *Arge jonasi* species group (Hymenoptera, Argidae). *Bulletin of the National Science Museum, Series A, Zoology*, **40**(2): 79–104.
- Hara H. and Shinohara A.** 2015. The *Gilpinia abieticola* species group (Hymenoptera, Diprionidae). *Bulletin of the National Science Museum, Series A, Zoology*, **41**(1): 21–41.
- Hara H. and Shinohara A.** 2017. Taxonomic notes and new distribution and host plant records for sawflies and woodwasps (Hymenoptera, Symphyta) of Japan II. *Bulletin of the National Science Museum, Series A, Zoology*, **43**(1): 53–69.
- Hara H., Shinohara A. and Ibuki Sh.-i.** 2012. *Arge obesa* n. sp. (Hymenoptera, Argidae) feeding on *Carpinus* and *Ostrya* in Japan, with taxonomic notes on *Hylotoma japonica*. *Bulletin of the National Science Museum, Series A, Zoology*, **38**(3): 125–143.
- Hara H. and Smith D.R.** 2012. *Nesodiprion orientalis* sp. nov., *N. japonicus*, and *N. biremis*, with a key to species of *Nesodiprion* (Hymenoptera, Diprionidae). *Zootaxa*, **3503**: 1–24.
- Haris A.** 2000. Study on the Palaearctic *Dolerus* Panzer, 1801 species (Hymenoptera: Tenthredinidae). *Folia Entomologica Hungarica*, **61**: 95–148.
- Haris A.** 2001. Six new *Dolerus* Panzer, 1801 species from Japan, Turkey and the United States (Hymenoptera: Tenthredinidae). *Folia Entomologica Hungarica*, **62**: 83–93.
- Haris A.** 2002. Symphyta (Hymenoptera) from Mongolia IV. *Folia Entomologica Hungarica*, **63**: 65–85.
- Haris A.** 2003. The World Nematinae collection of the Hungarian Natural History Museum with the description of three new species (Hymenoptera: Tenthredinidae). *Folia Entomologica Hungarica*, **64**: 99–113.
- Haris A.** 2006a. Sawflies from Sakhalin and the Kuril Islands (Hymenoptera, Tenthredinidae). *Natura Somogyiensis, Kaposvár*, **9**: 187–200.
- Haris A.** 2006b. Study on the Palaearctic *Pristiphora* species (Hymenoptera: Tenthredinidae). *Natura Somogyiensis, Kaposvár*, **9**: 201–277.
- Haris A.** 2009. Check list and distribution of Swedish Nematinae: Cladiini and Nematini (Hymenoptera: Tenthredinidae). *Entomologisk Tidskrift*, **130**(3–4): 209–222.
- Haris A. and Blank S.M.** 1996. Two new *Dolerus* species from Korea (Hymenoptera, Symphyta, Tenthredinidae). *Acta Zoologica Academiae Scientiarum Hungaricae*, **42**(1): 59–63.
- Haris A. and Zsolnai B.** 2007. New Nematinae species (Hymenoptera: Symphyta, Tenthredinidae) from Japan and Korea. *Zoologische Mededelingen*, **81**(7): 137–147.
- Hartig T.** 1837. *Die Aderflügler Deutschlands mit besonderer Berücksichtigung ihres Larvenzustandes und ihres Wirkens in Wäldern und Gärten für Entomologen, Wald- und Gartenbesitzer. Die Familien der Blattwespen und Holzwespen nebst einer allgemeinen Einleitung zur Naturgeschichte der Hymenopteren. Erster Band.* Haude und Spener, Berlin, i–xiv + 1–416.
- Haupt H.** 1927. Monographie der Psammocharidae (Pompilidae) von Mittel-, Nord- und Osteuropa. *Beiheft der Deutsche Entomologische Zeitschrift*, **1926–1927**: 1–367.
- Haupt H.** 1938. Zur Kenntnis der Dryinidae I (Hymenoptera-Sphecoidea). *Zeitschrift für Naturwissenschaften*, **92**: 13–35.
- Haupt H.** 1941. Zur Kenntnis der Dryinidae II (Hymenoptera-Sphecoidea). *Zeitschrift für Naturwissenschaften*, **95**: 27–67.
- Haupt H.** 1944. Zur Kenntnis der Dryinidae III (Hymenoptera-Sphecoidea). *Stettiner Entomologische Zeitung*, **105**: 90–94.
- Haupt H.** 1959. Elemente einer systematischen Aufteilung der Macromerinae m. (Hymenoptera-Sphecoidea) Fam. Pompilidae, Subfam. Macromerinae. *Nova Acta Leopoldina*, **21**(141): i–xi + 5–74.
- Heibo E., Lønnve O.J., Barstad T.E., Blank S.M., Liston A., Prous M. and Taeger A.** 2014. Sawflies (Hymenoptera, Symphyta) newly recorded from Norway. *Norwegian Journal of Entomology*, **61**: 15–26.
- Heidemaa M., Nuorteva M., Hantula J. and Saarma U.** 2004. *Dolerus asper* Zaddach, 1859 and *Dolerus brevicornis* Zaddach, 1859 (Hymenoptera: Tenthredinidae), with notes on their phylogeny. *European Journal of Entomology*, **101**(4): 637–650.
- Heidemaa M. and Prous M.** 2006. The larvae of *Empria pumila* (Konow, 1896) and *E. pumiliooides* Lindqvist, 1968 (Hymenoptera: Tenthredinidae). In: Blank S.M., Schmidt S., Taeger A. (Eds). *Recent Sawfly Research – Synthesis and Prospects*. Goecke & Evers, Keltern: 97–104.
- Heidemaa M., Smith D.R. and Shinohara A.** 2012. Taxonomy of *Dolerus subfasciatus* auct. and *D. subfasciatus* F. Smith with notes on the sawfly subgenus *Equidolerus* (Hymenoptera, Tenthredinidae). *Zootaxa*, **3525**: 1–17.
- Heidemaa M. and Viitasaari M.** 2004. Taxonomy of the *Dolerus gibbosus* species group, with descriptions of *Dolerus zhelochovtsevi*, sp. nov., and males of *D. blanki* Liston, 1995, and *D. quadrinotatus* Biro, 1884, and redescription of the male of *D. gibbosus* Hartig, 1837 (Hymenoptera: Tenthredinidae). *Dissertationes Biologicae Universitatis Tartuensis, Tartu*, **99**(I): 1–25.
- Heidemaa M. and Zinovjev A.** 2004. *Dolerus anatolii*, n. sp., the first Palaearctic member of the subgenus *Neodolerus* Goulet (Hymenoptera: Tenthredinidae). *Proceedings of the Entomological Society of Washington*, **106**(1): 159–165.

- Hellén W. 1920(1919).** Über finländische Goldwespen. *Meddelanden af Societas pro Fauna et Flora Fennica*, **46**: 203–213.
- Hellén W. 1948.** Mitteilungen über einige Tenthredinoiden aus Ostfennoskandien. *Notulae Entomologicae*, **28**: 40–46.
- Hellén W. 1956.** Studien über paläarktische Dolerinen (Hym., Tenth.). *Notulae Entomologicae*, **35**(4): 97–107.
- Hermes M.G., Melo G.A.R. and Carpenter J.M. 2014.** The higher-level phylogenetic relationships of the Eumeninae (Insecta, Hymenoptera, Vespidae), with emphasis on *Eumenes* sensu lato. *Cladistics*, **30**(5): 453–484.
- Hoá D.T., Lien N.T.P. and Kojima J. 2012.** Taxonomic notes on the genus *Euodynerus* Dalla Torre (Hymenoptera: Vespidae: Eumeninae) from Northern Vietnam. *Tạp Chí Sinh Học*, **34**(4): 427–431.
- Holmgren A.E. 1883.** Insecta a viris doctissimis Norden-skiöld illum ducem sequentibus in insulis Waigatsch et Novaja Semlia anno 1875 collecta. Hymenoptera et Diptera. *Entomologisk Tidskrift*, **4**(3–4): 141–190.
- Horne C. and Smith F. 1872.** Hymenoptera from the North-West Provinces of India. *Transactions of the Zoological Society of London*, **7**(3): 161–196.
- Humala A.E. 1997.** On the fauna of Hymenoptera Apocrita of Nature Reserve "Kivach". In: Kravchenko A.V. (Ed.). *Flora and fauna of protected territories of Karelian Republic*. Karelian Research Centre of the Russian Academy of Sciences, Petrozavodsk: 50–72. [In Russian].
- Humala A.E. 2006.** On the insect fauna of "Kivach" Nature Reserve. *Trudy Karelskogo Nauchnogo Tsentr RAN*, **10**: 153–159. [In Russian].
- Humala A.E. and Polevoi A.V. 2009.** On the insects fauna of South-East Karelia. *Trudy Karelskogo Nauchnogo Tsentr RAN*, **4**: 53–75. [In Russian].
- Humala A.E. and Polevoi A.V. 2011.** Records of new and remarkable insect species (Insecta) in Northern Ladoga area. *Trudy Karelskogo Nauchnogo Tsentr RAN*, **2**: 142–144. [In Russian].
- Humala A.E. and Polevoi A.V. 2015.** Records of rare and noteworthy insect species (insecta) in the republic of Karelia. *Trudy Karelskogo Nauchnogo Tsentr RAN*, **6**: 19–46. [In Russian].
- Ikudome S. 1989.** A revision of the family Colletidae of Japan (Hymenoptera, Apoidea). *Bulletin of the Institute of Minamikyūshū Regional Science*, **5**: 43–314.
- Inomata R. 1970.** Taxonomic and biological studies on *Parachyprotasis* (Hymenoptera, Tenthredinidae). I. Descriptions of 16 new species from Japan. *Mushi*, **43**(1): 1–27.
- International Commission on Zoological Nomenclature (ICZN). 1943.** Opinion 144. On the status of the names *Crabro* Geoffroy, 1762, *Crabro* Fabricius, 1775, and *Cimbex* Olivier, 1790 (Insecta, Hymenoptera). *Opinions Rendered by the International Commission on Zoological Nomenclature*, **2**: 91–96.
- International Commission on Zoological Nomenclature (ICZN). 1961.** *International Code of Zoological Nomenclature adopted by the XV International Congress of Zoology. International Trust for Zoological Nomenclature*. London, xvii + 176 p.
- International Commission on Zoological Nomenclature (ICZN). 1978.** Opinion 1106. Conservation of the generic name *Rhopalum* Stephens, 1829 (Insecta, Hymenoptera). *The Bulletin of Zoological Nomenclature*, **34**: 237–239.
- International Commission on Zoological Nomenclature (ICZN). 1985.** *International Code of Zoological Nomenclature. Third edition adopted by the XX General Assembly of the International Union of Biological Sciences. The International Trust for Zoological Nomenclature*. London, and University of California Press, Berkeley, Los Angeles, xx + 338 p.
- International Commission on Zoological Nomenclature (ICZN). 1994.** Opinion 1754. Histoire abrégée des insectes qui se trouvent aux environs de Paris (Geoffroy, 1762): some generic names conserved (Crustacea, Insecta). *The Bulletin of Zoological Nomenclature*, **51**: 58–70.
- International Commission on Zoological Nomenclature (ICZN). 1998.** Opinion 1906. *Euchroeus* Latreille, 1809 (Insecta, Hymenoptera): conserved; *Chrysis purpurata* Fabricius, 1787 (currently *Euchroeus purpuratus*): specific name conserved; and *Chrysis gloriosa* Fabricius, 1793: specific name suppressed. *Bulletin of Zoological Nomenclature*, **55**: 194–196.
- International Commission on Zoological Nomenclature (ICZN). 1999.** *International Code of Zoological Nomenclature. Fourth Edition. ITZN*. London, i–xxx + 1–306 p.
- Invrea F. 1964.** *Mutillidae – Myrmosidae*. Calderini, Bologna, xii + 303 p.
- Ito M. 1985.** Additional notes on the bumblebee fauna of North Korea (Hymenoptera, Apidae). *Folia Entomologica Hungarica*, **46**(1): 5–22.
- Iuga V.G. 1958.** Hymenoptera Apoidea, Fam. Apidae, Sub-fam. Anthophorinae. *Fauna Republicii Populare Romîne*, **9**(3): 1–270 p.
- Ivanov S.P., Fатеряга A.V. and Filatov M.A. 2009.** Retrospective assessment of species diversity of wild bees and wasps (Hymenoptera, Aculeata) of the Botanical Garden of Vernadskiy Taurida National University. *Scientific Notes of Taurida V.Vernadsky National University. Series: Biology, chemistry*, **22**(61/3): 40–51. [In Russian].
- Ivanov S.P., Filatov M.A., Fатеряга A.V., Amolin A.V., Procenko Yu.V. and Drozdovskaya A.V. 2011.** Hymenoptera insects (Insecta, Hymenoptera) in Red book of the Crimea: project. In: *The Nature reservations of Crimea. Biodiversity and Conservation in Sea of Azov and Black Sea Region. Materials of the 6th International Scientific-Practical Conference*. Simferopol: 287–292. [In Russian].

- Ivie M.A.** 2001. On the geographic origin of the wheat stem sawfly (Hymenoptera: Cephidae): A new hypothesis of introduction from Northeastern Asia. *American Entomologist*, **47**(2): 84–97.
- Ivie M.A. and Zinovjev A.G.** 1996. Discovery of the wheat stem sawfly (*Cephus cinctus* Norton) (Hymenoptera: Cephidae) in Asia, with the proposal of a new synonymy. *The Canadian Entomologist*, **128**: 347–348.
- Iwata K.** 1938. Die Crabronen aus den Kurilen, Sachalin und Hokkaido. *Insecta Matsumurana*, **12**: 81–88.
- Iwata M.** 2003. Comparisons of Scopal Hairs between *Tetralonia mitsukurii* and *Tetralonia nipponensis* (Hymenoptera: Apidae). *Japanese Journal of Entomology (N.S.)*, **6**(2): 74–88.
- Jakovlev A.** 1888. Quelques nouvelles espèces des mouches à scie de l'Empire Russe. *Trudy Russkago Entomologicheskago Obshchestva v S. Peterburge*, **22**: 368–375.
- Jakovlev A.** 1891. Diagnoses Tenthredinidarum novarum ex Rossia Europaea, Sibiria, Asia Media et confinum. *Trudy Russkogo Entomologicheskago Obshchestva*, **26**: 1–62.
- Jakovlev A.** 1900. Faune du district de Walouyki du gouvernement de Woronège (Russie). Fascicula 2. Tenthredinidae. In: Velitchkovsky V. (Ed.). *Faune du district de Walouyki du gouvernement de Woronege (Russie)*, Charkov, **2**: 1–5.
- Jakovlev J., Polevoi A. and Humala A.** 2015. Insect fauna of Zaonezhye Peninsula and adjacent islands. In: Lindholm T., Jakovlev J., Kravchenko A. (Eds.). *Biogeography, landscapes, ecosystems and species of Zaonezhye Peninsula, in Lake Onega, Russian Karelia*. Finnish Environment Institute, Helsinki: 257–338.
- Johansson N. and Paukkunen J.** 2017. *Osmia disjuncta* Tkalcú, 1995 – a bee species new to the Western Palaearctic (Hymenoptera: Megachilidae). *Entomologisk Tidskrift*, **138**(1): 25–32.
- Józán Z.** 2011. Checklist of Hungarian Sphecidae and Apidae species (Hymenoptera, Sphecidae and Apidae). *Natura Somogyiensis*, **19**: 177–200.
- Kajmuk E.L.** 1986. A new species of the Tenthredinid genus *Nematus* Panzer (Hymenoptera, Tenthredinidae) from Yakutia. *Entomologicheskoe Obozrenie*, **65**(1): 138–141. [In Russian].
- Kajmuk E.L.** 1988. The conifer-chewing sawflies of subfamily Nematinae of Central Yakutia. In: Revin Yu.V. (Ed.). *Nasemomye lugovo-taezhnykh biotsenozov Yakutii*, Yakutskiy filial SO AN SSSR, Yakutsk: 81–87. [In Russian].
- Kajmuk E.L. and Popov A.A.** 2003. The entomofauna of wild and cultivated berries of the near Yakutsk. In: Isaev A.P. (Ed.). *Entomologicheskie issledovaniya v Yakutii*, CP Li V.G., Yakutsk: 162–170. [In Russian].
- Kajmuk E.L. and Popov A.A.** 2006. On the fauna of sawflies (Hymenoptera, Symphyta) – inhabitants of trees and shrubs in Yakutia. Isaev A.P. (Ed.). *Lesnye issledovaniya v Yakutii: itogi, sostoyanie i perspektivu. Tom 1. Merzlotnoe lesovedenie i lesovedstvo. Lesnaya ekologiya*, Yakutskiy gosudarstvenny universitet, Yakutsk: 160–163. [In Russian].
- Karawajew W.** 1926. Beiträge zur Ameisenfauna des Kaukasus, nebst einigen Bemerkungen über andere paläarktische Formen. *Konowia*, **5**(1–3): 94–199.
- Kasperek M.** 2015. The cuckoo bees of the genus *Stelis* Panzer, 1806 in Europe, North Africa and the Middle East. *Entomofauna, Suppl.* **18**: 1–144.
- Kawall J.H.** 1864. Beiträge zur Kenntnis der Hymenopteren Fauna Russlands. *Bulletin de la Société des Naturalistes de Moscou, Section biologique, Nouvelle Série*, **37**: 293–303.
- Kazenas V.L.** 1978. *The digger wasps of Kazakhstan and Middle Asia (Hymenoptera, Sphecidae). The determinant*. Izdatelstvo Nauka Kazakhskoy SSR, Alma Ata, 172 p. [In Russian].
- Kazenas V.L.** 1980. Contributions to the digger wasp fauna (Hymenoptera, Sphecidae) of the Soviet Far East. In: Lehr P.A. (Ed.). *Taxonomiya nasekomykh Dal'nego Vostoka*. DVNTS AN SSSR, Vladivostok: 80–94. [In Russian].
- Kazenas V.L.** 1984. *The digger wasps Cerceris of Central Asia and Kazakhstan*. "Nauka" Kazakhskoy SSR, Alma Ata, 232 p. [In Russian].
- Kazenas V.L.** 1993. Digger wasps of the genus *Dinetus* Panzer in the fauna of USSR. In: *Zoologicheskie Issledovaniya v Kazakhstane (k 50-letiyu Instituta Zoologii NAN RK)*. Gylym, Almaty: 58–65. [In Russian].
- Kazenas V.L.** 2001. *Fauna and biology of sphecid wasps (Hymenoptera, Sphecidae) of Kazakhstan and Central Asia*. Kazgos INTI, Almaty, 334 p. [In Russian].
- Kazenas V.L.** 2004. Digger wasps of the genus *Pseudoscolia* Radoszkowski (Hymenoptera: Crabronidae) of Kazakhstan and Central Asia. *Tethys Entomological Research*, **10**: 117–156.
- Kemper H. and Döhring E.** 1967. *Die sozialen Faltenwespen Mitteleuropas*. Paul Parey Verlag, Berlin-Hamburg, 180 p.
- Kerenskij I.P.** 1919. *Eucera* (Hymenoptera, Apidae) from the environs of Rostov-on-Don. *Protokoly Zasedaniy Obshchestva Estestvoispytateley pri Donskom Universitete*, **2**(1): 46–56. [In Russian].
- Kerenskiy I.P.** 1919. Contribution to the knowledge of Hymenoptera of Russia. 1. Hymenoptera of Kerensk and Chembar Districts, Penza Government. *Izvestiya Donskogo Universiteta*, **1**: 1–48. [In Russian].
- Kholodov V.V.** 1999. Records of *Eucera kullenbergi* (Hymenoptera, Anthophoridae) from the Crimea. *Vestnik Zoologii*, **33** (4/5): 118.
- Khomentovskij P.A.** 1983. *The insects-xylophagous of conifers of Kamchatka*. DVNC AN SSSR, Vladivostok, 175 p. [In Russian].
- Kieffer J.-J.** 1914. *Bethylidae. Das Tierreich*, 41. R. Friedländer und Sohn, Berlin, 595 s.
- Kieffer J.J. and Marshall T.A.** 1904–1906. Proctotrypidae. In: André E. (Ed.). *Species des Hyménoptères d'Europe & d'Algérie*. A. Hermann: Paris, **1904**: 1–64 + pls. 1–8, **1905**: 65–288 + pls. 9–16, **1906**: 289–552 + pls. 17–21.

- Kilimnik A.N.** 1993. Chrysidae of the genus *Hedychrum* Latreille, 1806 (Hymenoptera) of Eastern Europe. *Entomologicheskoe Obozrenie*, **71**(2): 388–399. [In Russian].
- Kim J.-K.** 2012. Taxonomic review of the genus *Euodynerus* (Hymenoptera: Vespidae: Eumeninae) in the Korean Peninsula. *Animal Systematics, Evolution and Diversity*, **28**(3): 161–167.
- Kim J.-K.** 2013. Biogeographic characteristics of Far Eastern wasps: A case study with eumenine wasps (Hymenoptera: Vespidae: Eumeninae). *Entomological Research*, **43**(1): 1–11.
- Kim J.-K., Moon T.-Y. and Yoon I.B.** 1994. Systematics of vespine wasps from Korea, I. Genus *Vespa* Linnaeus (Vespidae, Hymenoptera). *Korean Journal of Entomology*, **24**: 107–115.
- Kim J.-W. and Shinohara A.** 1997. Occurrence of *Praia ussuriensis* (Hymenoptera, Cimbicidae) of Korea. *Korean Journal of Applied Entomology*, **36**(3): 203–205.
- Kim J.W., Shinohara A. and Woo K.-S.** 1997. The first record of the family Blasticotomidae (Hymenoptera: Symphyta) from Korea. *Korean Journal of Applied Entomology*, **36**(2): 126–128.
- Kimsey L.S.** 1991. Relationships among the tiphiid wasp subfamilies (Hymenoptera). *Systematic Entomology*, **16**: 427–438.
- Kimsey L.S. and Bohart R.M.** 1991. *The Chrysid Wasps of the World*. Oxford Press, New York, 652 p.
- Kirby W.F.** 1882. *List of Hymenoptera with descriptions and figures of the typical specimens in the British Museum. 1. Tenthredinidae and Siricidae Vol. 1*. By order of the Trustees, London, 450 p.
- Kirchner L.** 1867. *Catalogus Hymenopterorum Europae. Societatis Zoologico-Botanicae. Vindobonae [=Vienna]*, 285 p.
- Knyazev S.A., Fateryga A.V. and Danilov Yu.N.** 2015. First records of *Onychopterocheilus pallasi* (Klug, 1805) (Hymenoptera, Vespidae, Eumeninae) in the Russian part of the West-Siberian Plain. *Euroasian Entomological Journal*, **14**(1): 35–36. [In Russian].
- Kobayashi K.** 1931. On a collection from Shikotan, Kuriles. *The transactions of the Kansai Entomological Society*, **2**: 59–66.
- Koch F.** 1982. Ein Beitrag zur Symphytenfauna des Gebietes um Kursk (UdSSR) (Insecta, Hymenoptera). *Faunistische Abhandlungen Staatliches Museum für Tierkunde Dresden*, **9**(2): 57–60.
- Koch F.** 1988a. Die Gattung *Sterictiphora* Billberg (Insecta, Hymenoptera, Symphyta: Argidae). *Entomologische Abhandlungen*, **52**(2): 29–61.
- Koch F.** 1988b. Die palaearktischen Arten der Gattung *Apethymus* Benson, 1939 (Hymenoptera, Symphyta, Alantinae). *Mitteilungen der Münchener Entomologischen Gesellschaft*, **78**: 155–178.
- Koch F.** 1988c. Zur Differentialdiagnose einiger *Allantus*-Arten (Insecta, Hymenoptera, Symphyta: Tenthredinidae). *Reichenbachia*, **26**(9): 43–53.
- Koch F.** 1989. Eine neue *Pristiphora*-Art aus der Verwandtschaft der *Pristiphora kamtchatica* Malaise (Insecta, Hymenoptera, Symphyta: Tenthredinidae). *Reichenbachia*, **26**(25): 145–148.
- Koch F. and Goergen G.** 2008. Annotations to the Palaearctic sawfly genus *Pseudarge* Gussakovskij, 1935 gen. rev. with a contribution to the genus *Arge* Schrank, 1802 in the Afrotropical region (Hymenoptera: Symphyta: Argidae: Arginae). *Mitteilungen der Münchener Entomologischen Gesellschaft*, **98**: 43–57.
- Kochetkov D.N.** 2017. Spider wasps (Hymenoptera, Pompilidae) of the Khinganskiy nature reserve, Amur Province. *A.I. Kurentsov's Annual Memorial Meetings*, **28**: 67–81. [In Russian].
- Kohl F.F.** 1885 (1884). Die Gattungen und Arten der Lariden Autorum. *Verhandlungen der kaiserlich-königlichen Zoologisch-Botanischen Gesellschaft in Wien*, **34**: 171–268, 327–454.
- Kohl F.F.** 1888. Die Crabronen der Section *Thyreopus* Lep. (Schieldwespen, Siebbienen). *Zoologische Jahrbücher. Abtheilung für Systematik, Geographie und Biologie der Thiere*, **3**: 543–590.
- Kohl F.F.** 1905. Palaearktische Crabronen (Hym.). *Zeitschrift für Systematische Hymenopterologie und Dipterologie*, **5**: 217–227.
- Kohl F.F.** 1913. *Fascicula 11. Hymenoptera in V. Velitchkovsky. Faune du District de Walouky du gouvernement de Woronéje (Russie)*. Johann Zellmayer, Stockerau, 21 p.
- Kohl F.F.** 1915. Die Crabronen (Hymenopt.) der paläarktischen Region. Monographisch bearbeitet. *Annalen des kaiserlich-königlichen Naturhististorischen Hofmuseums*, **29**: 1–453.
- Kohl F.F.** 1923–1924. Die Hymenopteren-Gattung *Belomisrus* A. Costa (s.l.). *Konowia*, **2**: 98–122 (1923), 180–202 (1923), 258–278 (1924).
- Kokujev N.R.** 1902. Materials for the hymenopterous fauna of Russia. *Russkoe Entomologicheskoe Obozrenie*, **2**: 4–12. [In Russian].
- Kokujev N.R.** 1906. About Russian representatives of the subfamily Alysonini Dalla Torre (Hymenoptera, Crabronidae). *Horae Societatis Entomologicae Rossicae*, **37**: 209–219. [In Russian].
- Kokujev N.R.** 1910. On the distribution of the Hymenoptera of the subfamily Cephini Konow (Hymenoptera Chalastogastra Konow) in Russia, and the description of new species. *Russkoe Entomologicheskoe Obozrenie*, **10**(3): 127–139. [In Russian].
- Kokujev N.R.** 1927. Hymenoptera, collected by V.V. Sovinskij on the shores of the Lake Baikal in 1902. *Trudy Komissii po izucheniyu ozera Baykala*, **2**: 63–76. [In Russian].
- Kolesnikov V.A.** 1977. Sphecid wasps (Hymenoptera, Sphecidae) of the Bryansk region and their role as entomophagous insects. *Entomologicheskoe Obozrenie*, **56**: 315–325. [In Russian].

- Konovalova I.B.** 2010. The bumble bees of Ukraine: species distribution and floral preferences. *Psyche*, 2010: Article ID 819740, 10 pages.
- Konow F.W.** 1886. Sieben neue *Allantus*-Arten. *Wiener Entomologische Zeitung*, 5: 17–21.
- Konow F.W.** 1891. Neue Blattwespen. *Wiener Entomologische Zeitung*, 10(2): 41–48.
- Konow F.W.** 1895. Analytische und kritische Bearbeitung der Gattung *Amauronematus* Knw. *Természetrajzi Füzetek*, 18: 166–187.
- Konow F.W.** 1897. Neue palaearctische Tenthrediniden. *Wiener Entomologische Zeitung*, 16(6): 173–187.
- Konow F.W.** 1900. Neue Sibirische Tenthrediniden. *Entomologische Nachrichten*, 26: 119–126.
- Konow F.W.** 1902a. Eine neue Holcocneme. (Hym. Chalastog.). *Zeitschrift für systematische Hymenopterologie und Dipterologie*, 2(2): 104.
- Konow F.W.** 1902b. Novum genus et novae species Chalastogastrorum Rossiae. *Trudy Jaroslavskogo Estestvenno-Istoricheskogo Obshchestva*, 1: 1–4.
- Konow F.W.** 1902c. Über eine neue Varietät und eine neue Art der Gattung *Dolerus* Jur. *Trudy Jaroslavskogo Estestvenno-Istoricheskogo Obshchestva*, 1: 8–9.
- Konow F.W.** 1903. Zwei neue *Pamphilius*. (Hym.). *Zeitschrift für systematische Hymenopterologie und Dipterologie*, 3(1): 37–38.
- Konow F.W.** 1904. Neue paläarctische Chalastogastra. (Hym.). (Fortsetzung). *Zeitschrift für systematische Hymenopterologie und Dipterologie*, 4(5): 260–270.
- Konow F.W.** 1906. Ueber einige Tenthrediniden der alten Welt. (Hym.). *Zeitschrift für systematische Hymenopterologie und Dipterologie*, 6(2): 122–127.
- Konow F.W.** 1907. Ueber die Ausbeute der russischen Polar-Expedition an Blattwespen im arctischen Sibirien. *Zapiski Imperatorskoy Akademii Nauk po fiziko-matematicheskому otdeleniyu*, Ser. 8, 18(3): 1–26.
- Konow F.W.** 1908. Über die Ausbeute der Expeditionen der Kaiserlich–Russischen Geographischen Gesellschaft an Blattwespen aus Tibet 1893–1895 und 1899–1901. *Ezhegodnik Zoologicheskago Muzeya Imperatorskoy Akademii Nauk*, 13: 9–25.
- Konusova O.L.** 2012. *Psithyrus sylvestris* (Lepeletier de Saint-Fargeau, 1832) as social parasite in a colony of *Bombus schrencki* Morawitz, 1881 (Hymenoptera, Apidae). *Euroasian Entomological Journal*, 11(6): 537–543. [In Russian].
- Konusova O.L. and Yanushkin V.V.** 2000. Ecological characteristics of bees (Hymenoptera, Apoidea) of the southern taiga in the Tomsk part of the Ob River basin. *Sibirskiy Ekologicheskiy Zhurnal*, 7(3): 283–286. [In Russian].
- Kopelke J.-P.** 2007a. The European species of the genus *Phyllocolpa*, part I: the *leucosticta*-group (Insecta, Hymenoptera, Tenthredinidae, Nematinae). *Senckenbergiana Biologica*, 87(1): 75–109.
- Kopelke J.-P.** 2007b. The European species of the genus *Phyllocolpa*, part II: the *leucapsis*-group (Insecta, Hymenoptera, Tenthredinidae, Nematinae). *Senckenbergiana Biologica*, 87(2): 149–161.
- Kopelke J.-P.** 2007c. The European species of the genus *Phyllocolpa*, part III: the species-groups *crassispina*, *scotaspis*, and *piliserra* (Insecta, Hymenoptera, Tenthredinidae, Nematinae). *Senckenbergiana Biologica*, 87(2): 163–183.
- Kostjukov V.V., Kosheleva O.V., Golovan' A.V., Gunesheva Z.M. and Nakonechnaya I.V.** 2014. Species of Hymenoptera parasitica in the entomological reservation of Institute biological plant protection – parasitoids of Gelechiidae including *Tuta absoluta* Meyrick (Lepidoptera). *Biological Plant Protection as the Basis of Ecosystem Stabilization* Vol. 8. All-Russian Research Institute of Biological Plant Protection RAAS, Krasnodar: 174–177. [In Russian].
- Kostylev G.** 1928. Materialien zur Kenntnis der Vespidenfauna der östlichen Krim. *Entomologische Mitteilungen*, 17(6): 398–407.
- Kostylev G.** 1929. Materialien zur Kenntnis der Vespiden-Fauna Armeniens. *Revue Russe d'Entomologie*, 23(1–2): 76–82.
- Kostylev G.** 1940a. Espèces novielles et peu connues de Vespides, d'Euménides et de Masarides paléarctiques (Hymenoptera) I. *Bulletin de la Société des Naturalistes de Moscou. Section Biologique*, 49(3–4): 137–154.
- Kostylev G.** 1940b. Espèces novielles et peu connues de Vespides, d'Euménides et de Masarides paléarctiques (Hymenoptera) II. *Bulletin de la Société des Naturalistes de Moscou. Section Biologique*, 49(5–6): 24–42.
- Kostyunin A.E.** 2013. Sawflies of the family Diprionidae (Hymenoptera, Symphyta) from West Siberia. *Euroasian Entomological Journal*, 12(5): 501–505. [In Russian].
- Kostyunin A.E.** 2015. Sawfly fauna (Hymenoptera: Symphyta) of the south-eastern part of Western Siberia and its regional features. *Proceedings of the Russian Entomological Society*, 86(2): 57–65. [In Russian].
- Kostyunin A.E. and Efimov D.A.** 2014. New records of sawflies of the family Cephidae (Hymenoptera, Symphyta) in steppes of southeast part of West Siberia, Russia. *Euroasian Entomological Journal*, 13(1): 33–34. [In Russian].
- Krasnobayev Yu.P., Antropov A.V., Lyubvina I.V. and Zabelin S.I.** 1995. Invertebrates of Zhiguli Hills, European Russia. V. Order Hymenoptera (Insecta). *Samarskaya Luka*, 6: 123–144. [In Russian].
- Kraus M.** 1998. Die Orussidae Europas und des Nahen Ostens (Hymenoptera: Orussidae). In: Taeger A., Blank S.M. (Eds). *Pflanzenwespen Deutschlands (Hymenoptera, Symphyta)*. Goecke & Evers, Keltern: 283–300.
- Krombein K.V.** 1979a. Family Sapygidae. In: Krombein K.V., Hurd P.D., Smith D.R., Burks B.D. (Eds). *Catalog of Hymenoptera in America North of Mexico*. Vol. 2. Smithsonian Institution Press, Washington, D.C.: 1319–1321.

- Krombein K.V. 1979b.** Superfamily Pompiloidea. In: Krombein K.V., Hurd P.D.Jr., Smith D.R., Burks B.D. (Eds). *Catalog of Hymenoptera in America North of Mexico. Vol. 2. Apocrita (Aculeata)*. Smithsonian Institution Press, Washington D.C.: 1523–1571.
- Kuhlmann M. 1993.** Kritisches Verzeichnis ausgewählter Stechimmenfamilien Westfalens (Hym., Aculeata). I. Chrysidae, Tiphidae, Mutilidae, Sapygidae, Pompilidae, Eumenidae, Sphecidae und Apidae (excl. Apinae). *Mitteilungen der Arbeitsgemeinschaft Ostwestfälisch-Lippischer Entomologen*, 6: 69–85.
- Kuhlmann M., Ascher J.S., Dathe H.H., Ebmer A.W., Hartmann P., Michez D., Müller A., Patiny S., Pauly A., Praz C., Rasmont P., Risch S., Scheuchl E., Schwarz M., Terzo M., Williams P.H., Amiet F., Baldock D., Berg Ø., Bogusch P., Calabuig I., Cederberg B., Gogala A., Gusenleitner F., Josan Z., Madsen H.B., Nilsson A., Ødegaard F., Ortiz-Sánchez J., Paukkunen J., Pawlikowski T., Quaranta M., Roberts S.P.M., Sáropataki M., Schwenninger H.-R., Smit J., Söderman G. and Tomozei B. 2015.** Checklist of the Western Palaearctic Bees (Hymenoptera: Apoidea: Anthophila). <http://westpalbees.myspecies.info> (accessed 30 April 2017).
- Kuhlmann M. and Proshchalykin M.Yu. 2011.** Bees of the genus *Colletes* Latreille 1802 of the Asian part of Russia, with keys to species (Hymenoptera: Apoidea: Colletidae). *Zootaxa*, 3068: 1–48.
- Kuhlmann M. and Proshchalykin M.Yu. 2013a.** The genus *Colletes* (Hymenoptera: Apoidea: Colletidae) in Central Asia. *Zootaxa*, 3750(5): 401–449.
- Kuhlmann M. and Proshchalykin M.Yu. 2013b.** The bees of the genus *Colletes* Latreille 1802 of Mongolia (Hymenoptera, Apoidea: Colletidae). *Beiträge zur Entomologie*, 63(2): 255–269.
- Kuhlmann M. and Proshchalykin M.Yu. 2014.** The bees of the genus *Colletes* Latreille 1802 of the European part of Russia, with keys to species (Hymenoptera: Apoidea: Colletidae). *Zootaxa*, 3878(3): 201–247.
- Kuhlmann M. and Proshchalykin M.Yu. 2016.** The genus *Colletes* (Hymenoptera: Apoidea: Colletidae) in Caucasus. *Zootaxa*, 4161(4): 367–385.
- Kuhlmann M. and Quest M. 2006.** A new species of the bee genus *Colletes* Latreille, 1802 (Hymenoptera: Colletidae) from the Russian Far East and Mongolia. *Far Eastern Entomologist*, 157: 1–4.
- Kulagin A.Yu. and Simonenkova V.A. 2014.** Ecological features of main phylophagous insects of broad-leaved and coniferous trees of South Predural Region. *Izvestiya Samarskogo nauchnogo tsentra Rossiyskoy akademii nauk*, 16(1): 127–133. [In Russian].
- Kumar G.P., Carpenter J.M., Castro L. and Sureshan P.M. 2017.** A taxonomic review of the Indian species of the genus *Eumenes* Latreille (Hymenoptera: Vespidae: Eumeninae). *Zootaxa*, 4317(3): 469–498.
- Kupianskaya A.N., Proshchalykin M.Yu. and Lelej A.S. 2014.** Contribution to the fauna of bumble bees (Hymenoptera, Apidae: *Bombus* Latreille, 1802) of the Republic of Tyva, Eastern Siberia. *Euroasian Entomological Journal*, 13(3): 290–294.
- Kupianskaya A.N. 1990.** *Ants of the Far Eastern USSR*. DVO AN SSSR, Vladivostok, 258 p. [In Russian].
- Kupianskaya A.N. 1995.** Family Formicidae – Ants. In: Lehr P.A. (Ed.). *Key to the insects of the Russian Far East*. Vol. IV, pt 1. Nauka, St Petersburg: 325–368. [In Russian].
- Kupianskaya A.N. 2012.** Family Formicidae – Ants. In: Lelej A.S. (Ed.). *Annotated Catalogue of the Insects of Russian Far East. Vol. I. Hymenoptera*. Dalnauka, Vladivostok: 423–433. [In Russian].
- Kurzenko N.V. 1977a.** A review of wasps of the family Eumenidae (Hymenoptera, Vespoidea) of the fauna of the USSR. Genera *Paravespa* Rad., *Paragymnomerus* Blüthg., *Tropidodynerus* Blüthg., *Gymnomerus* Blüthg. and *Odynerus* Latr. *Entomologicheskoe Obozrenie*, 56(3): 676–690. [In Russian].
- Kurzenko N.V. 1977b.** Eumenid wasps (Hymenoptera, Eumenidae) of the Mongolian People's Republic and adjacent regions of China and Southern Siberia. In: Kerzhner I.M. (Ed.). *Insects of Mongolia*. Vol. 5. Nauka, Leningrad: 537–582. [In Russian].
- Kurzenko N.V. 1978a.** Solitary vespid wasps of the family Eumenidae (Hymenoptera) of the fauna of the USSR. 326 p. (PhD thesis, not published). [In Russian].
- Kurzenko N.V. 1978b.** Solitary wasps of the family Eumenidae (Hymenoptera, Vespoidea) of the fauna of the USSR. 1. Subfamilies Raphiglossinae and Discoeliinae. *Zoologicheskiy Zhurnal*, 62(6): 867–871. [In Russian].
- Kurzenko N.V. 1981.** Review of the genera of the vespid wasps of the family Eumenidae (Hymenoptera, Vespoidea) of the fauna of the USSR. In: Lehr P.A. (Ed.). *Hymenopterous Insects of Far East*. DVNTS AN SSSR, Vladivostok: 81–112. [In Russian].
- Kurzenko N.V. 1986.** The wasps of the family Sapygidae (Hymenoptera, Aculeata) of the USSR fauna. In: Lehr P.A. (Ed.). *Hymenoptera of the Eastern Siberia and Russian Far East*. DVNTS AN SSSR, Vladivostok: 64–80. [In Russian].
- Kurzenko N.V. 1995a.** Family Vespidae – vespid wasps. In: Lelej A.S. (Ed.). *Keys to the insects of Russian Far East*. Vol. IV, pt 1. Nauka, St Petersburg: 295–324. [In Russian].
- Kurzenko N.V. 1995b.** Family Sapygidae – Sapygids. In: Lehr P.A. (Ed.). *Key to the insects of the Russian Far East*. Vol. IV, pt 1. Nauka, St Petersburg: 190–193. [In Russian].
- Kurzenko N.V. 1996.** A new Nearctic genus of Sapygidae with a key to the Nearctic and Palearctic genera (Hymenoptera, Sapygidae). *Memoires of the Entomological Society of Washington*, 17: 89–94.

- Kurzenko N.V. 2004a.** Solitary vespid wasps of the genus *Onychopterocheilus* Blüthgen, 1955 (Hymenoptera: Vespidae, Eumeninae) of the Russia and adjacent countries. Subgenus *Onychopterocheilus* s. str. *Proceedings of the Russian Entomological Society*, **75**(1): 242–252. [In Russian].
- Kurzenko N.V. 2004b.** The wasps of the families Vespidae and Sapygidae (Hymenoptera) from Sakhalin Island. In: Storozhenko S.Yu. (Ed.). *Flora and fauna of Sakhalin Island. Part 1.* Dalnauka, Vladivostok: 193–208. [In Russian].
- Kurzenko N.V. 2009.** Family Sapygidae – Sapygids. In: Storozhenko S.Yu. (Ed.). *The insects of the Lazovskiy Reserve.* Dalnauka, Vladivostok: 224–225. [In Russian].
- Kurzenko N.V. 2012a.** Family Sapygidae – Sapygids. In: Lelej A.C. (Ed.). *Annotated Catalogue of the Insects of Russian Far East. Vol. 1. Hymenoptera.* Dalnauka, Vladivostok: 403–404. [In Russian].
- Kurzenko N.V. 2012b.** Family Vespidae – vespid wasps. In: Lelej A.S. (Ed.). *Annotated Catalogue of the Insects of Russian Far East. Vol. 1. Hymenoptera.* Dalnauka, Vladivostok: 415–423. [In Russian].
- Kurzenko N.V. and Guseleinert J. 1994.** Sapygidae from Turkey, with a key to Palaearctic species of Sapyginae (Hymenoptera). *Linzer Biologische Beiträge*, **26**(2): 583–632.
- Kurzenko N.V. and Lelej A.S. 2007.** Family Chrysidae. In: Lelej A.S. (Ed.). *Key to the insects of Russian Far East. Vol. IV, pt 5.* Dalnauka, Vladivostok: 998–1006. [In Russian].
- Kurzenko N.V., Lelej A.S. and Taeger A. 1995.** Data to the fauna of the Aculeata of the Ussuri area (Hymenoptera, Sapygidae, Pompilidae, Vespidae). *Beiträge zur Entomologie*, **45**: 299–305.
- Kuwayama S. 1967.** *Insect fauna of the southern Kurile Islands.* Sapporo, 225 p.
- Kuznetsova V.T. 1990.** The Hymenoptera of the “Galich’ya Gora” Nature Reserve. Informational Report. All-Union Institute for Information in Science and Technology of the Academy of Sciences of the USSR, Moscow, 85 p. [In Russian].
- Kuznetsova V.T. 2009.** *The bees of the Lipetsk Province. A cadastre of species.* IPTS VGU, Voronezh, 120 p. [In Russian].
- Kuznetsov-Ugamsky N.N. 1929.** Die Ameisenfauna Dagestan. *Zoologische Anzeiger*, **83**: 34–45.
- Kuznetsov-Ugamskij N.N. 1927.** Two new Asiatic species of the genus *Cleptes*. Hymenoptera, Cleptidae. *Acta Societas Entomologicae Stauropolitanae*, **3**: 25–30.
- Kuznetsov-Ugamskij N.N. 1927.** Beiträge zur Blattwespenfauna des Süd-Ussuri-Gebietes. *Zoologischer Anzeiger*, **71**(9–10): 224–238, tab. 14.
- Kuznetsov-Ugamskij N.N. 1929.** Tenthredinologische Notizen. (Hymenoptera, Tenthredinoidea). *Zoologischer Anzeiger*, **80**(3–4): 49–55.
- Lacourt J. 1999.** Répertoire des Tenthredinidae ouest-paléarctiques (Hymenoptera, Symphyta). *Mémoires de la SEF*, **3**: 1–432.
- Lavrov S.D. 1927.** Contributions to the study of the insect fauna in the environs of Omsk. *Trudy Sibirskogo Instituta Sel’skogo Khoziaystva i Lesovodstva*, **8**(3): 51–99.
- Lebedev A.G. 1932.** Einige neue *Andrena*-Arten aus S.S.R. *Konowia*, **11**(1): 65–73.
- Leclercq J. 1950a.** Sur quelques Crabroniens (Hymenoptera Sphecidae) du groupe *Lindenius-Entomognathus*. *Bulletin de l’Institut Royal des Sciences Naturelles de Belgique*, **26**(6): 1–8.
- Leclercq J. 1950b.** Contribution à l’étude des Crabroniens asiatiques appartenant au genre *Ectemnius* (Dahlbom, 1845) (Hymenoptera, Sphecidae). *Bulletin de l’Institut Royal des Sciences Naturelles de Belgique*, **26**(23): 1–12.
- Leclercq J. 1950c.** Remarques sur les *Crossocerus* du sous-genre *Blepharipus* Lepeletier et Brullé, 1834 (Syn. *Acanthocrabro* Perkins, 1913) (Hym. Sphecidae Crabroninae). *Bulletin de l’Institut Royal des Sciences Naturelles de Belgique*, **26**(28): 1–10.
- Leclercq J. 1954.** *Monographie systématique, phylogénétique et zoogéographique des Hyménoptères Crabroniens.* Les Presses de “Lejeunia”, Liège, 371 p.
- Leclercq J. 1974.** Pour la connaissance des *Crossocerus* asiatiques du sous-genre *Cuphopterus* Morawitz (Hymenoptera Sphecidae Crabroninae). *Bulletin de la Société Royale des Sciences de Liège*, **43**(11/12): 659–668.
- Leclercq J. 1988.** Crabroniens du genre *Crossocerus* Lepeltier & Brullé trouvés en Sibérie notamment dans le territoire de Primorskii (Hymenoptera Sphecidae Crabroninae). *Bulletin de la Société Royale des Sciences de Liège*, **57**(1): 15–27.
- Leclercq J. 1989.** En vue d’une monographie du genre *Lindenis* Lepeletier & Brullé (Hymenoptera Sphecidae Crabroninae). *Bulletin de la Société Royale des Sciences de Liège*, **58**(6): 419–444.
- Leclercq J. 1991.** Crabroniens paléarctiques du genre *Odonotocrabro* Tsuneki (Hymenoptera, Sphecidae). *Bulletin et Annales de la Société royale belge d’Entomologie*, **127**: 133–139.
- Leclercq J. 1996.** Pour les Hyménoptères Sphécides Crabroniens palearctiques du genre *Entomognathus* Dahlbom. *Bulletin et Annales de la Société royale belge d’Entomologie*, **132**: 261–264.
- Leclercq J. 2009.** Hyménoptères Crabroniens d’Asie du genre *Crossocerus* Lepeletier & Brullé 1835 (Hymenoptera: Crabronidae Crabroninae). *Entomologie Faunistique*, **61**: 157–192.
- Leclercq J. 2011.** Hyménoptères Crabroniens d’Asie du genre *Rhopalum* Stephens 1829 (Hymenoptera: Crabronidae: Crabroninae). *Entomologie faunistique*, **63**(1): 3–21.

- Lecocq T., Dellicour S., Michez D., Dehon M., Dewulf A., De Meulemeester T., Brasero N., Valterova I., Rasplus J.-Y. and Rasmont P.** 2015. Methods for species delimitation in bumblebees (Hymenoptera, Apidae, *Bombus*): towards an integrative approach. *Zoologica Scripta*, **44**(3): 281–297.
- Lecocq T., Dellicour S., Michez D., Lhomme P., Vanderplanck M., Valterova I., Rasplus J.-Y. and Rasmont P.** 2013. Scent of a break-up: phylogeography and reproductive trait divergences in the red-tailed bumblebee (*Bombus lapidarius*). *BMC Evolutionary Biology*, **13**: 263.
- Lecocq T., Lhomme P., Michez D., Dellicour S., Valterova I. and Rasmont P.** 2011. Molecular and chemical characters to evaluate species status of two cuckoo bumblebees: *Bombus barbutellus* and *Bombus maxillosus* (Hymenoptera, Apidae, Bombini). *Systematic Entomology*, **36**: 453–469.
- Lee H.-S., Tadauchi O. and Lyu D.** 2015. *Arthropoda: Insecta: Hymenoptera: Andrenidae*. Insect Fauna of Korea. Vol. 13, No. 8. National Institute of Biological Resources, Icheon, 131 p.
- Lee J.-W. and Jung J.-C.** 1999. Taxonomic review of the family Cimbicidae (Hymenoptera, Symphyta) from Korea. *Insecta Koreana*, **16**(2): 197–223.
- Lee J.-W., Ruy S.-M., Quan Y.T. and Jung J.-C.** 2000. Economic Insects of Korea. 2. Hymenoptera (Symphyta: Tenthredinidae). *Insecta Koreana*, **9**: 1–223.
- Lee J.-W. and Ryu S.-M.** 1996. A systematic study on the Tenthredinidae (Hymenoptera: Symphyta) from Korea II. Ten new species of the Tenthredinidae. *Entomological Research Bulletin*, **22**: 17–34.
- Lee T.-S.** 1986. Notes on the genus *Dolichovespula* from China (Hymenoptera: Vespidae). *Sinozoologia*, **4**: 195–200.
- Lelej A.S.** 1978. Superfamily Mutilloidea. In: Tobias V.I. (Ed.). *Keys to the Insects of the European Part of the USSR*. Vol. III, pt 1. Nauka, Leningrad: 71–84. [In Russian].
- Lelej A.S.** 1985. *The velvet ants (Hymenoptera, Mutillidae) of the USSR and neighbouring countries*. Nauka, Leningrad, 268 p. [In Russian].
- Lelej A.S.** 1990. New and little known species of the spider wasps (Hymenoptera, Pompilidae) from Far East of USSR. In: Lelej A.S., Storozheva N.A., Storozhenko S.Yu. (Eds). *News of Systematic of the Insects of USSR Far East*. DVO AN SSSR, Vladivostok: 71–78. [In Russian].
- Lelej A.S.** 1992. Fam. Pompilidae. In: Chistyakov Yu.A. (Ed.). *The Insects of the Khabarovskii Natural Reserve*. Part 2. Dalnauka, Valdivostok: 239–243. [In Russian].
- Lelej A.S.** 1995. Family Pompilidae – Spider wasps. In: Lelej A.S. (Ed.). *Key to the Insects of the Russian Far East*. Vol. IV, pt 1. Nauka, St Petersburg: 211–264. [In Russian].
- Lelej A.S.** 2000. Family Pompilidae – Spider wasps. Addition. In: Lelej A.S. (Ed.). *Key to the Insects of the Russian Far East*. Vol. IV, pt 4. Dalnauka, Vladivostok: 615–624. [In Russian].
- Lelej A.S.** 2002. *Catalogue of the Mutillidae (Hymenoptera) of the Palaearctic region*. Dalnauka, Vladivostok, 171 p.
- Lelej A.S.** 2003. Review: G. Pagliano. Revisione della sottofamiglia Apterogyninae (Hymenoptera: Bradyno-baenidae) Museo Regionale di Scienze Naturali. Torino, 2002, 387 p. *Entomologicheskoe Obozrenie*, **82**: 526–527. [In Russian].
- Lelej A.S.** 2005a. *Catalogue of the Mutillidae (Hymenoptera) of the Oriental Region*. Dalnauka, Vladivostok, 252 p.
- Lelej A.S.** 2005b. Spider wasps (Hymenoptera, Pompilidae) of Sakhalin Island. In: Storozhenko S.Yu. (Ed.). *Flora and Fauna of Sakhalin Island (Materials of International Sakhalin Island Project)*. Part 2. Dalnauka, Vladivostok: 122–140. [In Russian].
- Lelej A.S.** 2007. Biogeography of mutillid wasps (Hymenoptera, Mutillidae). In: Rasnitsyn A.P., Gokhman V.E. (Eds). *Studies on Hymenopterous Insects. Collection of scientific papers*. Scientific Press Ltd., Moscow: 82–111. [In Russian].
- Lelej A.S.** 2012a. Family Bethylidae – Bethylids. In: Lelej A.S. (Ed.). *Annotated Catalogue of the Insects of the Russian Far East. Vol. 1. Hymenoptera*. Dalnauka, Vladivostok, 398–400. [In Russian].
- Lelej A.S.** 2012b. Family Mutillidae – Velvet ants, or Mutilids. In: Lelej A.S. (Ed.). *Annotated Catalogue of the Insects of Russian Far East. Vol. I. Hymenoptera*. Dalnauka: Vladivostok, 406–407. [In Russian].
- Lelej A.S. and Brothers D.J.** 2008. The genus-group names of Mutillidae (Hymenoptera) and their type species, with a new genus, new name, new synonymies, new combinations and lectotypifications. *Zootaxa*, **1889**: 1–79.
- Lelej A.S., Fateryga A.V. and Ivanov S.P.** 2016. The velvet ants (Hymenoptera: Mutillidae) of the Crimean Peninsula. *Far Eastern Entomologist*, **314**: 1–24.
- Lelej A.S. and Kurzenko N.V.** 2006. Wasps of the families Dryinidae, Bethylidae Pompilidae and Vespidae (Hymenoptera) from Moneron Island. In: Storozhenko S.Yu. (Ed.). *Flora and Fauna of Moneron Island (Materials of International Sakhalin Island Project)*. Dalnauka, Vladivostok: 243–246. [In Russian].
- Lelej A.S. and Kurzenko N.V.** 2012. Family Chrysididae – chrysidids]. In: Lelej A.S. (Ed.) *Annotated Catalogue of the Insects of Russian Far East. Vol. I. Hymenoptera*. Dalnauka, Vladivostok: 400–402. [In Russian].
- Lelej A.S. and Loktionov V.M.** 2012. Family Pompilidae (Psammocharide) – Spider wasps. In: Lelej A.S. (Ed.). *Annotated Catalogue of the Insects of Russian Far East. Vol. I. Hymenoptera*. Dalnauka, Vladivostok: 407–414. [In Russian].
- Lelej A.S., Loktionov V.M., Mokrousov M.V. and Savranskaya Zh.V.** 2017. New records of the velvet ants (Hymenoptera: Mutillidae) and spider wasps (Hymenoptera: Pompilidae) from the Russia. *Proceedings of the Russian Entomological Society*, **88**(2). [In press].

- Lelej A.S. and Mokrousov M.M.** 2015. *Sierolomorpha sogiana* spec. nov. from Central Asia, and a World catalogue of Sierolomorphidae (Hymenoptera). *Zootaxa*, **4018**(4): 593–599.
- Lelej A.S. and Mokrousov M.M.** 2016. To the taxonomy and behavior of *Apterogyna mlokosevitzi* Radoszkowski, 1879 (Hymenoptera: Bradynobaenidae). *Far Eastern Entomologist*, **324**: 178–21.
- Lelej A.S. and Mokrousov M.V.** 2017. The types of Scoliidae (Hymenoptera) described by Eduard Eversmann, with some taxonomic notes and checklist of Russian Scoliidae. *Far Eastern Entomologist*, **340**: 1–17.
- Lelej A.S. and Nemkov P.G.** 1997. Phylogeny, evolution and classification of Mutillidae (Hymenoptera). *Far Eastern Entomologist*, **46**: 1–24.
- Lelej A.S., Proshchalykin M.Yu. and Laktionov V.M.** 2014. To the knowledge of velvet ants (Hymenoptera, Mutillidae) of East Siberia. *Euroasian Entomological Journal*, **13**(4): 388–394. [In Russian].
- Lelej A.S. and Schmid-Egger Ch.** 2005. The velvet ants (Hymenoptera, Mutillidae) of Central Europe. *Linzer Biologische Beiträge*, **37**(2): 1505–1543.
- Lelej A.S. and Shlyakhtenok A.S.** 2015. The velvet ants (Hymenoptera: Mutillidae) of Belarus, with analysis of the fauna of Eastern European countries. *Far Eastern Entomologist*, **297**: 1–15.
- Lelej A.S. and Taeger A.** 2007a. Family Siricidae – Wood wasps. In: Lelej A.S. (Ed.). *Key to the Insects of the Russian Far East*. Vol. IV, pt 5. Dalnauka, Vladivostok: 958–959. [In Russian].
- Lelej A.S. and Taeger A.** 2007b. Family Tenthredinidae – Tenthredinid sawflies. In: Lelej A.S. (Ed.). *Key to the Insects of the Russian Far East*. Vol. IV, pt 5. Dalnauka, Vladivostok: 947–958. [In Russian].
- Leontiev V.V.** 2015. Overview on the entomofauna of Chrysidae and Mutillidae in the north-east Republic of Tatarstan. In: International scientific-practical conference “Protection of environment and environmental sciences”, Yelabuga: 131–137. [In Russian].
- Lesieur V., Martin J.-F., Weaver D.K., Hoelmer K.A., Smith D.R., Morrill W.L., Kadiri N., Peairs F.B., Cockrell D.M., Randolph T.L., Waters D.K. and Bon M.-C.** 2016. Phylogeography of the wheat stem sawfly, *Cephus cinctus* Norton (Hymenoptera: Cephidae): Implications for pest management. *PLoS ONE*, **11**(12): 1–19.
- Levchenko T.V.** 2007. A contribution to the bee fauna (Hymenoptera, Apoidea) of the Moscow province: a preliminary list of species of the family Megachilidae. In: Rasnitsyn A.P., Gokhman V.E. (Eds). *Studies on hymenopterous insects. Collection of scientific papers*. Scientific Press Ltd., Moscow: 153–161. [In Russian].
- Levchenko T.V.** 2009. Contributions to the fauna of bees (Hymenoptera: Apoidea) of Moscow Province. 1. Families Melittidae and Colletidae. *Eversmannia*, **17/18**: 27–43. [In Russian].
- Levchenko T.V.** 2011. Contribution to the fauna of bees (Hymenoptera: Apoidea) of Moscow Province. 2. Family Apidae. Subfamilies Apinae (excluding *Bombus* Latr.) and Xylocopinae. *Eversmannia*, **27/28**: 87–103. [In Russian].
- Levchenko T.V.** 2012. Contributions to the fauna of bees (Hymenoptera: Apoidea) of Moscow Province. 3. Family Apidae. Genus *Bombus* Latreille, 1802. *Eversmannia*, **31/32**: 72–88. [In Russian].
- Levchenko T.V.** 2013. Contribution to the fauna of bees (Hymenoptera: Apoidea) of Moscow Province. 4. Family Apidae. Subfamilies Apinae (conclusive part) and Nomadinae. *Eversmannia*, **34**: 40–56. [In Russian].
- Levchenko T.V.** 2014. Contributions to the fauna of bees (Hymenoptera, Apoidea) of Moscow Province. 5. Family Halictidae (excluding *Lasioglossum* s. l.). *Eversmannia*, **40**: 18–40. [In Russian].
- Levchenko T.V.** 2015. Contributions to the fauna of bees (Hymenoptera, Apoidea) of Moscow Province. 6. Family Halictidae Genus *Lasioglossum* Curtis, 1833 and *Evylaeus* Robertson, 1902. *Eversmannia*, **43/44**: 20–40. [In Russian].
- Levchenko T.V.** 2017. Contributions to the fauna of bees (Hymenoptera: Apoidea) of Moscow Province. 7. Family Andrenidae. Genus *Andrena* Fabricius, 1778. *Eversmannia*, **50**: 11–43. [In Russian].
- Levchenko T.V. and Tomkovich K.P.** 2014. Contribution to the bee fauna (Hymenoptera: Apiformes) of the Khanty-Mansi Autonomous Region, Western Siberia, Russia. *Entomofauna*, **26**(18): 313–348.
- Levchenko T.V. and Yuferev G.I.** 2013. Additions and corrections to the checklist of bees (Hymenoptera: Apoidea: Apoformes) of the Kirov Province. *Trudy Gosudarstvennogo prirodnogo zapovednika “Nurgush”*, **2**: 99–108. [In Russian].
- Levchenko T.V., Schwarz M. and Byvaltsev A.M.** 2017. New data and corrections to the fauna of bees of the family Apidae (Hymenoptera) of Russia. *Proceedings of the Russian Entomological Society*, **88**(2). [In press].
- Levi É.K., Sysolitina L.G. and Shernin A.I.** 1974. Chapter 9. Order Hymenoptera. In: Boikov B.V., Koroleva V.A., Levi É.K. (Eds). *Animal world of Kirov Oblast’*. Vol. 2. Kirov: 236–288. [In Russian].
- Li T., Cai W. and Li Q.** 2009. Three new species and one new record of *Liris* Fabricius (Hymenoptera: Crabronidae) from China, with a key to the Palearctic species. *Entomologica Fennica*, **20**: 111–120.
- Li T.-J. and Chen B.** 2014. The taxonomic accounts of the genus *Symmorphus* Wesmael (Hymenoptera, Vespidae, Eumeninae) from China, with descriptions of three new species. *ZooKeys*, **389**: 9–26.
- Li Z., Liu M. and Wei M.** 2016. A new species of *Macrophyia* Dahlbom (Hymenoptera: Tenthredinidae) with a key to species of the *Macrophyia sibirica* group from China. *Entomotaxonomia*, **38**(1): 44–52.

- Li Z.-J., Liu M.-M. and Wei M.-C.** 2014. Four new species of *sanguinolenta*-group of the genus *Macrophya* (Hymenoptera: Tenthredinidae) from China. *Zoological Systematics*, **39**(4): 520–533.
- Li Z.-J., Xiao W. and Wei M.-C.** 2011. A new species of *Emphytus* Klug (Hymenoptera: Tenthredinidae) from Mountain Helanshan of Inner Mongolia. *Entomotaxonomia*, **33**(4): 312–316.
- Lieftinck M.A.** 1968. A review of Old World species of *Thyreus* Panzer (= *Crocisa* Jurine) (Hym., Apoidea, Anthophoridae) Part 4. Palearctic species. *Zoologische Verhandelingen*, **98**: 1–139.
- Lieftinck M.A.** 1974. Review of Central and East Asiatic *Habropoda* F. Smith with *Habrophorula*, a new genus from China (Hymenoptera, Anthophoridae). *Tijdschrift voor Entomologie*, **117**(5): 157–224.
- Lieftinck M.A.** 1980. Prodrome to a monograph of the Palaearctic species of the genus *Melecta* Latreille 1802 (Hymenoptera, Anthophoridae). *Tijdschrift voor Entomologie*, **123**(6): 129–349.
- Lindqvist E.** 1968. Die *Empria*-Arten Finnlands (Hymenoptera, Symphyta). *Notulae Entomologicae*, **48**: 23–33.
- Lindqvist E.** 1970. Neue Nematinen aus dem Pribäikal-gebiet, Sibirien (Hymenoptera, Tenthredinidae). *Notulae Entomologicae*, **50**: 97–104.
- Lindqvist E.** 1971. Einige von R. Malaise aus Kamtschatka beschriebene Nematinen (Hymenoptera, Tenthredinidae). *Notulae Entomologicae*, **51**: 124–129.
- Lindqvist E.** 1972. Zur Nomenklatur und Taxonomie einiger Blattwespen (Hymenoptera, Symphyta). *Notulae Entomologicae*, **52**: 65–77.
- Lindqvist E.** 1975. Neue Nematinen aus Sibirien (Hymenoptera, Tenthredinidae). *Notulae Entomologicae*, **55**: 13–20.
- Lindqvist E.** 1977. Einige neue Blattwespen (Hymenoptera, Tenthredinidae). *Annales Entomologici Fennici*, **43**(3): 92–95.
- Linsenmaier W.** 1959. Revision der Familie Chrysidae (Hymenoptera) mit besonderer Berücksichtigung der europäischen Spezies. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, **32**(1): 1–232.
- Linsenmaier W.** 1968. Revision der Familie Chrysidae (Hymenoptera). Zweiter Nachtrag. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, **41**(1–4): 1–144.
- Linsenmaier W.** 1969. The chrysidid wasps of Palestine (Hym., Chrysidae). A faunistic catalogue with descriptions of new species and forms. *Israel Journal of Entomology*, **4**: 343–375.
- Linsenmaier W.** 1987. Revision der Familie Chrysidae (Hymenoptera). 4 Teil. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, **60**(1/2): 133–158.
- Linsenmaier W.** 1997. Altes und Neues von den Chrysiden. (Hymenoptera, Chrysidae). *Entomofauna*, **18**(19): 245–300.
- Linsenmaier W.** 1999. Die Goldwespen Nordafrikas (Hymenoptera, Chrysidae). *Entomofauna*, Suppl., **10**: 1–210.
- Liston A., Knight G., Sheppard D., Broad G. and Livermore L.** 2014a. Checklist of British and Irish Hymenoptera – Sawflies, ‘Symphyta’. *Biodiversity Data Journal*, **2**: 1–140.
- Liston A.D., Heibo E., Prous M., Vårdal H., Nyman T. and Vikberg V.** 2017. North European gall-inducing *Euura* sawflies (Hymenoptera, Tenthredinidae, Neminae). *Zootaxa*, **4032**(1): 1–115.
- Liston A.D.** 1992. Taxonomy of European *Arge* related to *nigripes* (Retzius) (Hymenoptera, Symphyta: Argidae). *Entomologische Berichte Luzern*, **27**: 135–140.
- Liston A.D.** 1994. *Kaliofenusa* species from Asia and Asia Minor, including one new to science (Hym., Tenthredinidae). *The Entomologist's Monthly Magazine*, **130**: 81–83.
- Liston A.D.** 1995. *Compendium of European Sawflies. List of species, modern nomenclature, distribution, foodplants, identification literature*. Chalastos Forestry, Gottfrieding, 190 p.
- Liston A.D.** 2012. On West Palaearctic *Mesoneura* species (Hymenoptera: Tenthredinidae). *Entomologist's Monthly Magazine*, **148**: 187–194.
- Liston A.D. and Prous M.** 2014. Sawfly taxa (Hymenoptera, Symphyta) described by Edward Newman and Charles Healy. *ZooKeys*, **398**: 83–98.
- Liston A.D., Savina H., Nagy Z.T., Sonet G. and Boevé J.-L.** 2014b. Taxonomy, phylogeny and host plants of some *Abia* sawflies (Hymenoptera, Cimbicidae). *Zootaxa*, **3821**(1): 125–132.
- Liston A.D., Taeger A. and Blank S.M.** 2006. Comments on European sawflies (Hymenoptera: Symphyta). In: Blank S.M., Schmidt S., Taeger A. (Eds). *Recent Sawfly Research – Synthesis and Prospects*. Goecke & Evers, Keltern: 245–263.
- Lith J.P. van.** 1956. Notes on *Epeolus* (Hymenoptera Aculeata, Apidae). *Tijdschrift voor Entomologie*, **99**(1/2): 31–45.
- Litman J.R., Griswold T. and Danforth B.N.** 2016. Phylogenetic systematics and a revised generic classification of anthidiine bees (Hymenoptera: Megachilidae). *Molecular Phylogenetics and Evolution*, **100**: 183–198.
- Liu M., Chu B., Xiao W. and Li Z.** 2015. Two new species of *Macrophya* Dahlbom (Hymenoptera: Tenthredinidae) with a key to species of the *Macrophya annulitibia* group from China. *Entomotaxonomia*, **37**(1): 72–80.
- Løken A.** 1964. Social wasps in Norway (Hymenoptera, Vespidae). *Norsk Entomologisk Tidsskrift*, **12**: 195–218.
- Løken A.** 1973. Studies on Scandinavian bumble bees (Hymenoptera, Apidae). *Norsk Entomologisk Tidsskrift*, **20**: 1–218.
- Løken A.** 1984. Scandinavian species of the genus *Psithyrus* Lepeletier (Hymenoptera: Apidae). *Entomologica Scandinavica* (Suppl.), **23**: 1–45.

- Loktionov V.M.** 2010. *Anoplius eous* Yasumatsu, 1936 (Hymenoptera, Pompilidae) is newly recorded from Russia. *Far Eastern Entomologist*, **211**: 11–12.
- Loktionov V.M.** 2011. The fauna and geographical distribution of spider wasps (Hymenoptera: Pompilidae) of the Russia Far East. In: Lelej A.S. (Ed.). *Key to the Insects of the Russian Far East. Additional Volume. Analysis of the fauna and general index of the names*. Dalnauka, Vladivostok: 81–92. [In Russian].
- Loktionov V.M. and Lelej A.S.** 2010. Zoogeography of spider wasps (Hymenoptera: Pompilidae) of the Russian Far East. *Proceedings of the Russian Entomological Society*, **81**(2): 51–58. [In Russian].
- Loktionov V.M. and Lelej A.S.** 2014. *Spider wasps (Hymenoptera: Pompilidae) of the Russian Far East*. Dalnauka, Vladivostok, 472 p. [In Russian].
- Loktionov V.M. and Lelej A.S.** 2015. Keys to the genera of the spider wasps (Hymenoptera: Pompilidae) of Russia and neighbouring countries, with check-list of genera. *Zootaxa*, **4034**(1): 87–111.
- Loktionov V.M. and Lelej A.S.** 2017. An annotated catalogue of the spider wasps (Hymenoptera: Pompilidae) of Russia. *Zootaxa*, **4280**(1): 1–95.
- Lomholdt O.** 1982. On the origin of the bees (Hymenoptera: Apidae, Sphecidae). *Entomologica Scandinavica*, **13**: 185–190.
- Long K.D., Hue L.X., Hoa D.T. and Phong P.H.** 2012. A preliminary study on bees (Hymenoptera: Apoidea: Apiformes) from Northern and North Central Vietnam. *Tap Chí Singh Hoc*, **34**(4): 419–426.
- Lopatin A.V. and Dobrynin N.D.** 2005. Superfamily Apoidea. In: Negrobov O.P. (Ed.). *Inventory of invertebrates of the Voronezh Province*. Voronezh State University, Voronezh: 667–692. [In Russian].
- Lykov V.A.** 2000. A review of the fauna of bees of the families Melittidae, Megachilidae and Anthophoridae (Hymenoptera, Apoidea) in the Perm Province. *Vestnik Permskogo Universiteta*, **2**: 216–227. [In Russian].
- Ma Z.-X., Chen B. and Li T.-J.** 2016. A taxonomic account of the genus *Stenodynerus* from China, with descriptions of five new species (Hymenoptera, Vespidae, Eumeninae). *ZooKeys*, **595**: 17–48.
- Maa T.** 1949a. A synopsis of Asiatic Siricoidea with notes on certain exotic and fossil forms. *Notes d'Entomologie Chinoise*, **13**(2): 11–189.
- Maa T.** 1949b. A synopsis of Chinese sawflies of the superfamily Megalodontoidea (Hymenoptera). *The Chinese Journal of Zoology, Nanking*, **3**: 30–42.
- Macek J.** 2010. Taxonomy, distribution and biology of selected European *Dinax*, *Strongylogaster* and *Taxonus* species (Hymenoptera: Symphyta). *Acta Entomologica Musei Nationalis Pragae*, **50**(1): 253–271.
- Macek J.** 2012. Sawflies (Hymenoptera: Symphyta) of the Bílé Karpaty protected landscape area and biosphere reserve (Czech Republic). *Acta Musei Moraviae, Scientiae biologicae*, **96**(2): 819–896.
- Macek J., Strejček J. and Straka J.** 2007. Chrysidoidea: Bethylidae (hbítěnkovití). In: Bogusch P., Straka J., Kment P. (Eds). *Annotated checklist of the Aculeata (Hymenoptera) of the Czech Republic and Slovakia*. Národní museum, Praha: 21–40. (Acta Entomologica Musei Nationalis Pragae, Suppl. 11).
- Magis N.** 2003. Notes faunistiques sur les espèces du genre *Tenthredo* Linné, 1758 sensu lato dans la région franco-rhénane (Hymenoptera Symphyta: Tenthredinidae, Tenthredininae). *Notes fauniques de Gembloix*, **53**: 3–20.
- Maharramov M.M. and Bayramov A.B.** 2014. The fauna of true bees of the tribe Eucerini (Hymenoptera, Apoidea, Apidae) in the Nakhchivan autonomous republic of Azerbaijan. *Vestnik Altaiskogo Gosudarstvennogo Agrarnogo Universiteta*, **4**(14): 59–63.
- Mahmood K., Ullah M., Aziz A., Hasan S.A. and Inayatullah M.** 2012. To the knowledge of Vespidae (Hymenoptera) of Pakistan. *Zootaxa*, **3318**: 26–50.
- Maidl F.** 1912. Die Xylocopen (Holzbienen) des Wiener Hofmuseums. Ein Beitrag zu einer Monographie dieser Gattung. *Annalen des k. k. naturhistorischen Hofmuseums*, **26**(3/4): 249–330.
- Maidl F. and Klima A.** 1939. *Astatinae – Nyssoninae. Hymenopterorum Catalogus. Pars 8. Sphecidae 1.* – S-Dr. W. Junk, Verlag für Naturwissenschaften, Gravenhage, 150 p.
- Malaise R.** 1931a. Entomologische Ergebnisse der schwedischen Kamtchatka Expedition 1920–1922. (35. Tenthredinidae). [Separatum]. *Arkiv för Zoologi*, **23**(2[8]): 1–68.
- Malaise R.** 1931b. Blattwespen aus Wladiwostok und anderen Teilen Ostasiens. *Entomologisk Tidskrift*, **52**(2): 97–159.
- Malaise R.** 1939. The genus *Leptocimbex* Sem., and some other Cimbicidae. *Entomologisk Tidskrift*, **60**(1–2): 1–28.
- Malaise R.** 1943. A new *Pamphilius* (Hym. Tenth.) from Ussuri. *Entomologisk Tidskrift*, **4**(1–2): 126.
- Malaise R.** 1945. Tenthredinoidea of South-Eastern Asia with a general zoogeographical review. *Opuscula Entomologica*, Suppl. 4: 1–288.
- Malaise R.** 1947. Entomological results from the Swedish expedition 1934 to Burma and British India. Hymenoptera: Tenthredinoidea. Collected by René Malaise. The Tenthredinoidea of South Eastern Asia. Part III. The *Emphytus-Athlophorus* group. *Arkiv för Zoologi*, **39**(3[8]): 1–39.
- Malaise R.** 1963. Hymenoptera Tenthredinoidea, subfamily Selandriinae. Key to the genera of the World. *Entomologisk Tidskrift*, **84**(3–4): 159–215.
- Malaise R.** 1964. New genera and species of the subfamily Blennocampinae (Hym. Tenthred.). *Entomologisk Tidskrift*, **85**(1–2): 20–39.
- Malikova E.I.** 2012. Family Vespidae. In: *Zhivotnyi mir zapovednika "Bastak"*. Blagoveshchensk: 64–65. [In Russian].

- Malm T. and Nyman T.** 2015. Phylogeny of the symphytan grade of Hymenoptera: new pieces into the old jigsaw(fly) puzzle. *Cladistics*, **31**: 1–17.
- Marikovskaya T.P.** 1979. On the structure and zoogeography of the genus *Clisodon* Patton (Hymenoptera, Anthophoridae). *Izvestiya Akademii Nauk Kazakhstanskoy SSR, Biology*, **2**: 40–48.
- Marikovskaya T.P.** 1982. Bees – pollinators of Agricultural Plants. Nauka, Alma-Ata, 115 p. [In Russian].
- Marikovskaya T.P.** 2001. A contribution to the fauna of bees of the tribe Eucerini (Hymenoptera, Anthophoridae) in deserts of Kazakhstan. *Selevinia*, **2001**(1/4): 129–132.
- Marshakov V.G.** 1975. A review of the genera of the tribe Crabrononi (Hymenoptera, Sphecidae) from the USSR. The genus *Lestica* Billberg, 1820. *Entomologicheskoe Obozrenie*, **54**: 151–163. [In Russian].
- Marshakov V.G.** 1976a. Review of genera of the tribe Crabronini (Hymenoptera, Sphecidae) in the fauna of the USSR. III. Genus *Entomognathus* Dahlbom. *Zoologicheskiy Zhurnal*, **55**(4): 614–618. [In Russian].
- Marshakov V.G.** 1976b. Review of the genera of the tribe Crabronini (Hymenoptera, Sphecidae) from the USSR. The genus *Rhopalum* Stephens, 1829. *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, **67**: 100–112. [In Russian].
- Marshakov V.G.** 1976c. Digger wasps of the genera *Eremiasphecium* Kohl, *Ammoplanus* Gir., *Ammoplanops* Guss. and *Anomiopteryx* Guss. (Hymenoptera, Sphecidae) of the fauna of the USSR and Mongolia. *Entomologicheskoe Obozrenie*, **55**: 668–682. [In Russian].
- Marshakov V.G.** 1977. Review of digger wasps of the tribe Crabronini (Hymenoptera, Sphecidae) of the USSR. Genus *Crabro* Fabricius, 1775. *Entomologicheskoe Obozrenie*, **56**(4): 854–872. [In Russian].
- Marshakov V.G.** 1979. Fossiliferous wasps of the genera *Crossocerus* Lepeletier and Brullé, 1834, *Ectemnius* Dahlbom, 1845, and *Towada* Tsuneki, 1970 (Hymenoptera, Sphecidae) of the Soviet Far East. In.: Lehr P.A. (Ed.). *Nazemnyie Chlenistonogiye Dal'nego Vostoka*. BPI DVNTS AN SSSR, Vladivostok: 90–107. [In Russian].
- Matsumoto R., Fukushima H. and Morimoto K.** 2007. Discovery of Sapygidae (Hymenoptera) in Japan. *Bulletin of the Osaka Museum of Natural History*, **61**: 15–18.
- Matsumura S.** 1911. Erster Beitrag zur Insektenfauna Sachalins. *The journal of the College of Agriculture, Tohoku Imperial University*, **4**: 1–145.
- Mauss V., Fateryga A.V. and Prosi R.** 2016. Taxonomy, distribution and bionomics of *Celonites tauricus* Kostylev, 1935, stat. n. (Hymenoptera, Vespidae, Masarinae). *Journal of Hymenoptera Research*, **48**: 33–66.
- Melo G.A.R.** 1999. Phylogenetic relationships and classification of the major lineages of Apoidea (Hymenoptera), with emphasis on the crabronid wasps. *Scientific Papers, Natural History Museum, The University of Kansas*, **14**: 1–55.
- Menke A.S. and Pulawski W.J.** 2000. A review of the *Sphex flavidipennis* species group (Hymenoptera: Apoidea: Sphecidae: Sphecinii). *Journal of Hymenoptera Research*, **9**: 324–346.
- Mercet R.G.** 1907. El género “*Prosopigastra*”. *Boletín de la Real Sociedad Española de Historia Natural*, **7**: 292–304.
- Mercet R.G.** 1910. Especies europeas del género “*Anastromma*”. *Boletín de la Real Sociedad Española de Historia Natural*, **10**: 125–130.
- Merisuo A.K. and Valkeila E.** 1972. Beiträge zur Kenntnis der paläarktischen Arten der Gattung *Pemphredon* Latreille (Hym., Sphecoidea). *Annales Entomologici Fennici*, **38**: 7–24.
- Michener C.D.** 2007. The bees of the World. Second edition. John Hopkins University Press, Baltimore, xvi + [i] + 953 p. + 20 pls.
- Michez D. and Eardley C.** 2007. Monographic revision of the bee genus *Melitta* Kirby 1802 (Hymenoptera: Apoidea: Melittidae). *Annales de la Société entomologique de France (n.s.)*, **43**(4): 379–440.
- Michez D., Kuhlmann M., Ivanov S.P. and Radchenko V.G.** 2012. Description of four new species in the bee genus *Melitta* Kirby, 1802 (Hymenoptera: Melittidae). *Zootaxa*, **3337**: 57–67.
- Michez D. and Patiny S.** 2005. World revision of the oil-collecting bee genus *Macropis* Panzer 1809 (Hymenoptera: Apoidea: Melittidae) with a description of a new species from Laos. *Annales de la Société entomologique de France (n.s.)*, **41**(1): 15–28.
- Michez D., Terzo M. and Rasmont P.** 2004. Révision des espèces ouest-paléarctiques du genre *Dasyprocta* Latreille 1802 (Hymenoptera, Apoidea, Melittidae). *Linzer biologische Beiträge*, **36**(2): 365–430.
- Mingo E.** 1994. *Hymenoptera Chrysidae*. Fauna Iberica. Vol. 6. Museo Nacional de Ciencias Naturales Consejo Superior de Investigaciones Científicas, Madrid, 256 p.
- Minoranskiy V.A. and Shkuratov A.V.** 1996. On the sphecid fauna (Hymenoptera, Sphecidae) of Rostov Oblast'. *Izvestiya Vysshikh Uchebnykh Zavedeniy. Severo-Kavkazskiy Region. Estestvennye nauki*, **1996**(4): 80–83. [In Russian].
- Minoransky V.A., Kharchenko V.I. and Fomitschев A.L.** 1970. Some data on *Sceliphron destillatorium* Kl. (Sphecidae, Hymenoptera). *Vestnik Zoologii*, **5**: 15–20. [In Russian].
- Mitai K. and Tadauchi O.** 2008. The Genus *Nomada* (Hymenoptera, Apidae) from Kazakhstan and Kyrgyzstan collected by the Kyushu University expedition (1). *Esakia*, **48**: 25–35.
- Mitai K. and Tadauchi O.** 2013. Taxonomic study of the Japanese species of the genus *Sphecodes* (Hymenoptera, Halictidae). *Esakia*, **53**, 21–78.
- Mocsáry A.** 1882. *Chrysidae Faunae Hungaricae*. Academia Hungarica Scientiarum, Budapest, 94 p.

- Mocsáry A.** 1889. *Monographia Chrysidiadarum Orbis Terrarum Universi*. Hungarian Academy of Science, Budapest, 643 p.
- Mocsáry A.** 1909. Chalastogastra nova in collectione Musei nationalis Hungarici. *Annales historico-naturales Musei Nationalis Hungarici*, 7: 1–39.
- Mocsáry A. and Szépligeti V.** 1901. Hymenopteren. In: Horváth G. (Ed.). *Zoologische Ergebnisse der dritten asiatischen Forschungsreise des Grafen Eugen Zichy*. Vol. 2. V. Hornýanský et K.W. Hiersemann, Budapest-Leipzig: 121–169.
- Móczár L.** 1958. Die ungarischen Vertreter der Tribus Oxybelini (Hymenoptera, Sphecidae) unter Berücksichtigung der westpaläarktischen Arten. *Annales Historico-Naturales Musei Nationalis Hungarici. Series Nova*, 9, 50: 281–299.
- Móczár L.** 1964. Ergebnisse der Revision der Goldwespenfauna des Karpatenbeckens (Hymenoptera: Chrysidae). *Acta Zoologica*, 10: 433–450.
- Móczár L.** 1967. *Femdarásalkatúak – Chrysidoidea*. Magyarország Állatvilága. Fauna Hungariae, 86(2). Budapest, 118 p.
- Móczár L.** 1984. New and little known Mesitiinae from Southern Europe and Africa (Hymenoptera: Bethylidae). *Tijdschrift Voor Entomologie*, 127: 101–113.
- Mokrousov M.V.** 2000. Rare and interesting Hymenoptera Vespolomorpha of the Nizhniy Novgorod Oblast'. *Trudy Biologicheskogo Fakulteta Nizhegorodskogo Gosudarstvennogo Universiteta*, 3: 22–25. [In Russian].
- Mokrousov M.V.** 2002. Wasps (Hymenoptera, Vespolomorpha) of Kerzhenskiy Nature Reserve. *Trudy Gosudarstvennogo Prirodnoy Zapovednika "Kerzhenskiy"*, 2: 141–151. [In Russian].
- Mokrousov M.V.** 2003. Sphecidae. In: Anufriev G.A., Bakka S.V. (Eds). *Red book of Nizhniy Novgorod oblast'*. Volume I. Animals. Mezhdunarodnyi Sotsyalno-Ekologicheskiy Soyuz, Nizhniy Novgorod: 220–223. [In Russian].
- Mokrousov M.V.** 2004. New species of digger wasps (Hymenoptera, Crabronidae, Miscophini) from Eastern Europe. *Zoologicheskiy Zhurnal*, 83: 503–507. [In Russian].
- Mokrousov M.V.** 2009. To wasps fauna (Hymenoptera, Vespolomorpha) of the Republic of Mary El. *Nauchnyie Trudy Gosudarstvennogo Prirodnoy Zapovednika "Bolshaya Kokshaga"*, 4: 265–299. [In Russian].
- Mokrousov M.V.** 2010. Fauna of digger wasps (Hymenoptera: Ampulicidae, Sphecidae, Crabronidae) of the Upper and Middle Volga basin. *Proceedings of the Russian Entomological Society*, 81: 59–66. [In Russian].
- Mokrousov M.V.** 2015. New and little known species of digger wasps (Hymenoptera, Apoidea, Sphecidae, Crabronidae) in the fauna of southeastern Europe. In: Chibilyov A.A. (Ed.). *Stepi severnoi Evrazii. Materialy Sed'mogo Mezhdunarodnogo Simpoziuma*. Rossiyskaya Akademiya Nauk, Russkoe Geograficheskoe Obshchestvo, Orenburg: 537–539. [In Russian].
- Mokrousov M.V.** 2017a. New and little-known crabronid wasps of the subfamily Philanthinae (Hymenoptera: Crabronidae) from Russia. *Far Eastern Entomologist*, 335: 12–19.
- Mokrousov M.V.** 2017b. To the knowledge of digger wasps of subfamily Pemphedoninae (Hymenoptera: Crabronidae) of Russia. *Far Eastern Entomologist*, 337: 1–16.
- Mokrousov M.V.** 2017c. Digger wasps (Hymenoptera: Crabronidae) of sandy deserts in the south-east of European Russia. In: Zryanin V.A., Riss A.R. (Eds). Abstracts of the 12th International Symposium of the Russian Society of Nematologists "Nematodes and other Ecdysozoa under the growing ecological footprint on ecosystems" July 31 – August 6, 2017, Nizhny Novgorod, Russia: 89.
- Mokrousov M.V.** 2018. Tiphiid wasps (Hymenoptera: Scoliidae: Tiphidae) of Russia. *Entomological Review*. [In press.]
- Mokrousov M.V., Berezin A.Yu. and Egorov L.V.** 2011. The digger wasps (Hymenoptera: Ampulicidae, Sphecidae, Crabronidae) of the Chuvash Republic. *Eversmannia*, 27–28: 62–86. [In Russian].
- Mokrousov M.V. and Danilov Yu.N.** 2017. Additions and corrections to the fauna and distribution of digger wasps (Hymenoptera: Crabronidae) of Russia. *Proceedings of the Russian Entomological Society*, 88(2). [In press].
- Mokrousov M.V. and Popov I.B.** 2016. Digger wasps (Hymenoptera, Apoidea: Ampulicidae, Sphecidae, Crabronidae) of the Black Sea coast of Krasnodar Territory, Abkhazia, and adjacent areas. *Entomological Review*, 96: 559–599.
- Mokrousov M.V., Ruchin A.B. and Egorov L.V.** 2013. The wasp fauna of Mordovian State Nature Reserve and adjacent territories. *Trudy Mordovskogo Gosudarstvennogo Zapovednika Imeni P.G. Smidovicha*, 11: 193–205. [In Russian].
- Mokrousov M.V. and Selivanova O.V.** 2005. Superfamily Sphecoidea. In: Selivanova O.V. (Ed.). *A list of invertebrates of Voronezh Oblast'*. Voronezhskiy Gosudarstvennyi Universitet, Voronezh: 753–755. [In Russian].
- Mokrousov M.V., Shlyakhtenok A.S. and Grebennikov K.A.** 2016. New and little known crabronid wasps (Hymenoptera: Crabronidae) from Russia. *Far Eastern Entomologist*, 325: 18–32.
- Mokrousov M.V. and Vafin A.R.** 2014. Digger wasps (Hymenoptera: Sphecidae, Crabronidae) of the Republic of Tatarstan, Russia. *Trudy Kazanskogo Otdelenyia Russkogo Entomologicheskogo Obshchestva*, 3: 52–61. [In Russian].
- Monsevičius V.A.** 1995. A checklist of wild bee species (Hymenoptera, Apoidea) of Lithuania with data to their distribution and bionomics In: Jonaitis V. (Ed.). *New and rare for Lithuania insect species. Records and descriptions of 1994–1995*. Institute Ecology, Vilnius: 7–144.
- Morawitz A.** 1864. Verzeichniss der um St.-Petersburg aufgefundenen Crabroninen. *Bulletin de l'Académie Impériale des Sciences de St.-Pétersbourg*, 7: 451–465.

- Morawitz A. 1865.** Über eine neue, oder vielmehr verkannte Form von Männchen unter den Mutillen, nebst einer Übersicht der in Europa beobachteten Arten. *Bulletin de l'Académie Impériale des Sciences de St.-Pétersbourg*, 8: 82–141. (Reprinted in *Mélanges Biologiques tirés du Bulletin de l'Académie Impériale des Sciences de St.-Pétersbourg*, 4: 671–756).
- Morawitz A. 1866.** Einige Bemerkungen über die *Crabro*-artigen Hymenopteren. *Bulletin de l'Académie Impériale des Sciences de St.-Pétersbourg*, 9: 243–273.
- Morawitz F. 1865.** Ueber einige Andrenidae aus der Umgegend von St.-Petersburg. *Horae Societatis Entomologicae Rossicae*, 3(1): 61–79.
- Morawitz F. 1867.** Übersicht der im Gouvernement von Saratow und um St. Petersburg vorkommenden *Odynerus*-Arten. *Horae Societatis Entomologicae Rossicae*, 4: 109–144.
- Morawitz F. 1869.** Die Bienen des Gouvernements von St.-Petersburg. *Horae Societatis Entomologicae Rossicae*, 6(2): 27–71.
- Morawitz F. 1870.** Beitrag zur Bienenfauna Russlands. *Horae Societatis Entomologicae Rossicae*, 7(2/3): 305–320.
- Morawitz F. 1874.** Die Bienen Daghestans. *Horae Societatis Entomologicae Rossicae*, 10(2/4): 129–189.
- Morawitz F. 1890.** Hymenoptera Fossoria Transcaspia Nova. *Horae Societatis Entomologicae Rossicae*, 24: 570–645.
- Morawitz F. 1891.** Ueber Astrachan'sche Fossoria. *Horae Societatis Entomologicae Rossicae*, 25: 175–233.
- Morawitz F. 1892.** Hymenoptera Aculeata rossica nova. *Horae Societatis Entomologicae Rossicae*, 26: 132–181.
- Morawitz F. 1893.** Kareliens Fossoria. *Horae Societatis Entomologicae Rossicae*, 27: 95–115.
- Morawitz F. 1894.** Beitrag zur Raubwespen fauna Turkmeniens. *Horae Societatis Entomologicae Rossicae*, 28: 327–365.
- Morawitz F. 1895.** Materialien zu einer Vespidenfauna des Russischen Reiches. *Horae Societatis Entomologicae Rossicae*, 29: 407–493.
- Möschler A. 1938.** Ein Beitrag zur Bienenfauna in Ostpreußen, insbesondere der Kurischen Nehrung. *Schriften der Physikalisch-ökonomischen Gesellschaft zu Königsberg*, 70(2): 243–288.
- Muche W.H. 1965a.** 51. Tenthredininae. Ergebnisse der zoologischen Forschungen von Dr. Z. Kaszab in der Mongolei (Hymenoptera). *Reichenbachia*, 7(14): 117–126.
- Muche W.H. 1965b.** Mongolische Dolerini (Tenthredinidae, Dolerinae). *Reichenbachia*, 5(20): 181–189.
- Muche W.H. 1968a.** Beitrag zur Kenntnis der Blattwespen des Zentralkaukasus. *Faunistische Abhandlungen Staatliches Museum für Tierkunde Dresden*, 2(11): 85–93.
- Muche W.H. 1968b.** Tenthredinidae der Mongolisch-Deutschen biologischen Expedition 1964 (Hymenoptera). *Reichenbachia*, 10(25): 215–218.
- Muche W.H. 1969.** 3. Beitrag zur Kenntnis der Symphyten des Kaukasus (Hymenoptera). *Faunistische Abhandlungen Staatliches Museum für Tierkunde Dresden*, 2(22): 153–171.
- Muche W.H. 1971a.** Beitrag zur Kenntnis transkaukasischer Hymenopteren. *Entomologische Nachrichten (Dresden)*, 15(9–10): 99–103.
- Muche W.H. 1971b.** Die entomologisch fast unbekannte Umgebung von Ordshonikidse (Kaukasus). *Entomologische Nachrichten (Dresden)*, 15(6): 53–55.
- Muche W.H. 1972.** Beitrag zur Kenntnis der Argidae-Gattungen *Sterictophora* Billberg und *Aprosthemma* Konow (Hymenoptera). *Entomologische Nachrichten (Dresden)*, 16(11): 137–144.
- Muche W.H. 1973a.** 4. Beitrag zur Kenntnis der Symphyten des Kaukasus (Hymenoptera). *Faunistische Abhandlungen Staatliches Museum für Tierkunde Dresden*, 4(10): 77–97.
- Muche W.H. 1973b.** Beitrag zur Blattwespenfauna des Elbrus-Gebietes (Hymenoptera Symphyta). *Faunistische Abhandlungen Staatliches Museum für Tierkunde Dresden*, 4(23): 219–226.
- Muche W.H. 1973c.** Die Blattwespen meiner Mittelasienausbeute 1972. *Entomologische Nachrichten (Dresden)*, 17(11/12): 161–170.
- Muche W.H. 1973d.** Zwei neue kaukasische Symphyten aus der Verwandtschaft Schweizer Endemismen. *Entomologische Nachrichten (Dresden)*, 17(6): 86–90.
- Muche W.H. 1975a.** Eine neue Unterart von *Aglaostigma langei* (Konow) aus dem Kaukasus (Hymenoptera Symphyta). *Faunistische Abhandlungen Staatliches Museum für Tierkunde Dresden*, 5(9): 239–240.
- Muche W.H. 1975b.** Die Symphyten meiner Mittelasienreise 1975 (Hymenoptera, Symphyta). *Faunistische Abhandlungen Staatliches Museum für Tierkunde Dresden*, 5(13): 289–295.
- Muche W.H. 1977.** Die Argidae von Europa, Vorderasien und Nordafrika (mit Ausnahme der Gattung *Aprosthemma*) (Hymenoptera, Symphyta). *Entomologische Abhandlungen*, 41(Suppl.): 23–59.
- Müller A. 2017.** Palaearctic Osmiine Bees, ETH Zürich, <http://blogs.ethz.ch/osmiini> (accessed 18 July 2017).
- Murao R. 2015.** Notes on the distribution of the genus *Lasioglossum* (Hymenoptera: Halictidae) in Eastern Asia. *Japanese Journal of Systematic Entomology*, 21(1): 1–3.
- Murao R. 2017.** Bees of the *Hemihalictus* series of *Lasioglossum* Curtis (Hymenoptera, Halictidae) in South Korea. *Zootaxa*, 4268(4): 451–488.
- Murao R. and Tadauchi O. 2007.** Revision of the subgenus *Evylaeus* of the genus *Lasioglossum* in Japan (Hymenoptera, Halictidae). Part I. *Esakia*, 47: 169–254.
- Muskovits J. and György Z. 2011.** *Velvet ants of Hungary / Magyarország hangyadarazsai*. Kaposvár, Budapest, 198 p. (Natura Somogyensis, 18).

- Nagy C.G.** 1967. Systematisches Studium der Tiphiden Rumäniens (Hymenoptera, Tiphidae). *Reichenbachia*, 8: 175–204.
- Nagy C.G.** 1968. A new record of Mesitinae (Hymenoptera, Bethylidae). *Societá Entomologica Italiana. Memoire*, 47: 168–176.
- Nagy C.G.** 1971. First record of the Old World species of *Sierolomorpha* Ashm. (Hym., Heterogynoidea). *Reichenbachia*, 13 (27): 247–249.
- Naito T.** 1971a. A revision of the genus *Hemitaxonius* in the Old World, I (Hymenoptera, Tenthredinidae). *Kontyû*, 39(1): 19–28.
- Naito T.** 1971b. A revision of the genus *Hemitaxonius* in the Old World, II (Hymenoptera, Tenthredinidae). *Kontyû*, 39(2): 77–82.
- Naito T.** 1980. Studies on the Japanese sawflies of the genus *Strongylogaster* Dahlbom (Hymenoptera, Tenthredinidae). *Kontyû*, 48(3): 390–401.
- Naito T.** 1990. Sawflies of the genus *Strongylogaster* (Hymenoptera, Tenthredinidae) from the Baikal region, USSR, with description of two new species. *Japanese Journal of Entomology*, 58(1): 75–80.
- Naito T.** 2004. Species diversity of sawflies in Hyogo Prefecture, Central Japan. *Monograph of Nature and Human Activities, Hyogo*, 1: 1–87.
- Naito T., Smith D.R. and Huang F.-S.** 1998. Records of Cephidae (Hymenoptera) from China and Southeastern Asia with two new species of *Janus* Stephens. *Japanese Journal of Systematic Entomology, Matsuyama*, 4(2): 237–242.
- Nakonechnaya I.V., Ivchenko V.M. and Kostyukov V.V.** 2014. Noctuids of soybean (*Glycine max*): species and parasite complex in the Krasnodar territory. *Biological Plant Protection as the Basis of Ecosystem Stabilization. Vol. 8*. All-Russian Research Institute of Biological Plant Protection RAAS, Krasnodar: 458–465. [In Russian].
- Nemkov P.G.** 1986. On the digger wasp fauna (Hymenoptera, Sphecidae) of Lake Baikal area. In: Lehr P.A. (Ed.). *Hymenopterous Insects of Eastern Siberia and Far East*. BPI DVNTS AN SSSR, Vladivostok: 92–110. [In Russian].
- Nemkov P.G.** 1988. Review of the Holarctic species of the genus *Tracheliodes* A. Morawitz with descriptions of two new species from the Far East of USSR. *Trudy Vsesoyuznogo Entomologicheskogo Obshchestva*, 70: 116–124. [In Russian].
- Nemkov P.G.** 1989. Digger wasps of the genus *Gorytes* of Siberia and Far East of CCCP. *Zoologicheskiy Zhurnal*, 68: 66–73. [In Russian].
- Nemkov P.G.** 1990a. New and little known species of digger wasps (Hymenoptera, Sphecidae) of Siberia and Far East. In: Lelej A.S. (Ed.). *Novosti sistematiki nasekomykh Dalnego Vostoka*. BPI DO AN SSSR, Vladivostok: 79–85. [In Russian].
- Nemkov P.G.** 1990b. Digger wasps of the tribe Gorytini (Hymenoptera, Sphecidae) of the fauna of the USSR. Genera *Gorytes* Latreille, *Pseudoplatus* Ashmead, *Kohlia* Handlirsch. *Entomologicheskoe Obozrenie*, 69(3): 675–690. [In Russian].
- Nemkov P.G.** 1992a. Digger wasps of the tribe Gorytini (Hymenoptera, Sphecidae) of the fauna of Russia and neighbouring countries. The genera *Lestiphorus* Lepeletier, *Oryttus* Spinola and *Olgia* Radoszkowski. *Entomologicheskoe Obozrenie*, 71(4): 935–949. [In Russian].
- Nemkov P.G.** 1992b. Sphecidae. In: Chistyakov Yu.A. (Ed.). *Insects of Khingan Nature Reserve. Part 2*. Dalnauka, Vladivostok: 243–251. [In Russian].
- Nemkov P.G.** 1995a. Sphecid wasps of the tribe Gorytini (Hymenoptera, Sphecidae) of the fauna of the CIS. The genera *Argogorytes* Ashmead, *Hoplisoides* Gribodo, *Psammaecius* Lepeletier. *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, 258: 128–137. [In Russian].
- Nemkov P.G.** 1995b. Digger wasps of the tribe Gorytini (Hymenoptera, Sphecidae) of the fauna of Russia and neighbouring countries. Genera *Sphecius* Dahlbom and *Ammatomus* A. Costa. *Entomologicheskoe Obozrenie*, 74(1): 177–185. [In Russian].
- Nemkov P.G.** 1996. Digger wasps of the tribe Gorytini (Hymenoptera, Sphecidae) in the fauna of Russia and neighbouring countries. Genus *Harpactus* Shuckard. *Zoologicheskiy Zhurnal*, 75(8): 1204–1213. [In Russian].
- Nemkov P.G.** 2001. Review of the digger wasps of the tribe Nyssonini (Hymenoptera: Sphecidae, Bembicinae) of Russia and neighboring countries. 1. Genus *Synneurus* A. Costa. *Far Eastern Entomologist*, 98: 1–11.
- Nemkov P.G.** 2003. Review of the digger wasps of the genus *Brachystegus* A. Costa (Hymenoptera, Crabronidae, Bembicinae) of Russia and neighboring countries. *Far Eastern Entomologist*, 131: 1–5.
- Nemkov P.G.** 2004. Contribution to the knowledge of the species of *Crossocerus* Lepeletier et Brullé (Hymenoptera; Crabronidae, Crabroninae) described by K. Tsuneki. *Proceedings of the Russian Entomological Society*, 75: 263–269. [In Russian].
- Nemkov P.G.** 2005a. Digger wasp fauna (Hymenoptera, Sphecidae, Crabronidae) of Sakhalin Island. In: *Flora and fauna of Sakhalin Island. Part 2*. Dalnauka, Vladivostok: 141–167. [In Russian].
- Nemkov P.G.** 2005b. Review of the digger wasps of the genus *Palarus* Latreille in Russia and neighbouring countries (Hymenoptera, Crabronidae, Larrinae). *Entomofauna*, 26: 241–252.
- Nemkov P.G.** 2006a. Contribution to the fauna of digger wasps (Hymenoptera: Sphecidae, Crabronidae) of Kedrovaya Pad' Nature Reserve. In: Makarchenko A. (Ed.). *Flora and Fauna of Kedrovaya Pad Nature Reserve*. Dalnauka, Vladivostok: 166–170. [In Russian].

- Nemkov P.G. 2006b.** Digger wasps (Hymenoptera, Crabronidae) of Moneron Island. In: Storozhenko S.Yu. (Ed.). Flora and Fauna of Moneron Island (Materials of International Sakhalin Island Project). Dalnauka, Vladivostok: 247–249. [In Russian].
- Nemkov P.G. 2007.** The digger wasp fauna (Hymenoptera: Sphecidae, Crabronidae) of Kuril Islands. *Euroasian Entomological Journal*, 6: 67–76. [In Russian].
- Nemkov P.G. 2008a.** The digger wasp fauna (Hymenoptera: Sphecidae, Crabronidae) of the Asiatic part of Russia. *A.I. Kurentsov's Annual Memorial Meetings*, 19: 15–34. [In Russian].
- Nemkov P.G. 2008b.** A key to the digger wasps of the genus *Mellinus* (Hymenoptera, Crabronidae, Mellininae) of the Russia. *Far Eastern Entomologist*, 185: 19–20.
- Nemkov P.G. 2008c.** New records of digger wasps (Hymenoptera: Sphecidae, Crabronidae) from the Asian part of Russia. *Far Eastern Entomologist*, 187: 10–11.
- Nemkov P.G. 2009.** Annotated catalogue of digger wasps (Hymenoptera: Sphecidae, Crabronidae) of Asian part of Russia. Dalnauka, Vladivostok, 193 p. [In Russian].
- Nemkov P.G. 2012a.** Section Spheciiformes – Digger wasps. In: A. Lelej (Ed.). *Annotated catalog of insects of the Far East of Russia. Vol. I. Hymenoptera*. Dalnauka, Vladivostok: 433–447. [In Russian].
- Nemkov P.G. 2012b.** Digger wasps of the genus *Stizus* Latreille, 1802 (Hymenoptera, Crabronidae, Bembicinae) of the fauna of Russia and neighbouring countries. *Euroasian Entomological Journal*, 11(1): 55–62.
- Nemkov P.G. 2014a.** Annotated catalog of digger wasps (Hymenoptera: Sphecidae, Crabronidae) of Asian part of Russia. Appendix I. *Euroasian Entomological Journal*, 13: 286–289. [In Russian].
- Nemkov P.G. 2014b.** The digger wasps of the genus *Alysosson* Panzer (Hymenoptera: Crabronidae: Bembicinae) of Russia and adjacent territories, with a key to species and new synonymies. *Zootaxa*, 3838: 276–286.
- Nemkov P.G. 2015.** The digger wasps of the genus *Didineis* Wesmael (Hymenoptera: Crabronidae: Bembicinae) of Russia and adjacent territories, with a key to species and new synonymies. *Zootaxa*, 3941(3): 391–400.
- Nemkov P.G. 2016.** Digger wasps of the genus *Bembix* Fabricius, 1775 (Hymenoptera: Crabronidae, Bembicinae) of Russia and adjacent territories. *Far Eastern Entomologist*, 313: 1–34.
- Nemkov P.G., Kazenas V.L., Budrys E.R. and Antropov A.V. 1995.** Superfamily Sphecoidea. Family Sphecidae – Digger wasps. In: Lehr P.A. (Ed.). *Key to the insects of Russian Far East. Vol. IV, pt 1*. Nauka, St Petersburg: 368–480. [In Russian].
- Nesterov M.A. 1992.** Fossilial wasps (Hymenoptera, Sphecidae) of Daurian Reserve. In: Amirkhanov A.M. (Ed.). *Insects of Dauria and adjacent territories. Vol. 1*. Izdatel'stvo Tsentral'noy Nauchnoissledovatel'skoy Laboratorii Okhotnich'ego Khozaystva i Zapovednikov, Moscow: 83–85. [In Russian].
- Neumeyer R., Baur H., Guex G.-D. and Praz Ch. 2014.** A new species of the paper wasp genus *Polistes* (Hymenoptera, Vespidae, Polistinae) in Europe revealed by morphometrics and molecular analyses. *ZooKeys*, 400: 67–118.
- Neumeyer R., Gereys B. and Castro L. 2015.** New data on the distribution of *Polistes bischoffi* Weyrauch, 1937 and *Polistes helveticus* Neumeyer, 2014, a synonym of *Polistes albellus* Giordany Soika, 1976 n. stat. (Hymenoptera: Vespidae). *Boletín de la Sociedad Entomológica Aragonesa*, 57: 205–216.
- Neumeyer R. and Praz C. 2015.** DNA barcoding of the potter wasp genus *Eumenes* Latreille, 1802 (Hymenoptera: Vespidae: Eumeninae) in Switzerland. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, 88(3–4): 307–320.
- Nie H. and Wei M. 2001.** Taxonomical studies on the genus *Eriocampa* (Hymenoptera: Tenthredinidae: Tenthredininae) from China. *Entomotaxonomia*, 23(4): 283–294.
- Nie H. and Wei M. 2004a.** Taxonomic study on the genus *Adamas* Malaise (Hymenoptera: Tenthredinidae). *Entomotaxonomia*, 26(3): 200–210.
- Nie H. and Wei M. 2004b.** Two new species of *Taxoble-nus* from Northwestern China (Hymenoptera: Tenthredinidae). *Journal of Central South Forestry University*, 24(1): 18–22.
- Nie H., Wei M. and Taeger A. 2012.** Revision of the *Siobla metallica* group (Hymenoptera: Tenthredinidae). *Zootaxa*, 3196: 1–49.
- Nikiforuk K.S. 1957.** Bees of the Bashkir ASSR. *Zapiski Bashkirskogo filiala Geographicheskogo obshchestva SSSR*, 1: 139–162. [In Russian].
- Nikol'skaya M.N. 1978.** Chryridoidea. In: Medvedev G.S. (Ed.). *Key to the insects of the European part of the USSR. Vol. III, pt 1*. Nauka, Leningrad: 58–71. [In Russian].
- Niu G. and Wei M. 2013.** Review of *Aneugmenus* Hartig (Hymenoptera: Tenthredinidae) with description of a new species from China and a key to world species (excluding Neotropical). *Entomotaxonomia*, 35(3): 221–232.
- Niu G., Xiao W. and Wei M. 2012.** Seven new species and a key to species of *Siobla* (Hymenoptera: Tenthredinidae) from Shaanxi, China. *Entomotaxonomia*, 34(2): 399–422.
- Niu G., Xiao W. and Wei M. 2015.** Two new species of *Hartigia* Schiødte from China (Hymenoptera: Cephidae). *Zoological Systematics*, 40(2): 223–227.
- Niu Z., Wu Y. and Huang D. 2004.** A taxonomic study on the subgenus *Seladonia* (Hymenoptera: Halictidae) in China with a description of a new species. *Zoological Studies*, 43(4): 647–670.
- Niu Z.Q., Zhu C.D. and Kuhlmann M. 2014.** The bees of the genus *Colletes* (Hymenoptera: Apoidea: Colletidae) from China. *Zootaxa*, 3856(4): 451–483.
- Noblecourt T. 2016.** *Liste systématique des Hyménoptères Symphytes de France*. Office National des Forêts, Laboratoire National d'Entomologie Forestière, Quillan, 94 p.

- Nordström A. and Forsius R.** 1928. Über die Psamnochoridae (Pompilidae) Finlands. *Notulae Entomologicae*, 8: 1–13.
- Noskiewicz J.** 1936. *Die Palearktischen Colletes-Arten*. Wydawnictwo Towarzystwa Naukowego we Lwowie, Lwow, v + 532 p.
- Novgorodova T.A. and Biryukova O.B.** 2011. Some ethological aspects of the trophobiotic interrelations between ants (Hymenoptera: Formicidae) and larvae of the sawfly *Blasticotoma filiceti* (Hymenoptera: Blasticotomidae). *European Journal of Entomology*, 108: 47–52.
- Nylander W.** 1847. Mutillidae, Scoliidae et Sapygidae boreales. *Notiser ur Sällskaps Pro Fauna et Flora Fennica Förhandlingar*, 1: 7–31 + tab. 2.
- Nylander W.** 1859. A list of Hymenoptera and Diptera species collected by G. Appelberg and W. Pipping in Karelia in 1841. *Notiser ur Sällskaps Pro Fauna et Flora Fennica Förhandlingar*, 4: 109–118.
- Ohl M.** 1999. A revision of *Stizoides* Guérin-Méneville, 1844: taxonomy, phylogenetic relationships, biogeography, and evolution (Hymenoptera: Apoidea: "Sphecidae"). *Mitteilungen aus dem Museum für Naturkunde in Berlin, Zoologische Reihe*, 75: 63–169.
- Okutani T.** 1972. A new genus and a key to Japanese genera of the subfamily Blennocampinae (Hym. Tenth.). Studies on Symphyta XXVIII. *Entomological Review of Japan*, 24(1–2): 57–61.
- Olmi M.** 1984. A revision of the Dryinidae (Hymenoptera). *Memoirs of the American Entomological Institute*, 37: I–XII + 1–1913.
- Olmi M.** 1996. A revision of the world Embolemidae (Hymenoptera: Chrysidoidea). *Frustula Entomologica. N.S.*, 1: 85–146.
- Olmi M.** 2003. A contribution to the knowledge of Dryinidae. *Frustula Entomologica. N.S.*, 24(37): 21–49.
- Olmi M.** 2004. Dryinidae (Hym.) from the Kuril Islands, Kamchatka and Sakhalin (Russia), with the description of a new species of Aphelopus Dalman. *The Entomologist's Monthly Magazine*, 140: 301–306.
- Olmi M., Belokobylskij S.A. and Guglielmino A.** 2014a. Revision of the family Embolemidae of Russia and Ukraine (Hymenoptera: Chrysidoidea), with description of a new species. *Annales Zoologici*, 64(1): 97–108.
- Olmi M., Mita T. and Guglielmino A.** 2014b. Revision of the Embolemidae of Japan (Hymenoptera: Chrysidoidea), with description of a new genus and two new species. *Zootaxa*, 3793: 423–440.
- Olmi M. and Xu Z.** 2015. Dryinidae of the Eastern Palearctic region (Hymenoptera: Chrysidoidea). *Zootaxa*, 3996(1): 1–253.
- Ono M.** 1997. *Science of Japanese hornets (Vespa spp.)*. Kaiyusha, Tokyo, 174 p.
- Ortiz-Sánchez F.J. and Roberts P.M.S.** 2007. A new record for *Eucera* (*Eucera*) *dalmatica* Lepeletier, 1841 in the Iberian Peninsula (Hymenoptera, Anthophoridae). *Entomofauna*, 28(5): 61–64.
- Osten T.** 1999. Kritische Liste der palaearktischen Scoliiden (Hymenoptera, Scoliidae). *Entomofauna*, 20(27): 422–428.
- Osten T.** 2005. Checkliste der Dolchwespen der Welt (Insecta: Hymenoptera, Scoliidae). *62. Bericht der Naturforschenden Gesellschaft Augsburg*, 220: 1–62.
- Osytshnjuk A., Romasenko L., Banaszak J. and Cierzniak T.** 2005. *Andreninae of the Central and Eastern Palaearctic. Part 1*. Polish Entomological Monographs. Vol. 2. Polish Entomological Society, Poznań, Bydgoszcz, 235 p.
- Osytshnjuk A., Romasenko L., Banaszak J. and Motyka E.** 2008. *Andreninae of the Central and Eastern Palaearctic. Part 2*. Polish Entomological Monographs. Vol. 5. Polish Entomological Society, Poznań, Bydgoszcz, 233 p.
- Osytshnjuk A.Z.** 1970. *Fauna of the Ukraine. Vol. 12. Bees. No. 4. Family Colletidae*. Naukova Dumka, Kiev: 158 p. [In Ukrainian].
- Osytshnjuk A.Z.** 1977. *Fauna of the Ukraine. Vol. 12. Bees. No. 5. Family Andrenidae*. Naukova Dumka, Kiev, 328 p. [In Ukrainian].
- Osytshnjuk A.Z.** 1982. A new species of the genus *Andrena* F. (Hymenoptera, Apoidea, Andrenidae) from the Primorski territory. In: Lerh P.A. (Ed.). *Hymenopterous Insects of Far East*. DVNTS AN SSSR, Vladivostok: 113–116. [In Russian].
- Osytshnjuk A.Z.** 1983. Three new species of the genus *Andrena* (Hymenoptera, Andrenidae) from Kazakhstan. *Vestnik Zoologii*, 1: 20–27. [In Russian].
- Osytshnjuk A.Z.** 1984. A new Palaearctic subgenus and a new species of the genus *Andrena* (Hymenoptera, Andrenidae). *Vestnik Zoologii*, 18(2): 23–30. [In Russian].
- Osytshnjuk A.Z.** 1986a. New species of the subgenus *Euandrena* Hed. (Hymenoptera, Andrenidae, *Andrena* F.) from the Far East. In: Lerh P.A. (Ed.). *Hymenopterous Insects of Eastern Siberia and Far East*. BPI DVNTS AN SSSR, Vladivostok: 111–116. [In Russian].
- Osytshnjuk A.Z.** 1986b. New Palaearctic species of the subgenus *Euandrena* Hed. (Hymenoptera, Andrenidae, *Andrena* F.). *Entomologicheskoe Obozrenie*, 65(2): 407–418. [In Russian].
- Osytshnjuk A.Z.** 1993a. New Palaearctic subgenera and species of the genus *Andrena* (Hymenoptera, Andrenidae). Report 2. *Vestnik Zoologii*, 5: 60–66. [In Russian].
- Osytshnjuk A.Z.** 1993b. New Palaearctic species of the bee subgenus *Micrandrena* Ashmead (Hymenoptera, Andrenidae, *Andrena*). *Entomologicheskoe Obozrenie*, 72(2): 401–409. [In Russian].
- Osytshnjuk A.Z.** 1994a. New subspecies of the Palaearctic *Andrena* bees (Hymenoptera, Andrenidae). *Vestnik Zoologii*, 1: 30–36. [In Russian].
- Osytshnjuk A.Z.** 1994b. New Palaearctic subgenera and species of the genus *Andrena* (Hymenoptera, Andrenidae). Report 3. *Vestnik Zoologii*, 4/5: 17–23, 34. [In Russian].

- Osytshnjuk A.Z.** 1995. Family Andrenidae. In: Lehr P.A. (Ed). *Key to the insects of Russian Far East*. Vol. 4, pt 1. Nauka, St. Petersburg: 489–527. [In Russian].
- Osytshnjuk A.Z., Panfilov D.V. and Ponomareva A.A.** 1978. Superfamily Apoidea. In: Medvedev G.S. (Ed.). *Key to the insects of the European part of the USSR*. Vol. III, pt 1. Nauka, Leningrad: 279–519. [In Russian].
- Osytshnjuk A.Z. and Romankova T.G.** 1995. Family Colletidae. In: Lehr P.A. (Ed). *Key to the insects of Russian Far East*. Vol. 4, pt 1. Nauka, St. Petersburg: 480–489. [In Russian].
- Pagliano G.** 1994. Catalogo degli Imenotteri italiani. (IV Apoidea: Colletidae, Megachilidae, Anthophoridae, Apidae). *Memorie della Società Entomologica Italiana*, **72**: 331–467.
- Pagliano G.** 2002. Revisione della sottofamiglia *Apterogyninae* (Hymenoptera: Bradynobaenidae). Museo Regionale di Scienze Naturali di Torino, Monografie, 34, 387 p.
- Pagliano G. and Strumia F.** 2007. Contributo alla conoscenza dei Mutilidae italiani (Hymenoptera, Scolioidea). *Bulletino del Museo Regionale di Scienze Naturali, Torino*, **24**(1): 25–110.
- Pallas P.S.** 1771. *Reise durch verschiedene Provinzen des Russischen Reichs. Erster Theil*. Kayserlichen Akademie der Wissenschaften, St. Petersburg, x + 3–504 p. + pls. 1–11.
- Pallas P.S.** 1773. *Reisen durch verschiedene Provinzen des Russischen Reichs. Zweiter Theil*. Zweites Buch vom Jahr 1771. Kayserlichen Academie der Wissenschaften, St. Petersburg, [12] + 744 s.
- Panfilov D.V.** 1951. Bumblebees of the subgenus *Cullumanobombus* Vogt (Hymenoptera, Apoidea). *Trudy Vsesoyuznogo entomologicheskogo obshchestva*, **43**: 115–128. [In Russian].
- Panfilov D.V.** 1954. Apterogins (Hymenoptera: Apterogynidae) of the USSR fauna. *Proceedings of the Zoological Institute of the USSR Academy of Sciences*, **15**: 146–153. [In Russian].
- Panfilov D.V.** 1956. Contribution to the taxonomy of bumblebees (Hymenoptera, Bombinae), including the description of new forms. *Zoologicheskiy Zhurnal*, **35**(9): 1325–1334. [In Russian].
- Panfilov D.V.** 1957. On the geographical distribution of bumblebees (*Bombus*) in China. *Acta Geographica Sinica*, **23**(3): 221–239.
- Panfilov D.V.** 1981. Map 91: Genus *Bombus* Latreille, 1802. Map 92: *Bombus lapidarius* (Linnaeus, 1758). Map 93: *Bombus hortorum* (Linnaeus, 1761). Map 94: *Bombus agrorum* (Fabricius, 1787). Map 95: *Bombus terrestris* (Linnaeus, 1758). Map 96: *Bombus lucorum* (Linnaeus, 1761). Map 97: *Bombus pratorum* (Linnaeus, 1761). In: Gorodkov K.B. (Ed.). *Provisional Atlas of the Insects of the European Part of U.S.S.R. Maps 73–125*. Nauka, Leningrad: 22–28. [In Russian].
- Panfilov D.V.** 1982. Map 147: *Bombus silvarum* (Linnaeus, 1761). Map 148: *Bombus equestris* (Fabricius, 1793). Map 149: *Bombus distinguendus* Morawitz, 1869. Map 150: *Bombus jonellus* (Kirby, 1802). In: Gorodkov K.B. (Ed.). *Provisional Atlas of the Insects of the European Part of U.S.S.R. Maps 126–178*. Nauka, Leningrad: 25–28. [In Russian].
- Panfilov D.V.** 1984. Map 186: *Bombus hypnorum* (Linnaeus, 1758). Map 187: *Bombus muscorum* (Fabricius, 1775). Map 188: *Bombus balteatus* Dahlbom, 1832. Map 189: *Bombus proteus* Gerstaecker, 1869. Map 190: *Bombus confusus* Schenck, 1859. Map 191: *Bombus ruderatus* (Fabricius, 1775). Map 192: *Bombus fragrans* (Pallas, 1771). *Provisional Atlas of the Insects of the European Part of U.S.S.R. Maps 179–221*. Nauka, Leningrad: 28–35. [In Russian].
- Panfilov D.V., Rossolimo O.L. and Syroechkovskii E.E.** 1961. Contribution to the bumblebees fauna (Bombinae) of Tyva. *Izvestiya Sibirskogo Otdeleniya AN SSSR. Seriya Biologicheskikh nauk*, **6**: 106–113. [In Russian].
- Pankov G.I.** 2013. Digger wasps of Ivanovsk Oblast', Russia. *Krayevedcheskiye Zapiski (Ivanovo)*, **14**: 226–232. [In Russian].
- Pate V.S.** 1946. The generic names of the spider wasps (Psammocharidae olim Pompilidae) and their type species. *Transactions of the American Entomological Society*, **72**: 65–137.
- Pate V.S.** 1947a. Neotropical Sapygidae with conspectus of the family (Hymenoptera: Aculeata). *Acta Zoologica Lilloana*, **4**: 393–426.
- Pate V.S.** 1947b. The generic names of Sapygidae and their type species (Hymenoptera: Aculeata). *Entomological News*, **57**(9): 219–221.
- Paukkunen J., Berg A., Soon V., Ødegaard F. and Rosa P.** 2015. An illustrated key to the cuckoo wasps (Hymenoptera, Chrysididae) of the Nordic and Baltic countries, with description of a new species. *ZooKeys*, **548**: 1–116.
- Paukkunen J. and Kozlov M.V.** 2015. Stinging wasps, ants and bees (Hymenoptera: Aculeata) of the Murmansk region, Northwest Russia. *Entomologica Fennica*, **26**: 53–73.
- Paukkunen J., Rosa P., Soon V., Johansson N. and Ødegaard F.** 2014. Faunistic review of the cuckoo wasps of Fennoscandia, Denmark and the Baltic countries (Hymenoptera: Chrysididae). *Zootaxa*, **3864**(1): 1–67.
- Pavlusenko I.N.** 2006. A historical review of the research on Cephid sawflies (Hymenoptera: Symphyta: Cephidae) in Ukraine. *The Kharkov Entomological Society Gazette*, **13**(1–2): 168–171.
- Perkins J.F.** 1976. Hymenoptera, Bethyloidea (excluding Chrysididae). *Handbook for the identification of British Insects*, **6**(3a): 1–38.
- Perrard A., Grimaldi D. and Carpenter J.M.** 2017. Early lineages of Vespidae (Hymenoptera) in Cretaceous amber. *Systematic Entomology*, **42**(2): 379–386.

- Pesarini P., Campadelli G. and Grudele G.** 1995. Imenotteri sinfitti delle foreste Demaniali Casentinesi a aree circostanti. (Materiali per una sinfitofauna appenninica. I). *Bollettino dell'Istituto di entomologia G. Grandi dell'Università di Bologna*, **50**: 29–55.
- Pesenko Yu.A.** 1971. Contributions to the fauna and ecology of bees (Hymenoptera, Apoidea) in steppes of the Lower Don basin. Report I. The family Megachilidae. *Entomologicheskoe Obozrenie*, **50**(1): 66–78. [In Russian].
- Pesenko Yu.A.** 1972. Contributions to the fauna and ecology of bees (Hymenoptera, Apoidea) in steppes of the Lower Don basin. Report II. The family Halictidae. *Entomologicheskoe obozrenie*, **51**(2): 282–295. [In Russian].
- Pesenko Yu.A.** 1974. Contributions to the fauna and ecology of bees (Hymenoptera, Apoidea) in the Lower Don basin. Report IV. The family Anthophoridae. *Entomologicheskoe Obozrenie*, **53**(2): 324–333. [In Russian].
- Pesenko Yu.A.** 1985. The taxonomy of bees of the genus *Halictus* Latreille (Hymenoptera, Halictidae) with description of the 7th and 8th metasomal sterna of males: subgenus *Monilapis* Cockerell. *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, **132**: 77–105. [In Russian].
- Pesenko Yu.A.** 1986. An annotated key to the Palaearctic species of bees of the genus *Lasioglossum* sensu stricto (Hymenoptera, Halictidae) for females, with descriptions of new subgenera and species. *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, **159**: 113–151. [In Russian].
- Pesenko Yu.A.** 2005a. New data on the taxonomy and distribution of the Palaearctic halictids: genus *Halictus* Latreille (Hymenoptera: Halictidae). *Entomofauna*, **26**(18): 313–348.
- Pesenko Yu.A.** 2005b. Contributions to the halictid fauna of the Eastern Palaearctic Region: subfamily Nomioidinae (Hymenoptera: Halictidae). *Far Eastern Entomologist*, **152**: 1–12.
- Pesenko Yu.A.** 2006a. Contributions to the halictid fauna of the Eastern Palaearctic Region: genus *Seladonia* Robertson (Hymenoptera: Halictidae, Halictinae). *Esakia*, **46**: 53–82.
- Pesenko Yu.A.** 2006b. Contributions to the halictid fauna of the Eastern Palaearctic Region: genus *Lasioglossum* Curtis (Hymenoptera: Halictidae, Halictinae). *Zoosystematica Rossica*, **15**(1): 133–166.
- Pesenko Yu.A.** 2007a. A taxonomic study of the bee genus *Evylaeus* Robertson of Eastern Siberia and Far East of Russia (Hymenoptera: Halictidae). *Zoosystematica Rossica*, **16**(1): 79–123.
- Pesenko Yu.A.** 2007b. Subfamily Halictidae. Introduction; Subfamily Halictinae. In: Lelej A.S. (Ed.). *Key to the insects of Russian Far East*. Vol. IV, pt 5. Dalnauka, Vladivostok: 745–754, 760–773; 824–878. [In Russian].
- Pesenko Yu.A., Banaszak J., Radchenko V.G. and Cierznia T.** 2000. *Bees of the family Halictidae (excluding Sphecodes) of Poland: taxonomy, ecology, bionomics*. Pedagogical University, Bydgoszcz, 348 p.
- Pesenko Yu.A. and Situdikov A.A.** 1988. The classification and phylogenetic relationships of the genera of the bee tribe Eucerini (Hymenoptera, Anthophoridae) with two submarginal cells. *Entomologicheskoe Obozrenie*, **67**(4): 846–860. [In Russian].
- Petersen B.** 1988. The Palaearctic Mutillidae of I.C. Fabricius and some related material (Insecta, Hymenoptera, Aculeata). *Steenstrupia*, **14**(6): 129–224.
- Petrov D.L. and Sautkin F.V.** 2013. Insect gall – pests of shrubs of green spaces of Belarus. *Vestnik Belarusskogo gosudarstvennogo universiteta, Seriya 2*, **1**: 65–71. [In Russian].
- Pfeiffer M., Schultz R., Radchenko A., Yamane S., Woyciechowski M., Aibek U. and Seifert B.** 2007. A critical checklist of the ants of Mongolia (Hymenoptera: Formicidae). *Bonner Zoologische Beiträge*, **55**: 1–8.
- Pham P.H. and Kumar P.G.** 2016. Taxonomic notes on the genus *Rhynchium* Spinola (Hymenoptera: Vespidae: Eumeninae) from Vietnam, with a first checklist of the species worldwide. *Ecologica Montenegrina*, **7**: 530–537.
- Piotrowski W., Rozwałka R. and Sołytk D.** 2006. Data on the occurrence of some species of Xyelidae, Pamphilidae, and Megalodontesidae (Hymenoptera: Symphyta) in Poland. *Wiadomości Entomologiczne*, **25**(2): 89–96.
- Pitts J.P., Wasbauer M.S. and von Dohlen C.D.** 2006. Preliminary morphological analysis of relationships between the spider wasp subfamilies (Hymenoptera: Pompilidae): revisiting an old problem. *Zoologica Scripta*, **35**: 63–84.
- Plavil'shchikov N.N.** 1964. List of insects found in Mordovian State Nature Reserve. *Trudy Mordovskogo Gosudarstvennogo Zapovednika imeni P.G. Smidovicha*, **2**: 105–134. [In Russian].
- Polaszek A.** 2011. Fauna Europaea: Bethylidae // de Jong Y. (ed.). Fauna Europaea: Hymenoptera. Fauna Europaea version 2.4. <http://www.faunaeur.org> (accessed 15 February 2017).
- Polumordvinov O.A. and Shibaev S.V.** 2015. To the Steppe fauna of insects in the Penza Region. In: Steppes of the North Eurasia. Materials of the 7th international symposium. Dimur, Orenburg: 669–672. [In Russian].
- Ponomarenko N.G.** 1965. New parasitic Hymenoptera of the subfamily Gonatopodinae (Hymenoptera, Dryinidae) in the USSR fauna. *Entomologicheskoe Obozrenie*, **44**: 622–631. [In Russian].
- Ponomarenko N.G.** 1966. New Gonatopodinae of the USSR fauna (Hymenoptera, Dryinidae). *Zoologicheskiy Zhurnal*, **45**(10): 1578–1580. [In Russian].
- Ponomarenko N.G.** 1970. New species of Dryinids (Hymenoptera, Dryinidae) in the Europeans areas of the USSR. *Entomologicheskoe Obozrenie*, **49**: 423–433. [In Russian].

- Ponomarenko N.G.** 1972. On the fauna of dryinids (Hymenoptera, Dryinidae) of the Mongolian People's Republic. In: Kerzner I.M. (Ed.). *Insects of Mongolia*. Vol. 1. Nauka, Leningrad, 1: 673–680. [In Russian].
- Ponomarenko N.G.** 1975. On the fauna of dryinids (Hymenoptera, Dryinidae) of the Mongolian People's Republic, II. In: Kerzner I.M. (Ed.). *Insects of Mongolia*. Vol. 3. Nauka, Leningrad: 314–319. [In Russian].
- Ponomarenko N.G.** 1978. Family Dryinidae – Dryinids. In: Medvedev G.S. (Ed.). *Key to the Insects of the European part of USSR*. Vol. III, pt 2. Nauka, Leningrad: 16–27. [In Russian].
- Ponomarenko N.G.** 1979. On the fauna of dryinids (Hymenoptera, Dryinidae) of the Mongolian People's Republic, III. In: Kerzner I.M. (Ed.). *Insects of Mongolia*. Vol. 6. Nauka, Leningrad: 354–361. [In Russian].
- Ponomarenko N.G.** 1992. On the fauna of Dryinids (Hymenoptera, Dryinidae) of Siberia and Far East. *Entomologicheskoe Obozrenie*, 71: 929–934. [In Russian].
- Ponomarenko N.G.** 1995. Family Dryinidae – Dryinids. In: Lehr P.A. (Ed.). *Key to the Insects of the Russian Far East*. Vol. IV, pt 1. Nauka, St Petersburg: 153–175. [In Russian].
- Ponomarenko N.G.** 2000. A new species of the genus *Mystrophorus* Förster (Hymenoptera, Dryinidae) from Russia. *Entomologicheskoe Obozrenie*, 79: 204–206. [In Russian].
- Ponomarenko N.G. and Olmi M.** 2007. New records of Dryinidae (Hymenoptera: Chrysidoidea) from Russia, Ukraine, Moldova, Georgia, Armenia and Azerbaijan. *Entomologist's Gazette*, 58: 51–58.
- Ponomareva A.A.** 1966. On some little-known species of the genus *Anthophora* s. l. (Hymenoptera, Apoidea) from the USSR. *Entomologicheskoe Obozrenie*, 45(1): 155–167. [In Russian].
- Ponomareva A.A.** 1967. On trophic links of some bees of the subfamily Anthophorinae and of main pollinators (Hymenoptera, Apoidea) of leguminous plants in central Kazakhstan and Middle Asia. *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, 38: 330–365. [In Russian].
- Popov A.A.** 2011a. Synopsis of dendro-tamnobiont sawflies of family Tenthredinidae (Hymenoptera, Symphyta) of Yakutia. *Proceedings of the Russian Entomological Society*, 82: 77–88. [In Russian].
- Popov A.A.** 2011b. A review of sawflies of the genus *Alantus* Panzer, 1801 (Hymenoptera, Tenthredinidae) for Yakutia (Eastern Siberia). *Euroasian Entomological Journal*, 10(2): 191–196. [In Russian].
- Popov A.A.** 2013. The sawflies of the family Diprionidae (Hymenoptera, Symphyta) of Yakutia (Eastern Siberia). *Vestnik Mordovskogo Universiteta, Seria: Biologicheskie nauki*, 3–4: 125–128. [In Russian].
- Popov A.A.** 2015. Sawflies of the Family Pamphiliidae (Hymenoptera, Symphyta) of Yakutia. *Nauka i Obrazovanie*, 4(80): 133–137. [In Russian].
- Popov A.A.** 2016. Survey of the fauna and ecology of sawflies (Hymenoptera, Symphyta) of the Yakut botanical garden. *Nauka i Obrazovanie*, 4(84): 129–133. [In Russian].
- Popov A.A. and Kajmuk E.L.** 2007. About sawflies of family Argidae (Symphyta, Tenthredinoidea) fauna of Yakutia. *Nauka i Obrazovanie*, 2(46): 31–34. [In Russian].
- Popov A.A. and Kajmuk E.L.** 2008. The fauna of sawflies of tribe Dineurini (Hymenoptera, Tenthredinidae) of Yakutia. *Trudy Kemerovskogo Otdeleniya Russkogo Entomologicheskogo Obshchestva*, 6: 95–99. [In Russian].
- Popov A.A. and Kajmuk E.L.** 2009a. Materials on the fauna of sawflies (Hymenoptera, Tenthredinoidea) of resource reserve "Pil'ka" (Southwest Yakutia). In: Bagashov A.N. (Ed.). *Chelovek i Sever: Antropologiya, arkeologiya, ekologiya: Materialy vserossiyskoy konferentsii*. IPOS SO RAN, Tyumen: 269–271. [In Russian].
- Popov A.A. and Kajmuk E.L.** 2009b. The sawfly of tribes Macrophyini (Hymenoptera, Symphyta) of Yakutia. *Sovremennye problemy bioraznoobraziya. Materialy Mezhdunarodnoy nauchnoy konferentsii*, IPC VGU, Voronezh: 315–318. [In Russian].
- Popov A.A. and Kajmuk E.L.** 2010a. On the sawflies of the Cimbicidae family (Hymenoptera, Symphyta) of the fauna of Yakutia. *Entomologicheskie issledovaniya v Severnoy Azii. Materialy VIII Mezhdunarodnogo soveshchaniya entomologov Sibiri i Dal'nego Vostoka s uchastiem zarubezhnykh uchenykh*, Novosibirsk: 173–174. [In Russian].
- Popov A.A. and Kajmuk E.L.** 2010b. The sawflies of the family Tenthredinidae (Hymenoptera: Symphyta) of Yakutia. *Proceedings of the Russian Entomological Society*, 80(4): 67–73. [In Russian].
- Popov V.B.** 1931. Zur Kenntnis der paläarktischen Schmarotzerhummeln (*Psithyrus* Lep.). *Eos*, 7(2): 131–209.
- Popov V.B.** 1933a. Notes on the parasitic bees allied to the genus *Biastes* Panz. (Hymenoptera, Nomadidae). *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, 2(1): 51–75. [In Russian].
- Popov V.B.** 1933b. Palaearctic forms of the tribe Stelidini Roberts. (Hymenoptera, Megachilidae). *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, 1(3/4): 375–414. [In Russian].
- Popov V.B.** 1935. Beiträge zur Bienenfauna von Tadzhikistan (Hymenoptera, Apoidea [sic]). *Travaux de la Filiale de l'Académie des Sciences de l'URSS au Tadjikistan*, 5: 351–408.
- Popov V.B.** 1940. A contribution to the knowledge of the Palaearctic species of the genus *Andrena* F. (Hymenoptera, Apoidea). *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, 6(1/2): 252–262. [In Russian].
- Popov V.B.** 1947. Zоogeographical characteristics of the Palaearctic species of the genus *Xylocopa* Latr. (Hymenoptera, Apoidea) and their associations with melittophilous plants. *Izvestiya Akademii Nauk SSSR, Biology*, 1947(1): 29–52. [In Russian].

- Popov V.B.** 1948. The geographical distribution of bees of the genus *Habropoda* F. SM. (Hymenoptera, Anthophoridae). *Doklady Akademii Nauk SSSR* (N.S.), **59**(9): 1673–1676. [In Russian].
- Popov V.B.** 1949. The subgenus *Plastandrena* Hedicke and its new forms (Hymenoptera, Apoidea). *Entomologicheskoe Obozrenie*, **30**(3/4): 389–404. [In Russian].
- Popov V.B.** 1950. On the genus *Amegilla* Friese (Hymenoptera, Apoidea). *Entomologicheskoe Obozrenie*, **31** (1/2): 257–261. [In Russian].
- Popov V.B.** 1951. Parasitic bees of the genus *Ammobates* Latr. (Hymenoptera, Anthophoridae). *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR R*, **9**(3): 895–949. [In Russian].
- Popov V.B.** 1954. On the fauna of bees (Hymenoptera, Apoidea) in the southern part of the Western Kazakhstan Province. *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, **16**: 351–373. [In Russian].
- Popov V.B.** 1956. The geographical distribution and trophic links of the genus *Ctenoplectra* F. Sm. (Hymenoptera, Melittidae). *Doklady Akademii Nauk SSSR*, **108**(5): 969–972. [In Russian].
- Popov V.B.** 1958a. Hymenoptera of the Crimea. In: Vinogradov B.S. (Ed.). *Animal World of the USSR. Vol. 5. Mountain Regions of the European Part of the USSR*. Academy of Sciences of the USSR, Moscow, Leningrad: 100–115. [In Russian].
- Popov V.B.** 1958b. On three subgeneric andrenid groupings (Hymenoptera, Andrenidae). *Trudy Vsesoyuznogo entomologicheskogo obshchestva*, **46**: 109–161. [In Russian].
- Popov V.B.** 1967a. The bees (Hymenoptera, Apoidea) of Middle Asia and their associations with angiosperm plants. *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, **38**: 11–329. [In Russian].
- Popov V.B.** 1967b. The bees (Hymenoptera, Apoidea) of Iran. *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, **43**: 184–216. [In Russian].
- Popov V.B. and Ponomareva A.A.** 1961. On the fauna of the Xylocopini (Hymenoptera, Apoidea) of the Soviet Union. *Entomologicheskoe Obozrenie*, **40**(2): 393–404. [In Russian].
- Popov V.B. and Yasumatsu K.** 1935. Notes on the bee-genus *Pasites* Jurine (Hymenoptera, Nomadidae) with description of a new subspecies of *P. maculatus* Jurine from South Manchuria. *Mushi*, **8**(2): 97–104.
- Praz C.J.** 2017. Subgeneric classification and biology of the leafcutter and dauber bees (genus *Megachile* Latreille) of the western Palearctic (Hymenoptera, Apoidea, Megachilidae). *Journal of Hymenoptera Research*, **55**: 1–54.
- Praz C.J., Müller A., Danforth B.N., Griswold T.L., Widmer A. and Dorn S.** 2008. Phylogeny and biogeography of bees of the tribe Osmiini (Hymenoptera: Megachilidae). *Molecular Phylogenetics and Evolution*, **49**(1): 185–197.
- Priesner H.** 1957. A review of the *Anthophora*-species of Egypt (Hymenoptera: Apidae). *Bulletin de la Société entomologique d'Égypte*, **41**: 1–115.
- Prisniy A.V.** 2012. Digger wasps (Hymenoptera: Sphecidae, Crabronidae) of the Belgorod Oblast'. *Euroasian Entomological Journal*, **11**: 44–54. [In Russian].
- Prokofiev A.M. and Skomorokhov M.O.** 2010. *Sceliphron curvatum* (F. Smith, 1870), a new in the fauna of Russia and invasive species of digger wasps (Hymenoptera: Sphecidae). *Russian Entomological Journal*, **19**: 67–70. [In Russian].
- Proshchalykin M.Yu.** 2004. A check list of the bees (Hymenoptera, Apoidea) of the southern part of the Russian Far East. *Far Eastern Entomologist*, **143**: 1–17.
- Proshchalykin M.Yu.** 2006. Bees of the genus *Coelioxys* Latreille, 1809 (Hymenoptera, Apoidea, Megachilidae) from the Russian Far East. *Euroasian Entomological Journal*, **5**(4): 318–324. [In Russian].
- Proshchalykin M.Yu.** 2007a. Family Colletidae. In: Lelej A.S. (Ed.). *Key to the insects of Russian Far East*. Vol. IV, pt 5. Dalnauka, Vladivostok: 878–883. [In Russian].
- Proshchalykin M.Yu.** 2007b. Family Andrenidae. In: Lelej A.S. (Ed.). *Key to the insects of Russian Far East*. Vol. IV, pt 5. Dalnauka, Vladivostok: 884–886. [In Russian].
- Proshchalykin M.Yu.** 2007c. Family Melittidae. In: Lelej A.S. (Ed.). *Key to the insects of Russian Far East*. Vol. IV, pt 5. Dalnauka, Vladivostok: 886–889. [In Russian].
- Proshchalykin M.Yu.** 2007d. Family Megachilidae. In: Lelej A.S. (Ed.). *Key to the insects of Russian Far East*. Vol. IV, pt 5. Dalnauka, Vladivostok: 889–897. [In Russian].
- Proshchalykin M.Yu.** 2007e. Family Apidae. In: Lelej A.S. (Ed.). *Key to the insects of Russian Far East*. Vol. IV, pt 5. Dalnauka, Vladivostok: 897–908. [In Russian].
- Proshchalykin M.Yu.** 2007f. The bees of family Megachilidae (Hymenoptera, Apoidea) of Transbaikalia. *Far Eastern Entomologist*, **175**: 1–18.
- Proshchalykin M.Yu.** 2008a. The bees of family Colletidae (Hymenoptera, Apoidea) of Transbaikalia. *Far Eastern Entomologist*, **187**: 1–9.
- Proshchalykin M.Yu.** 2008b. The bees of family Melittidae (Hymenoptera, Apoidea) of Transbaikalia. *Far Eastern Entomologist*, **191**: 8–9.
- Proshchalykin M.Yu.** 2009a. The long-tongued bees (Hymenoptera, Apoidea: Megachilidae, Apidae) of Transbaikalia. *Vestnik Kharkovskogo Natsionalnogo Universiteta im. V.N. Karazina. Seriya: biologiya*, **856**(9): 136–142. [In Russian].
- Proshchalykin M.Yu.** 2009b. *Ctenonomia blakistoni* (Sakagami et Munakata, 1990) is newly recorded bee species (Hymenoptera, Apoidea, Halictidae) from Russia. *Far Eastern Entomologist*, **202**: 7–8.
- Proshchalykin M.Yu.** 2010. The short-tongued bees (Hymenoptera, Apoidea: Colletidae, Andrenidae, Halictidae, Melittidae) of Transbaikalia. *Euroasian Entomological Journal*, **9**(3): 508–514.

- Proshchalykin M.Yu.** 2012a. Section Apiformes. In: Lelej A.S. (Ed.). *Annotated Catalogue of the Insects of Russian Far East. Vol. 1. Hymenoptera*. Dalnauka, Vladivostok: 448–473. [In Russian].
- Proshchalykin M.Yu.** 2012b. Additional data on the long-tongued bee fauna (Hymenoptera, Apoidea: Megachilidae, Apidae) of Eastern Siberia. *Far Eastern Entomologist*, **253**: 24–27.
- Proshchalykin M.Yu.** 2013a. New records of bees (Hymenoptera, Apoidea, Apiformes) from Siberia. *A.I. Kurentsov's Annual Memorial Meetings*, **24**: 135–148. [In Russian].
- Proshchalykin M.Yu.** 2013b. The bees of the tribe Anthidiini Ashmead, 1899 (Hymenoptera: Apoidea: Megachilidae) of Siberia and the Russian Far East. *Caucasian Entomological Bulletin*, **9**(1): 147–158. [In Russian].
- Proshchalykin M.Yu.** 2014a. The species-group names of bees (Hymenoptera: Apoidea, Apiformes) described from the Russian Far East. Part I. Families Colletidae, Andrenidae and Melittidae. *Euroasian Entomological Journal*, **13**(6): 481–488.
- Proshchalykin M.Yu.** 2014b. The species-group names of bees (Hymenoptera: Apoidea, Apiformes) described from the Russian Far East. Part II. Families Halictidae, Megachilidae and Apidae. *Euroasian Entomological Journal*, **13**(6): 527–534.
- Proshchalykin M.Yu.** 2017. The bees of the genus *Colletes* Latreille (Hymenoptera, Colletidae) of the Palaearctic Region: taxonomic diversity and distribution patterns. *Meetings in memory of N.A. Cholodkovsky*, **68**(2): 1–81. [In Russian].
- Proshchalykin M.Yu. and Astafurova Yu.V.** 2012. Halictid bees (Hymenoptera, Apoidea: Halictidae) of Ukraine: fauna and zonal distribution. *A.I. Kurentsov's Annual Memorial Meetings*, **23**: 93–113. [In Russian].
- Proshchalykin M.Yu. and Astafurova Yu.V.** 2016. The species-group names of bees (Hymenoptera: Apoidea, Apiformes) described from Crimea, North Caucasus, European part of Russia and Ural. Part I. Families Colletidae and Halictidae. *Far Eastern Entomologist*, **312**: 1–20.
- Proshchalykin M.Yu., Astafurova Yu.V. and Osytshnjuk A.Z.** 2017a. The species-group names of bees (Hymenoptera: Apoidea, Apiformes) described from Crimea, North Caucasus, European part of Russia and Ural. Part II. Families Andrenidae and Megachilidae. *Far Eastern Entomologist*, **328**: 1–34.
- Proshchalykin M.Yu., Astafurova Yu.A., Schwarz M., Levchenko T.V. and Byvaltsev A.M.** 2017b. New records to the bee fauna of Russia (Hymenoptera, Apiformes). *Far Eastern Entomologist*, **337**: 17–24.
- Proshchalykin M.Yu., Astafurova Yu.V. and Shlyakhtenok A.S.** 2016. A review of the genus *Melitturga* Latreille, 1809 (Hymenoptera, Apoidea, Andrenidae) of the fauna of Russia and adjacent territories. *Euroasian Entomological Journal*, **15**(6): 566–571. [In Russian].
- Proshchalykin M.Yu. and Dathe H.H.** 2012. The bees of the genus *Hylaeus* Fabricius 1793 of the Asian part of Russia, with a key to species (Hymenoptera: Apoidea: Colletidae). *Zootaxa*, **3401**: 1–36.
- Proshchalykin M.Yu. and Dathe H.H.** 2016a. Additional records of the genus *Hylaeus* Fabricius, 1793 (Hymenoptera: Apoidea: Colletidae) from Siberia, with description of a new species. *Zootaxa*, **4105**(4): 301–320.
- Proshchalykin M.Yu. and Dathe H.H.** 2016b. A new species of the genus *Hylaeus* Fabricius, 1793 (Hymenoptera: Colletidae) from Siberia. *Euroasian Entomological Journal*, **15**(Suppl. 1): 113–115.
- Proshchalykin M.Yu. and Kuhlmann M.** 2012. The bees of the genus *Colletes* Latreille 1802 of the Ukraine, with a key to species (Hymenoptera: Apoidea: Colletidae). *Zootaxa*, **3488**: 1–40.
- Proshchalykin M.Yu. and Kuhlmann M.** 2015a. Type specimens of *Colletes* Latreille (Hymenoptera, Colletidae) deposited in the Zoological Institute of the Russian Academy of Sciences, St. Petersburg, with description of a new species. *Zootaxa*, **3949**(4): 540–554.
- Proshchalykin M.Yu. and Kuhlmann M.** 2015b. Additional records of the genus *Colletes* Latreille (Hymenoptera: Apoidea: Colletidae) from Siberia, with a checklist of Russian species. *Zootaxa*, **3949**(3): 323–344.
- Proshchalykin M.Yu. and Kupianskaya A.N.** 2005. The bees (Hymenoptera, Apoidea) of the northern part of the Russian Far East. *Far Eastern Entomologist*, **153**: 1–39.
- Proshchalykin M.Yu. and Kupianskaya A.N.** 2009. The bees of family Apidae (Hymenoptera, Apoidea) of Transbaikalia. *Eurasian Entomological Journal*, **8**(1): 59–68. [In Russian].
- Proshchalykin M.Yu. and Lelej A.S.** 2004. Bees of the subgenus *Alloceroioxys* Tkalcu of the genus *Ceroioxys* Latreille (Hymenoptera: Apoidea: Megachilidae) from the Russian Far East. *Zootaxa*, **517**: 1–6.
- Proshchalykin M.Yu. and Lelej A.S.** 2010. Review of the *Nomada roberjetiana* species-group (Hymenoptera: Apidae) of Russia, with description of new species. *Zootaxa*, **2335**: 1–15.
- Proshchalykin M.Yu. and Lelej A.S.** 2013. The species-group names of bees (Hymenoptera: Apoidea, Apiformes) described from Siberia. *Euroasian Entomological Journal*, **12**(4), 315–327.
- Proshchalykin M.Yu. and Lelej A.S.** 2014. Review of the genus *Ammobatoides* Radoszkowski, 1867 (Hymenoptera: Apidae, Nomadinae) from Russia and neighbouring countries. *Zootaxa*, **3852**(4): 445–460.
- Proshchalykin M.Yu., Novomodnyi E.V., Bezborodov V.G. and Koshkin E.S.** 2014. First records of wax bee *Apis cerana* Fabricius, 1793 (Hymenoptera, Apidae) in Khabarovskii Krai. *Euroasian Entomological Journal*, **13**(3): 295–298. [In Russian].

- Proshchalykin M.Yu. and Schwarz M.** 2017. The genus *Nomada* (Hymenoptera: Apidae) in Siberia. *Linzer Biologische Beiträge*, **49**(1): 985–1006.
- Protsenko Yu.V., Fateryga A.V., Ivanov S.P. and Puzanov D.V.** 2012. Digger wasps (Hymenoptera: Apoidea: Ampulicidae, Sphecidae) of the collection of V.I. Vernadskiy Taurida National University. *Ekosistemy, ikh Optimizatsiya i Okhrana*, **6**: 50–61. [In Russian].
- Prous M., Blank S.M., Goulet H., Heibo E., Liston A., Malm T., Nyman T., Schmidt S., Smith D.R., Vårdal H., Viitasaari M., Vikberg V. and Taeger A.** 2014. The genera of Nematinae (Hymenoptera, Tenthredinidae). *Journal of Hymenoptera Research*, **40**: 1–69.
- Prous M., Heidemaa M., Shinohara A. and Soon V.** 2011a. Review of the sawfly genus *Empria* (Hymenoptera, Tenthredinidae) in Japan. *ZooKeys*, **150**: 347–380.
- Prous M., Heidemaa M. and Soon V.** 2011b. *Empria longicornis* species group: taxonomic revision with notes on phylogeny and ecology (Hymenoptera, Tenthredinidae). *Zootaxa*, **2756**: 1–39.
- Prous M., Kramp K., Vikberg V. and Liston A.** 2017. North-Western Palaearctic species of *Pristiphora* (Hymenoptera, Tenthredinidae). *Journal of Hymenoptera Research*, **59**: 1–190.
- Prous M., Vikberg V., Liston A. and Kramp K.** 2016. North-Western Palaearctic species of the *Pristiphora ruficornis* group (Hymenoptera, Tenthredinidae). *Journal of Hymenoptera Research*, **51**: 1–54.
- Pulawski W.J.** 1956. Europejskie gatunki rodzaju *Astata* Latr. (Hym., Sphecid.). *Polskie Pismo Entomologiczne*, **25**: 33–71.
- Pulawski W.J.** 1962. Les *Tachytes* Panz. de la région paléarctique occidentale et centrale (Hym., Sphecidae). *Polskie Pismo Entomologiczne*, **32**: 311–475.
- Pulawski W.J.** 1963. *Klucze do oznaczenia owadów Polski. Część 24. Błonkówki – Hymenoptera*. Zeszyt 57–62. *Sapygidae, Scoliidae, Tiphiidae, Methocidae, Myrmosidae, Mutillidae*. Państwowe Wydawnictwo Naukowe, Warszawa, 66 p.
- Pulawski W.J.** 1971. Les *Tachysphex* (Hym., Sphecidae) de la région paléarctique occidentale et centrale. Państwowe Wydawnictwo Naukowe, Wrocław, 464 p.
- Pulawski W.J.** 1973. Les *Ammatomus* A. Costa (Hym., Sphecidae) de la région paléarctique occidentale et centrale. *Polskie Pismo Entomologiczne*, **43**: 273–288.
- Pulawski W.J.** 1977. A revision of the Old World *Parapagetia* Kohl (Hymenoptera, Sphecidae). *Polskie Pismo Entomologiczne*, **47**: 601–669.
- Pulawski W.J.** 1978. Superfamily Sphecoidea. In: Medvedev G.S. (Ed.). *Keys to the Identification of insects of European USSR*. Vol. III, pt 1. Nauka, Leningrad: 173–279. [In Russian].
- Pulawski W.J.** 1979. A revision of the World *Prosopigastra* Costa (Hymenoptera, Sphecidae). *Polskie Pismo Entomologiczne*, **49**: 3–134.
- Pulawski W.J.** 1984. The status of *Trypoxylon figulus* (Linnaeus, 1758), *medium* de Beaumont, 1945, and *minus* de Beaumont, 1945 (Hymenoptera: Sphecidae). *Proceedings of the California Academy of Sciences*, **43**: 123–140.
- Pulawski W.J.** 2016. Catalog of Sphecidae sensu lato (= Apoidea excluding Apidae). California Academy of Sciences, Golden Gate Park, San Francisco, California, USA. Available from: http://researcharchive.calacademy.org/research/entomology/entomology_resourcess/hymenoptera/sphecidae/ (accessed 15 December 2016).
- Pulawski W.J. and Prentice M.A.** 2008. A revision of the wasp tribe Palarini Schrottky, 1909 (Hymenoptera: Apoidea: Crabronidae). *Proceedings of the California Academy of Sciences (Series 4)*, **59**: 307–479.
- Qi L., Niu G., Liu T. and Wei M.** 2015. New species and new synonyms of *Conaspidia* Konow (Hymenoptera: Tenthredinidae) with keys to species of the *Conaspidia bicuspis* group and Japanese species. *Entomotaxonomia*, **37**(3): 221–233.
- Quest M.** 2009. Arbestand, Ökologie und Habitatwahl von Bienen ausgewählter Offenlebensräume im Lazovski Zapovednik (Ferner Osten Russland) (Hymenoptera, Apiformes). *Entomofauna, Suppl.* **15**: 1–357.
- Radchenko A.G.** 1991. Ants of the genus *Strongylognathus* (Hymenoptera, Formicidae) of the USSR fauna. *Zoologicheskiy Zhurnal*, **70**(10): 84–90. [In Russian].
- Radchenko A.G.** 1992. Ants of the genus *Tetramorium* (Hymenoptera, Formicidae) of the USSR fauna. Report 1. *Zoologicheskiy Zhurnal*, **71**(8): 39–49. [In Russian].
- Radchenko A.G.** 1996a. Ants of the genus *Plagiolepis* Mayr (Hymenoptera, Formicidae) of Central and Southern Palearctic. *Entomologicheskoe Obozrenie*, **75**: 178–187. [In Russian].
- Radchenko A.G.** 1996b. A review of the ant genus *Leptothorax* (Hymenoptera, Formicidae) of central and eastern Palearctic. Communication 4. Group *congruus*. Species of uncertain position. Zoogeographical characteristics. *Vestnik Zoologii*, **1/2**: 16–22. [In Russian].
- Radchenko A.G.** 1997a. Review of ants of the genus *Camponotus* (Hymenoptera, Formicidae) of the Palearctic. The subgenus *Camponotus* s. str. *Zoologicheskiy Zhurnal*, **76**: 554–564. [In Russian].
- Radchenko A.G.** 1997b. Review of ants of the subgenus *Myrmentoma* genus *Camponotus* (Hymenoptera, Formicidae) of the Asian Palearctic. *Zoologicheskiy Zhurnal*, **76**: 703–711. [In Russian].
- Radchenko A.G.** 1997c. Review of ants of the subgenera *Tanaemyrmex*, *Colobopsis*, *Myrmamblis*, *Myrmosericus*, *Orthonotomyrmex* and *Paramyrmamblis* of the genus *Camponotus* (Hymenoptera, Formicidae) in the Asian Palearctic. *Zoologicheskiy Zhurnal*, **76**: 806–815. [In Russian].

- Radchenko A.G.** 1997d. Review of ants of the genus *Castaglyphis* Foerster (Hymenoptera, Formicidae) of Asia. *Entomologicheskoe Obozrenie*, **76**: 424–442.
- Radchenko A.G.** 2004. A review of the ant genera *Lepthorax* Mayr and *Temnothorax* Mayr (Hymenoptera, Formicidae) of the Eastern Palaearctic. *Acta Zoologica Academiae Scientiarum Hungaricae*, **50**(2): 109–137.
- Radchenko A.G.** 2005a. A review of the ants of the genus *Lasius* Fabricius, 1804, subgenus *Dendrolasius* Ruzsky, 1912 (Hymenoptera: Formicidae) from East Palearctic. *Annales Zoologici*, **55**(1): 83–94.
- Radchenko A.G.** 2005b. Monographic revision of the ants (Hymenoptera: Formicidae) of North Korea. *Annales Zoologici*, **55**(2): 127–221.
- Radchenko A.G.** 2016. *Ants (Hymenoptera, Formicidae) of Ukraine*. Kiev, 495 p.
- Radchenko A.G. and Elmes G.W.** 2010. *Myrmica ants of the Old World*. Fauna Mundi, Vol. 3. Natura Optima Dux Foundation, Warsaw, 789 p.
- Radchenko A.G. and Scupola A.** 2015. Taxonomic revision of the *strativentre* species group of the genus *Tetramorium* (Hymenoptera, Formicidae). *Vestnik Zoologii*, **49**: 219–244.
- Radchenko A.G. and Yusupov Z.M.** 2012. A new peculiar *Myrmica* species (Hymenoptera, Formicidae) from the North Caucasus. *Annales Zoologici*, **62**(4): 593–598.
- Radchenko A.G. and Yusupov Z.M.** 2017. New data on the *Myrmica* ants (Hymenoptera, Formicidae) from the North Caucasus. *Proceedings of the Russian Entomological Society*, **88**(2). [In press].
- Radchenko A.G., Yusupov Z.M. and Fedoseeva E.B.** 2015. Taxonomic notes for some Caucasian *Temnothorax* Mayr species (Hymenoptera, Formicidae), with a description of three new species. *Caucasian Entomological Bulletin*, **11**(1): 161–167.
- Radchenko A.G., Yusupov Z.M. and Komarov Y.E.** 2016. New data on the distribution and ecology of *Myrmica ravasinii* Finzi, 1923 (Hymenoptera, Formicidae). *Asian Myrmecology*, **8**: 1–7.
- Radchenko V.G. and Pesenko Yu.A.** 1989. A key to the bees of the genus *Dasypoda* Latreille (Hymenoptera, Melittidae) of the European part of the USSR, with a designation of lectotypes. *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, **188**: 114–121. [In Russian].
- Radchenko V.G.** 2009. *Dasypoda spinigera* Kohl, 1905. *Data Book of Ukraine. Animal kingdom*. Globalconsulting, Kiev: 253. [In Ukrainian].
- Radchenko V.G.** 2016. A new widespread European bee species of the genus *Dasypoda* Latreille (Hymenoptera, Apoidea). *Zootaxa*, **4184**(3): 491–504.
- Radoszkowski O.** 1865. Les mutilles russes. *Bulletin de la Société Impériale des Naturalistes de Moscou*, **38**(2): 422–467, tab. 7–9.
- Radoszkowski O.** 1866. Enumération des espèces de Chrysides de Russie. *Horae Societatis Entomologicae Rossicae*, **3**: 295–310.
- Radoszkowski O.** 1866. Supplément aux descriptions des mutilles russes. *Bulletin de la Société Impériale des Naturalistes de Moscou*, **39**(2): 299–306, tab. 9, figs 1–3.
- Radoszkowski O.** 1879. Les Chrysides et Sphégides du Caucase. *Horae Societatis Entomologicae Rossicae*, **15**: 140–156.
- Radoszkowski O.** 1880(1879). Les Chrysides et Sphégides du Caucase. *Horae Societatis Entomologicae Rossicae*, **15**: 140–156.
- Radoszkowski O.** 1888. Études hyménoptérologiques. I. Revision des armures copulatrices des mâles. II. Description de nouvelles espèces russes. *Horae Societatis Entomologicae Rossicae*, **22**(3/4): 315–337.
- Radoszkowski O.** 1891. Révision des armatures copulatrices des males du genre *Colletes*. *Horae Societatis Entomologicae Rossicae*, **25**: 249–260.
- Radoszkowski O.** 1893. Revue des armures copulatrices des mâles des genres *Crocisa* Jur., *Melecta* Latr., *Pseudomelecta* Rad., *Chrysanthes* Pert., *Mesocheira* Lep., *Aglae* Lep., *Melissa* Smit, *Euglossa* Lat., *Eulema* Lep., *Acanthopodus* Klug. *Bulletin de la Imperiale Society d'Naturalistes de Moscou*, (N. S.), **7**(2/3): 163–190.
- Radoszkowski O.I.** 1887a. Insecta in itinere cl. N. Przewalskii in Asia Centrali novissime lecta. III. *Horae Societatis Entomologicae Rossicae*, **21**: 41–52.
- Radoszkowski O.I.** 1887b. Faune hyménoptérologique Transcaspienne. *Horae Societatis Entomologicae Rossicae*, **21**: 88–101.
- Radoszkowski O.I.** 1888. Révision des armures copulatrices des mâles de la famille Pompilidae. *Bulletin de la Société Impériale des Naturalistes de Moscou*, **3**: 462–493 + pl. 12–15.
- Radoszkowski O.** 1889(1888). Révision des armures copulatrices des mâles de la tribu des Chrysides. *Horae Societatis Entomologicae Rossicae*, **23**(1–2): 3–40, pls 1–6.
- Rasmont P.** 1995. Les Anthophores de France du sous-genre *Lophanthophora* Brooks avec la redescription de trois espèces au statut confus (Hymenoptera, Apoidea, Anthophoridae). *Annales de la Société Entomologique de France* (N.S.), **31**(1): 3–20.
- Rasmont P., Devalez J., Pauly A., Michez D. and Radchenko V.G.** 2017. Addition to the checklist of IUCN European wild bees (Hymenoptera: Apoidea). *Annales de la Société entomologique de France* (N.S.), **53**(1): 17–32.
- Rasmont P., Ebner A., Banaszak J. and Zanden G.** 1995. Hymenoptera Apoidea Gallica. Liste taxonomique des abeilles de France, de Belgique, de Suisse et du Grand Duché du Luxembourg. *Bulletin de la Société Entomologique de France*, **100**: 1–98.

- Rasmont P., Franzén M., Lecocq T., Harpke A., Roberts S.P.M., Biesmeijer J.C., Castro L., Cederberg B., Dvorák L., Fitzpatrick Ú., Gonseth Y., Haubrûge E., Mahé G., Manino A., Michez D., Neumayer J., Ødegaard F., Paukkunen J., Pawlikowski T., Potts S.G., Reemer M., Settele J., Straka J. and Schweiger O.** 2015. Climatic risk and distribution atlas of European bumblebees. *Biorisk*, 10(Special Issue): 1–246.
- Rasnitsyn A.P.** 2002. Superorder Vespoidea Laicharting, 1781. Order Hymenoptera Linne, 1758 (=Vespidae Laicharting, 1781). In: Rasnitsyn A.P., Quicke D.L.J. (Eds). *History of Insects*. Kluwer Academic Publishers: 242–255.
- Rasnitsyn A.P. and Matveev D.G.** 1989. First Palaearctic representative of the genus *Ampulicomorpha* Ashmead. *Entomologicheskoe Obozrenie*, 68: 657–661. [In Russian].
- Richards O.W.** 1962. A revisional study of the masarid wasps (Hymenoptera: Vespidae). British Museum (Natural History), London, 302 p.
- Richards O.W.** 1973. The subgenera of *Polistes* Latreille (Hymenoptera, Vespidae). *Revista Brasileira de Entomologia*, 17: 85–104.
- Richards O.W. and Richards M.J.** 1951. Observations on the social wasps of South America (Hymenoptera Vespidae). *Transactions of the Royal Entomological Society of London*, 102: 1–170.
- Rightmyer M.G.** 2008. A review of the cleptoparasitic bee genus *Triepeolus* (Hymenoptera: Apidae). Part I. *Zootaxa*, 1710: 1–170.
- Risch S.** 1999. Neue und wenig bekannte Arten der Gattung *Eucera* Scopoli 1770 (Hymenoptera, Apidae). *Linzer Biologische Beiträge*, 31(1): 115–145.
- Risch S.** 2001. Die Arten des Genus *Eucera* Scopoli 1770 (Hymenoptera, Apidae) Untergattung *Pareucera* Tkalcú, 1979. *Entomofauna*, 22(15): 365–376.
- Rohwer S.A.** 1910. Japanese sawflies in the collection of the United States National Museum. *Proceedings of the United States National Museum*, 39(1777): 99–120.
- Rohwer S.A.** 1925. Sawflies from the Maritime Province of Siberia. *Proceedings of the United States National Museum*, 68(2609): 1–12.
- Roller L. and Haris A.** 2008. Sawflies of the Carpathian Basin, history and current research. *Natura Somogyensis*, 11: 1–259.
- Romankova T.G.** 1995a. Family Megachilidae. In: Lehr P.A. (Ed.). *Key to the insects of the Russian Far East*. Vol. IV, pt 1. Nauka, St Petersburg: 530–547. [In Russian].
- Romankova T.G.** 1995b. Family Anthophoridae. In: Lehr P.A. (Ed.). *Key to the insects of the Russian Far East*. Vol. IV, pt 1. Nauka, St Petersburg: 547–551. [In Russian].
- Romankova T.G. and Astafurova Yu.V.** 2011. Bees of the genus *Panurginus* in Siberia, Far East of Russia, and allies areas (Hymenoptera: Andrenidae, Panurginae). *Zootaxa*, 3112: 1–35.
- Romanova V.P.** 1969. On the fauna of sphecid wasps (Hymenoptera, Sphecidae) from the North Caucasus. *Entomologicheskoe Obozrenie*, 48: 132–137. [In Russian].
- Rosa P.** 2004. Alcune osservazioni sulle relazioni tra Vegetazione e Crisidi (Hymenoptera, Chrysidae) in Italia. *Giornale Italiano di Entomologia*, 11: 79–90.
- Rosa P.** 2006. *I Crisidi della Valle d'Aosta (Hymenoptera, Chrysidae)*. Monografie del Museo Regionale di Scienze Naturali, 6, St.-Pierre, Aosta, 368 p., xvi + 32 pls
- Rosa P., Antropov A. and Xu Z-f.** 2015. A catalogue of the Chrysidae (Insecta, Hymenoptera) types deposited in the Zoological Museum, Moscow Lomonosov State University, Russia. *Zootaxa*, 3990(1): 1–31.
- Rosa P., Belokobylskij S.A. and Zaytseva L.A.** 2017a. The Chrysidae types described by Semenov-Tian-Shanskij and deposited at the Zoological Institute of the Russian Academy of Sciences, Saint Petersburg (Insecta, Hymenoptera). *Proceedings of the Zoological Institute RAS*, Suppl. 5: 1–266.
- Rosa P., Proshchalykin M.Yu., Lelej A.S. and Loktionov V.M.** 2017b. Contribution to the Siberian Chrysidae (Hymenoptera). Part 1. *Far Eastern Entomologist*, 341: 1–44.
- Rosa P., Proshchalykin M.Yu., Lelej A.S. and Loktionov V.M.** 2017c. Contribution to the Siberian Chrysidae (Hymenoptera). Part 2. *Far Eastern Entomologist*, 342: 1–42.
- Rosa P., Lelej A.S., Proshchalykin M.Yu., Loktionov V.M. and Mokrousov M.** 2017d. New records of Chrysidae (Hymenoptera) from Russia with description of five new species. *Far Eastern Entomologist*, 345: 1–33.
- Ruchin A.B. and Antropov A.V.** 2014. On the wasp fauna (Hymenoptera, Vespoidea) of the Republic of Mordovia, Russia. *Aktual'nyie Problemy Gumanitarnykh i Estestvennykh Nauk*, 3(62): 29–36. [In Russian].
- Ruchin A.B. and Antropov A.V.** 2016. New and interesting finds of wasps (Insecta, Hymenoptera) in the Mordovian Reserve. *Trudy Mordovskogo gosudarstvennogo prirodno-zapovednika imeni P.G. Smidovicha*, 16: 399–403. [In Russian].
- Ruchin A.B., Antropov A.V. and Shibaev S.V.** 2009. Materials to the wasp fauna (Hymenoptera: Chrysidae, Scoliidae, Tiphiidae, Pompilidae, Vespidae, Sphecidae, Crabronidae, Trigonalyidae) of the Republic of Mordovia, Russia. *Vestnik Mordovskogo Universiteta. Seriya "Biologicheskiye Nauki"*, 1: 164–172. [In Russian].
- Rudoiskatel P.V.** 2010. Sphecidae and Crabronidae of central and southern Ural mountains. In: *Ecology from the southern mountains to the northern seas*. Rossiskaya Akademiya Nauk, Ural'skoye Otdeleniye, Institut Ekologii Rasteniy i Zhivotnykh, Yekaterinburg: 145–161. [In Russian].
- Rudoiskatel P.V., Nikolaenkova A.V. and Fadeev K.I.** 2011. Fauna and ecology of vespid wasps (Hymenoptera, Vespidae) of Sverdlovsk Province. *Agrarnyi Vestnik Urala*, 2011(2): 31–33. [In Russian].

- Ruzsky M.D.** 1905. The ants of Russia. (Formicariae Imperii Rossici). Systematics, geography and data on the biology of Russian ants. Part I. *Trudy Obshchestva Estestvoispytatelei pri Imperatorskom Kazanskom Universitete*, **38**(4–6): 1–800. [In Russian].
- Ruzsky M.D.** 1907. The ants of Russia. (Formicariae Imperii Rossici). Systematics, geography and data on the biology of Russian ants. Part II. *Trudy Obshchestva Estestvoispytatelei pri Imperatorskom Kazanskom Universitete*, **40**(4): 1–122 + 3 p.
- Saarinen A.** 1950a. *Amauronematus hartigi* n.sp., eine neue Blattwespenart (Hym., Symphyta) aus der Sammlung Th. Hartigs in München. *Annales Entomologici Fennici*, **16**(1): 18–24.
- Saarinen A.** 1950b. Drei neue paläarktische *Trichiosoma*-Arten (Hym., Symphyta). *Entomologica Fennica*, **16**(1): 1–13.
- Sahlberg J.** 1910. Lyhyt katsaus Suomen kulta-ampiaisiin (Chrysidae) (!). *Luonnon Ystävä*, **14**: 94–99.
- Saini M.S., Blank S.M. and Smith D.R.** 2006. Checklist of the sawflies (Hymenoptera: Symphyta) of India. In: Blank S.M., Schmidt S., Taeger A. (Eds). *Recent Sawfly Research – Synthesis and Prospects*. Goecke & Evers, Keltern: 575–612.
- Sapaev E.A.** 2004. Order Hymenoptera. In: S.M. Shafiqullina (Ed.). *Cadastre of useful insects of Tatarstan Republic*. Innovation technology center, Kazan: 123–212. [In Russian].
- Schaefer H. and Renner S.S.** 2008. A phylogeny of the oil bee tribe Ctenoplectriini (Hymenoptera: Anthophila) based on mitochondrial and nuclear data: Evidence for Early Eocene divergence and repeated out-of-Africa dispersal. *Molecular Phylogenetics and Evolution*, **47**: 799–811.
- Schedl W.** 2009. Hymenoptera (Insecta). In: Winkler H., Stuessy T. (Eds). *Checklisten der Fauna Österreichs*, Wien, **4**: 8–40.
- Scheuchl E.** 2008. Illustrierte Bestimmungstabellen der Wildbienen Deutschlands und Österreichs. Bd. 1: Anthophoridae. 2. erweiterte Auflage. Velden, 175 s.
- Scheuchl E.** 2010. Beitrag zur Kenntnis westpaläarktischer bienen der Gattung *Andrena* (Hymenoptera: Apidae: Andreninae). *Linzer Biologische Beiträge*, **42**(2): 1445–1455.
- Schiff N.M., Goulet H., Smith D.R., Boudreault C., Wilson D.A. and Scheffler B.E.** 2012. Sircidae (Hymenoptera: Symphyta: Siricoidea) of the Western Hemisphere. *Canadian Journal of Arthropod Identification*, **21**: 1–305.
- Schmid-Egger C.** 2005. *Proxiandrena* subgen. nov. und Revision der west- und zentralpaläarktischen Arten der *Andrena proxima* Gruppe (Hymenoptera, Apidae). *Revue Suisse de Zoologie*, **112**(4): 1029–1044.
- Schmid-Egger Ch.** 2002. Key and new records for the western Palaearctic species of *Gorytes* Latreille 1804 with description of a new species (Hymenoptera, Sphecidae, Bembicinae). *Linzer Biologische Beiträge*, **34**: 167–190.
- Schmid-Egger Ch.** 2004. Revision of *Bembecinus* (Hymenoptera, Crabronidae) of the Palearctic Region. *Notes Fauniques de Gembloux*, **54**: 3–69.
- Schmid-Egger Ch.** 2005. The genus *Olgia* Radoszkowski 1877 with description of a new species (Hymenoptera, Crabronidae, Bembicinae). *Linzer Biologische Beiträge*, **37**: 1545–1553.
- Schmidt K.** 2000. Bestimmungstabelle der Gattung *Cerceris* Latreille, 1802 in Europa, dem Kaukasus, Kleinasien, Palästina und Nordafrika (Hymenoptera, Sphecidae, Philanthinae). *Stapfia*, **71**: 1–325.
- Schmidt S.** 1997. *Amauronematus* Konow, 1890 – Ökologie und Taxonomie der nordeuropäischen Arten des *fallax*-Komplexes (Hymenoptera, Tenthredinidae). *Beiträge zur Entomologie*, **47**(2): 227–326.
- Schmidt S., Taeger A., Morinière J., Liston A., Blank S.M., Kramp K., Kraus M., Schmidt O., Heibo E., Prous M., Nyman T., Malm T. and Stahlhut J.** 2016. Identification of sawflies and horntails (Hymenoptera, ‘Symphyta’) through DNA barcodes: successes and caveats. *Molecular Ecology Resources*, **17**(4): 1–16.
- Schmiedeknecht O.** 1882. Apidae Europaeae (Die Bienen Europas) per genera, species et varietates dispositae atque descriptae. 1. *Nomada, Bombus, Psithyrus et Andrena*. R. Friedländer & Sohn, Berlin, 866 s.
- Schneider H.** 2016. The ghost of the Cretaceous terrestrial revolution in the evolution of fern-sawfly associations. *Journal of Systematics and Evolution*, **54**(2): 93–103.
- Schtundjuk A.V. and Zhelochovtsev A.N.** 1974. To fauna of Symphyta (Hymenoptera) of middle Amur region. In: Kulik S.A. (Ed.). *Fauna nasekomykh Vostochnoy Sibiri i Dal'nego Vostoka*. Irkutskiy gosudarstvenny universitet, Irkutsk: 244–252. [In Russian].
- Schwarz M.** 1965. Zwei neue *Nomada*-Arten aus Griechenland (Hymenoptera, Apoidea). *Nachrichtenblatt der bayrischen Entomologen*, **14**: 81–87.
- Schwarz M.** 1967a. Ergebnisse der Albanien-Expedition 1961 des Deutschen Entomologischen Institutes. 68. Beitrag Hymenoptera: Apidae III (Genus *Nomada* Scopoli). *Beiträge zur Entomologie*, **17**: 529–535.
- Schwarz M.** 1967b. Die Gruppe der *Nomada cinctiventris* Fr. (= *stigma* auct. nec. F.). *Polskie Pismo Entomologiczne*, **37**: 263–339.
- Schwarz M.** 1980. Beitrag zur Kenntnis weiterer von F. Morawitz beschriebenen *Nomada*-Arten (Hymenoptera, Apoidea). *Entomofauna*, **1**(9): 103–118.
- Schwarz M.** 1981. 1. Beitrag zur *Nomada*-Fauna der Türkei (Hymenoptera, Apoidea). *Entomofauna*, **2**(28): 357–414.
- Schwarz M. and Gusenleitner F.** 2001. Beitrag zur Kenntnis paläarktischer Anthophorini und Habropodini (Hymenoptera: Apidae). *Entomofauna*, **22**(6): 53–92.

- Schwarz M. and Guseinleitner F.** 2013. Zur Kenntnis der *Nomada*-Fauna Spaniens, mit Klärung der *Nomada dolosa* Mocsay (= *Nomada centenarii* Dusmet) (Hymenoptera: Apidae). *Linzer Biologische Beiträge*, **45**(1): 971–993.
- Schwarz M. and Guseinleitner F.** 2015. Über den Verbleib der von Schmiedeknecht 1882 beschriebenen *Nomada*-Arten und Festlegung von Lectotypen der aufgefundenen Arten. Mit zusätzlichen Ergänzungen und Beschreibungen zu einigen Arten (Hymenoptera, Apidae, Nomadinae). *Linzer Biologische Beiträge*, **47**(1): 1003–1044.
- Schwarz M. and Guseinleitner F.** 2017. *Nomada pilosa* nov. sp., eine neue paläarktische Wespenbiene (Hymenoptera, Apidae). *Linzer Biologische Beiträge*, **49**(1): 979–983.
- Scobiola-Palade X.** 1989. L'Ordre Hymenoptera (Sous-ordre Symphyta, Superfamilles Vespoidea, Pompiloidea, Sphecoidea) du cordon de sable Sărăturile – Sf. Gheorghe (Delta du Danube, Roumanie). *Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa"*, **30**: 85–89.
- Sedykh K.F.** 1974. *The animal world of the Komi ASSR. Invertebrates*. Book Publishing House of the Komi Republic, Syktyvkar, 191 p.
- Seifert B.** 1992. A taxonomic revision of the Palaeartic members of the ant subgenus *Lasius* s.str. (Hymenoptera: Formicidae). *Abhandlungen und Berichte des Naturkundemus Görlitz*, **66**(5): 1–67.
- Seifert B.** 2000. A taxonomic revision of the ant subgenus *Cryptoformica* Mueller, 1923 (Hymenoptera: Formicidae). *Zoosystema*, **22**: 517–568.
- Seifert B.** 2003. The ant genus *Cardiocondyla* (Insecta: Hymenoptera: Formicidae) - a taxonomic revision of the *C. elegans*, *C. bulgarica*, *C. batesii*, *C. nuda*, *C. shuckardi*, *C. stambuloffii*, *C. wroughtonii*, *C. emeryi*, and *C. minutior* species groups. *Annalen des Naturhistorischen Museum in Wien*, **104**(B): 203–338.
- Seifert B.** 2007. *Die Ameisen Mittel- und Nordeuropas*. Iutra Verlags und Vertriebsgesellschaft, Tauer, 368 p.
- Seifert B.** 2012. A review of the West Palaearctic species of the ant genus *Bothriomyrmex* Emery, 1869 (Hymenoptera: Formicidae). *Myrmecological News*, **17**: 91–104.
- Seifert B. and Schultz R.** 2009. A taxonomic revision of the *Formica subpilosa* Ruzsky, 1902 group (Hymenoptera: Formicidae). *Myrmecological News*, **12**: 67–83.
- Seiyama Y.** 1981. A revision of the Japanese species of the genus *Eutomostethus* Enslin (Hymenoptera: Tenthredinidae). *Transactions of the Shikoku Entomological Society*, **15**(3–4): 155–171.
- Semenov A.** 1921. Praecursoriae Siricidarum novorum diagnoses (Hymenoptera). *Russkoe Entomologicheskoe Obozrenie*, **17**[1917]: 81–95.
- Semenov A. and Gussakovskij V.V.** 1935. Siricides nouveaux ou peu connus de la faune paléarctique (Hymenoptera). *Annales de la Société Entomologique de France*, **104**: 117–126.
- Semenov A. and Gussakovskij V.V.** 1937. Note sur les représentants paléarctiques de la tribu des Abiini (Hymenoptera, Tenthredinidae). *Konowia*, **16**(1): 1–9.
- Semenov-Tian-Shanskij A.** 1967. New species of gold wasps (Hymenoptera, Chrysididae). *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, **43**: 118–184. [In Russian].
- Sergeev M.E. and Lelej A.S.** 2011. On the parasitism of velvet ant *Physetopoda halensis* (Hymenoptera: Mutilidae) on the larva of leaf beetle *Labidostomis hummeralis* (Coleoptera, Chrysomelidae, Clytrinae) in South-East Ukraine. *Vestnik Zoologii*, **45**(2): 144. [In Russian].
- Setrakova E.M.** 2014. The current state of knowledge of the taxonomic composition of phylophagous insects of oak (*Quercus robur* L.) in Belarus. *Trudy Belorusskogo gosudarstvennogo universiteta*, **9**(2): 236–245. [In Russian].
- Shcherbakov D.E.** 2006. Fern sawfly larvae *Blasticotoma filiceti* Klug, 1834 (Hymenoptera: Blasticotomidae) are visited by ants: a new kind of trophobiosis. *Russian Entomological Journal*, **15**(1): 67–72.
- Shcherbakov D.E.** 2008. New records of Hymenoptera from the Moscow Region and other parts of Russia, with notes on synonymy of *Konowia* species. *Russian Entomological Journal*, **17**: 209–212.
- Shchurov V.I., Gninenco Yu.I., Lengesova N.A. and Gninenco M.Yu.** 2012. *Aproceros leucopoda* in the European part of Russia. *Zashchita i Karantin Rasteniy*, **2**: 37–38. [In Russian].
- Shebl M. and Li Q.** 2013. New record of *Panurginus niger* Nylander (Hymenoptera: Andrenidae: Panurginae) from China. *Entomotaxonomia*, **35**(2): 146–154.
- Shebl M.A. and Tadauchi O.** 2009. The genus *Andrena* from Kazakhstan and Kyrgyzstan (Hymenoptera, Andrenidae) (3). *Esakia*, **49**: 21–62.
- Shestakov A.V.** 1925. Materials to the knowledge of digger wasps (Crabronidae) of Yaroslavl province. *Trudy Yaroslavl'skogo Yestestvenno-Istoricheskogo Obshchestva*, **4**: 35–37. [In Russian].
- Shibaev S.V.** 2006. Addition to the fauna and ecology of some of chrysidid wasps (Hymenoptera: Chrysididae) of the Penza region. *News of Penza State Pedagogical University*, **5**: 108–112. [In Russian].
- Shibaev S.B. and Polumordvinov O.A.** 2012. A review of species diversity of hymenoptera (Insecta, Hymenoptera) in Penza region. *Izvestiya Penzenskogo gosudarstvennogo pedagogicheskogo universiteta imeni V. G. Belinskogo, Estestvennye Nauki*, **29**: 274–279. [In Russian].
- Shilenkov V.G. and Tolstonogova E.V.** 2006. Phytophagous insects on cannabis in Baikal Region. *Byulleten' VSNTs SShch RAMN*, **2**(48): 159–160. [In Russian].
- Shimizu A.** 1994. Phylogeny and classification of the family Pompilidae. *Tokyo Metropolitan University, Bulletin of Natural History*, **2**: 1–142.

- Shimizu A. and Wahis R.** 2009. Systematic studies on the Pompilidae occurring in Japan: genus *Agenioideus* Ashmead (Hymenoptera), supplement. *Entomological Science*, **12**: 238–251.
- Shinohara A. and Hara H.** 2015. Taxonomic notes and new distribution and host plant records for sawflies and woodwasps (Hymenoptera, Symphyta) of Japan. *Bulletin of the National Science Museum, Series A, Zoology*, **41**(3): 171–184.
- Shinohara A.** 1979. A study of the *sulphureipes* complex of the genus *Pamphilius* (Hymenoptera: Pamphiliidae). *Transactions of the Shikoku Entomological Society*, **14**(3–4): 151–161.
- Shinohara A.** 1980. East Asian species of the genus *Neurotoma* (Hymenoptera: Pamphiliidae). *Transactions of the Shikoku Entomological Society*, **15**(1–2): 87–117.
- Shinohara A.** 1987. Systematic studies on the genus *Onycholyda* (Hymenoptera, Pamphiliidae) of the Far East I. *Kontyû*, **55**(4): 644–653.
- Shinohara A.** 1988a. Descriptions of two *Pamphilius*, with a checklist of the Pamphiliidae (Hymenoptera) from Caucasia. *Bulletin of the National Science Museum, Series A, Zoology*, **14**(2): 105–111.
- Shinohara A.** 1988b. *Pamphilius stramineipes* (Hymenoptera, Pamphiliidae) and its close relatives. *Bulletin of the National Science Museum, Series A, Zoology*, **14**(4): 179–197.
- Shinohara A.** 1990. New faunal records of the three East Asian species of the sawfly genus *Abia* (Hymenoptera, Cimbicidae). *Memoirs of the National Science Museum*, **23**: 113–117.
- Shinohara A.** 1992a. Records of *Neurotoma atrata* and *N. sibirica* (Hymenoptera, Pamphiliidae) from the Russian Far East. *Japanese Journal of Entomology*, **60**(4): 826.
- Shinohara A.** 1992b. The sawfly genus *Megaxyela* (Hymenoptera, Xyelidae) in East Asia. *Japanese Journal of Entomology*, **60**(4): 783–796.
- Shinohara A.** 1993. *Pamphilius sulphureipes* (Hymenoptera, Pamphiliidae) and its close relatives. *Memoirs of the National Science Museum*, **26**: 111–123.
- Shinohara A.** 1994a. New record of *Tenthredo xanthotarsus* (Hymenoptera, Tenthredinidae) from Honshu and Kunashir Island. *Japanese Journal of Entomology*, **62**(3): 482.
- Shinohara A.** 1994b. *Tenthredo viridatrix nippon* n. subsp. (Hymenoptera, Tenthredinidae) from Japan and Sakhalin. *Proceedings of the Japanese Society of Systematic Zoology*, **51**: 51–58.
- Shinohara A.** 1995a. *Pamphilius histrio* (Hymenoptera, Pamphiliidae) and its close relatives. *Bulletin of the National Science Museum, Series A, Zoology*, **21**(1): 37–70.
- Shinohara A.** 1995b. The sawfly genus *Pleroneura* (Hymenoptera, Xyelidae) in East Asia. *Japanese Journal of Entomology*, **63**(4): 825–840.
- Shinohara A.** 1997a. New distribution records and taxonomic notes on eight Pamphiliid sawflies (Hymenoptera) in the Western Palaearctic Region. *Japanese Journal of Systematic Entomology*, **3**(2): 231–237.
- Shinohara A.** 1997b. New or noteworthy distribution records of six Palaearctic species of Pamphiliid sawflies (Hymenoptera). *Bulletin of the National Science Museum, Series A, Zoology*, **23**(1): 69–72.
- Shinohara A.** 1997c. The type material of Japanese *Tenthredo* and *Macrophya* sawflies (Hymenoptera, Tenthredinidae) described by A. Mocsáry and R. Malaise. *Bulletin of the National Science Museum, Series A, Zoology*, **23**(3): 165–175.
- Shinohara A.** 1998. *Pamphilius albopictus* (Hymenoptera, Pamphiliidae) and its close relatives. *Bulletin of the National Science Museum, Series A, Zoology*, **24**(4): 225–252.
- Shinohara A.** 1999. A study on stem boring sawflies (Hymenoptera, Cephidae) of the tribe Hartigiini from Japan and Korea. *Japanese Journal of Systematic Entomology*, **5**(1): 61–77.
- Shinohara A.** 2000. Pine-feeding webspinning sawflies of the *Acantholyda posticalis* group (Hymenoptera, Pamphiliidae). *Bulletin of the National Science Museum, Series A, Zoology*, **26**(2): 57–98.
- Shinohara A.** 2001a. Conifer-feeding webspinning sawflies of the genus *Acantholyda* (Hymenoptera, Pamphiliidae) of Japan. *Species Diversity*, **6**: 23–63.
- Shinohara A.** 2001b. The group of *Pamphilius sylvaticus* (Hymenoptera, Pamphiliidae): two new species, new collection records, and a key to Palearctic species. *Japanese Journal of Systematic Entomology*, **7**(1): 99–116.
- Shinohara A.** 2002a. Sawflies of the subgenus *Tenthredina* (Hymenoptera, Tenthredinidae) in Northeastern Asia. *Bulletin of the National Science Museum, Series A, Zoology*, **28**(3): 151–157.
- Shinohara A.** 2002b. Systematics of the leaf-rolling or web-spinning sawfly subfamily Pamphiliinae (Hymenoptera): a preliminary overview. In: Viitasaari M. (Ed.). *Sawflies (Hymenoptera, Symphyta) I. A review of the suborder, the Western Palaearctic taxa of Xyeloidea and Pamphilioidea*. Tremex, Helsinki: 359–438.
- Shinohara A.** 2003. Leaf-rolling sawflies of the *Pamphilius vafer* complex (Hymenoptera, Pamphiliidae) in Europe. *Insect Systematics & Evolution*, **34**: 453–480.
- Shinohara A.** 2005a. Leaf-rolling sawflies of the *Pamphilius vafer* complex (Hymenoptera, Pamphiliidae). *National Science Museum Monographs*, **27**: 1–116.
- Shinohara A.** 2005b. Taxonomic species of the changes and new distribution records of four sawfly genus *Tenthredo* (Hymenoptera, Tenthredinidae) in Japan. *Bulletin of the National Science Museum, Series A, Zoology*, **31**(4): 183–189.

- Shinohara A.** 2015. Japanese sawflies of the genus *Macrophyia* (Hymenoptera, Tenthredinidae), taxonomic notes and key to species *Bulletin of the National Science Museum, Series A, Zoology*, 41(4): 225–251.
- Shinohara A. and Byun B.K.** 1993. Pamphiliid sawfly genera *Neurotoma* and *Onycholyda* (Hymenoptera, Symphyta) of Korea. *Insecta Koreana*, 10: 75–91.
- Shinohara A. and Byun B.K.** 1996. Conifer-feeding webspinning sawflies of the genus *Acantholyda* (Hymenoptera, Pamphiliidae) from Korea. *Insecta Koreana*, 13: 91–104.
- Shinohara A. and Hara H.** 2000. Notes on a spruce-feeding webspinning sawfly, *Acantholyda aglaia* (Hymenoptera, Pamphiliidae), with description of a new subspecies from Hokkaido, Japan. *Bulletin of the National Science Museum, Series A, Zoology*, 26(1): 1–11.
- Shinohara A. and Hara H.** 2007. Type material of Japanese sawflies of the genus *Arge* (Insecta, Hymenoptera, Argidae) described by Snellen van Vollenhoven (1860), Smith (1874) and Kirby (1882). *Bulletin of the National Science Museum, Series A, Zoology*, 33(3): 127–132.
- Shinohara A., Hara H. and Kim J.-W.** 2009. The species-group of *Arge captiva* (Insecta, Hymenoptera, Argidae). *Bulletin of the National Museum of Nature and Science, Series A*, 35(4): 249–278.
- Shinohara A., Hara H. and Prous M.** 2015. Host plants of *Empria* sawflies (Hymenoptera, Tenthredinidae) in Japan include *Rhododendron* (Ericaceae). *Zootaxa*, 4007(1): 143–148.
- Shinohara A., Ibuki Sh.-i. and Hara H.** 2011. *Arge meliosmae* n. sp. (Hymenoptera, Argidae) feeding on *Meliosma myriantha* in Japan. *Bulletin of the National Museum of Nature and Science, Series A*, 37(2): 99–112.
- Shinohara A. and Kato Yu.** 2017. The genus *Acantholyda* (Hymenoptera, Pamphiliidae) of Japan: A revised key and a description of the previously unknown female of *Acantholyda flaviventris*. *Zootaxa*, 4221(2): 215–219.
- Shinohara A. and Lee J.-W.** 1997. Collection records of Pamphiliid sawflies (Hymenoptera) from Korea. *Bulletin of the National Science Museum, Series A, Zoology*, 23(4): 213–220.
- Shinohara A. and Lelej A.S.** 2007. Family Pamphiliidae. In: Lelej A.S. (Ed.). *Key to the insects of Russian Far East*. Vol. IV, pt 5. Dalnauka, Vladivostok: 922–942. [In Russian].
- Shinohara A., Saito A. and Kuranishi R.B.** 2000a. Records of the genus *Tenthredo* (Insecta: Hymenoptera: Tenthredinidae) from Kamchatka and the North Kuril Islands. *Natural History Research, Special issue*, 7: 295–300.
- Shinohara A. and Taeger A.** 2007. Description of a new species and collection data of Pamphiliid sawflies (Insecta, Hymenoptera, Symphyta) mainly from the Russian Far East. *Bulletin of the National Museum of Nature and Science, Series A, Zoology*, 33(1): 31–40.
- Shinohara A. and Vasilenko S.** 2005. Some Pamphiliid sawflies (Hymenoptera) in the collection of the Siberian Zoological Museum, Novosibirsk. *Japanese Journal of Systematic Entomology*, 11(1): 31–37.
- Shinohara A., Vikberg V., Zinovjev A.G. and Yamagami A.** 2000b. *Fagineura crenativora*, a new genus and species of sawflies (Hymenoptera, Tenthredinidae, Nematinae) injurious to beech trees in Japan. *Bulletin of the National Science Museum, Series A, Zoology*, 26(3): 113–124.
- Shinohara A., Wei M. and Niu G.** 2013. Revision of *Siobla* (Hymenoptera, Tenthredinidae) from Japan. *Zootaxa*, 3746(1): 1–40.
- Shinohara A. and Yoshida H.** 2015. *Macrophyia togashii* n. sp. (Hymenoptera, Tenthredinidae) from Japan. *Bulletin of the National Museum of Nature and Science, Series A, Zoology*, 41(2): 123–129.
- Shinohara A. and Zhou H.** 2006. Leaf-rolling sawflies of the *Pamphilus komonensis* complex (Insecta, Hymenoptera, Pamphiliidae). *Bulletin of the National Science Museum, Series A, Zoology*, 32(4): 153–189.
- Shinohara A. and Zinovjev A.G.** 1996. New distribution records of pamphiliid sawflies (Hymenoptera) from Russia, Ukraine, Kazakhstan and Sweden. *Bulletin of the National Science Museum, Series A, Zoology*, 22: 107–112.
- Shinohara A. and Zombori L.** 2003. Records of pamphiliid sawflies (Hymenoptera) from the Ukrainian Carpathians, Poland and western Russia. *Folia Entomologica Hungarica*, 63: 223–226.
- Shkuratov A.V.** 2000. Sphecid wasps in the village Vyoshenskaya area. In: *Gosudarstvennyi Muzey-Zapovednik M.A. Sholokhova. Rostovskiy Gosudarstvennyi Universitet. Priroda Gosudarstvennogo Muzeya-Zapovednika M.A. Sholokhova. Sbornik Statyei*. RostIzdat, Rostov-na-Donu: 54–60. [In Russian].
- Shkuratov A.V.** 2004. Digger wasps (Hymenoptera: Sphecidae) of Rostov Region and some neighbouring territories. *Izvestiya Khar'kovskogo Entomologicheskogo Obshchestva*, 11: 70–85. [In Russian].
- Shlyahetenok A.S., Lelej A.S. and Loktionov V.M.** 2012. A review of the genus *Anoplius* Dufour, 1834 (Hymenoptera, Pompilidae) of European fauna. *Euroasian Entomological Journal*, 11(5): 449–464. [In Russian].
- Shorenko K.I.** 2003. New data on the digger wasp fauna (Apoidea: Sphecidae, Crabronidae) of Ukraine. *Izvestiya Kharkovskogo Entomologicheskogo Obshchestva*, 10: 96–98. [In Russian].
- Shorenko K.I.** 2005. On the digger wasp fauna (Hymenoptera: Ampulicidae, Sphecidae, Crabronidae) of Crimea. *Caucasian Entomological Bulletin*, 1: 161–170. [In Russian].
- Shorenko K.I.** 2007. First find of *Dolichurus haemorrhous* (Hymenoptera: Ampulicidae) in Ukraine. *Vestnik Zoologii*, 41: 554. [In Russian].

- Shorenko K.I.** 2009. Dimensions of species ranges of fosorial wasps (Hymenoptera: Ampulicidae, Sphecidae, Crabronidae) of the Crimean peninsula. In: *Zapovedniki Kryma. Teoriya, Praktika i Perspektivy Zapovednogo Dela v Chernomorskem Regeone. Materiyaly V Mezhdunarodnoy Nauchno-Prakticheskoy Konferentsii*. Simferopol: 366–371. [In Russian].
- Shorenko K.I. and Konovalov S.V.** 2010. New data on digger wasps (Hymenoptera: Ampulicidae, Sphecidae, Crabronidae) in the fauna of Ukraine. *Ukrains'ka Entomofaunistyka*, 1: 9–32. [In Russian].
- Shumakova P.I., Babenko Z.S. and Zolotarenko G.S.** 1982. Bee pollinators (Hymenoptera, Apoidea) of leguminous plants in the Kulunda. In: Zolotarenko G.S. (Ed.). *Beneficial and Harmful Insects of Siberia*. Nauka, Novosibirsk: 157–174. [In Russian].
- Sichel J. and Radoszkowski O.** 1869–1870. Essai d'une monographie des Mutilles de l'ancien continent. *Horae Societatis Entomologicae Rossicae*, (1869) 6(3): 139–172 + pl. 6–8; (1870) 6(4): 173–309 + pl. 9–11.
- Sidorov D.A. and Eremeeva N.I.** 2008. Fauna of bees of some genera of the family Apidae in the Kuznetsk-Salair Mountain Province. *Trudy Kemerovskogo otdeleiniya Russkogo Entomologicheskogo Obchhestva*, 6: 110–116. [In Russian].
- Sidorov D.A., Luzyanin S.L. and Yakovleva S.N.** 2016. First records of bees (Hymenoptera, Apoiformes) in the relict lime forest of the Mountain Shoria. *Sovremenneye Problemy Nauki i Obrazovaniya*, 6 (<http://www.science-education.ru/article/view?id=25798>).
- Sidorov D.A. and Proshchalykin M.Yu.** 2017a. A contribution to the knowledge of the bee genus *Andrena* Fabricius (Hymenoptera, Andrenidae) of the Republic of Khakassia. *Acta Biologica Sibirica*, 3(2): 13–19. [In Russian].
- Sidorov D.A. and Proshchalykin M.Yu.** 2017b. New records on bees of the genus *Andrena* Fabricius (Hymenoptera, Apoidea: Andrenidae) in the south of Eastern Siberia. *Euroasian Entomological Journal*, 16(2): 173–179. [In Russian].
- Sidorov D.A., Proshchalykin M.Yu., Konusova O.L., Belova K.A. and Byvaltsev A.M.** 2017. The bees of the genus *Andrena* Fabricius (Hymenoptera, Apoidea: Andrenidae) of Western Siberia. *Proceedings of the Russian Entomological Society*, 88(2). [In press].
- Sitdikov A.A.** 1986. The fauna of bees (Hymenoptera, Apoidea) of Udmurtia, with description of *Melitta udumurica* sp. n. *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, 159: 103–112. [In Russian].
- Sitdikov A.A.** 1988. The taxonomy of the bee genus *Eucera* Scopoli (Hymenoptera, Anthophoridae) of the fauna of the USSR and neighbouring countries: the subgenus *Pteneucera* Tkalcu. *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, 175: 102–111. [In Russian].
- Sitdikov A.A. and Pesenko Yu.A.** 1988. The subgeneric classification of bees of the genus *Eucera* Scopoli (Hymenoptera, Anthophoridae), with a scheme of the phylogenetic relationships between the subgenera. *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, 175: 75–101. [In Russian].
- Skhirtladze I.A.** 1981. The Bees of the Transcaucasus (Hymenoptera, Apoidea). Metsniereba, Tbilisi, 148 p. [In Russian].
- Skorikov A.S.** 1922. Bumblebees of the Petrograd Province. In: *Faunae Petropolitanae catalogus. Vol. 2*. Petrogradskii agronomicheskii institut, Petrograd, 51 p.
- Skorikov A.S.** 1927. Sur la sousfam. des Myrmillini (Myrmecidae, Hymenoptera) dans la région Paléarctique. *Ezhegodnik Zoologicheskogo Muzeya*, 28(1): 33–47.
- Skorikov A.S.** 1931. Die Hummelfauna Turkestans und ihre Beziehungen zur zentralasiatischen Fauna (Hymenoptera, Bombidae). In: Lindholm V.A. (Ed.). *Abhandlungen der Pamir-Expedition 1928. Vol. 8*. Academy of Sciences of the USSR, Leningrad: 175–247.
- Skorikov A.S.** 1935. Zur Mutilliden-Fauna Zentralasiens. *Trudy Tadzhikskoi Basy Akademii Nauk SSSR*, 5: 257–349.
- Smith D.R.** 1975. The sawfly genus *Hemicroa* Stephens: A review of species (Hymenoptera: Tenthredinidae). *Entomologica Scandinavica*, 6: 297–302.
- Smith D.R.** 1981. Studies on the leaf-mining sawflies of the tribe Fenusini in Asia (Hymenoptera: Tenthredinidae). *Proceedings of the Entomological Society of Washington*, 83(4): 763–771.
- Smith D.R.** 2008. Xiphydriidae of the Philippines, Insular Malaysia, Indonesia, Papua New Guinea, New Caledonia, and Fiji (Hymenoptera). *Beiträge zur Entomologie*, 58(1): 15–95.
- Smith D.R. and Altenhofer E.** 2011a. A new elm leafmining sawfly (Hymenoptera: Tenthredinidae) from Russia. *Proceedings of the Entomological Society of Washington*, 113(1): 50–56.
- Smith D.R. and Altenhofer E.** 2011b. The second record of a sawfly leafminer on *Rosa*, a new species of *Fenusia* (Hymenoptera: Tenthredinidae). *Proceedings of the Entomological Society of Washington*, 113(1): 57–60.
- Smith D.R. and Shinohara A.** 2002. The stem-boring sawfly genus *Cephus* Latreille (Hymenoptera: Cephidae) in Japan. *Proceedings of the Entomological Society of Washington*, 104(2): 479–484.
- Smith D.R. and Shinohara A.** 2011. Review of the Asian wood-boring genus *Euxiphydria* (Hymenoptera, Symphyta, Xiphydriidae). *Journal of Hymenoptera Research*, 23: 1–22.
- Smith D.R., Tripotin P. and Shinohara A.** 2011. Xiphydriid woodwasps (Hymenoptera: Xiphydriidae) of Korea. *Proceedings of the Entomological Society of Washington*, 113(1): 61–70.

- Smith D.R. and Wei M.** 2015. A new Asian *Monophadnoides* Ashmead (Hymenoptera: Tenthredinidae) with high antennal crests. *Proceedings of the Entomological Society of Washington*, **117**(2): 203–208.
- Smith L. and Gourlay H.** 2016. What's happening with old man's beard? *What's new in biological control of weeds?*, **76**: 4–5.
- Sokolov G.I.** 2014. The pine web-spinning sawfly (*Acantholyda posticalis*) in Chelyabinsk Region and measures to control its population. *Forestry Bulletin*, **6**: 110–118. [In Russian].
- Soon V. and Rosa P.** 2015. Fauna Europaea: Hymenoptera, Chrysidae. Fauna Europaea version 2.6.2. Available from: <http://www.faunaeur.org> (accessed 15 February 2017).
- Spradberry J.P.** 1973. *Wasps: an account of the biology and natural history of solitary and social wasps*. University of Washington Press, Seattle, 408 p.
- Standfuss K. and Schwarz M.** 2007. Zur aktuellen Bienenaufauna der Ölbaumzone in SO-Thessalien / Griechenland (Hymenoptera: Apoidea: Apiformes). 2. Die parasitischen Bienen (pro parte: Apidae, Megachilidae, Halictidae). *Entomofauna*, **28**: 293–320.
- Steinberg D.M.** 1962. *Family Scoliidae*. Nauka, Leningrad, 189 p. (Fauna of the USSR. N. S., 84, Hymenoptera, 13.) [In Russian].
- Stepanov A.D., Nogovitsyna S.N., Popov A.A. and Sivtseva L.V.** 2007. List of insects and spiders of protected areas of the Republic Sakha (Yakutia). In: Labutin Yu.V. (Ed.). *Raznoobrazie nasekomykh i paukov osobo okhranyaemykh prirodnnykh territoriy Yakutii*. IBPK SO RAN, Yakutsk: 90–159. [In Russian].
- Stöckhert E.** 1944. Über die Gruppe der *Nomada furva* Panz. (Hym. Apid.). *Deutsche Entomologische Zeitschrift*, **3/4**: 89–126.
- Straka J., Bogusch P. and Přidal A.** 2007. Apoidea: Apiformes (včely). *Acta Entomologica Musei Nationalis Pragae*, Suppl., **11**: 241–299.
- Straka J. and Rozen J.G.** 2012. First observations on nesting and immatures of the bee genus *Ancyla* (Apoidea: Apidae: Apinae: Ancylini). *American Museum Novitates*, **3749**: 1–24.
- Stroganova V.K.** 1960. To the fauna of the sawflies (Tenthredinidae) of Western Siberia. *Materialy III nauchno-tehnicheskoy konferentsii molodykh uchenykh. SO AN SSSR*, Novosibirsk: 35–38. [In Russian].
- Stroganova V.K.** 1966. A new species of sawflies of the genus *Arge* Schr. (Tenthredinidae, Hymenoptera). *Trudy Biologicheskogo instituta SO AN SSSR*, **22**: 105–108. [In Russian].
- Stroganova V.K.** 1968. *Woodwasps of Siberia*. Nauka, Novosibirsk, 148 p. [In Russian].
- Stroganova V.K.** 1970. Materials to the fauna of Argidae of Western Siberia. *Trudy Biologicheskogo instituta SO AN SSSR*, **26**: 123–129. [In Russian].
- Stroganova V.K.** 1973. Little-known sawflies living on the false hellebore. In: Cherepanov A.I. (Ed.). *Novye i maloizvestnye vidy fauny Sibiri*, **7**: 86–97 + 147. [In Russian].
- Stroganova V.K.** 1976. New and little-known species of the family Cephidae (Hymenoptera) in the fauna of West Siberia. In: Cherepanov A.I. (Ed.). *Novosti fauny Sibiri. Novye i maloizvestnye vidy fauny Sibiri*, **10**: 168–174. [In Russian].
- Stroganova V.K.** 1978a. To fauna of Symphyta (Hymenoptera) of Barguzinsky Reserve. In: Zolotarenko G.S. (Ed.). *Trudy Biologicheskogo instituta SO AN SSSR*, **34**: 101–103. [In Russian].
- Stroganova V.K.** 1978b. New species of sawflies (Hymenoptera, Tenthredinoidea) from Siberia. In: Cherepanov A.I. (Ed.). *Taksonomiya i ekologiya chlenistono-gikh Sibiri. Novye i maloizvestnye vidy fauny Sibiri*, **12**: 144–148. [In Russian].
- Stroganova V.K.** 1980. New data on the morphology and distribution of some sawflies of the genus *Tenthredo* L. In: Cherepanov A.I. (Ed.). *Sistemmatika i ekologiya zhivotnykh. Novye i maloizvestnye vidy fauny Sibiri*, **14**: 95–104. [In Russian].
- Stroganova V.K.** 1981. New species of stem sawfly of the genus *Cephus* Latreille (Hymenoptera, Symphyta). In: Cherepanov A.I. (Ed.). *Nasekomye i kleshchi Sibiri. Novye i maloizvestnye vidy fauny Sibiri*, **15**: 50–56. [In Russian].
- Stroganova V.K.** 1982. New to the fauna of stem sawflies (Hymenoptera, Tenthredinoidea) of Western Siberia. In: Cherepanov A.I. (Ed.). *Fauna Sibiri. Gel'minty, kleshchi i nasekomye*, Nauka: Novosibirsk: 47–53. [In Russian].
- Stroganova V.K.** 1987. The use of new morphological characters in taxonomy of stem sawflies. In: Cherepanov A.I. (Ed.). *Nasekomye i kleshchi Sibiri. Novye i maloizvestnye vidy fauny Sibiri*, **19**: 4–9. [In Russian].
- Stroganova V.K.** 1988. A new species of the genus *Rhogogaster* Konow (Hymenoptera, Symphyta) from Siberia. In: Zolotarenko G.S. (Ed.). *Taksonomiya zhivotnykh Sibiri. Novye i maloizvestnye vidy fauny Sibiri*, **20**: 98–102. [In Russian].
- Strumia F.** 2008. Order Hymenoptera, family Chrysidae. In: van Harten A. (Ed.). *Arthropod fauna of the UAE, Vol. 1*. Dar Al Ummah Printing, Publishing, Distribution & Advertising, Abu Dhabi: 375–387.
- Suárez F.J.** 1988. Mirmosidos de la Peninsula Iberica (Hymenoptera, Myrmecidae). *Graellsia*, **44**: 81–158.
- Sugihara Y. and Kim H.K.** 1936. A new species of Sapygidae from Korea (Hym.). *Insecta Matsumurana*, **10**(3): 123–126.
- Sundukov Yu.N.** 2007. *Arge stroganovae* nom. n. – a new replacement name for *Arge altaica* Stroganova, 1977 (Hymenoptera, Symphyta, Argidae). *Far Eastern Entomologist*, **178**: 11.
- Sundukov Yu.N.** 2009. Suborder Symphyta – Sawflies. In: Storozhenko S.Yu. (Ed.). *Nasekomye Lazovskogo zapovednika*, Dalnauka, Vladivostok: 212–220. [In Russian].

- Sundukov Yu.N. 2010.** A new species of the genus *Apethymus* Benson, 1939 (Hymenoptera, Tenthredinidae) from Sikhote-Alin Mountains, Russian Far East. *Far Eastern Entomologist*, **212**: 1–6.
- Sundukov Yu.N. 2011a.** The new record of the *Xiphydriola amurensis* Semenov, 1921 (Hymenoptera, Symphyta, Xiphydriidae) for the fauna of Russia. *Amurian Zoological Journal*, **3**(4): 378–380. [In Russian].
- Sundukov Yu.N. 2011b.** A preliminary list of the insects (Insecta) and spiders (Aranei) of National Park "Zov tigra". In: Bersenev Yu.I. (Ed.). *Fauna natsionalnogo parka "Zov tigra". Annotirovannye spiski vidov*. Afrika, Vladivostok: 97–131. [In Russian].
- Sundukov Yu.N. 2013.** List of insects and spiders of Baikal-sky Reserve. In: Boichenko V.S. (Ed.). *Rezul'taty izuchenija prirodnnykh kompleksov Baikal'skogo zapovednika. Trudy Baikal'skogo gosudarstvennogo prirodnogo biosfernogo zapovednika*, **4**: 95–147. [In Russian].
- Sundukov Yu.N. 2014.** First records of the *Orussus coreanus* Takeuchi, 1938 and *O. rufipes* Tsuneki, 1963 (Hymenoptera: Orussidae) from Russia. *Amurian Zoological Journal*, **6**(1): 81–84. [In Russian].
- Sundukov Yu.N. 2015.** To the sawfly fauna (Hymenoptera, Symphyta) of Southern Kuriles. *A.I. Kurentsov's Annual Memorial Meetings*, **26**: 241–258. [In Russian].
- Sundukov Yu.N. 2017.** A new data on the Symphyta fauna (Hymenoptera) from Russia]. *Proceedings of the Russian Entomological Society*, **88**(2). [In press]. [In Russian].
- Sundukov Yu.N. and Lelej A.S. 2009.** Sawflies (Hymenoptera, Symphyta) of the Russian Far East. Additions and corrections. *Far Eastern Entomologist*, **200**: 1–12.
- Sundukov Yu.N. and Lelej A.S. 2012.** Suborder Symphyta – Sawflies. In: Lelej A.S. (Ed.). *Annotated Catalogue of the Insects of Russian Far East. Vol. 1. Hymenoptera*. Dalnauka, Vladivostok: 62–119. [In Russian].
- Sung I.-H., Dubitzky A., Eardley C. and Yamane S. 2009.** Descriptions and biological notes of *Ctenoplectra* bees from Southeast Asia and Taiwan (Hymenoptera: Apidae: Ctenoplectrini) with a new species from North Borneo. *Entomological Science*, **12**(3): 324–340.
- Szentgyörgyi H., Grześ I.M., Woyciechowski M., Blinov A., Eremeeva N. and Luzyanin S. 2011.** Bumblebees (Bombidae) along pollution gradient – heavy metal accumulation, species diversity, and *Nosema bombi* infection level. *Polish Journal of Ecology*, **59**(3): 599–610.
- Tadauchi O. 2006.** The genus *Andrena* from Kazakhstan and Kyrgyzstan collected by the Kyushu University expedition (Hymenoptera, Andrenidae) (1). *Esakia*, **46**: 1–19.
- Tadauchi O. 2008.** The genus *Andrena* from Kazakhstan and Kyrgyzstan collected by the Kyushu University expedition (Hymenoptera, Andrenidae) (2). *Esakia*, **48**: 1–18.
- Tadauchi O. and Murao R. 2014.** An Illustrated Guide to Japanese Bees. Bun-ichi Sogo Shuppan Co, Tokyo, 479 p.
- Tadauchi O. and Xu H-I. 1995.** A revision of the subgenus *Simandrena* of the genus *Andrena* of Eastern Asia with a key to Palaearctic species (Hymenoptera, Andrenidae). *Esakia*, **35**: 201–222.
- Tadauchi O. and Xu H-I. 1999.** Subgeneric positions and redescriptions of Cockerell's Siberian *Andrena* preserved in the British Museum (Natural History) (Hymenoptera, Andrenidae). *Esakia*, **39**: 13–30.
- Taeger A. 1985.** Zur Systematik der Blattwespengattung *Tenthredo* (s. str.) L. (Hymenoptera, Symphyta, Tenthredinidae). *Entomologische Abhandlungen*, **48**(8): 83–148.
- Taeger A. 1986.** Beitrag zur Taxonomie und Verbreitung paläarktischer Allantinae (Hymenoptera, Symphyta). *Beiträge zur Entomologie*, **36**(1): 107–118.
- Taeger A. 1988.** Dritter Beitrag zur Kenntnis der Blattwespengattung *Tenthredo* L. (Hymenoptera: Symphyta: Tenthredinidae). *Beiträge zur Entomologie*, **38**(2): 337–359.
- Taeger A. 1989.** Die Gattung *Macrophya* Dahlbom in der DDR (Insecta, Hymenoptera, Symphyta: Tenthredinidae). *Entomologische Abhandlungen*, **53**(5): 57–69.
- Taeger A. 1991.** Vierter Beitrag zur Systematik der Blattwespengattung *Tenthredo* Linnaeus. Die Untergattung *Zonuledo* Zhelochovtsev, 1988 (Hymenoptera, Tenthredinidae). *Entomofauna*, **12**(23): 373–398.
- Taeger A. 1992.** Fünfter Beitrag zur Systematik der Blattwespengattung *Tenthredo* L. (Hymenoptera, Symphyta). *Beiträge zur Entomologie*, **42**(1): 3–53.
- Taeger A. 1998.** Die Megalodontesidae Europas (Hymenoptera: Symphyta). In: Taeger A., Blank S.M. (Eds.). *Pflanzenwespen Deutschlands (Hymenoptera, Symphyta)*. Goecke & Evers, Keltern: 175–192.
- Taeger A. 2002.** The Megalodontesidae of Europe (Hymenoptera, Symphyta). In: Viitasaari M. (Ed.). *Sawflies (Hymenoptera, Symphyta) I. A review of the suborder, the Western Palaearctic taxa of Xyeloidea and Pamphilioidea*, Vol. 1. Tremex, Helsinki: 461–480.
- Taeger A. 2005.** Catalogue of types in the Hymenoptera collection of the DEI – Symphyta. *Leibniz-Zentrum für Agrarlandschafts- und Landnutzungsforschung (ZALF) e. V., Deutsches Entomologisches Institut (DEI)*: 1–92.
- Taeger A. 2013.** The type specimens of *Tenthredo* Linnaeus, 1758 (Hymenoptera: Tenthredinidae) deposited in the Hungarian Natural History Museum. *Zootaxa*, **3626**(2): 201–244.
- Taeger A. and Blank S.M. 2011.** ECatSym – Electronic World Catalog of Symphyta (Insecta, Hymenoptera). Program version 3.9, data version 38 (07.12.2011). http://www.zalf.de/home_zalf/institute/dei/php_e/ecatsym/index.html.
- Taeger A., Blank S.M. and Liston A.D. 2006.** European sawflies (Hymenoptera: Symphyta) – A species checklist for the countries. In: Blank S.M., Schmidt S., Taeger A. (Eds.). *Recent Sawfly Research – Synthesis and Prospects*. Goecke & Evers, Keltern: 399–504.

- Taeger A., Blank S.M. and Liston A.D.** 2010. World catalog of Symphyta (Hymenoptera). *Zootaxa*, **2580**: 1–1064.
- Taeger A. and Kramp K.** 2017. *Tianmuthredo davei*, a new species from the Russian Far East (Hymenoptera: Tenthredinidae). *Proceedings of the Washington Entomological Society*. [In press].
- Taeger A., Shinohara A. and Kramp K.** 2017. *Pachyprotasis kojimai* sp. nov. – the “*Pachyprotasis nigronotata*” of Japanese authors (Hymenoptera: Tenthredinidae). *Zootaxa*, **4227**(4): 583–592.
- Taeger A. and Viitasaari M.** 2015. European *Rhogogaster* s. str., with notes on several Asian species (Hymenoptera: Tenthredinidae). *Zootaxa*, **4013**(3): 369–398.
- Taeger A., Wei M. and Shinohara A.** 2016. Sibling species in *Tenthredo* Linné (Hymenoptera: Tenthredinidae): the status of some East Asian taxa. *Euroasian Entomological Journal*, **15**(Suppl. 1): 156–164.
- Takeuchi K.** 1927. Some Chalastogastra from Corea. *Transactions of the Natural History Society of Formosa*, **17**(93): 378–387.
- Takeuchi K.** 1931. Some sawflies from the Island of Shikotan in the southern Kuriles. *The Transactions of the Kansai Entomological Society*, **2**: 30–34.
- Takeuchi K.** 1932. A revision of the Japanese Argidae. *The Transactions of the Kansai Entomological Society*, **3**: 27–42.
- Takeuchi K.** 1933. Four species of sawflies from the Northern Kuriles. *Bulletin of the Biogeographical Society of Japan*, **4**(2): 86–90.
- Takeuchi K.** 1936. Tenthredinoidea of Saghalien (Hymenoptera). *Tenthredo. Acta Entomologica*, **1**(1): 53–108.
- Takeuchi K.** 1937. A study on the Japanese species of the genus *Macrophyia* Dahlbom (Hymenoptera Tenthredinidae). *Acta Entomologica*, **1**(4): 376–454.
- Takeuchi K.** 1938. A systematic study on the suborder Symphyta (Hymenoptera) of the Japanese Empire (I). *Tenthredo. Acta Entomologica*, **2**(2): 173–229.
- Takeuchi K.** 1939. A systematic study on the suborder Symphyta (Hymenoptera) of the Japanese Empire (II). *Tenthredo. Acta Entomologica*, **2**(4): 393–439.
- Takeuchi K.** 1940. A systematic study on the suborder Symphyta (Hymenoptera) of the Japanese Empire (III). *Tenthredo. Acta Entomologica*, **3**(2): 187–199.
- Takeuchi K.** 1941. A systematic study on the suborder Symphyta (Hymenoptera) of the Japanese Empire (IV). *Tenthredo. Acta Entomologica*, **3**(3): 230–274.
- Takeuchi K.** 1955. Sawflies of the Kurile Islands (I). *Insecta Matsumurana*, **19**(1–2): 9–22.
- Takeuchi K.** 1956. Sawflies of the Kurile Islands (II). *Insecta Matsumurana*, **19**(3–4): 71–81.
- Tan J.-L., van Achterberg K., Duan M.-J. and Chen X.-X.** 2014. An illustrated key to the species of subgenus *Gyrostoma* Kirby, 1828 (Hymenoptera, Vespidae, Polistinae) from China, with discovery of *Polistes* (*Gyrostoma*) *tenuispunctia* Kim, 2001. *Zootaxa*, **3785**(3): 377–399.
- Terayama M.** 2003a. Phylogenetic systematics of the family Bethylidae (Insecta: Hymenoptera). Part I. Higher classification. *Academic Reports. Faculty of Engineering Tokyo Polytechnic University*, **26**(1): 1–15.
- Terayama M.** 2003b. Phylogenetic systematics of the family Bethylidae (Insecta: Hymenoptera). Part 2. Key to subfamilies, tribes and genera in the world. *Academic Reports. Tokyo Polytechnic University, Faculty of Engineering* **26**(1): 16–29.
- Terayama M.** 2006. *Bethylidae (Hymenoptera)*. *The Insect of Japan*. 1. Touka Shobo Co. Ltd., Fukuoka, 319 p.
- Terayama M.** 2016a. Bethylidae. In: Terayama M., Suda H. (Eds). *A Guide to the Aculeate Wasps of Japan*. Tokai University Press, Minamiyama: 335–387, pls. 79–81 + 608–611. [In Japanese].
- Terayama M.** 2016b. Tiphiidae. In: Terayama M., Suda H. (Eds). *A Guide to the Aculeate Wasps of Japan*. Tokai University Press, Minamiyama: 264–289, pls. 59–61. [In Japanese].
- Terayama M. and Nagase H.** 2016. Scoliidae. In: Terayama M., Suda H. (Eds). *A Guide to the Aculeate Wasps of Japan*. Tokai University Press, Minamiyama: 248–259, pls. 54–57 + 599–601. [In Japanese].
- Terayama M., Suda H., Tano T. and Murota T.** 2016. Chrysididae. In: Terayama M., Suda H. (Eds). *A Guide to the Aculeate Wasps of Japan*. Tokai University Press, Minamiyama: 388–415, pls 81–85 + 611–613. [In Japanese].
- Terzo M. and Rasmont P.** 1997. Révision des *Xylocopa* Latreille du sous-genre *Copoxyla* Maa des pays circum-méditerranéens (Hymenoptera, Apoidea). *Bulletin de la Société entomologique de France*, **102**(4): 367–377.
- Terzo M. and Rasmont P.** 2004. Biogéographie et systématique des abeilles rubicoles du genre *Ceratina* Latreille au Turkestan (Hymenoptera, Apoidea, Xylocopinae). *Annales de la Société Entomologique de France*, **40**(2): 109–130.
- Tkalcù B.** 1961. Zur Hummelfauna der Umgebung Kukunors (Hymenoptera, Bombinae). *Casopis Československé Společnosti Entomologické*, **58**: 344–379.
- Tkalcù B.** 1974a. Ergebnisse der 1. und 2. mongolisch-tschechoslowakischen entomologisch-botanischen Expedition in der Mongolei. Nr. 29: Hymenoptera, Apoidea, Bombinae. *Sborník faunistických prací Entomologického oddělení Národního muzea v Praze*, **15**: 25–57.
- Tkalcù B.** 1974b. Eine Hummel-Ausbeute aus dem Nepal-Himalaya (Insecta, Hymenoptera, Apoidea, Bombinae). *Senckenbergiana Biologica*, **55**: 311–349.
- Tkalcù B.** 1978. Beiträge zur Kenntnis der Fauna Afghanistans. *Melitturga* Latr., *Eucera* Scop., Apidae; *Lithurge* Latr., *Stelis* Pz., *Creightonella* Cockll., Megachilidae, Apoidea, Hym. *Acta Musei Moraviae*, **63**: 153–181.
- Tkalcù B.** 1979. Revision der europäischen Vertreter der Artengruppe von *Tetralonia ruficornis* (Fabricius) (Hymenoptera, Apoidea). *Acta Musei Moraviae*, **64**: 127–152.

- Tkalcú B.** 1984a. Systematisches Verzeichnis der westpaläarktischen *Tetralonia* und *Eucera* Arten, deren Männchen als Blütenbesucher verschiedener *Ophrys*-Arten festgestellt wurden. Mit Beschreibung neuer Taxa (Hymenoptera: Apoidea). *Nova Acta Regiae Societatis Scientiarum Upsaliensis*, 5(3): 57–77.
- Tkalcú B.** 1984b. Revision der Gattung *Cubitalia* Friese, 1911 (Hymenoptera, Apoidea). *Annotationes Zoologicae et Botanicae*, 161: 1–15.
- Tobias V.I.** 1965. To the knowledge of the family Fedtschenkiidae (Hymenoptera, Sapygoidea). *Zoologicheskij Zhurnal*, 44: 706–715. [In Russian].
- Tobias V.I.** 1978a. Superfamily Scolioidea. In: Tobias V.I. (Ed.). *Keys to the Insects of the European Part of the USSR*. Vol. III. Pt. 1. Nauka, Leningrad: 47–56. [In Russian].
- Tobias V.I.** 1978b. Superfamily Pompiloidea. In: Tobias V.I. (Ed.). *Keys to the Insects of the European Part of the USSR*. Vol. III. Pt. 1. Nauka, Leningrad: 83–147. [In Russian].
- Tobias V.I.** 1978c. Superfamily Sapygoidea. In: Tobias V.I. (Ed.). *Keys to the Insects of the European Part of the USSR*. Vol. III. Pt. 1. Nauka, Leningrad: 56–58. [In Russian].
- Tobias V.I. and Kurzenko N.V.** 1978. Family Eumenidae. In: Tobias V.I. (Ed.). *Keys to the Insects of the European Part of the USSR*. Vol. III. Pt. 1. Nauka, Leningrad: 152–173. [In Russian].
- Togashi I.** 1973. On the species of *Megalodontes* Latreille (Hym., Symphyta) from Inner Mongolia. *Kontyû*, 41(3): 294–297.
- Togashi I.** 1989. Japanese sawflies of the family Blastocotomidae (Hymenoptera: Symphyta). *Proceedings of the Entomological Society of Washington*, 91(3): 406–413.
- Togashi I.** 1998. Symphyta (Hymenoptera) of Hokkaido, Japan. *Bulletin of the Biogeographical Society of Japan*, 53(1): 39–47.
- Togashi I.** 2000. Japanese sawflies of the *japonicus* group of the genus *Taxonus* Hartig (Hymenoptera: Tenthredinidae). *Proceedings of the Entomological Society of Washington*, 102(2): 313–321.
- Togashi I.** 2008. An additional species of the genus *Orussus* Konow (Hymenoptera, Symphyta, Orissidae) from Japan. *Biogeography*, 10: 23–25.
- Tolstova Yu.S.** 1962. On the fauna of the inhabitants of the raspberries stems in the vicinity of Leningrad. *Entomologicheskoe Obozrenie*, 61: 285–293. [In Russian].
- Tomilova V.N.** 1974. The mined insects of Eastern Siberia and Far East. In: Kulik S.A. (Ed.). *Fauna nasekomykh Vostochnoy Sibiri i Dal'nego Vostoka*. Irkutskiy gosudarstvennyy universitet, Irkutsk: 253–259. [In Russian].
- Trjapitsin V.A.** 1978. Family Bethylidae – Bethylids. In: Trjapitsin V.A. (Ed.). *Key to the insects of the European part of the USSR*. Vol. III, pt 2. Nauka, Leningrad: 6–16. [In Russian].
- Trjapitzin V.A.** 1978. Family Embolemidae – Embolemids. In: Trjapitsin V.A. (Ed.). *Key to the insects of the European part of the USSR*. Vol. III, pt 2. Nauka, Leningrad: 27–28. [In Russian].
- Trunz V., Packer L., Vieu J., Arrigo N. and Praz C.J.** 2016. Comprehensive phylogeny, biogeography and new classification of the diverse bee tribe Megachilini: Can we use DNA barcodes in phylogenies of large genera? *Molecular Phylogenetics and Evolution*, 103: 245–259.
- Tsuneki K.** 1947. Chrysidae from North China and Inner Mongolia. *Mushi*, 17(9): 43–60.
- Tsuneki K.** 1948. Chrysidae from Shansi, North China (Hymenoptera). *Mushi*, 18(19): 115–131.
- Tsuneki K.** 1950. Descriptions of new species and subspecies of the Chrysidae from East Asia (Hymenoptera). *Mushi*, 21(8): 21–81.
- Tsuneki K.** 1952. The genus *Rhopalum* Kirby (1829) of Japan, Korea, Saghalien and the Kuriles, with a suggested reclassification of the subgenera and descriptions of four new species (Hym. Sphecidae, Crabroninae). *Journal of the Faculty of Sciences, Hokkaido Imperial University (Series VI, Zoology)*, 11: 110–125.
- Tsuneki K.** 1953a. Chrysidae of Manchuria (Hymenoptera). *Mushi*, 25(8): 53–61.
- Tsuneki K.** 1953b. Chrysidae of Korea (Hymenoptera). *Kontyû*, 20(1–2): 22–28.
- Tsuneki K.** 1954. The genus *Crossocerus* Lepeletier et Brûlé (1834) of Japan, Korea, Saghalien and the Kuriles. *Memoirs of the Faculty of Liberal Arts, Fukui University (Series II, Natural Science)*, 3: 57–78.
- Tsuneki K.** 1955a. *Chrysis (Pentachrysis)* of North-Eastern Asia (Hymenoptera, Chrysidae). *Memoirs of the Faculty of Liberal Arts, Fukui University, Series II, Natural Science*, 5(11): 35–46.
- Tsuneki K.** 1955b. Crabroninae of the Kurile Islands (Hymenoptera, Sphecidae) with remarks on new specific synonymy. *Bulletin of the Biogeographical Society of Japan*, 16–19: 212–219.
- Tsuneki K.** 1956. Die Trypoxylonen der nordöstlichen Gebiete Asiens (Hymenoptera, Sphecidae, Trypoxyloninae). *Memoirs of the Faculty of Liberal Arts, Fukui University (Series II, Natural Science)*, 6: 1–42.
- Tsuneki K.** 1959. Contributions to the knowledge of the Cleptinae and Pseninae Faunae of Japan and Korea (Hymenoptera, Chrysidae and Sphecidae). *Memoirs of the Faculty of Liberal Arts, Fukui University, Series II, Natural Science*, 9: 1–78.
- Tsuneki K.** 1961. Studies on *Cerceris* of North Eastern Asia (Hymenoptera, Sphecidae). *Memoirs of the Faculty of Liberal Arts, Fukui University (Series II, Natural Science)*, 11: 1–72.
- Tsuneki K.** 1963. *Chrysis (Hexachrysis)* of Eastern Asia. *Etzienia*, 3: 1–9.
- Tsuneki K.** 1963. The tribe Gorytini of Japan and Korea (Hymenoptera, Sphecidae). *Etzienia*, 1: 1–20.

- Tsuneki K.** 1971a. Ergebnisse der zoologischen Forschungen von Dr. Z. Kaszab in der Mongolei. 239. Sphecidae (Hymenoptera). I-II. *Acta Zoologica Academiae Scientiarum Hungaricae*, 17: 139–217.
- Tsuneki K.** 1971b. Sphecididae aus der Inneren Mongolei und dem nördlichen China (Hym.). *Etzenzia*, 58: 1–38.
- Tsuneki K.** 1972. Studies on the Mutillidae of Japan (Hymenoptera). *Etzenzia*, 61: 1–26.
- Tsuneki K.** 1979. Sphecidae and Chrysididae collected by Dr. K. Baba in northern part of Japan proper and Hokkaido, with descriptions of two new species. In: *Insects of Niigata Prefecture, Japan*: 9–14.
- Tsuneki K.** 1981. Revision of the *Trypoxylon* species of Japan and northeastern part of Asiatic continent, with comments on some species of Europe (Hymenoptera, Sphecidae). *Special Publications of the Japan Hymenopterists Association*, 17: 1–92.
- Trautmann W.** 1927. *Die Goldwespen Europas*. Uschman, Weimar, 194 p. + 2 pls.
- Turillazzi S.** 1991. The Stenogastrinae. In: Ross K.G., Matthews R.W. (Eds). *The social biology of wasps*. Ithaca, Comstock: 74–98.
- Uchida T.** 1927. Eine uebersicht der Chrysididen Japans und mit den beschreibungen der neuen Arten und Varietaeten. *Insecta Matsumurana*, 1(3): 149–157.
- van der Vecht J. and Carpenter J.M.** 1990. A catalogue of the genera of the Vespidae. *Zoologische Verhandelingen*, 260: 1–62.
- van der Vecht J. and Fischer F.C.J.** 1972. *Hymenopterorum Catalogus. Pars 8. Palaearctic Eumenidae*. 's- NV. Junk, Gravenhage, vi + 199 p.
- Vargas-Rojas J.M. and Terayama M.** 2006. Familia Bethylidae. In: Fernández F., Sharkey M.J. (Eds). *Introducción a los Hymenoptera de la Región Neotropical*. Sociedad Colombiana de Entomología y Universidad Nacional de Colombia, Bogotá, D.C.: 427–442.
- Varrone R. and Olmi M.** 2012. First record of host of *Embolemus ruddii* Westwood (Hymenoptera Embolemidae). *Frustula Entomologica. N.S.*, 33: 91–95.
- Vasilenko S.V.** 2004. Suborder Symphyta. In: Mordkovich V.V. (Ed.). *Biodiversity of the Sokhondo Nature Reserve. Arthropoda*, Novosibirsk–Chita: 348–350. [In Russian].
- Vasilenko S.V.** 2006. The sawflies (Hymenoptera, Symphyta: Tenthredinidae, Tenthredininae) of Novosibirsk Area. *Entomologicheskie issledovaniya v Severnoy Azii. Materialy VII Mezhdunarodnogo soveshchaniya entomologov Sibiri i Dal'nego Vostoka v ramkakh Sibirskoy zoologicheskoy konferentsii. Novosibirsk, 20–24 sentyabrya 2006 g.*, Novosibirsk: 39–40. [In Russian].
- Vasilenko S.V.** 2007. About two type species of saw-flies (Hymenoptera, Symphyta: Tenthredinidae, Cephidae) described by V.K. Stroganova. *Euroasian Entomological Journal*, 6(1): 34–35. [In Russian].
- Vasilenko S.V.** 2008. *Tenthredo pamiriensis* Jakovlev, 1888 (Hymenoptera, Symphyta, Tenthredinidae), a new species for the fauna of Russia. *Euroasian Entomological Journal*, 7(4): 356. [In Russian].
- Vasilenko S.V.** 2009a. Two new sawfly species of the genus *Aprosthemia* Konow, 1899 (Hymenoptera, Symphyta, Argidae) from West Siberia. *Euroasian Entomological Journal*, 8(1): 27–30. [In Russian].
- Vasilenko S.V.** 2009b. Data on the fauna of sawflies (Hymenoptera, Symphyta) of the Bolshekhekhtsirsky Nature Reserve. *Amurian Zoological Journal*, 1(1): 83–87. [In Russian].
- Vasilenko S.V.** 2010a. The sawflies of the *Tenthredo marginella* Fabricius, 1793-group (Hymenoptera, Symphyta, Tenthredinidae) in West Siberia. *Euroasian Entomological Journal*, 9(3): 492–494. [In Russian].
- Vasilenko S.V.** 2010b. Data on the fauna of sawflies (Hymenoptera, Symphyta) in environs of Teletskoye Lake. *Altayskiy Zoologicheskiy Zurnal*, 4: 3–15. [In Russian].
- Vasilenko S.V.** 2010c. A new species of the genus *Arge* Schrank, 1802 (Hymenoptera, Argidae) from West Siberia. *Euroasian Entomological Journal*, 9(4): 681–682. [In Russian].
- Vasilenko S.V.** 2010d. The review of the cimbicid species of subgenus *Zaraea* Leach, 1817 (Hymenoptera: Cimbicidae: *Abia* Leach) of Siberia. *Amurian Zoological Journal*, 2(4): 338–340. [In Russian].
- Vasilenko S.V.** 2011a. Species of the genus *Athalia* Leach, 1817 (Hymenoptera, Tenthredinidae, Allantiinae) in the Asian part of Russia. *Euroasian Entomological Journal*, 10(2): 197–200. [In Russian].
- Vasilenko S.V.** 2011b. Data on the fauna of sawflies (Hymenoptera, Symphyta) of Novosibirsk Oblast. 2. Cephidae, Siricidae, Xyphidiidae, Blasticotomidae, Diprionidae. *Euroasian Entomological Journal*, 10(1): 113–116. [In Russian].
- Vasilenko S.V.** 2011c. *Trichiosoma nanae* Vikberg et Viita-saari, 1991 (Hymenoptera, Symphyta: Cimbicidae) – a new species for the fauna of Russia. *Amurian Zoological Journal*, 3(2): 199–200. [In Russian].
- Vasilenko S.V. and Dolgov I.S.** 2005. A contribution to the fauna of saw-flies (Hymenoptera, Symphyta) of the Novosibirsk Oblast'. Part 1. Pamphiliidae, Megalodontesidae, Argidae, Cimbicidae. *Euroasian Entomological Journal*, 4(1): 57–62. [In Russian].
- Vasilenko S.V. and Korshunov A.V.** 2012. On the sawfly fauna (Hymenoptera, Symphyta) of Kemerovskaya Oblast. *Euroasian Entomological Journal*, 11(3): 271–275. [In Russian].
- Verzhbitskii B.N.** 1973. *Keys to the larvae of the wood-wasps and sawflies of Siberia and the Soviet Far East*. Nauka, Moscow. 140 p. [In Russian].

- Verzhutskii B.N.** 1974. Fauna of Symphyta (Hymenoptera) of Eastern Siberia. In: Kulik S.A. (Ed.). *Fauna nasekomykh Vostochnoi Sibiri i Dal'nego Vostoka*. Irkutskiy gosudarstvennyy universitet, Irkutsk: 194–243. [In Russian].
- Verzhutskii B.N.** 1981. *Herbivorous insects in ecosystems of East Siberia (sawflies and wood-wasps)*. Nauka, Novosibirsk, 303 p. [In Russian].
- Vilhelmsen L., Blank S.M., Liu Z. and Smith D.R.** 2014. Discovery of new species confirms Oriental origin of *Orussus* Latreille (Hymenoptera: Orussidae). *Insect Systematics & Evolution*, 45(1): 51–91.
- Viitasaari M.** 1989. Taxonomic notes on the genus *Trichiosoma* Leach (Hymenoptera, Cimbicidae). II. *Annales Entomologici Fennici*, 55(3): 111–129.
- Viitasaari M.** 2002. The suborder Symphyta of the Hymenoptera. In: Viitasaari M. (Ed.). *Sawflies (Hymenoptera, Symphyta) I. A review of the suborder, the Western Palaearctic taxa of Xyloidea and Pamphilioidea*. Tremex, Helsinki: 11–174.
- Viitasaari M. and Varama M.** 1987. Sahapistäiset 4. Havupistäiset (Diprionidae). *Helsingin yliopisto, Maatalous- ja metsäläintieteen laitos, Julkaisuja*, 10: 1–78.
- Viitasaari M. and Zinovjev A.G.** 1991. *Taxonus zhe-lochovtsevi* sp. n. and *Apethymus parallelus* (Eversmann, 1847) from the Soviet Far East (Hymenoptera, Tenthredinidae). *Entomologica Fennica*, 2(3): 175–178.
- Vikberg V.** 2003. Taxonomy of the species of *Eupontania crassipes*- and *aquilonis*-groups (Hymenoptera: Tenthredinidae: Nematinae). *Entomologica Fennica*, 14(3): 129–172.
- Vikberg V.** 2004. Seasonal head dimorphism and taxonomy of some European species of *Aprosthemma* (Hymenoptera: Symphyta: Argidae). *Beiträge zur Entomologie*, 54(1): 107–125.
- Vikberg V.** 2009. On the taxonomy of *Phyllocolpa scotaspis* (Förster, 1854) and *Phyllocolpa anglica* (Cameron, 1877) and notes on the species groups of *Phyllocolpa* (Hymenoptera: Tenthredinidae: Nematinae). *Sahlbergia*, 15: 3–13.
- Vikberg V.** 2010. European species of *Tubpontania* gen. nov., a new genus for species of the *Pontania crassispina* group. *Zootaxa*, 2620: 1–28.
- Vikberg V.** 2013. Hemp sawfly found in European Russia (Hymenoptera: Tenthredinidae, Nematinae, Cladiini). *Entomologica Fennica*, 24: 172–178.
- Vikberg V.** 2017. Two sawflies new for the fauna of Russia, one being new for Europe and the western Palaearctic region (Hymenoptera: Symphyta: Heptamelidae and Tenthredinidae). *Sahlbergia*, 23(1): 17–19.
- Vikberg V. and Liston A.D.** 2009. Taxonomy and biology of European Heptamelini (Hymenoptera, Tenthredinidae, Selandriinae). *Zootaxa*, 2112: 1–24.
- Vikberg V. and Viitasaari M.** 1991. *Trichiosoma nanae* sp. n., a monophag on *Betula nana* from Finland (Hymenoptera, Cimbicidae). *Entomologica Fennica*, 2: 67–77.
- Vikberg V. and Zinovjev A.G.** 2006. On the taxonomy and host plants of North European species of *Eupontania* (Hymenoptera: Tenthredinidae: Nematinae). *Beiträge zur Entomologie*, 56: 239–268.
- Vikberg V. and Zinovjev A.G.** 2014. Three new species of the *Pontania dolichura* group from Europe and Siberia (Hymenoptera: Tenthredinidae: Nematinae). *w-album*, 18: 3–15.
- Vinokurov N.B.** 2004. Chrysidid wasps (Hymenoptera, Chrysidiidae) in specially protected eco-resort region of Caucasian Mineral Waters. In: *Actual problems of preserving the stability of living systems. Materials VIII International scientific ecological conference*. Belgorod, 27–29 September 2004. Belgorod: 33–34. [In Russian].
- Vinokurov N.B.** 2006. Daily activity and seasonal dynamics of flight in cuckoo wasps (Hymenoptera, Chrysidiidae) in Central Ciscaucasia. In: *Problems of Ecology of Mountain Territories: Collection of scientific papers*. Moscow: 19–21. [In Russian].
- Vinokurov N.B.** 2007. Laccoith mountains and cretaceous ridges of the Caucasian Mineral spas as refugium for the fauna of cuckoo wasps from the subfamily Hedychriniae in foothills of the Central Caucasus. *Proceedings of the International Conference "Mountain ecosystems and their components". Association of scientific editions*, 13–18 August 2007. Moscow: 138–142. [In Russian].
- Vinokurov N.B.** 2009. Cuckoo wasps (Hymenoptera, Chrysidiidae) in steppe biocenoses of the Middle Kama region (North Caucasus). *Steppes of Northern Eurasia: Materials of the V International Symposium*. Orenburg: 206–208. [In Russian].
- Vinokurov N.B.** 2010. Zoogeographical complexes of cuckoo wasps (Hymenoptera: Chrysidiidae) in the fauna of high mountains of the Central Caucasus. *Proceedings of the Russian Entomological Society*, 81(2): 39–42. [In Russian].
- Vinokurov N.B.** 2012. The review of the cuckoo wasps species (Hymenoptera, Chrysidiidae, *Chrysura*) from the group *dichroa* from the central part of the North Caucasus. In: *Ecology, Evolution and Systematics of Animals. Materials of IV International Scientific Conference*, 13–16 November 2012. Ryazan: 53–54. [In Russian].
- Vinokurov N.B.** 2013. Peculiarities in ecology of cuckoo wasps (Hymenoptera, Chrysidiidae) in soil biocenoses from the northern macroslope of the Central Caucasus. *Proceedings of the Samara Scientific Center of the Russian Academy of Sciences*, 15[3(3)]: 1105–1109. [In Russian].
- Vorontsovskij P.A.** 1930. Materials for the study of the Orenburg District Chrysidiidae (Hymenoptera). *Proceedings of the Society for the Study of the Orenburg District*, 1(2): 1–72. [In Russian].
- Wahis R.** 1986. Catalogue systematique et codage des Hyménoptères Pompiliides de la région ouest-européenne. *Notes fauniques de Gembloux*, 12: 1–91.

- Wahis R.** 2013. Fauna Europaea: Hymenoptera, Pompilidae. *Fauna Europaea version 2.6.2*. Available from <http://www.faunaeur.org> (accessed 15 February 2017).
- Waichert C., Rodriguez J., Wasbauer M.S., von Dohlen C.D. and Pitts J.P.** 2015. Molecular phylogeny and systematics of spider wasps (Hymenoptera: Pompilidae): redefining subfamily boundaries and the origin of the family. *Zoological Journal of the Linnean Society*, **175**: 271–287.
- Wang M., Rasnitsyn A.P., Li H., Shih Ch., Sharkey M.J. and Ren D.** 2016. Phylogenetic analyses elucidate the inter-relationships of Pamphilioidea (Hymenoptera, Symphyta). *Cladistics*, **32**: 239–260.
- Ward P.S., Blaimer B.B. and Fisher B.L.** 2016. A revised phylogenetic classification of the ant subfamily Formicinae (Hymenoptera: Formicidae), with resurrection of the genera *Colobopsis* and *Dinomyrmex*. *Zootaxa*, **4072**: 343–357.
- Warncke K.** 1972. Beitrag zur Systematik und Verbreitung der Bienengattung *Prosopis* F. in der Westpaläarktis (Hymenoptera, Apoidea, Colletidae). *Bulletin des Recherches agronomique de Gembloux, N.S.*, **5**: 746–768.
- Warncke K.** 1973. Die westpaläarktische Arten der Bienen Familie Melittidae (Hymenoptera). *Polskie Pismo Entomologiczne*, **43**: 97–126.
- Warncke K.** 1975. Beitrag zur Systematik und Verbreitung der Furchenbienen in der Türkei (Hymenoptera, Apoidea, *Halictus*). *Polskie Pismo Entomologiczne*, **45**(1): 81–128.
- Warncke K.** 1978. Über die westpaläarktischen Arten der Bienengattung *Colletes* Latr. (Hymenoptera, Apoidea). *Polskie Pismo Entomologiczne*, **48**: 329–370.
- Warncke K.** 1979. Beiträge zur Bienenfauna des Iran: 10. Die Gattung *Ancyla* Lep., mit einer Revision der Bienengattung *Ancyla* Lep. *Bollettino del Museo civico di storia naturale di Venezia*, **30**: 183–195.
- Warncke K.** 1980. Die Bienengattung *Anthidium* Fabricius, 1804 in der Westpaläarktis und im turkestanischen Becken. *Entomofauna*, **1**(10): 119–210.
- Warncke K.** 1982a. Beitrag zur Bienenfauna des Iran 14. - Die Gattung *Halictus* Latr., mit Bemerkungen über bekannte und neue *Halictus*-Arten in der Westpaläarktis und Zentralasien. *Bollettino del Museo Civico di Storia Naturale di Venezia*, **32**: 67–166.
- Warncke K.** 1982b. Die Holzbienen des Vorderen Orients (Hym., Apidae). *Linzer Biologische Beiträge*, **14**(1): 23–37.
- Warncke K.** 1983. Zur Kenntnis der Bienengattung *Pasites* Jurine, 1807, in der Westpaläarktis (Hymenoptera, Apidae, Nomadinae). *Entomofauna*, **4**(21): 261–334.
- Warncke K.** 1992. Die westpaläarktischen Arten der Bienengattung *Sphecodes* Latr. *Bericht der Naturforschenden Gesellschaft Augsburg*, **52**: 9–64.
- Wei M.** 1997a. Further studies on the tribe Fenusini (Hymenoptera: Tenthredinidae). *Acta Zootaxonomica Sinica*, **22**(3): 286–300.
- Wei M.** 1997b. Revision of the genus *Caliroa* O. Costa (Hymenoptera, Heterarthridae) from China. *Entomotaxonomia*, **19**(Suppl.): 51–59.
- Wei M.** 1998a. Study on the tribe Heterarthrini of China (Hymenoptera: Tenthredinidae). *Journal of Jishou University (Natural Science Edition)*, **19**(3): 6–10.
- Wei M.** 1998b. Two new genera of Hoplocampinae from China with a key to known genera of the subfamily in the world (Hymenoptera: Nematidae). *Journal of Central-South Forestry University*, **18**(4): 12–18.
- Wei M.** 2006. On the genus *Casipteryx* Taeger (Hymenoptera, Tenthredinidae) with description of a new species from China. *Acta Zootaxonomica Sinica*, **31**(3): 617–620.
- Wei M., Lee S.-B., Choi J.-K. and Lee J.-W.** 2017. Taxonomy of *Sterictiphora* Billberg (Hymenoptera: Argidae: Sterictiphorinae) in South Korea with descriptions of two new species. *Entomological Research*, **47**(2): 63–73.
- Wei M. and Nie H.** 1997a. Studies on Chinese Cephidae IV – Four new species of *Janus* Stephens with a list of Chinese Cephidae (Hymenoptera). *Entomotaxonomia*, **19**(2): 146–152.
- Wei M. and Nie H.** 1997b. Studies on the genus *Conaspidia* Konow (Hymenoptera: Tenthredinidae) from China with a key to the known species of the world. *Entomotaxonomia*, **19**(Suppl.): 95–117.
- Wei M. and Nie H.** 1999. New species of Blasticotomidae (Hymenoptera: Blasticotomorpha) from China with keys to known genera and species of the family. *Entomotaxonomia*, **21**(1): 51–59.
- Wei M. and Nie H.** 2008. A new species of *Megadineura* Malaise (Hymenoptera: Tenthredinidae) and a key to the known species of the genus. *Zootaxa*, **1920**: 51–60.
- Wei M., Nie H. and Taeger A.** 2006. Sawflies (Hymenoptera: Symphyta) of China – checklist and review of research. In: Blank S.M., Schmidt S., Taeger A. (Eds). *Recent Sawfly Research – Synthesis and Prospects*. Goecke & Evers, Keltern: 505–574.
- Wei M. and Niu G.** 2010. Review of the genus *Eriocampopsis* Takeuchi (Hymenoptera: Tenthredinidae) with description of a new species from China. *Proceedings of the Entomological Society of Washington*, **112**(2): 295–301.
- Wei M. and Smith D.R.** 2010. Review of *Syrista* Konow (Hymenoptera: Cephidae). *Proceedings of the Entomological Society of Washington*, **112**(2): 302–316.
- Wei M. and Wen J.** 2002. Argidae. In: Li Z., Jin D. (Eds). *Insects from Maolan Landscape*, Guizhou Science and Technology Publishing House: 422–427.
- Wen J., Wei M. and Nie H.** 1998. Taxonomical study of *Sterictiphora* Billberg (Hymenoptera: Argidae) of China with descriptions of nine new species. *Journal of Guangxi Agricultural University*, **17**(1): 61–70.
- Westerlund A.** 1892. Hymenopteroologisia havainnoita Laatokan pohjois-rannikolta. Title. *Acta Societatis pro Fauna et Flora Fennica*, **9**(2): 3–30.

- Williams P.H., Byvaltsev A., Sheffield C. and Rasmont P.** 2013. *Bombus cullumanus* – an extinct European bumblebee species? *Apidologie*, **44**: 121–132.
- Williams P.H.** 1991. The bumble bees of the Kashmir Himalaya (Hymenoptera: Apidae, Bombini). *Bulletin of the British Museum (Natural History) (Entomology)*, **60**: 1–204.
- Williams P.H.** 1998. An annotated checklist of bumble bees with an analysis of patterns of description (Hymenoptera: Apidae, Bombini). *Bulletin of The Natural History Museum (Entomology)*, **67**: 79–152.
- Williams P.H., An J. and Huang J.** 2011. The bumblebees of the subgenus Subterraneobombus: integrating evidence from morphology and DNA barcodes (Hymenoptera, Apidae, Bombus). *Zoological Journal of the Linnean Society*, **163**: 813–862.
- Williams P.H., Brown M.J.F., Carolan J.C., An J., Goulson D., Aytekin A.M., Best L.R., Byvaltsev A.M., Cederberg B., Dawson R., Huang J., Ito M., Monfared A., Raina R.H., Schmid-Hempel P., Sheffield C.S., Sima P. and Xie Z.** 2012. Unveiling cryptic species of the bumblebee subgenus *Bombus* s. str. world-wide with COI barcodes (Hymenoptera: Apidae). *Systematics and Biodiversity*, **10**: 21–56.
- Williams P.H., Byvaltsev A.M., Cederberg B., Berezin M.V., Ødegaard F., Rasmussen C., Richardson L.L., Huang J.-X., Sheffield C.S. and Williams S.T.** 2015. Genes suggest ancestral colour polymorphisms are shared across morphologically cryptic species in arctic bumblebees. *PLoS ONE*, **10**(12): e0144544.
- Williams P.H., Cameron S.A., Hines H.M., Cederberg B. and Rasmont P.** 2008. A simplified subgeneric classification of the bumblebees (genus *Bombus*). *Apidologie*, **39**: 46–74.
- Williams P.H., Huang J., Rasmont P. and An J.** 2016. Early-diverging bumblebees from across the roof of the world: the high-mountain subgenus Mendacibombus revised from species' gene coalescents and morphology (Hymenoptera, Apidae). *Zootaxa*, **4204**: 1–72.
- Williams P.H., Tang Y., Yao J. and Cameron S.** 2009. The bumblebees of Sichuan (Hymenoptera: Apidae, Bombini). *Systematics & Biodiversity*, **7**(2): 101–190.
- Wiśniowski B.** 2009. *Spider-hunting wasps (Hymenoptera: Pompilidae) of Poland. Diversity, identification, distribution.* Ojców National Park, Ojców, 432 p.
- Wnukowski W.W.** 1927. Verzeichnis der Hymenopteren des Bezirks Kamenj (südwestliches Sibirien, früheres Gouvernement Tomsk). *Konowia*, **6**(1): 31–34.
- Wnukowski W.W.** 1936. Beiträge zur Insekten-Fauna des Bezirkes von Tomsk (West-Sibirien). *Konowia*, **15**(1/2): 113–128.
- Woldstedt F.W.** 1875. Förteckning over några finska guld- och rofsteklar samt getingar. *Notiser ur Sällskapets Pro Fauna et Flora Fennica Förfhandlingar*, **14**: 344–349.
- Wolf H.** 1963(1960). Monografie der Westpalaeartischen Priocnemis-Arten (Hym. Pompilidae). *Bollettino del Museo Civico di Storia Naturale di Venezia*, **1963**: 21–181.
- Wolf H.** 1967. Wegwespen (Hym., Pompiloidea) Finnlands. *Acta Entomologica Fennica*, **23**: 1–46.
- Wolf H.** 1972. *Insecta Helvetica Fauna. 5. Hymenoptera Pompilidae.* Zurich, 176 p.
- Wolf H. and Sorg M.** 2007. *Die Wegwespen (Pompilidae) von Nordrhein-Westfalen.* UWM-Verlag, Krefeld, 133 p.
- Wu X.** 2010. A list of genera and species of Chinese stem sawflies with a proposal of a new tribe for an Australian stem sawflies (Hymenoptera: Cephidae: Australcephinae). *Journal of Northwest Forestry University*, **25**(6): 107–114.
- Wu X., Liu M. and Wei M.** 2016. Review of *Propodea* Malaise (Hymenoptera: Tenthredinidae) with descriptions of two new species, three new records from China and a new record from Nepal. *Entomotaxonomia*, **38**(4): 297–314.
- Wu Y.-R.** 2000. *Fauna Sinica - Insecta, Vol. 20: Melittidae and Apidae.* Sciense Press, Beijing, 412 p. [In Chinese].
- Xiao G.** 2000. A revisional list of the Chinese Pamphiliids (Hymenoptera: Pamphiliidae). *Forest Pests and Diseases*, **6**: 3–5. [In Chinese].
- Xu Z., Olmi M. and He J.** 2013. Dryinidae of the Oriental region (Hymenoptera: Chrysidoidea). *Zootaxa*, **3614**(1): 1–460.
- Yakovleva S.N.** 2015. Species diversity and ecology of the bees of the family Megachilidae (Hymenoptera, Apoidea) of in Kuznetsk-Salair region. Abstract of the Candidate of Biological Sciences thesis. Kemerovo State University, Kemerovo, 20 p. [In Russian].
- Yamane Sk.** 1975. Taxonomic notes on the subgenus *Bo-reovespula* Blüthgen (Hymenoptera, Vespidae) of Japan, with notes on specimens from Sakhalin. *Kontyû*, **43**: 343–355.
- Yamane Sk. and Terayama M.** 2016. Vespidae. In: Terayama M., Suda H. (Eds). *A Guide to the Aculeate Wasps of Japan.* Minamiyama, Tokai University Press: 290–331, pls. 62–78 + 602–608. [In Japanese].
- Yamane Sk., Wagner R.E. and Yamane Sô.** 1980. A tentative revision of the subgenus *Paravespula* of Eastern Asia (Hymenoptera: Vespidae). *Insecta Matsumurana (N.S.)*, **19**: 1–46.
- Yang J.** 1995. Revision of the Chinese pear stem girdlers with description of a new species (Hymenoptera: Cephidae). *Journal of Hubei University (Natural Science Edition)*, **17**(1): 7–13.
- Yasumatsu K.** 1939. Sur deux Sapygides nouveaux pour la faune de Chine et de Mongolie (Hymenoptera). *Notes d'Entomologie Chinoise, Shanghai*, **6**(7): 143–145.
- Yasumatsu K. and Sugihara Y.** 1938. Etude sur les Sapygides de Corée (Hymenoptera). *Mushi*, **11**: 75–77.
- Yegorenkova E.N., Lengesova N.A. and Yefremova Z.A.** 2010. Some biological data of *Fenusula ulmi* Sundevall, 1847 (Hymenoptera: Tenthredinidae) in Ulyanovsk region (Russia). *Caucasian Entomological Bulletin*, **6**(2): 199–202.

- Yıldırım E. and Gusenleitner J.** 2012. Contribution to the knowledge of the Vespidae (Hymenoptera, Aculeata) of Turkey, with a checklist of the Turkish species. *Turkish Journal of Zoology*, **36**(3): 361–374.
- Yusupov Z.M.** 2009. An ant species of the genus *Pyramica* (Hymenoptera, Formicidae) new for the fauna of Russia and the Northern Caucasus. *Entomological Review*, **89** (1): 123–124.
- Yusupov Z.M.** 2014. Two ant species (Hymenoptera, Formicidae) from the Northern Caucasus, new to the fauna of Russia. *Entomological Review*, **94** (2): 256–257.
- Zabroda V.V. and Sirenko A.N.** 2007. About spreading *Dolerus* and *Loderus* (Dolerinae, Tenthredinidae, Hymenoptera, Insecta) in different biotops and ekotops of Prekarpathian Region. *Naukovyi visnyk Volyn'skogo derzhavnogo universytetu imeni Lesi Ukrainky. II. Zoologiya*, **5**: 111–115. [In Russian].
- Zabroda V.V. and Sirenko A.N.** 2008. The sawflies of the genus *Macrophyia* (Tenthredinidae, Hymenoptera) in northern megaslope of the Ukrainian Carpathians. *Naukovyi visnyk Volyn'skogo derzhavnogo universytetu imeni Lesi Ukrainky. II. Zoologiya*, **3**: 133–135. [In Russian].
- Zaddach G.** 1859. Beschreibung neuer oder wenig bekannter Blattwespen aus dem Gebiet der preussischen Fauna. In: *Zur öffentlichen Prüfung der Schüler des Königlichen Friedrichs-Collegiums am Montag den 3. und am Dienstag den 4. October 1859*, Schultzsche Hofbuchdruckerei, Königsberg: 3–40.
- Zamprogno L.N. and Azevedo C.O.** 2014. Phylogeny and reclassification of *Pristocera* Klug (Hymenoptera: Bethylidae). *Insect Systematics & Evolution*, **45**: 1–49.
- Zerova M.D., D'jakonchuk L.A. and Ermolenko V.M.** 1988. *Gall-inducing insect of cultivated and wild plants of the European part of the USSR*. Naukova Dumka, Kiev, 159 p. [In Russian].
- Zhelochovtsev A.** 1941. On the sawflies of Armenia. *Archives of the Zoological Museum of the M. V. Lomonosov Moscow State University*, **6**: 225–238. [In Russian].
- Zhelochovtsev A.N.** 1928. Über paläarktische Dolerinae. *Zoologischer Anzeiger*, **79**(3–4): 105–112.
- Zhelochovtsev A.N.** 1935a. Neue oder wenig bekannte Tenthredinoidea aus Ost-Sibirien. *Archives of the Zoological Museum of the M. V. Lomonosov Moscow State University*, **1**: 150–154.
- Zhelochovtsev A.N.** 1935b. Notes sur les Dolerinae (Hym.) paléarctiques. *Archives of the Zoological Museum of the M. V. Lomonosov Moscow State University*, **2**: 79–84.
- Zhelochovtsev A.N.** 1939. Sawflies notes (Hym.). *Archives of the Zoological Museum of the M. V. Lomonosov Moscow State University*, **5**: 155–160.
- Zhelochovtsev A.N.** 1951. Review of the Palaearctic sawflies of the subfamily Selandriinae (Hym., Tenth.). *Archives of the Zoological Museum of the M. V. Lomonosov Moscow State University*, **7**: 123–153.
- Zhelochovtsev A.N.** 1952. Review of the sawflies of the subfamily Cladiinae (Hymenoptera, Tenthredinidae) of the fauna of the USSR. *Zoologicheskiy Zhurnal*, **31**(2): 257–269.
- Zhelochovtsev A.N.** 1968. New species of Symphyta (Hymenoptera) of the fauna of the USSR. *Archives of the Zoological Museum of the M. V. Lomonosov Moscow State University*, **11**: 47–56.
- Zhelochovtsev A.N.** 1976. The fauna of sawflies and wood-wasps (Hymenoptera Symphyta) of the southern Magadan area. *Archives of the Zoological Museum of the M. V. Lomonosov Moscow State University*, **15**: 74–96.
- Zhelochovtsev A.N.** 1988. Suborder Symphyta (Chalastogastra) – Sawflies and woodwasps. In: Tobias V.I., Zinovjev A.G. (Eds). *Keys to the Insects of the European Part of the USSR*. Vol. III, pt 6. Nauka, Leningrad: 21–234. [In Russian].
- Zhelochovtsev A.N. and Prochorova L.G.** 1976. Sawflies and woodwasps Moscow region. *Archives of the Zoological Museum of the M. V. Lomonosov Moscow State University*, **15**: 97–110. [In Russian].
- Zhelochovtsev A.N. and Zinovjev A.G.** 1981. New species of sawflies (Hymenoptera, Symphyta). *Archives of the Zoological Museum of the M. V. Lomonosov Moscow State University*, **19**: 15–17. [In Russian].
- Zhelochovtsev A.N. and Zinovjev A.G.** 1992. Suborder Symphyta. In: Chistyakov Yu.A. (Ed.). *Insects of Khingan Nature Reserve. Part 2*. Dalnauka, Vladivostok: 199–221. [In Russian].
- Zhelochovtsev A.N. and Zinovjev A.G.** 1994. A new sawfly species, *Nematus mandshuricus* sp. n. (Hymenoptera, Tenthredinidae) from Transbaikalia and Far East. *Bulletin of the Moscow Society of Naturalists, Section of Biology (N.S.)*, **99**(3), 37–39. [In Russian].
- Zhelochovtsev A.N. and Zinovjev A.G.** 1995. A list of the sawflies and horntails (Hymenoptera, Symphyta) of the fauna of Russia and Adjacent territories. I. *Entomologicheskoe Obozrenie*, **74**(2): 395–415. [In Russian].
- Zhelochovtsev A.N. and Zinovjev A.G.** 1996. A list of the sawflies and horntails (Hymenoptera, Symphyta) of the fauna of Russia and Adjacent territories. II. *Entomologicheskoe Obozrenie*, **75**(2): 357–379. [In Russian].
- Zhou X., Chen B. and Li T.-J.** 2012. Two new species and a key to species of the genus *Eumenes* Latreille (Hymenoptera: Vespidae: Eumeninae) from southwestern China. *Entomotaxonomia*, **34**(2): 467–474.
- Zimmermann K.** 1930. Zur Systematik der palaearktischen Polistes (Hym. Vespi.). *Mitteilungen aus dem Zoologischen Museum in Berlin*, **15**: 609–621.
- Zinovjev A.G.** 1978. Description of a new subgenus *Paranematus* subgen. n. for the group of species related to *Nematus wahlbergi* Thomson (Hymenoptera, Tenthredinidae) with a review of species from the European part of the USSR. *Entomologicheskoe Obozrenie*, **57**(3), 625–635.

- Zinovjev A.G. 1981.** On the study of gall mining Hymenoptera of the genus *Pontania* Costa (Tenthredinidae). In: Ler P.A. (Ed.). *Hymenopterous Insects of Far East*. DVNC AN SSSR, Vladivostok: 18–24. [In Russian].
- Zinovjev A.G. 1985.** On the taxonomy of the sawfly genus *Pontania* O. Costa (Hymenoptera, Tenthredinidae). Subgenus *Eupontania* subg. n. *Proceedings of the Zoological Institute of the Academy of Sciences of the USSR*, **132**: 3–16. [In Russian].
- Zinovjev A.G. 1986.** Sawflies of the genus *Platycampus* Schiødte (Hymenoptera, Tenthredinidae) from the Far East of the USSR. In: Ler P.A. (Ed.). *Hymenopterous Insects of Eastern Siberia and Far East*. BPI DVNC AN SSSR Vladivostok: 3–14. [In Russian].
- Zinovjev A.G. 1993a.** Pristicampini – a new tribe for a new genus of sawflies from Northern Europe and Siberia (Hymenoptera: Tenthredinidae). *Zoosystematica Rossica*, **1**: 78–85.
- Zinovjev A.G. 1993b.** Subgenera and Palaearctic species groups of the genus *Pontania*, with notes on the taxonomy of the *viminalis*-group (Hymenoptera: Tenthredinidae). *Zoosystematica Rossica*, **2**: 145–154.
- Zinovjev A.G. 1993c.** Two new species of Nematinae (Hymenoptera Tenthredinidae) from the Eastern Palaearctic. *Russian Entomological Journal*, **2**(2): 31–35.
- Zinovjev A.G. 1993d.** A new species of *Pikonema* Ross (Hymenoptera, Tenthredinidae) from Japan. *Bulletin of the National Science Museum, Series A, Zoology*, **19**(1): 21–25.
- Zinovjev A.G. 1994.** Revision of sawflies of the genus *Aglao-stigma*, subgenus *Neurosiobla* (Hymenoptera: Tenthredinidae). *Zoosystematica Rossica*, **3**(1): 115–127.
- Zinovjev A.G. 2000.** Supplements and corrections to the list of sawflies (Hymenoptera, Symphyta) of the fauna of Russia and adjacent territories. *Entomologicheskoe Obozrenie*, **79**(2): 450–457. [In Russian].
- Zinovjev A.G. 2010.** Palaearctic sawflies of the genus *Pontania* Costa (Hymenoptera: Tenthredinidae) and their host specificity. In: Csoka G., Mattson W.J., Stone G.N., Price P.W. (Eds). *The biology of Gall-inducing Arthropods. U.S. Department of Agriculture, Forest Service, General Technical Report NC*: 204–225.
- Zinovjev A.G. and Schmidt S. 1994.** *Amauronematus compactus* Bogacheva and *A. harpicola* Bogacheva (Hymenoptera, Tenthredinidae) from the Polar Urals. *Entomologica Fennica*, **5**: 135–138.
- Zinovjev A.G. and Vikberg V. 1999.** The sawflies of the *Pontania crassispina*-group with a key to the genera of the subtribe Euurina (Hymenoptera: Tenthredinidae: Nematinae). *Entomologica Scandinavica*, **30**: 281–298.
- Zombori L. 1971.** Symphyta (Hymenoptera) from Mongolia with the description of two new species I. *Ergebnisse der Zoologischen Forschungen von Dr. Z. Kaszab in der Mongolei* (No. 229). *Acta Zoologica Academiae Scientiarum Hungaricae*, **17**(1–2): 233–241.
- Zombori L. 1973.** Symphyta (Hymenoptera) from Mongolia III. *Acta Zoologica Academiae Scientiarum Hungaricae*, **19**(3–4), 445–460.
- Zombori L. 1974.** Sawflies from Korea (Hymenoptera: Symphyta). *Acta Zoologica Academiae Scientiarum Hungaricae*, **20**(3–4), 453–461.
- Zombori L. 1978.** New sawfly species from Korea (Hymenoptera: Symphyta). *Acta Zoologica Academiae Scientiarum Hungaricae*, **24**(1–2): 253–268.
- Zombori L. 2013.** *Pseudoryssus henschii* (Mocsáry, 1910) new in the Hungarian fauna (Hymenoptera: Orussidae). *Folia Entomologica Hungarica*, **74**: 185–187.
- Zvantsov A.B. 1988.** Fauna and geographical distribution of the chrysidid wasps (Hymenoptera, Chrysidae) in the Moscow region. In: Rasnitsyn A.P. (Ed.). *Insects of the Moscow region. Problems of inventory and protection*, Nauka, Moscow: 80–97. [In Russian].

INDEX OF LATIN NAMES OF HYMENOPTERA

(**bold** is given for family-group names, *italic* for: synonyms, nomina nuda, wrong names, and generic names, when current combination is different from original combination).

- 14-notatus**, Oxybelus 243
2-cincta, Tenthredo 97
3-cincta, Tenthredo 97
4-cincta, Tenthredo 97
4-fasciata, Tenthredo 30
4-maculata, Vespa 196
4-notatus, Oxybelus 243
4punctatus, Crabro 230
5 maculatus, Blepharipus 227
5-macutatus, Crabro 230
6-maculatus, Crabro 232
8 maculata, Crabro 231
8-notatus, Solenius 232
- aaliensis, Tenthredo, *Pachyprotasis* 91
abbreviata, Pristiphora, *Nematus* 70
abdominalis, Ammobatooides,
 Phileremus 318
abdominalis, Cimbex femorata 29
abdominalis, Dolerus 40
abdominalis, Hylaeus 282
abdominalis, Lipotriches 291
abdominalis, Monostegia, *Tenthredo* 43
abdominalis, Tenthredo,
 Tenthredopsis 97
abdominalis, Tipha 218
abdominis, Taxonus 43
abeillei, Chrysida 142
abeillei, Ponera 209
aberrans, Andrena 274
aberrans, Anoplus 172
aberrans, Anoplus 172
aberrans, Crabro 234
aberrans, Nomada 315
aberrans, Stelis 318
aberrans, Stizus 224
aberrans, Thyreus, *Crocisa* 326
Abia 28
abieticola, Gilpinia, *Lophyrus* 33
abietina, Lyda 102
abietina, Pristiphora, *Tenthredo* 70
abietina, Tenthredo 70
abietinus, Nematus 70
abietinus, Orussus, *Sphex* 117
abietis, Cephalcia, *Tenthredo* 102
abietis, Lophyra 33
abietum, Tenthredo 70
Abiinae 28
abjecta, Tenthredo 93
ablatus, Tachysphex 239
Ablepharipus 227
abnormis, Arachnospila, *Pompilus* 168
abnormis, Euura, *Nematus* 55
abnormis, Odontocrabro 235
abnormis, Polemistus, *Passaloecus* 249
abnormis, Sierolomorpha, *Tipha* 151
abnormis, Strongylogaster 39
- aborigenica*, Myrmica 205
aborigensis, Euura, *Eupontania* 55
aborigensis, Hylaeus 259
abortiva, Euura 56
Abrupta subg. 259
absens, Fenusa 48
Absentia 90
absimilis, Pteronidea 57
absoluta, Andrena marginata var. 268
abyssinicus, Sphecodes hispanicus 289
Acanthocnema 101
Acanthocrabro 227
Acantholepis 202
Acantholyda 101
Acantholyda subg. 101
Acanthonomada 311
acanthophorus, Crossocerus, *Crabro* 231
Acanthoptenos 22
Acarepiona 178
accola, Cerceris 255
accuratus, Loderus 79
Aceratosmia 297
acerbum, Lasioglossum, *Halictus*
 obscuratum 285
acerbus, Loderus 77
aceris, Heterarthrus, *Phyllotoma* 49
aceris, *Phyllotoma* 49
aceris, Pteronidea 64
aceroides, Heterarthrus, *Phyllotoma* 49
Aceropales 160
acerosus, Nematus 64
acerrima, Tenthredo 96
acervolum, Lasioglossum 281
acervorum, Apis auct. 323
acervorum, Formica 204
acervorum, Leptothorax, *Formica* 204
Achaetopriion 77
Achaetopriion subg. 77
achaues, Leptochilus limbiferus 185
Achrysis 141
Aciandrena subg. 263
aciculata, Andrena 263
aciculatus, Crabro 233
aciculatus, Halictus brevicornis 282
aciculatus, Messor, *Aphaenogaster* 204
Aclastocera 145
Acolpus 256
Acrepyris 123
acriculus, Loderus 78
acris, Apis 267
acrobates, Tachytes 238
Acrodontochelys 120
Actenopoda 166
Aculeata 118
aculeata, Stelis 302
acuminata, Apis 305
acuminata, Coelioxys 304
- acuminata*, Euura 64
acuminatus, Nematitus, *Nematus* 68
acuta, Ceratina 310
acuta, Coelioxys 305
acuta, Euura, *Amauronematus* 55
acutangula, Tetralonia 322
acuticornis, Hoplitis, *Osmia* 296
acuticornis, Oxybelus 243
acuticrista, Lasioglossum 281
acutidens, Pristiphora 70
acutifoliae, Euura, *Pontania* 55
acutinodis, Leptothorax tuberum 208
acutiserra, Euura, *Pontania* 56
Acutopterocheilus subg. 187
adamantis, Metallus 50
Adamas 40
adelpha, Ammophila 212
adelpha, Cerceris 252
adelungi, Pristiphora 72
ademonia, Myrmica 205
adi, Camponotus fedtschenkoi var. 199
Adipogon 164
Adirostes 161
Adirus 110
adjikenticus, Halictus 277
Adonta subg. 161
adulterina, Dolichovespula, *Vespa* 194
adumbrata, Tenthredo 45
adunca, Hoplitis, *Apis* 297
aduncoides, Osmia 297
aduncta, Apis 296
Adungia 89
adusta, Amasis obscura 32
adusta, Lyda 102
adusta, Tenthredo 91
adusta, Tetralonia 322
adustus, Microdynerus timidus 186
adventicium, Hedychrillum 128
adventor, Agrobombus 331
adversa, Dasylabris 153
adversa, Dasylabris italica var. 153
adzhharicum, Trypoxylon 245
aeger, Euura, *Amauronematus* 56
aeginus, Halictus 284
aegra, Tenthredo 96
aegyptiaca, Coelioxys 305
aegyptiaca, Vespa 194
aegyptiaca, Vespa orientalis var. 194
aegyptiacus, Halictus quadricinctus 277
aegyptica, Megachile 305
aegypticola, Halictus 277
aegypticellum, Lasioglossum,
 Halictus 280
aegyptius, Lindenius, *Crabro* 234
aegyptorum, Anthophora atricilla 324
aello, Chrysis 133
aemula, Andrena 269

aemula, Dolerus tremulus 77
aemulans, Hoplammophila,
 Ammophila 213
aemulus, Amauronematus 65
aemulus, Hylaeus 261
Aenandrena subg. 263
aenea, Abia, *Cimbex* 28
aenea, Apis 298
aenea, Ceratina 310
aenea, Chrysis 131
aeneidorsum, Halictus 285
aeneipennis, Vespa 179
aeneiventris, Andrena 263
aeneiventris, Anthophora 324
aeneopaca, Chrysis longula 136
aeneovarius, Pachycephus 110
aeneoviridescens, Arge dimidiata 23
aenescens, Cataglyphis, *Formica* 199
aenescens, Crabro 234
aenescens, Hylotoma 23
aenescens, Nomioides 279
aenescens, Rhogogastera 93
aenescens, Trichiosoma 30
aeneus, Cladius 54
aeneus, Dolerus 79
aeneus, Omalus, *Chrysis* 131
Aenoabia 28
aequalis, Cladius 54
aequisculptus, Stenodynerus,
 Odysnerus 189
aerarius, Halictus 278
aeratum, Lasioglossum, *Melitta* 282
aereum, Hedychrum simile 130
aericeps, Dolerus 77
aerosus, Cleptes aerosus 126
aeruginosus, Anoplus, *Pompilus* 172
aestiva, Andrena 267
aestiva, Chrysis 133
aestivalis, Andrena nitida var. 269
aestivalis, Anthophora, *Apis* 325
aestivalis, Nomada fabriciana var. 312
aestivus, Nematus 64
aethiops, Andrena 270
aethiops, Camponotus, *Formica* 199
aethiops, Endelomyia, *Tenthredo* 46
Aethomegachile subg. 305
afasciata, Smicromyrme orientalis f. 156
afer, Cleptes 142
afer, Coelioxys 304
afer, Sphecodes 216
affine, Lasioglossum, *Halictus* 282
affinis, Anthophora 324
affinis, Cephus 110
affinis, Crabro 227
affinis, Crossocerus 227, 229
affinis, Halidamia, *Hylotoma* 48
affinis, Harpactus, *Gorytes* 223
affinis, Hartigia 111
affinis, Hylaeus 260, 284
affinis, Hylotoma 48
affinis, Lasius 201

affinis, Lasius auct. 201
affinis, Melitta 268
affinis, Odynerus 179
affinis, Osmia 298
affinis, Oxybelus 242
affinis, Pachynematus 58
affinis, Palarus 243
affinis, Podalonia, *Ammophila* 213
affinis, Pompilus 161, 173
affinis, Pristiphora, *Lygaeonematus* 70
affinis, Psen 251
affinis, Sphecodes 288
affinis, Temnothorax, *Leptothorax* 208
affinis, Thyreus, *Crocisa* 326
affinissimus, Eumenes 181
affinis, Megachile 305
afghaniensis, Sphecodes 215
afra, Anthophora albigena 323
africana, Empria 41
africanus, Holepyris 123
africanus, Metrionotus 124
africanus, Sphecodes 287
africanus, Tachysphex julliani 238
Afrisobrachium 123
afromontana, Pseudepipona
 herrichii 188
afzeliella, Melitta 274
agelastum, Lasioglossum 280
Agenia 164
agenia, Priocnemis, *Salius* 163
Ageniellina 165
Ageniellini 165
Agenioideus 167
Agenioideus subg. 167
Agenioxenus 160
Agenocimber 29
Agenoideus 167
agilis, Apis 305
agilis, Astata 218
agilis, Nematus 70
agilis, Phylloecus, *Cephus* 110
agilis, Priocnemis, *Pompilus* 163
agilis, Tenthredo 38, 39
agilis, Tiphiia 150
aglaia, Acantholyda aglaia 101
Aglaopapis 303
Aglaostigma 83
Aglaostigma subg. 83
agnani, Macrophyta albicincta 84
agnolii, Elampus 127
agnolii, Pseudomalus 132
Agnosicrabro 225
agona, Coelioxys rufescens var. 304
Agonatopoides 120
Agonatopus 120
Agraptus 224
agraria, Andrena bicolor ssp. 267
agrestis, Crabro 232
agrestis, Tenthredo 104
agrimoniae, Fenella 48
agrimoniae, Macrocephus 110
agrorum, Apis 331
agrorum, Taxonus, *Tenthredo* 43
agyo, Crossocerus dimidiatus 230
aheneum, Hedychridium,
 Hedychrum 128
ahngeri, Athalia 44
ahngeri, Myrmica scabrinodis var. 206
aihinoensis, Heterarthrus 49
aino, Andrena 263
aino, Anoployda viriditibialis 105
aino, Crabro 227
aino, Gorytes 222
aino, Sphecodes 289
Ainocrabro 227
ainu, Arachnospila 169
aiunensis, Gastrosericus 236
ajnu, Paramasaakia 51
akazui, Xiphydria 115
akitanus, Sphecodes 289
aksuense, Lasioglossum 280
alabastipes, Eriocampa 39
alagesianus, Bombus 329
alaica, Anthophora aestivalis 325
alaicus, Leptothorax tuberum 208
alamannicus, Evagetes, *Pompilus* 170
alashanica, Cephalcia, *Cephaleia* 102
alaskana, Tenthredo, *Allantus* 96
Alastor 178
Alastor subg. 178
Alastorellus subg. 178
alastoroides, Microdynerus 186
Alastorynerus subg. 186
alata, Lestica, Crabro 234
alatulus, Crabro 232
alatus, Coelioxys 304
albamacula, Tenthredo 85
albanica, Chrysis 133
albanica, Chrysis succinata var. 133
albarius, Halictus 277
albata, Tenthredo 91
albata, Tenthredopsis 100
albata, Tenthredopsis austriaca 99
albellus, Polistes 192
albellus, Polistes foederatus 192
albeola, Mutilia 153
albertii, Dolerus 79
albescens, Anthophora
 quadrifasciata var. 322
albibarbis, Andrena 273
albicans, Apis auct. 275
albicarpus, Nematus 63
albicaudata, Andrena 265
albiceps, Stromboceros delicatulus 82
albicilla, Andrena flavobila ssp. 271
albicilla, Megachile 307
albicincta, Ceropales, *Evania*
 albicincta 160
albicincta, Macrophyta, *Tenthredo* 84
albicincta, Tarpa 107
albicinctus, Allantus, *Dolerus* 38
albiclypeata, Deuteragenia,
 Dipogon 164
albiclypeata, Tetralonia
 salicariae var. 322
albicollis, Andrena ephippium var. 269

albicolon, *Tenthredo coryli* 95
albicolor, *Cerceris* 252
albicornis, *Siobla ruficornis* 89
albicornis, *Tenthredo* 91, 95
albicurvis, *Hylotoma* 244
albicus, *Melitta* 268
albida, *Anthophora albigena* var. 323
albida, *Megachile analis* var. 307
albida, *Megachile lefebvrei* var. 305
albida, *Periclista*, *Tenthredo* 51
albidemaculata, *Nomada* 311
albidicollis, *Anteon* 118
albidopleuralis, *Agrobombus schrencki* 332
albidulum, *Anthidium* 301
albidulus, *Gorytes*, *Hoplitus* 222
albidus, *Euodynerus disconotatus* 183
albidus, *Neogonatopus* auct. 121
albidus, *Polistes atrimandibularis* 192
albifrons, *Anthophora* 324
albifrons, *Apis* 267
albifrons, *Auplopus*, *Pompilus* 165
albifrons, *Dolerus*, *Dosytheus* gilvipes 79
albifrons, *Lyda* 103
albifrons, *Melecta*, *Apis* 325
albigena, *Amegilla*, *Anthophora* 322
albilabris, *Apis* 272
albilabris, *Crabro* 234
albilabris, *Euspongus* 222
albilabris, *Hylaeus* 310
albilabris, *Lindenius*, *Crabro* 234
albilabris, *Nematus* 70
albilabris, *Pristiphora*, *Nematus* 70
albilabris, *Sphecodes*, *Nomada* 287
albilabris, *Tenthredo* 85
albimacula, *Macrophyia* 85
albimacula, *Tenthredella balteata* 94
albimana, *Tenthredo* 86
albinasata, *Prosopis* 262
albipalpis, *Tenthredo* 84
albipennis, *Elampus*, *Ellampus* 127
albipennis, *Euura*, *Nematus* 56
albipennis, *Periclista*, *Selandria* 51
albipes, *Ametastegia*, *Taxonus* 39
albipes, *Andrena* 267
albipes, *Aphelinus* 118
albipes, *Armitarsus* 84
albipes, *Ceraphron* 118
albipes, *Cladius* 54
albipes, *Eriocampa* 42
albipes, *Lasioglossum*, *Apis albipes* 282
albipes, *Metallus*, *Phaenus* 50
albipes, *Podalirius acervorum* var. 323
albipes, *Tenthredo* 50
albipes, *Tenthredo* 50, 54, 76, 84
albipleuris, *Tenthredopsis* 100
albipuncta, *Macrophyia*, *Tenthredo* 84
albiscopa, *Osmia fulviventris* 298
albisecta, *Megachile*, *Anthophora* 305
albisectus, *Sphecius* 215
albisternus, *Emphytus* 43
albitarsis, *Hylaeus* 283

albitarsis, *Macrophyia* 87
albitarsis, *Macrophyia sanguinolenta* 86
albitibia, *Pristiphora*, *Nematus* 70
albiventre, *Anthidium* 300
albiventris, *Tenthredo*, *Allantus* albiventris 96
alboannulata, *Tenthredo*, *Tenthredella* 93
albociliata, *Crocisa ramosa* var. 327
albociliatus, *Hortobombus consobrinus* 328
albocincta, *Andrena* 268
albocincta, *Nomia* 291
albocinctus, *Bombus lucorum* 327
albocineta, *Megachile* 306
alboclypea, *Filixungulua* 42
albofasciata, *Andrena* 274
albofasciata, *Cerceris*, *Vespa* 252
albofasciata, *Eucera* 320
albofasciata, *Melitta* 294
albofasciatum, *Anthidium punctatum* var. 301
albofimbriata, *Andrena* 274
alboguttata, *Nomada* 311
albohirta, *Megachile lagopoda* 307
albolapidaria, *Macrophyia* 85
albomaculata, *Anthophora* 324
albomaculata, *Megachile lefebvrei* var. 305
albomaculata, *Poecilotiphia* 150
albomaculatus, *Colletes*, *Halictus* 257
albomaculatus, *Nomada* 319
albomaculatus, *Tenthredopsis* 99
albonotata, *Cerceris* 254
albonotata, *Tenthredopsis stigma* 100
albonotatum, *Episyron*, *Pompilus* 166
albopicta, *Euura*, *Pontania* 56
albopicta, *Tenthredo* 93
albopictus, *Odynerus* 187
albopictus, *Pamphilus*, *Lyda* 104
albopilosum, *Trichosoma* 30
albopilosus, *Psen* 250
alboplagiata, *Tenthredella cunyi* 94
alboplagiata, *Tenthredopsis tessellata* 100
albopunctata, *Andrena*, *Apis* 268
albopunctata, *Eucera* 320
albopunctata, *Tenthredo* 98
albopunctatus, *Oryssus* 117
alborufa, *Eucera*, *Tetralonia* 321
alborufa, *Osmia* 298
alboscutellata, *Crocisa histrionica* var. 327
albotricinctus, *Odynerus* 179
albovaria, *Ametastegia*, *Emphytina* 39
albovaria, *Melecta* 326
albrechti, *Dryinus*, *Richardsdryinus* 120
albufeirae, *Miscophus* 240
alces, *Tenthredo* 54
alchymista, *Tremex* 113
Alcidamea subg. 296
aldrichi, *Odynerus* 188
alecto, *Melecta* 326
alector, *Empria* 40
alemannia, *Andrena pectoralis* var. 269
Alepidaspis 242
alexandrina, *Andrena flavipes* ssp. 275
alexii, *Hedychrum* 130
alexoides, *Halictus* 278
Alfieria 180
Alfenella 322
alfkenella, *Andrena* 270
alfkeni, *Andrena rosae* var. 267
alfkeni, *Nomada obscura* var. 315
alfkeni, *Schizocera* 26
alga, *Andrena minutula* ssp. 270
algerica, *Amasis crassicornis* 32
algerica, *Fertonella* 297
algericoellus, *Halictus* 285
algeriensis, *Andrena nitida* var. 269
algeriensis, *Ceratina callosa* 310
algeriensis, *Coelioxys aurolimbata* var. 305
algeriensis, *Odynerus* 184
algeriensis, *Sphecodes reticulatus* 287
algira, *Andrena lucens* var. 271
algira, *Ferreola* 166
algira, *Tachysphex* 239
algitrus, *Sphecodes pellucidus* 289
Alicrabro 227
alienatus, *Hylaeus* 259
alienatus, *Nematus* 60
alienonigra, *Lasius niger* var. 202
alienus, *Lasius*, *Formica* 202
alienus, *Pompilus* 225
alinae, *Temnothorax*, *Leptothorax* 208
alinae, *Tracheliodes* 236
alinense, *Lasioglossum*, *Halictus* 280
alini, *Colletes* 257
alius, *Epyris* 123
alias, *Odynerus* 179
aljkeni, *Megachile lagopoda* 307
aljkeni, *Nomada* 312, 315
Allanteon 118
allantiformis, *Leptocimbex*, *Clavellaria* 30
Allantinae 37, 43
Allantus 37
Allantus 89
Allantus subg. 37
allegrus, *Ancistrocerus antilope* 178
Allepyris 123
alliaria, *Andrena* 273
Alliognathus 235
allobrogus, *Symporus*, *Odynerus* 190
Allocimbex 29
Allocelioxys subg. 304
allodalam, *Lasioglossum* 282
Allodynerus 178
Allogonatopus 120
Allomutilla 153
Alloridestus 167
allosa, *Andrena* 267
Alloscenia 22
Allosmia subg. 297

Allovespula 195
Alloxylocopa subg. 309
almasyana, Haba, *Holopyga* 128
almasyana, Holopyga 128
almasyanus, Allantus 97
alneti, Tenthredo 69, 99
alni, Psen 105
alni, Tenthredo 67, 69
alniastri, Nematus, *Tenthredo* 69
alnicola, Nematus 70
alnicola, Pamphilus 104
alnicola, Tenthredo 70
alnina, Tenthredo 68
alnivaga, Tenthredo 29
alnivora, Xiphidria 115
alnivorus, Heterarthrus 49
alnivorus, Nematus 70
alnivorus, Pamphilus 104
Alocandreninae 263
Alocochrysis 141
alpestris, Halictus 280
alpestris, Leptochilus, *Odynerus* 185
alpestris, Myrmica lobicornis 206
alpestris, Pristiphora, *Pachynematus* 70
alpestris, Sphecodes rufescens 288
Alpha 181
Alphadryinus 120
Alphastromboceros 75
alpicola, Andrena gallica ssp. 269
alpicola, Euura, *Amauronematus* 56
alpicola, Halictus albipes 282
alpicola, Lygaeonematus 72
alpicola, Megachile 306
alpicola, Odynerus parvulus var. 180
alpigena, Allantus arcuatus 96
Alpigenixyla 20
Alpigenobombus subg. 327
alpina, Andrena 264, 268
alpina, Athalia 44
alpina, Cephalcia, *Lyda* 102
alpina, Empria 40
alpina, Hoplocampa, *Tenthredo* 67
alpina, Nomada robertjeotiana var. 315
alpina, Perineura 83
alpina, Podalonia, *Ammophila* 213
alpina, Synairema 82
alpina, Tenthredo velox 96
Alpinematus 55
alpinobalticus, Anoplus 172
alpinum, Anthidium 301
alpinum, Anthidium 302
alpinum, Pseudoanthidium,
 Anthidium 302
alpinus, Bombus, *Apis* 327
alpinus, Colletes 258
alpinus, Crabro 226
alpinus, Epeolus 317
alpinus, Nematus 71
alpinus, Odynerus 186
alpinus, Odynerus spinipes var. 186
alpinus, Temnothorax, *Leptocephalus* 208
alpium, Polistes biglumis 192
alpotanini, Panurginus 276

alsia, Messa 48
alsius, *Amauronematus* 64
altaica, Chrysis 143
altaica, Andrena 273
altaica, Anthophora 325
altaica, Arge 22
altaica, Arge 25
altaica, Camponotus
 herculeanus var. 198
altaica, Tenthredo providus 98
altaica, Vespa crabro var. 194
altaica, Xylocopa 310
altaica, Xylocopa nitidiventris 310
altaicum, Camptopeum 276
altaicus, Bombus 329
altaicus, Crabro 226
altaicus, Halictus 278
altaicus, Hylaeus 259
altaicus, Lindenius 235
altaicus, Pamphilus 104
altaicus, Passaloecus 248
altaicus, Thyreus, *Crocisa* 326
altaicensis, Podalonia, *Ammophila* 213
altercator, Apis 293
altercator, Dasypoda auct. 293
alternans, Eucera, *Macrocerca* 321
alternans, Tenthredo 82
alternanus, Odynerus 191
alternata, Apis 313
alternatus, Pompilus 162
alternatus, Sphecodes 287
alternipes, Claremontia, *Tenthredo* 46
alticincta, Tetraloniella, *Macrocerca* 321
alticinctus, Loderus 77
alticola, Anthophora vulpina 323
alticola, Chrysura, *Chrysis* 139
alticola, Ciliissa 294
alticola, Pachynematus 57, 71
alticola, Psen 251
alticolus, Panurginus 276
alutacea, Andrena 273
alutus, Discolia 148
alvarabnormis, Arachnospila,
 Pompilus 168
Alyson 219
Alysson 219
Alysson 219
Alyssontini 219
amabilis, Scolia 148
amaculatus, Platycampus 70
amadanensis, Eustenancistrocerus,
 Odynerus 184
amaguella, Megachile melanopyga 307
amaguensis, Andrena 271
amaguensis, Osmia 297
amakusensis, Sphecodes 288
amala, Senoclidea 53
amanda, Melecta 326
amaniensis, Andrena furcata var. 267
amanusense, Anthidium 301
amasia, Andrena 264
Amasis 31
amaura, Tenthredo 49
Amauronematus 55
ambidens, Tachytes 240
ambigua, Andrena 272
ambigua, Chrysis 133
ambigua, Coelioxys 305
ambigua, Sierola 151
ambigua, Smicromyrme 155
ambigua, Tenthredo 73, 99
ambiguum, Tapinoma auct. 198
ambiguum, Tapinoma erraticum 198
ambiguum, Trypoxylon 244
ambiguus, Crabro 227
ambiguus, Elampus 128
ambiguus, Epeoloides 319
ambiguus, Epeoloides 319
ambiguus, Gorytes 222
ambiguus, Nematus 62
ambiguus, Oxybelus 243
ambiguus, Psen 252
Amblyellus 168
Amblys 297
amblystoma, Nomada 316
ameidei, Eumenes 184
Amegilla 322
Amegilla subg. 322
amelanchieris, *Amauronematus* 72
amentivora, Pontania 56
amentorum, Euura, *Nematus* 56
americana, Dineura 67
americana, Synairema 87
amerinae, Euura, *Cynips* 56
amerinae, Pseudoclavellaria,
 Tenthredo 30
amerinae, Tenthredo 30
Ametastegia 39
Ametastegia subg. 39
amethystina, Hylotoma 23
Ammatodus 220
Ammobates 318
Ammobates subg. 318
Ammobatini 318
Ammobatoides 318
Ammobatoidini 318
Ammophila 212
Ammophilinae 212
ammophiloides, Trypoxylon 245
Ammoplanellus 246
Ammoplanus 246
Ammosphecidium 241
Ammosphecidium subg. 168
ameris, Chrysis 133
amoena, Chrysis 140
amoena, Corynis, *Cimbex* 31
amoena, Pentachrysis, *Chrysis* 140
amoena, Tenthredo 93
amoena, Tenthredo 97
amoena, Tetracis 321
Amoeniformis 31
amoenita, Apis 300
amoenula, Holopyga 131
amoenula, Holopyga amoenula 131
amoenus, Pompilus 174
amoorense, Aglaostigma, *Tenthredo* 83

amphibola, Pristiphora, *Nematus* 70
 amphybolia, Euxiphydria 115
 ampla, Euura, *Amauronematus* 56
amplicapitata, Tenthredo 95
amplitarsis, Eucera 320
 Ampulex 211
Ampulicidae 211
Ampulicinae 211
 Ampulicomorpha 122
amurensis, Lasioglossum, *Halictus* 282
amurensis, Agenioideus,
Psammochares 167
amurensis, Andrena 271
amurensis, Anthidium 300
amurensis, Arachnospila, *Pompilus* 170
amurensis, Bombus 330
amurensis, Calameuta 108
amurensis, Chrysis 133
amurensis, Cimbex 29
amurensis, Crossocerus, *Crabro* 231
amurensis, Euura, *Pontania* 56
amurensis, Hoplomerus spinipes 187
amurensis, Macropis dimidiata 293
amurensis, Macropis fulvipes 293
amurensis, Nomada 311
amurensis, Nysson 220
amurensis, Platycampus 70
amurensis, Priocnemis 163
amurensis, Tenthredo 83, 93
amurensis, Xiphydriola 116
amurica, Andrena 271
amurica, Tenthredo 93
Anachoreta 33
anachoreta, Bombus, *Agrobombus* 331
anachoreta, Lophyrus 33
Anafenusia 45
analis, Allantus 93
analis, *Amauronematus* 65
analis, Andrena 267
analis, Astatus 108
analis, Athalia lineolata 44
analis, Chrysis analis 133
analis, Crabro 230
analis, Dolerus 80
analis, Halictus 282
analis, Lithurgus 295
analis, Megachile 307
analis, Oxybelus 242
analis, Periclista pubescens 51
analis, Philanthus 262
analis, Pontopristia 59
analis, Selandria 76
analis, Tenthredo 99
analis, Vespa 193
anastasiae, Bothriomyrmex 197
anathema, Larra, *Sphex* 236
anathema, Sphex 236
anatoliae, Odontodynerus
 ephippium 188
anatolica, Andrena 271
anatolica, Clavellaria amerinae 30
anatolica, Eoferreola 174
anatolica, Halictus euboeensis 283

anatolica, Pseudapis, *Nomia* 292
anatolica, Systropha planidens 291
anatolicus, Colletes tuberculatus 259
anatolicus, Eumenes mediterraneus 182
anatolicus, Heterarthrus vagans 49
anatolicus, Leptochilus limbiferus 185
anatolicus, Pasites oraniensis 318
anatolicus, Sphecodes 214
anatolii, Dolerus 79
anceps, Arachnospila, *Pompilus* 168
anceps, Colletes 257
anceps, Hoplitis 223
anceps, Hylotoma 25
anceps, Nematus 64
ancilla, Athalia 44
ancilla, Athalia ancilla 44
ancilla, Pontania 55
Ancistrocerus 178
Ancistrostromma 236
ancisus, Loderus 77
Ancyla 319
Ancylini 319
andalusica, Nomada 311
anderschi, Nematus 73
andinus, Halictus 285
andreae, Cephus 109
andreae, Eucera 320
andrej, Monoctenus 35
andrej, Tenthredo 92
Andrena 263
Andrena subg. 263
Andrenidae 263
Andreninae 263
andrenoides, Osmia 298
Andreas subg. 149
anellum, Lasioglossum, *Halictus* 282
anemones, Endophytus, *Pelmatopus* 55
anemones, Pelmatopus 55
Anemphytus 37
Anergates 203
Aneugmenus 75
angarensis, Andrena 274
angarensis, Bombus maculidorsis 331
angarensis, Megachile 307
angarensis, Miscophus 241
angarensis, Phiarus 318
angarensis, Sphecodes 288
angularis, Acantholyda 101
angularis, Lasioglossum, *Halictus* 282
angelica, Cerceris 252
anglica, Andrena 267
anglica, Hylotoma 24
anglica, Vespa 195
anglica, Vespa crabro var. 194
anglicus, Nematus 64
angolensis, Chrysis 133
angularis, Blennocampa subcana 46
angularis, Pemphredon 249
angularis, Tachysphex 239
angulata, Osmia 299
angulata, Pristiphora 70
angulata, Prosopis 261
angulata, Vespa 179
angulicollis, Psen 250
angulinodis, Myrmica 205
angulinodis, Myrmica scabrinodis 205
angulosa, Melitta 264
angusta, Tenthredo 80
angustata, Caliroa 45
angustata, Ephutomma,
Smicromyrme 155
angustata, Tenthredo 38
angustata, Thrinax, *Hemitaxonus* 83
angustatus, Hylaeus, *Prosopis* 259
angustatus, Pachynematus 65
angustatus, Symmorphus, *Odynerus* 191
angustatus, Tachysphex 237
angustifrons, Chrysis angustifrons 134
angustifrons, Philanthus 255
angustipes, Andrena 268
angustirostris, Cerceris 252
angustula, Anthophora 299
angustula, Chrysis angustula 134
angustula, Tenthredo 39
angustus, Nematus 56
angustus, Passaloecus 247
anhinga, Andrena florea ssp. 273
Anisoneura 40
anivskiensis, Pristiphora 71
ankarae, Andrena 270
Annosmia 296
annulare, Anthidium 301
annularis, Melitta 260
annularis, Strongylogaster 82
annularis, Tenthredo 91, 95
annulata, Euura, *Nematus* 56
annulata, Lyda 102
annulata, Macrophya, *Tenthredo* 84
annulata, Prosopis 259
annulata, Sphex 161
annulata, Sphex 261
annulata, Stelis, *Anthidium* 302
annulata, Tenthredo 44, 90
annulatus, Crabro 29
annulatus, Crossocerus 229
annulatus, Crossocerus elongatulus 229
annulatus, Cryptocheilus, *Sphex* 162
annulatus, Dolerus 80
annulatus, Harpactus 223
annulatus, Hylaeus, *Apis* 259
annulatus, Stizus, *Larra* 225
annulatus, Urocerus 116
annulicornis, Macrophya 85
annuligera, Tenthredopsis, *Tenthredo* 98
annulipes, Caliroa, *Tenthredo* 45
annulipes, Crossocerus, *Blepharipus*
 annulipes 227
annulipes, Dolerus 77
annulipes, Lasioglossum, *Halictus* 282
annulitarsis, Birka, *Selandria* 76
annulitibia, Macrophya 85
annulitibia, Xiphydria 115
annulus, Crabro 228
anomala, Pseudapis 292
anomala, Tenthredo 43
anomalipes, Halictus 277

anomaloptera, Euura, *Nematus* 56
 anomoneurae, Psenulus,
Nipponopsen 252
Anopliini 172
Anoplinelus 172
Anoplius 164
Anoplius 172
Anoplius subg. 172
Anoplochares subg. 168
Anoplolyda 104
Anoplonyx 53
Anopompilinus 168
Anospilus 168
Anospilus subg. 168
Anothyreus 226
Anoxus 125
Anoxybelus 242
Antalastor 178
Antaphelopus 118
antennalis, Belomicrus 242
antennalis, Pedinomma
 rufescens var. 122
antennalis, Polistes biglumis var. 192
antennalis, Selandria flavens 76
antennata, Eucera 321
antennata, Pachyprotasis, *Tenthredo* 87
antennata, Prosopis 259
antennata, Sciapteryx laeta 89
antennatus, Sphecius, *Larra* 224
antennatus, Urocerus, *Sirex* 113
Anteon 118
Anteoninae 118
Antepipona 180
Anthemois 305
Anthidiellum 300
Anthidiellum subg. 300
Anthidiini 300
Anthidium 300
Anthidium subg. 300
Anthocopa subg. 297
anthocopoides, Hoplitis, *Osmia* 297
Anthomegilla subg. 323
Anthophilus 255
anthophilus, Nematus 56
Anthophora 323
Anthophorini 322
anthracinum, Trichiosoma 30
anthracinus, Ceratophorus 247
anthracinus, Dolerus, *Tenthredo* 79
anticus, Dolerus, *Tenthredo* anticus 77
antigae, Sphecodes 287
antigana, Nomada 314
antilope, Ancistrocerus, *Vespa* 178
antipoda, Monostegia 45
antoniae, Megalodontes 107
Antronius 249
antropovi, Dipogon 164
antropovi, Laelius 124
antropovi, Odontocerabro 235
anxius, Crabro 229
Aomodyctium 39
Aoplopus 165
apakensis, Cerceris 252

apakensis, Megalodontes 107
Apareophora 45
apatellatus, Halictus 278
Apericlista 51
aperta, Corymbas, *Siobloides* 84
aperta, Tenthredo 76
Apethymus 40
Aphadnurus 48
Aphaenogaster 203
Aphanilopterus subg. 192
Aphanisus 51
aphantoneura, Pristiphora, *Nematus* 70
Aphelinidae 118
Aphelinus 118
aphidum, Crossocerus 229
Aphiloctenus 172
Apiarus 332
apicalis, Ammophila 213
apicalis, Anoplonyx, *Leptopus* 53
apicalis, Apethymus, *Tenthredo* 40
apicalis, Eutomostethus,
Monophadnus 47
apicalis, Haplogonatopus 121
apicalis, Hylaeus 284
apicalis, Lindenius 234
apicalis, Macrophyia 85
apicalis, Megachile 305, 306
apicalis, Nematus 55, 61
apicalis, Parna, *Blennocampa* 51
apicalis, Sciapteryx 82
apicalis, Sphecodes 215
apicalis, Tenthredo 99
apicalis, Tremex 113
apicalis, Trypoxylon 244
apicata, Andrena 263
apicata, Osmia 299
apicatus, Sphecodes 288
Apicerceris 252
apicornis, Monoplopus 109
apicifrons, Pontania 58
apiornatus, Symmorphus,
Ancistrocerus 191
apiciscerra, Amauronematus
viduatinus 66
Apicula 332
apiculata, Coelioxys 304, 305
Apidae 309
Apiformes 257
apiformis, Masaris 176
apilinguaria, Vespa 222
Apinae 319
Apini 332
Apiraptrix 252
Apis 332
Apius 244
Apobembex 221
Apocabro 227
Apocrita 118
Apoctemnius 231
Apoidea 211
Apoides 227
apollineus, Bombus cullumanus 328
Aporhopalum 235
Aporia 250
Aporina 166
Aporina 250
Aporinellus 168
Aporinellus subg. 168
Aporini 166
Aporinus 166
Aporoideus 167
Aporus 166
Aporus subg. 166
apostata, Astata 218
Apoxiphia 115
appendiculata, Ceropales pygmaea 160
appendiculata, Pristiphora, *Nematus* 70
appendiculata, Xylocopa 309
appia, Megachile 306
appropinquans, Myrmilla 154
appropinquans, Alpinobombus
 kirbyellus 327
appropinquans, Halictus 282
approximatus, Nematus 61
apraria, Andrena bicolor ssp. 267
apricoti, Pristiphora 71
apricus, Dolerus, *Dosytheus* 77
aprilia, Andrena 269
apristum, Lasiosclossum, *Halictus* 282
Aproceros 25
Aprosthemia 25
aptera, Apis 157
Apterodryinus 120
Apterogyna 159
Apterogninae 159
apterus, Mystrophorus 119
apunctata, Smicromyrme
 hasanensis f. 156
Apycnemia 212
aquilegiae, Nematus 74
aqulonia, Formica 199
aqulonius, Eumenes 181
aqulonius, Eumenes rubronotatus 181
aquilus, Megalodontes 107
aquisgranensis, Prosopis 261
arabica, Vespa orientalis 194
Arachnodaicter 172
Arachnophila 172
Arachnophroctonus subg. 172
Arachnospila 168
Arachnospila subg. 168
Arachnotheutes 173
aragonensis, Spatiariella hyalinata 262
arapahonum, Pteronus 64
Arapatka 149
arborea, Vespa 195
arbusculae, Euura, *Pontania* 56
Arbusia 82
arbustorum, Hylaeus 277
arbustorum, Katamenes, *Vespa* 184
arbustorum, *Tenthredo* 84
arbuti, Lyda 105
Arcesillas 224
archanensis, Andrena 265
archanensis, Osmia 296
archangelskii, Empria 41

Archarpactus 221
Archianthidium subg. 303
archiducalis, *Pamphilus* 104
arctica, *Anthophora* 323
arctica, *Euura* 56
arctica, *Mesoneura* 72
arctica, *Rhogogaster* 91
arctica, *Tenthredo* 91
arctica, *Tenthredo, Enicia* 96
arctica, *Trichiosoma* 30
arctica, *Vespa saxonica* var. 195
arcticola, *Amauronematus* 64
arcticola, *Dolerus* 78
arcticola, *Lygaeonematus* 71
arcticola, *Nematus* 64
articornis, *Euura, Pontania* 56
arcticum, *Trichiosoma* 30
arcticus, *Dolerus* 78
arcticus, *Emphytus* 39
arcticus, *Nematus* 64, 66
arcticus, *Odynerus* 191
Arctochrysis 139
arcuata, *Tenthredo* 96
arcuatooides, *Tenthredo* 96
arcuatum, *Anteon* 118
ardens, *Andrena* 273
ardens, *Bombus* 330
ardens, *Hedychridium* 128
ardens, *Hedychridium, Chrysis*
ardens 128
ardens, *Nematus* 67
Ardenthidium 300
ardeola, *Andrena humilis* ssp. 265
Ardis 45
arduinoi, *Polistes bimaculatus* var. 192
arenaria Psammophila 214
arenaria, *Cerceris, Sphex* 253
arenaria, *Mutilla* 153
arenarius, *Mellinus* 223
arenarius, *Panurginus* 276
arenarum, *Pheidole pallidula* var. 207
arenicola, *Formica sanguinea* 200
arenosus, *Halictus leucaheneus* 279
arensi, *Hedychridium* 128
Argalictus subg. 277
Arge 22
argentata, *Andrena* 268
argentata, *Dasylabris maura*
nostras var. 153
argentata, *Dasypoda* 293
argentata, *Enodia* 215
argentata, *Megachile* auct. 306
argentata, *Nomada* 311
argentata, *Tachytes* 240
argentatus, *Oxybelus* 242
argentatus, *Tachyagetes* 174
argentatus, *Tachysphex* 237
argenteofasciata, *Mutilla* 153
argenteus, *Crabro* 233
argenteus, *Dicyrtomellus* 170
argentifrons, *Cerceris* 253
Argidae 22
argillaceus, *Bombus, Apis* 328

Arginae 22
argiola, *Pyramica* auct. 207
argiola, *Strumigenys, Epitritus* 207
argivus, *Odynerus rufospinosus* var. 187
Argogorytes 221
argonautarum, *Urocerus, Sirex* 113
Argus, *Sphex Cribalaria* 226
Argyrammophila 212
argyrauxa, *Dasylabris maura*
nostras var. 153
argyrea, *Larrada* 240
argyrea, *Megachile* 306
argyreus, *Tachytes, Larrada* 240
argyrius, *Chilosphaex, sphex* 214
argyrius, *Sphex* 214
argropyga, *Notogonia* 237
Argyroselenis 317
argyrotrichus, *Tachysphex* 239
ariadne, *Chrysis* 138
ariadne, *Chrysis scutellaris* var. 138
ariasi, *Anthidium punctatum* var. 301
Aridopomilus 168
arietina, *Andrena* 267
arietina, *Anthophora* 323
armata, *Ammophila* 213
armata, *Andrena* 325
armata, *Nomada* 311
armata, *Nomiapis, Nomia* 291
armata, *Onycholyda, Pamphilus* 103
armata, *Pristiphora, Nematus* 71
armata, *Vespa* 222
armatus, *Sphex* 213
armeniaca, *Coelioxys brevis* 304
armeniaca, *Dasylabris, Mutilla*
maura 154
armeniaca, *Eucera, Tetralonia* 321
armeniaca, *Myrmilla calva* 154
armeniacus, *Bombus* 331
armeniacus, *Hoplomerus* 187
armeniacus, *Pamphilus* 104
armenica, *Formica cinerea* var. 200
armenius, *Ateuchopus* 111
armifera, *Pristocera* 123
armiger, *Oxybelus* 242
armillatus, *Dolerus* 77
armipes, *Crabro* 230
Armitarsus 84
arnaudi, *Ammophila* 212
arnoldi, *Halictus* 284, 285
arnoldii, *Chrysis* 133
arnoldii, *Myrmica* 205
arnoldii, *Pachygonatopus* 121
arnoldii, *Strongylognathus* 207
arnoldii, *Temnothorax* 208
Arpactus 222, 223
arpactus, *Mellinus* 221
arrectus, *Katamenes dimidiatus* 185
arrogans, *Episyron, Pompilus* 166
arrogans, *Nomada* 311
arrogans, *Tenthredopsis* 98
arsenievei, *Panurginus* 276
arsenjevi, *Colletes* 257
articulata, *Andrena* 272
articulata, *Methocha, Mutilla* 149
articulata, *Mutilla* 149
Artiepyris 123
Artiotilla 157
arundinis, *Cephus* 108
arvenica, *Sapuga*
quinquepunctata var. 145
arvensis, *Cephalcia* 102
arvensis, *Mellinus, Vespa* 246
arvensis, *Miscus* 212
arvensis, *Vespa* 246
arzhakanensis, *Eudynerus egregius* 191
asachalinense, *Trichiosoma* 30
Asaconoton 244
asahi, *Arge simillima* 25
asakura, *Sphecodes* 290
Asaropoda 322
aschabadensis, *Andrena*
lepeletieri var. 270
aschabadensis, *Eumenes* 181
aschabadensis, *Pasites*
maculatus var. 319
asensioi, *Hoplitis tridentata* 297
asericae, *Tiphia* 150
ashabadensis, *Crocisa* 327
ashmeadi, *Sierola* 125
asiaeminoris, *Halictus* 277
asianum, *Hedychridium* 128
asianum, *Hedychridium integrum* 128
Asiapterocheilus subg. 187
asiatica, *Ammophila* 212
asiatica, *Ancyla* 319
asiatica, *Bembix* 222
asiatica, *Chrysis rutilans* var. 135
asiatica, *Dolichovespula* 194
asiatica, *Eucera* 321
asiatica, *Holopyga generosa* 131
asiatica, *Holopyga gloriosa* var. 131
asiatica, *Larropsis, Ancistromma* 236
asiatica, *Podalonia* 213
asiaticum, *Diploplectron* 218
asiaticus, *Bombus* 330
asiaticus, *Crossocerus* 230
asiaticus, *Pompiloides sexmaculatus* 168
Asiemphytus 40
asioboreus, *Eumenes* 182
Asitrichiosoma 30
askhabadensis, *Meria* 149
asper, *Dolerus* 79
asperatus, *Tachysphex* 239
aspericeps, *Pristiphora* 73
asperlatus, *Pristiphora* 74
aspersa, *Myrmica* 205
asperula, *Andrena* 267
asperulus, *Halictus* 279
aspidiphorus, *Crabro* 233
assamensis, *Elampus, Ellampus* 127
assiduus, *Apis* 269
assimilis, *Blennocampa* 48
assimilis, *Chrysis* 133
assimilis, *Crossocerus, Crabro* 227
assimilis, *Hylobota* 24
associa, *Cerceris* 252

associa, Polistes 192
associus, Polistes 192
astarte, Nomada 311
Astata 218
Astatinae 218
Astatus 109, 111, 218
asterabadiae, Andrena 269
Asthenoctenia 170
Asthenoctenus 170
Asticta 42
Astochus 83
Astochus subg. 83
Astomyrme subg. 156
astrachanensis, Euodynerus caspicus 183
astragali, Anthophora 324
astragalina, Andrena 272
astrictus, Leptochilus 185
astutus, Nematus 71
asunica, Andrena 265
asuniensis, Andrena 270
aterrima, Melecta 326
Atelosphex 237
ater, Alyson 244
ater, Bethylus 150
ater, Crabro 227
ater, Crabro 251
ater, Miscophus 240
ater, Pasites 318
ater, Psen, Crabro 251
ater, Sphecodes 251
ater, Stigmus 250
ater, Philanthus 276
aterrima, Andrena 272
aterrima, Apis 302
aterrima, Apis 303
aterrima, Camponotus herculeanus var. 198
aterrima, Cataglyphis aenescens 199
aterrima, Osmia 298
aterrima, Phymatocera, Tenthredo 52
aterrima, Pristiphora 70
aterrima, Tenthredo 52
aterrimum, Diprion pallipes 34
aterrimus, Allantus 95
aterrimus, Camponotus 198
aterrimus, Crabro 228
aterrimus, Dolerus germanicus 78
aterrimus, Nematinus 68
Ateuchopus 111
athalassae, Andrena morio var. 269
Athalia 43
Athalinae 43
athaloides, Eurhadinoceraea, *Monophadnus* 47
athaloides, Poppia 81
athenensis, Andrena 271
athyrii, Thrinax, *Hemitaxonius* 83
atlantica, Osmia 299
atlantis, Paururus 112
Atomostethus 47
Atoposelandria 75
Atopostigmus 249

atra, Andrena 272
atra, Anthophora robusta 324
atra, Apis 268, 272
atra, Bembix 222
atra, Blasticotoma 27
atra, Crocisa 326
atra, Euura, *Pteronus* 56
atra, Myrmosa 152
atra, Myrmosa atra 152
atra, Osmia 298
atra, Pontania 56
atra, Selandria 45
atra, Tenthredo 93
atracornis, Monophadnus 50
atramentaria, Tenthredella mesomelas 91
atramentaria, Tenthredopsis parvula 99
atramentarium, Diprion virens 34
atramentata, Anthophora khambana var. 324
atrata, Andrena 271
atrata, Anthophora furcata f. 323
atrata, Calameuta 108
atrata, Hylotoma 23
atrata, Larra 237
atrata, Neurotoma 103
atrata, Nomada 311, 318
atrata, Pristiphora 72
atrata, Prosopis 260
atrata, Tenthredo 24
atratilobis, Tenthredopsis hungarica 99
atratina, Mimumesa, *Mimesa* 251
atratinus, Dicyrtomellus 168
atratula, Anthophora robusta 324
atratula, Eriocampa 46
atratula, Myrmica 203
atratula, Tenthredo 46
atratulus, Anergates, *Myrmica* 203
atratulus, Evagetes 171
atratulus, Hylaeus 259, 286
atratum, Trypoxylon 252
atratus, Allantus 91
atratus, Amauronematus 60
atratus, Ammobates apunctatus var. 318
atratus, Aphelopus, *Dryinus* 118
atratus, Dolerus dubius 78
atratus, Dryinus 118
atratus, Haplogonatopus 121
atratus, Liris, *Larra* 237
atratus, Psen 251
atrescens, Sphecodes 287
atricallosa, Prosopis moricella f. 260
atricapillus, Dolerus 79
atriceps, Eucera 320
atriceps, Melitta 272
atricilla, Anthophora 324
atricollaris, Tenthredella maculata 92
atricollis, Eucera 320
atricolor, Camponotus, *Formica* 198
atricornis, Emphytus viennensis 38
atricornis, Eumenes 181
atricornis, Halictus 286
atricornis, Monophadnus 50
atricornis, Sirex 112
atricornis, Tenthredo 82
atricula, Andrena 270
atrifacialis, Prosopis sinuata var. 262
atrifemoribus, Tenthredopsis parvula 99
atrilobis, Tenthredopsis parvula 99
atrimandibularis, Polistes 192
atripes, Cephus 108
atripes, Discoelius dufouri 181
atripes, Euodynerus quadrifasciatus 183
atripes, Pelopoeus 214
atripleuris, Tenthredopsis parvula 99
atriscutis, Tenthredopsis excisa 99
atritiventris, Megachile 308
atritiventris, Osmia 298
atritiventris, Tenthredopsis sordida 100
atropacalis, Tenthredo maculata 92
atrocaerulea, Osmia 299
atrocaudatus, Bombus ussurensis 328
atroclypeata, Tenthredo 97
atrocoerulea, Andrena 270
atrocoerulea, Hylotoma 23
atrocyanæa, Podalonia, *Psammophila* 213
atrofasciata, Tenthredo 93
atroferruginea, Anthophora 324
atroflava, Nomada pectoralis var. 315
atrophirtus, Sphecodes 215
atromicans, Hylaeus 261
atronotata, Tenthredopsis andrei 100
Atropancistrocerus 189
atropilosus, Sphecodes 215
atropis, Halictus 283
atropos, Alastor 178
atropos, Melecta 326
atropos, Odontomyrus 189
atropos, Sierolomorpha 151
atropyga, Chalicodoma 306
atrorubricata, Andrena 272
atroskopacea, Anthophora robusta 324
atrosutellaris, Nomada 311
atrosutellatum, Diprion simile 33
atrum, Pison Alyson 244
attelaboides, Formica 197
attenuatum, Trypoxylon 244
attila, Camponotus 199
auberti, Pontania 61
auctumnalis, Nomada ruficornis var. 316
auctus, Ancistrocerus, *Vespa* 178
aucupariae, Aglaostigma, Tenthredo 83
aucupariae, Pamphilus 104
augusta, Pseudepipona, *Odynerus* 188
aulatus, Amauronematus 55
aulica, Andrena 272
aulica, Tenthredo 90
Auplopodina 165
Auplopus 165
auraneus, Dolerus elderi 78
aurantiaca, Meria, *Myzine* 149
aurantiaca, Nomada 313
aurantiacus, Nematus 57
aurantiacus, Odynerus 188
aurantiacus, Oxybelus 242

aurantifer, Heriades crenulatus 296
aurata, Abia, *Zaraea* 28
aurata, Arge, *Hylotoma* 22
aurata, Dasypoda 293
auratus, Crabro 232
auratus, Nysson 219
auratus, Pseudomalus, *Sphex*
 auratus 132
auratus, Sphecodes 237
aureatensis, Tenthredo 86
aureicolle, Hedychrum 130
aureicollis, Pseudochrysis, *Chrysis* 140
aureocincta, Nomia 291
aureomaculata, Mutilla 158
aureopectus, Nematus 58
auriceps, Chrysis 134
auricomus, Crabro 233
auriculata, Tenthredopsis, *Perineura* 98
aurifrons, Psen 251
aurifrons, Tachytes 239
aurigena, Nomada lineola var. 313
auriginosus, Palarus 243
aurilabris, Crabro 231
auripennis, Arge 22
auripes, Anthidium 301
aurita, Cerceris 253
aurita, Lyda 104
auritae, Euura 56
Auroabia 28
aurolimbata, Nomada lineola var. 313
aurolimbatus, Coelioxys 305
aurora, Andrena 274
aurotecta, Chrysis 143
aurulenta, Osmia, *Apis* 298
aurulenta, Vespa orbata var. 196
aurulentus, Sphecodes 237
ausa, Arachnospila, *Pompilus* 168
ausonius, Smicromyrme 156
auster, Symmorphus mutinensis 191
austerus, Dolerus 77
australe, Anthidium oblongatum 301
australis, Hemichroa, *Tenthredo* 67
australis, Sirex 112
australis, Tenthredo 67
austriaca, Andrena 273
austriaca, Andrena auct. 267
austriaca, Chrysis 139
austriaca, Chrysis 140
austriaca, Chrysura, *Chrysis* 139
austriaca, Mutilla 153
austriaca, Nomada 316
austriaca, Pemphredon, *Diphlebius* 248
austriaca, Tenthredopsis 100
austriaca, Vespa 195
austriaca, Vespa, *Vespa* 195
austriacum, Aprosthemma, *Schizocera* 26
austriacum, Lasioglossum
 subfulvicorne 286
austriacum, Rhopalum, Crabro 236
australialis, Holopyga 131
austrinus, Sphecodes subovalis 290
Austrolasius subg. 201
Austronomia subg. 291

autumnalis, Andrena
 carbonaria var. 272
autumnalis, Emphytus 40
Avodryinus 120
awaensis, Sphecodes 289
axillare, Aprosthemma, *Schizocera* 26
azalearum, Andrena 270
azhendarovensis, Arge 22

babuganicus, Leptothonax
 pithyusae 208
baccarum, Nematus 63
baccata, Nomada 311
Bacconematus 55
bacillus, Osmia 299
backi, Trypoxylon 245
bactriana, Didineis 219
bactriana, Solierella 241
Bactroceros 104
baeri, Chrysis 139
baeri, Nomada 311
baerii, Eumenes 185
baerii, Melecta, *Pseudomelecta* 326
baeticus, Gonatopus, *Mercetia* 120
baguenai, Notogonidea 237
Baguenaia subg. 174
baicalensis, Andrena 274
baicalensis, Anthophora aestivalis 325
baicalensis, Eumenes
 arbustorum var. 185
baicalensis, Osmia 298
baicalica, Anthrena 265
baicalica, Megachile 307
baicalica, Megachile lapponica 307
baicaliella, Andrena 265
baigacuma, Smicromyrme 155
baikalensis, Myrmica 205
baikalensis, Strongylogaster 82
baikalica, Cephalcia 102
Baisobius 122
baja, Tenthredo 90
bajulus, Dolerus 77
bakurianica, Myrmica 205
bakurianica, Myrmica pilosiscapus 205
balassaglooi, Pteronus 64
balassoglooi, Anthophora,
 Habropoda 324
balcanicus, Eumenes lunulatus var. 181
balcanicus, Oplomerus reniformis 187
balcanicus, Sphecodes subovalis
 aestrinus 290
baldensis, Perineura scutellaris 100
balearicus, Cephus 110
baleicum, Lasioglossum, *Halictus*
 baleicum 282
baleina, Andrena 267
baleina, Megachile 307
ballioni, Mutilla 157
ballioni, Odynerus 180
ballioni, Taxonus 43
balneorum, Lasioglossum 284
balteata, Tenthredo 94
balteatus, Bombus 327

balteatus, Colletes 259
balteatus, Pamphilus, *Lyda* 104
baltica, Andrena nitida 269
baltica, Empria 41
baltica, Eumenes papillarius var. 182
baltica, Eupontania acutifoliae 55
baltica, Odynerus crassicornis var. 191
baltica, Pemphredon 248
balticus, Colletes 257
balticus, Halictus 284
Baltilla subg. 153
baltistanica, Anthophora parietina 324
baluchistanensis, Plesia 149
banatica, Nomada 317
bantaizana, Abia 28
baratonis, Sphecodes 288
barbara, Formica 204
barbareae, Andrena auct. 268
barbarum, Trypoxylon figulus 244
barbarus, Euodynerus egegius 191
barbata, Melitta 274
barbatula, Andrena 268
barbatula, Melitta 274
barbatulum, Monomorium 205
barbatulus, Eumenes pomiformis 181
barbatum, Anthidium 302
barbatus, Hylaeus 259
barbilabris, Andrena, *Melitta* 268
barbipes, Crossocerus 227
barbiventris, Bembix 221
barbouri, Vespa analis var. 193
barbutellus, Bombus, *Apis* 329
barchanica, Myrmica bergi var. 205
barkensis, Halictus 287
basalicincta, Nomia 292
basalis, Allantus, *Tenthredo basalis* 38
basalis, Andrena 274
basalis, Andrena parviceps var. 268
basalis, Crabro 234
basalis, Empria 41
basalis, Halictus morio 284
basalis, Lophyrus 34
basalis, Nematus fulvus 61
basalis, Nomada 311
basalis, Nomia 291
basalis, Odynerus 188
basalis, Psen 250
basalis, Sphecodes fuscipennis 287
basalis, Tetralonia 322
basicirus, Halictus 281
basilaris, Andrena 268
basilinea, Andrena 274
basimacula, Apis 283
basimacula, Tenthredo 98
basirufa, Mimesa 250
basizonata, Tenthredo 94
batai, Dasylabris italicica f. 154
batava, Andrena 263
batava, Nomada 313
bateiae, Andrena 269
batesiae, Andrena 269
Banthanthidium 301
Bathyblepta 22

Batozonellus 166
batsumerensis, Megachile analis 307
atumicum, Trypoxylon 244
bavarica, Vespa 195
beata, Spilomena 249
beaticola, Bombus, *Pyrobombus* 330
beumonti, Andrena 274
beumonti, Eopsis 42
beumonti, Mimumesa, *Psen* 251
beumonti, Palarus 243
beumonti, Tenthredo 97
beumonti, Trypoxylon 244
beckeri, Arge 22
beckeri, Metallus, *Entodecta* 50
beckeri, Mimesa 250
beckeri, Pseudepipona, *Odynerus* 188
beckeri, Pseudotiphia, *Tiphia* 150
bekkeri, Metrionotus 125
Belalastor 178
Beleses 40
Belesidea 40
belgica, Mimesa 251
belgicus, Arpactus 223
belicovi, Andrena 274
belikovi, Nomada 313
beljgovskii, Myrmilla sarmatica var. 154
bellicosus, Oxybelus 242
bellieri, Cephus 109
bellina, Andrena labiata ssp. 272
bellona, Vespa 194
bellus, Nematus 63
bellus, Oxybelus 243
belokobylskii, Arachnospila 168
belokobylskii, Priocnemis 163
belokobylskij, Hedychradium 128
belokobylskyi, Pseudisobrachium 123
Belomiricus 241
bembeciformis, Hemipterochilus,
Odynerus 184
Bembecinus 221
Bembex 221
Bembicinae 219
Bembicinae 220
Bembix 221
benefica, Andrena 263
bengasinus, Eumenes mediterraneus 182
bennetti, Aphelopus 118
bensonii, Metallus 50
bensonii, Rhadinoceraea 52
bequaerti, Anthidium
punctatum var. 301
berberiana, Chrysis mediata 143
berberidis, Arge 22
Berberidisformis 22
berberus, Halictus 287
berberus, Sphecodes monilicornis 289
berezovskii, Omalus, *Ellampus* 131
berezowskii, Abia 28
bergi, Chrysis 134
bergi, Dasylabris 153
bergi, Myrmica 205
bergi, Pseudomalus, *Ellampus* 132
bergiellus, *Ellampus* 132

bergmani, Euura, *Amauronematus* 56
bergmani, Siobla sibirica 89
bergmanni, Euura, *Nematus* 56
bergrothi, Conaspida 43
berlandi, Bethylus 125
Berlandi, Crabro elongatus 229
berlandi, Dolerus germanicus 78
berlandi, Macrophyia albicincta 84
berlandi, Prosopis 262
bernardi, Tenthredo 91
bernardiformis, Tenthredella stulta 92
berolinensis, Halictus 286
bertilpoppii, Euura 57
bertolini, Macrophyia 86
Bethylidae 123
Bethylinae 125
Bethylus 125
bettoh, *Psen* 251
betulae, Cimbex 29
betulae, Cynips 31
betulae, Euura, *Amauronematus* 57
betulae, Fenusia 52
betulae, Lasioglossum fratellum 283
betulae, Leptothorax muscorum var. 204
betulae, Nematus 74
betulae, Pamphilus, *Tenthredo* 104
betulae, Tenthredo 74
betulae, Xiphydria,
Pseudoxiphydria 115
betularius, Nematus 74
betuleti, Cimbex 31
betuleti, Scolioneura, *Tenthredo* 52
betuleti, Tenthredo 52
betulinus, Nematus 74
beuthini, Andrena 267
beybienkoi, Nematus 64
bezzii, Pterochilus 189
biaccinctus, Omalus, *Ellampus* 131
bianchii, Chrysis 134
bianchii, Hortobombus consobrinus 328
biannulatus, Nematus 62
biaruatus, *Odynerus* 180
biareolatus, Hylaeus 262
Biastes 318
Biastini 318
Biastoides 318
biblicus, Eumenes sichelii var. 185
bicallosum, Lasioglossum, *Halictus* 280
bicarinata, Andrena auct. 271
bicarinata, *Tiphia* 150
biciliata, Anthophora 324
biciliatus, Bombus tricornis 332
bicincta, Cerceris 253
bicincta, Tenthredo 93, 97
bicinctus, Allantus 98
bicinctus, Crabro 221
bicinctus, Crabro 224
bicinctus, Halictus 280
bicinctus, Lestiphorus, *Crabro* 224
bicinctus, Pamphilus hortorum 104
bicolor, Ammobates 318
bicolor, Andrena 267
bicolor, Aporus 166
bicolor, Athalia 44
bicolor, Bembix 221
bicolor, Blennocampa 51
bicolor, Chrysis 134
bicolor, Dasylabris, *Mutilla* 153
bicolor, Dolichurus 211
bicolor, Eremiasphecium
Shestakova 246
bicolor, Euura, *Amauronematus* 57
bicolor, Evagetes 170
bicolor, Gonatopus, *Dryinus* 120
bicolor, Larra 238
bicolor, Lyda 105
bicolor, Microphadnus 173
bicolor, Mimesa, *Psen* 250
bicolor, Miscophus 240
bicolor, Miscophus 241
bicolor, Myrmica 205
bicolor, Myrmosa 152
bicolor, Osmia, *Apis* 299
bicolor, Oxybelus 242
bicolor, Tachytes 238
bicolor, Tenthredo 23, 39
bicolor, Tenthredopsis litterata 99
bicoloripes, Ceropales bicoloripes 160
bicoloriventris, Megachile 306
bicornes, Calameuta 108
bicornis, Apis 297
bicornis, Lasius, *Formica* 201
bicornis, Osmia, *Apis* 299
bicornuta, Cerceris 254
bidens, Crabro 228
bidens, Elampus, *Notozus* 127
bidens, Lestodryinus 120
bidens, Osmia 298
bidens, Tachytes 240
bidens, Vespa 220
bidentata, *Apis* 305
bidentata, Bembix 221
bidentata, Chrysis 143
bidentata, Osmia 298
bidentatus, *Psen* 250
bidenticulata, Heriades 296
bidenticulatus, *Odynerus* 180
bidenticulatus, Smicromyrme 156
bidentulus, Philoctetes, *Hedychrum* 132
bieglebeni, Alastor 178
bifasciata, Deuteragenia,
Ichneumon 164
bifasciata, Larra 225
bifasciata, Nomada 311
bifasciata, Tenthredo 90, 97
bifasciatellus, Halictus 281
bifasciatus, Symmorphus, *Vespa* 191
bifida, Abia 28
bifida, Nomada 316
Bifidoceropales subg. 160
bifidum, Aprosthemia, *Hyloboma* 26
bifidus, Halictus 278
bifidus, Microdynerus 186
bifidus, Pompilus 172
bifoveolatum, Dianthidium 301
bifoveolatus, Rhophites 291

- bifrons*, Janus 110
bifurca, Aprosthemia, *Hylotoma* 26
bifurcata, Konowia 116
bigallae, Euura, *Pontania* 57
bigemmis, Tenthredo 95
bigens, Abia 28
biglumis, Polistes, *Vespa* 192
biguetina, Cimbex 29
biguttata, Artiotilla, *Mutilla* 157
biguttata, Mutilla 157
biguttata, Tenthredo 95
biguttata, Tenthredopsis dorsalis 99
biguttatus, Crabro 226
biguttatus, Dolerus 84
bihamata, Chrysis 143
bilineata, Trichiosoma 31
bilineatus, Anoplonyx 53
bilineatus, Nematinus, *Tenthredo* 68
bilineatus, Philanthus 262
biloba, Selandria 68
biloba, Vespa 195
bilobata, Chrysis 134
bilunatus, Pompilus 167
bilunulatus, Lestiphorus 224
bilunulatus, Lestiphorus
bilunulatus 224
bimacula, Vespa 192
bimaculata, Andrena, *Melitta* 272
bimaculata, Anthophora, *Apis* 323
bimaculata, Lyda 101
bimaculata, Mutilla 155
bimaculata, Nomada 311
bimaculata, Selandria 76
bimaculata, Tenthredo 94
bimaculata, Vespa 192
bimaculatus, Allantus 95
bimaculatus, Crossocerus 230
bimaculatus, Dolerus, *Tenthredo* 77
bimaculatus, Eumenes 182
bimaculatus, Sphecodes 219
binghami, Vespa 193
binominata, Xylocopa 310
binominatus, Nannodynerus 180
binotata, Anthophora 323
binotata, Thomsonia excisa 99
binotatus, Crossocerus 230
bipartitor, Pemphredon 249
bipartita, Andrena 269
bipartita, Eucera 320
bipartita, Euura, *Nematus* 57
bipartita, Tenthredo 51
bipartitus, Nematus 57
bipicta, Pteronidea 60
bipilosa, Formica cinerea var. 201
bipunctata, Bembix 221
bipunctata, Hartigia draconis 110
bipunctata, Melecta 326
bipunctata, Tenthredo 45
bipunctata, Tenthredo 45, 73, 95
bipunctata, Vespa 262
bipunctatus, Crabro 232, 246
bipunctatus, Hylaeus 282
bipunctatus, Mellinus 262
bipunctatus, Monophaenus 50
bipunctatus, Oxybelus 242
bipunctatus, Stizus, *Larra* 225
bipunctatus, Tachysphex 238
bipunctis, Eumenes 182
bipunctula, Tenthredo bipunctula 94
bipunetis, Nomada 311
bipustulata, Andrena 271
Birk 76
Birk subg. 76
Birkmania 276
biroi, Bombus 330
biroi, Vespa parallela var. 193
biroi, Tetralonia julliani 322
birulae, Pontania 63
birulai, Pseudovespa 196
birulai, Vespa crabro var. 194
bisbimaculatus, Hylaeus 283
bisbistrigatus, Hylaeus 280
biscalis, Pristiphora, *Nematus* 71
bischoffi, Chrysis ignita 143
bischoffi, Polistes 192
bisexmaculatus, Crabro 232
bisignata, Tenthredo velox 96
bisinuatus, Hylaeus 260
bison, Crabro 228
bispinosa, Nomada 311
bispinosa, Nomiapis, *Nomia* 291
bispinosa, Rophites 290
bisstrigatus, Odynerus 183
bistillatus, Oxybelus pugnax 243
bistriatus, Allantus 98
bistriatus, Nematus 73
bistrispinosus, Rophites 291
bisulcata, Andrena 263
bitineta, Nomada obscura var. 315
bituberculata, Tenthredella 94
bituberculatus, Sphecodes 289
Bivena 83
Bivena subg. 83
biverrucatum, Trichiosoma 31
bivinctus, Halictus 277
bivittatus, Odynerus 186
bizonata, Tenthredo 39
bizonatus, Sialus 162
blakistoni, Lasioglossum 280
blanchardi, Athalia 44
blanchardianus, Odynerus 184
blancoburgensis, Nomada
lateralis var. 312
blanda, Macrophyta, Tenthredo 85
blandus, Hylaeus 261
blanki, Dolerus 79
Blankia 90
Blasticotoma 27
Blasticotomidae 27
blattivorus, Tachysphex 237
Blennocampa 45
Blennocampinae 45
blepharipes, Nomada 311
Blepharipus 227
bleusei, Arge fuscipes 23
bleusei, Schizocera 26
bleyli, Amasis 32
blidahensis, Halictus 286
bluethgeni, Lasioglossum, *Halictus* 282
bluethgeni, Nomada 311
bluethgeni, Osmia 303
bluethgeni, Rophites 291
bluethgeni, Stenodynerus 189
blüthgeni, Andrena 272
Bnunius 227
Bocchinae 119
Bocchoides 120
bofillana, Nomada 317
bogdanovi, Cardiocondyla 203
bogdanovii, Ceropales bogdanovii 160
bogdanovii, Philoctetes, *Holopyga* 132
Bohea 27
bohemani, Euura, *Nematus* 57
bohemani, Odynerus 191
bohemanni, Tenthredo 39
bohemica, Rhogogastera olivacea 92
boemicus, Bombus 329
boemicus, Nysson 220
bojus, Crabro 227
bokhaica, Deuteragenia, *Dipogon* 164
bolivari, Athalia 44
Bombias subg. 327
bombiforme, Trichiosoma 30
Bombini 327
Bombomelecta 325
Bombus 327
Bombus subg. 327
bombycina, Megachile 306
bonivuri, Andrena 274
boops, Astata, *Sphex boops* 218
boops, Sphex 218
borea, Pristiphora, *Lygaeonematus* 71
borea, Tenthredo 94
boreale, Trichiosoma 31
borealis, Anthophora 324
borealis, Belomicrus 242
borealis, Chrysis 134
borealis, Crabro 232
borealis, Cyrtogonatopoides 121
borealis, Dolichovespula 195
borealis, Ectemnius, *Crabro* 232
borealis, Formica fusca var. 200
borealis, Gonatopus 120
borealis, Macrophyta sanguinolenta 86
borealis, Megachile ligniseca var. 307
borealis, Melitta 294
borealis, Mimesa 251
borealis, Nitela 241
borealis, Nomada 314
borealis, Passaloecus 247
borealis, Tachysphex ibericus 239
borealis, Tarpa 107
borealis, Tenthredo 71
borealis, Vespa 195
borealis, Vespa 195
borealis, Vespa crabro var. 193
Boreocoelioxys subg. 304
boreoibericus, Pterocheilus
phaleratus 189

boreophila, Euura, *Pteronidea* 57
Boreopomilus 168
Boreovespula 194
borisi, Euura, *Eupontania* 57
borisi, Pamphilius 104
borodini, Ellampus 133
Bothriomyrmex 197
bothriorhina, Andrena 267
boulderensis, Pachynematus 62
braccata, Dasypoda 293
braccata, Tenthredo 40
brachyacanthus, Nematus 57
Brachyandrena subg. 264
Brachycephalapis 294
brachycephalum, Lasioglossum,
Halictus 282
brachycera, Pteronidea 57
brachycera, Tenthredo, *Allantus* 93
brachycera, Tomostethus 47
brachycercus, Cephus 109
brachycercus, Nematus 59
brachycercus, Pseudocephaleia 106
brachyceros, Osmia 297
brachycerum, Anteon, *Dryinus* 118
brachycerus, Colletes 258
Brachycolumna 55
Brachycolus 55
brachygaster, Dolerus 79
Brachymerus 236
Brachyocampa 42
Brachyodynerus 180
brachyotus, Nematus 66
Brachypipona 180
Brachyponera 209
brachypterus, Cephus 111
Brachystegus 219
Brachythops 76
brachytomus, Odynerus dantici 183
bracteata, Cerceris 253
Bradynobaenidae 159
Bradynobaninae 159
branensis, Allantus scrophulariae 97
branickei, Bombus, *Psithyrus* 329
brassicae, Andrena 271
brauneri, Temnothorax
 erviculus var. 208
brauneri, Temnothorax,
Leptothorax 208
braunsiana, Andrena 271
braunsiana, Nomada 311
braunsi, Camponotus 53
braunsi, Thomsonia 98
breadalbanensis, Pristiphora,
 Nematus 71
bremensis, Andrena 266
bremensis, Nomada lineola var. 313
breve, Tapinoma 198
brevicauda, Poecilotiphia, *Meria*
 brevicauda 150
breviceps, Cyrtuchridium 129
breviceps, Eucera 319
breviceps, Prosopis auct. 259
brevicera, Euura 57

breviclypeatus, Sphecodes 289
brevicorne, Aprosthemma, *Hyloboma* 26
brevicorne, Lasioglossum, *Halictus*
 brevicorne 282
brevicornis, Biastes, *Tiphia* 318
brevicornis, Claremontia,
Blennocampa 46
brevicornis, Colletes 258
brevicornis, Cryptocampus 66
brevicornis, Dolerus 77
brevicornis, Dolerus 79
brevicornis, Gonatopus 119
brevicornis, Hylaeus 259
brevicornis, *Hyloboma* 26
brevicornis, Macrophyia blanda 85
brevicornis, Nematus 72
brevicornis, Nomada 311
brevicornis, Osmia, *Eucera* 299
brevicornis, Pamphilus brevicornis 104
brevicornis, Passaloecus 248
brevicornis, Sphecodes 289
brevicornis, Tenthredo 53
brevicornis, Tenthredo, *Allantus* 96
brevicornis, *Tiphia* 318
brevicuneatus, Hylaeus 261
brevigena, Pompilus 170
brevihirtscopa, Andrena 263
brevilabris, Diodontus 247
brevilabris, Macrophyia 87
brevilabris, Passaloecus 247
brevimaculata,
 Prosopis variegata var. 262
breviniger, Allantus amaenus 97
brevipalpis, Hylaeus 259
brevipennis, Aglaopis 303
brevipennis, Tachysphex 237
brevipennis, *Tiphia* 150
brevipetiolata, Pemphredon 248
brevis, Coelioxys 304
brevis, Crabro 229
brevis, Crabro 234
brevis, Dolerus 79
brevis, Entomognathus, *Crabro* 234
brevis, Hoplocampa, *Tenthredo* 67
brevis, Mimesa 250
brevis, Pachycephus 110
brevis, Sphecodes 290
brevis, Tenthredo 67
breviscapus, Pompilus 173
brevisetata, Euura, *Pteronidea* 57
brevispina, Cimbex 30
brevispina, Perineura 99
brevispinis, Nematus 57
brevispinosus, Mesitius 124
brevitarsis, Andrena 292
brevitarsis, Astata 218
brevitarsis, Chrysos 134
brevitarsis, Psenulus 252
brevitarsis, Dolerus 80
brevithorax, Halictus 286
breviuscula, Stelis, *Heriades* 302
breviuscula, Tenthredo 88
breviusculus, Nematus 61

brevivalva, Euura, *Nematus* 57
breviventris, Allantus 91
breviventris, Mimesa 250
bridgmaniana, Nomada 312
bridgmani, Euura, *Nematus* 57
brischkei, Nematus 69
briseis, Lasioglossum briseis 282
britannica, Vespa 195
Brodiei, Oxybelus 243
brodmannicus, Bombus 330
bruesi, Myrmica laevinodis var. 206
Brugmoia 140
brullei, Cladius, *Priophorus* 54
Brullei, Tachytes 238
brullii, Tachysphex, *Tachytes* 238
Brullii, Tachytes 238
bruneipes, Macrophyia 87
brunnea, Tenthredo 68
brunneipes, Liris 218
brunneipecta, Euura,
Amauronematus 57
brunneofasciata, Nomada
 lineola var. 313
brunneonitida, Formica 199
brunnescens, Halictus, *Hylaeus* 277
brunnescens, Tenthredopsis
 coqueberti 98
brunneus, Aneugmenus 81
brunneus, Lasius, *Formica* 202
brunneus, Pasites maculatus var. 319
brunni, Camponotus marginatus 198
brunniapex, Pristiphora 74
brunnicornis, Nematus 63
brunnipes, Macrophyia 87
brunnipes, Paramyrmosa, *Myrmosa* 153
brunniventris, Tenthredo 45
brutia, Mutilla 157
brutia, Ronisia, *Mutilla brutia* 157
bruxellensis, Mimesa 250
bubulca, Andrena niveata ssp. 270
bubulcus, Halictus 284
buccale, Lasioglossum, *Halictus* 282
buccatus, Nematus 56
Bucephalus, Sphex scutellata 233
Bucerceris 252
bucharensis, Polistes 193
bucharica, Didineis 219
bucoda, Pristiphora 72
buda, Chrysis 134
budashkini, Melitta 294
budensis, Melitta, *Cilissa* 294
budrissi, Stigmatodipogon 165
bufo, Pristiphora, *Nematus* 71
bulgarica, Athalia glabricollis 44
bulgarica, Prospigastra 237
bulgaricus, Odynerus 192
bulgariensis, Andrena 263
Bulsanensis, Crabro 233
bulsanensis, Leptochilus alpestris
 alpestris var. 185
bupresticida, Cerceris 253
burlinii, Eumenes arbustorum var. 184
bursakovi, Tetramorium 209

busaei, Dolerus 77
Busarbidea 76
butovitschi, Diprion 33
butuani, Ectemnius 233
buyoo, Nomada 314
buyssoni, Chrysis 141
buyssoni, Dolerus germanicus 78
buyssoni, Megachile 306
buyssoni, Odynerus 185
buyssoni, Psithyrus rupestris 329
buyssoni, Tenthredopsis nassata 99
buyssoni, Xiphydria 115
Buzatlana 147
byssina, Anthidium, *Apis* 303
Bytinskia 296

cabrerae, Macrophyia militaris 86
Cabrerai, Tachysphex 238
cadderensis, Euura, *Nematus* 57
cadusa, Myrmica 205
caeca, Ctenotilla, *Mutilla* 155
caeca, Mutilla 155
caementaria, Osmia 297
caementarium, Sceliphron, *Sphex* 214
Caenocephus 108
Caenolarra 236
Caenolyda 102
Caenoneura 36
Caenopen 251
caeruleocarpa, Euura 57
caeruleocephala, Dufourea 290
caerulescens, Osmia, *Apis* 298
caeruleum, Chalybion 214
caeruleus, Elampus 127
caeruleus, Sphecodes 214
caespitum, Formica 209
caespitum, Tetramorium, *Formica* 209
caja, Tenthredo persa 90
calabra, Andrena 267
calabra, Nomada 315
calabricus, Eumenes 182
calabricus, Odynerus 185
calabrina, Melecta 326
Calagenia 165
Calameuta 108
Calameuta subg. 108
calandra, Chrysis soror 138
Calarge 22
calcarata, Apis 276
calcaratus, Allantus 97
calcaratus, Hoplomerus 187
calcaratus, Panurgus, *Apis* 276
calceatum, Lasioglossum, *Apis* 282
calceatus, Allantus, *Tenthredo* 38
calceatus, Crabro 223
calceatus, Crabro 235
calceolata, Hoplocampa 50
Calceorhopalum 235
calderoni, Harpiphorus 41
caledonicus, Nematinus, *Nematus* 68
calendarum, Apis 259
calens, Chrysis 141
calens, Stilbum, *Chrysis* 141

Caliadurgus 162
Calicurgus 161, 162
calidus, Bombus 330
californicus, Gymnonychus 70
californicus, Pelopeus 214
caligator, Tenthredo 97
caligatus, Emphytus 40
caliginosa, Megilla 324
caliginosa, Tenthredo 99
caliginosum, Lasioglossum 282
calimorpha, Chrysis calimorpha 134
calimorpha, Nomada 311
Caliosysphinga 48
Caliroa 45
calliblepharus, Allantus, *Emphytus* 37
callicerus, Nematus 75
callopyrrha, Andrena 274
callosa, Ceratina auct. 310
callosum, Hedychrum 128
callosus, Odynerus 179
Calomelissa subg. 264
calopteryx, Astata 219
calopteryx, Tachysphex 239
Calosphe 215
Calotachytes 240
calvaria, Tenthredo 96
Calyoza 123
Calyozaella 123
calypso, Chrysis 135
camelinus, Gonatopus 120
camelogigas, Sirex 113
camelus, Ichneumon 115
camelus, Lestica, *Crabro* 234
camelus, Xiphydria, *Ichneumon* 115
cameroni, Alysson 219
cameroni, Amauronematus 65
cameroni, Anteon 119
cameroni, Mutilla 156
cameronii, Nematus 63
Cameronitus 231
camilla, Tenthredo 90
campanaria, Vespa 195
campanularum, Chelostoma, *Apis* 295
campestris, Ammophila 212
campestris, Andrena 280
campestris, Bombus, *Apis* 329
campestris, Gonatopus 120
campestris, Tenthredo 91
campestris, Vespa 221
Camponiscus 69
Camponotus 198
Camponotus subg. 198
Campsomerini 147
Campsoscolia 147
Camptopoeum 275
Camptopoeum subg. 276
Campylogaster subg. 264
Campylonyx 119
camtschatcalis, Cephus 109
camtschatcalis, Dolerus pratensis 78
camtschatcalis, Euura,
Amauronematus 57
camtschatcalis, Loderus gilvipes 79

camtschatcalis, Micronematus 73
camtschatcalis, Tenthredo,
Tenthredella 94
camtschatcalis, Tenthredopsis 98
camtschatica, Athalia lugens 44
camtschatica, Empria 41
camtschatica, Euura, *Pontania* 57
camtschatica, Tenthredo maculigera 91
camtschatica, Tenthredopsis
auriculata 98
camus, Aphelopus 118
cana, Eucera 320
cana, Megachile analis 307
cana, Rophites 291
canadensis, Emphytus 39
canaliculatus, Nematus 71
canariensis, Andrena 270
canariensis, Tachysphex costai 238
candens, Abia 28
candens, Andrena 273
candiae, Andrena 269
candida, Formica 200
candida, Tenthredopsis austriaca 99
candidata, Empria, *Tenthredo* 41
candidata, Macrophyia albicincta 84
candidum, Episyron 167
canescens, Andrena 274
canescens, Bembix, *Apis* 221
canescens, Dasypoda 293
canescens, Eucera 322
canescens, Tenthredo 80
canigica, Andrena allosa ssp. 267
caninae, Eriocampa 46
cannabis, Cladius, *Trichiocampus* 54
Canonarea 83
Canoniades 76
canonota, Dasylabris mauropis var. 153
cantabrica, Heriades 296
cantatrix, Eucera 320
canus, Apis 257
canus, Rhophitoides, *Rophites* 291
canuta, Xylocopa 309
caocinnus, Psen 251
capillosa, Andrena 267
capitalis, Andrena 275
capitata, Cerceris 254
capitata, Spilomena 249
capitatus, Acanthomyops
fuliginosus 201
capitatus, Lasiurus, *Acanthomyops* 201
Capito, Crabro 227
capito, Euura, *Pteronus* 57
capitosus, Crossocerus, *Crabro* 228
capra, Odynerus 178
caprae, Apis 313
caprae, Cimbex 30
caprae, Cynips salicis 64
caprae, Tenthredo 64
caprifolia, Psen 103
caprimulga, Andrena 268
capta, Andrena decipiens ssp. 267
captiva, Arge, *Hylotoma* 22
caputaureum, Hedyphridium 128

caputaureum, Hedychridium roseum var. 128
capverdensis, Sphecodes 289
Carabro 225
caraimica, Andrena similis ssp. 274
 Carandrena subg. 264
 Caranthophora subg. 323
 carantonica, Andrena 267
 carbo, Aglaostigma, *Macrophyopsis* 83
carbonaria, Anthophora 324
carbonaria, Apis auct. 272
carbonaria, Macrophyia 85
carbonaria, Mimesa 250, 251
carbonaria, Neocimber taukushi 29
carbonaria, Tenthredo 99
carbonarium, Poecilosoma 41
carbonarius, Auplopus carbonarius, *Sphecodes* 165
carbonarius, Cephus 109
carbonarius, Crabro 228
carbonarius, Dolerus 81
carbonarius, Gorytes 221
carbonarius, Halictus 281
carbonarius, Taxonus 43
carbonatus, Halictus 281
Carbonelis subg. 147
carbonicolor, Anopilus,
Psammochares 168
carceli, Arpactus 223
carcelii, Gerbekas, *Mesitius* 124
carcelii, Mesitius 124
cardinalis, Lionotus 183
cardinalis, Pompilus 173
Cardiocondyla 203
cardioscapus, Hylaeus 259
carelica, Euura, *Nematus* 57
carinaevarnis, Halictus 279
carinata, Coelioxys 304
carinata, Dasypoda 293
carinata, Dufourea, *Halictoides* 290
carinata, Pempredon 247
carinata, Pristiphora, *Nematus* 71
carinata, Tenthredo 82
carinata, Tenthredopsis carinata 98
carinatus, Pompilus 169
carinatus, Anteon 119
carinatus, Dolerus 80
carinatus, Psen 250
Carinellum 301
Carinevagetes 170
carinifrons, Aneugmenus, *Selandria* 76
carinifrons, Arge 23
carinifrons, Psenulus, *Psen* 252
cariniventris, Halictus pollinosus 279
Carinoscolia 147
Carinostigmus 246
carinthiaca, Macrophyta, *Tenthredo* 85
carinthiacus, Camponotus 53
carinula, Andrena 275
carinulata, Megachile 305
carinulatus, Cimbex 29
carnca, Nomada flavomaculata var. 311
carnea, Andrena rutila ab. 270

carnea, Chrysis 142
carneiventris, Osmia 298
carnifex, Chrysis 134
carniolica, Osmia 298
carniolicus, Lasius 201
carpathia, Chalicodoma parietina 305
carpatica, Tenthredo 88
carpentieri, Aprosthemma 26
carpentieri, Pseudosobrachium 123
carpinata, Tenthredo 93
carpini, Ametastegia, *Emphytus* 39
carpini, Lyda 104
carpini, Tenthredo 95
casia, Tenthredopsis 98
Casipteryx 90
caspia, Cerceris 255
caspia, Tenthredo, *Allantus* 90
caspica, Amasis 31
caspica, Andrena 273
caspica, Cerceris 253
caspica, Crematogaster sordidula var. 204
caspica, Eucera 320
caspica, Eumenes sesquicinctus viratus var. 185
caspica, Mutilla 153
caspica, Sterictiphora 26
caspica, Tenthredo 92
caspica, Vespa crabro var. 194
caspicum, Lasioglossum, *Halictus* 280
caspicus, Colletes 257
caspicus, Crabro, *Blepharipus* 226
caspicus, Euodynerus, *Leionotus* 183
caspicus, Jucancistrocerus 184
caspicus, Mesitius 124
caspicus, Pachycephalus 110
caspicus, Polistes 193
casiensis, Chrysis 134
casiensis, Myrmecocystus cursor var. 199
castanea, Cimbex 29
castanea, Macrophyta 89
castellana, Nomada 311
castigata, Chrysis 134
castilianus, Sphecodes 288
castiliensis, Macrophyta rufipes 86
casularum, Heriades 296
catachloris, Nematus 65
Cataglyphis 199
catagraphus, Bombus ikonnikovi 327
catanensis, Trogaspidia, *Mutilla* 157
catellata, Birka, *Selandria* 76
cathoraticus, Nematus 70
catocala, Gilpinia, *Lophyrus* 33
catulus, Nematus 73
caucasica, Poecilotiphia 150
caucasica, Anthophora 324
caucasica, Arge 23
caucasica, Cerceris 253
caucasica, Chrysis 138
caucasica, Chrysis analis var. 134
caucasica, Corynus, *Amasis* 31
caucasica, Crocisa 327
caucasica, Empria 42

caucasica, Eucera 321
caucasica, Euura, *Pachynematus* 57
caucasica, Ferreola 173
caucasica, Macrophyta 85
caucasica, Macrophyta, *Allantus* 85
caucasica, Megachile 305
caucasica, Melitturga 276
caucasica, Mimesa 250
caucasica, Myrmica rugulosa 205
caucasica, Myrmilla meda 154
caucasica, Myrmilla, *Mutilla* 154
caucasica, Nomada fucata var. 313
caucasica, Podalonia, *Psammophila* 213
caucasica, Pseudotiphia, *Tiphia* 150
caucasica, Rhadinoceraea 52
caucasica, Sciateryx 89
caucasica, Tenthredo caucasica 90
caucasica, Xiphydria 116
caucasicola, Anthophora furcata 323
caucasicola, Cataglyphis nodus 199
caucasicola, Chrysis 134
caucasicola, Chrysis analis 134
caucasicola, Hoplitis 297
caucasicola, Macrophyta hamata 85
caucasicum, Anthidium 300
caucasicum, Anthidium diadema var. 300
caucasicum, Hedychridium 129
caucasicum, Stenamma 207
caucasicus, Allantus 90
caucasicus, Ancistrocerus oviventris 179
caucasicus, Bombus 329
caucasicus, Camponotus herculeanus 198
caucasicus, Cleptes 127
caucasicus, Harpactus 223
caucasicus, Pamphilus 104
caucasicus, Podalirius furcata var. 323
caucasicus, Rophites 291
caucasicus, Strongylogaster 82
caucasicus, Tachytes 240
caucasiensis, Chrysis analis 134
caucasiun, Hedychridium sculpturatum var. 129
caudalis, Euura, *Nematus* 57
caudatus, Coelioxys 304
caudatus, Halictus aeratus 282
caularis, Hoplitis, *Osmia* 296
caulicola, Apis 296
Cautolasius subg. 201
cavata, Empria 39
cavifrons, Ectemnius, *Crabro* 231
caviventris, Anoplus, *Pompilus* 172
Ceballosia 246
cebrionicornis, Nematus 75
ceciliae, Bombus lapponicus 330
cecilioides, Bombus lapponicus 330
cecrop, Colletes 258
cedens, Halictus alkenellus 278
cedri, Lasioglossum leucozonium 281
Celia 249
cellularis, Nematus 70
Celonites 176

- celsia*, Tenthredella 92
celtica, Mimesa 251
celtica, Tenthredo 93
celticus, Colletes cunicularius 257
Cemonus 248
cenchrus, Dolerus 80
centaureae, Melitta 294
centifoliae, Tenthredo 44
Centrias 310
centronigrata, Tenthredopsis
 flavomaculata 100
centricularis, Apis 305
centricularis, Megachile, *Apis* 306
Cepha 111
Cephalcia 102
Cephalcinae 101
Cephaledo 89
Cephaledo subg. 90
Cephaleia 102
cephalica, Ceratina callosa 310
cephalica, Mutilla 154
Cephalictoides subg. 290
cephalicus, Euodynerus 183
cephalicus, Halictus 278
Cephaloderma 124
Cephalonomia 124
Cephalonomiini 124
cephalotes, Allantus 98
cephalotes, Bethylus, *Perisemus* 125
cephalotes, Crabro 233
cephalotes, Ectemnius, *Crabro* 233
cephalotes, Ephippionotus 110
cephalotes, Megalodontes,
 Tenthredo 107
cephalotes, Osmia 299
cephalotes, Sphecodes 289
cephalotes, Tenthredo 107
cephaloiformis, Sphecodes
 quadratus 289
Cephen 310
Cephidae 108
Cephinae 108
Cephoidea 108
Cephosoma 110
Cephus 109
cerana, Apis 332
Ceraplastes 302
cerasi, Caliroa, *Tenthredo* 45
cerasorum, Tenthredo 76
cerastes, Chrysis 134
Ceratina 310
Ceratina subg. 310
Ceratinidia 310
Ceratinidia subg. 310
Ceratinini 310
Ceratocnemis 162
Ceratocolus 234
Ceratocrabro 231
Ceratopales 160
Ceratophorus 247
Ceratopompilus 168
Ceratosmia 297
ceraunius, Crabro 245
- cerberus*, Anthophora semperi 324
Cercerini 252
Cerceris 252
cereipes, Selandria 76
Cerianthidium 300
cerinthidis, Osmia 299
Cerobactrus 110
Ceropales 160
Ceropales subg. 160
Ceropalinae 160
cervicornis, Prosopis 260
cetaria, Empria 39
cetii, Apis 268
cetratus, Crossocerus, Crabro 228
Ceylalictus 292
Ceylalictus subg. 292
ceylonica, Tachytes 239
ceylonicus, Oxybelus
chaharensis, Halictus 277
chalcites, Ceratina 310
chalcites, Spinolia, *Chrysis* 141
Chalcogonatopus 120
Chalcolamprus 234
Chalepoxenus 203
Chalicodoma subg. 305
chalybaeum, Hedydium 130
chalybea, Ceratina 310
chalybea, Ctenoplectra 319
chalybea, Podalonia, *Ammophila* 213
Chalybion 214
chambersi, Pachynematus 59
changaiensis, Diodontus 247
chappelli, Dolerus 77
Characopygus 110
Charitandrena subg. 264
charusini, Cerceris 253
chasanensis, Prosopis 259
chasanicus, Epyris 123
chatkalenis, Cataglyphis aenescens 199
chefketi, Tetramorium 209
chefketi, Tetramorium
 caespitum var. 209
cheilon, Nematus 72
Chelogynus 118
Chelostoma 295
Chelostoma subg. 295
chelostomae, Sapyga 146
Chelothelius 119
chengtehensis, Colletes 257
chersona, Andrena 263
chersonensis, Gonatopus 120
chevrieranus, Pterochilus 189
chevrieranus, Stenodynerus,
 Odynerus 190
chevrieri, Ceratina 310
chevrieri, Psenulus, *Psen* 252
Chevrieria 248
chevrolatii, Formica 122
chibaensis, Sphecodes 288
chichibuensis, Sphecodes 290
chichibuus, Sphecodes 290
chikuzenensis, Andrena
 taraxaci ssp. 265
- Chilocares* 161
Chilosphex 214
chimaera, Andrena 270
chinensis, Brachyponera, *Ponera* 209
chinensis, Chrysis 134
chinensis, Eucera, *Tetralonia* 321
chinensis, Odynerus 189
chinensis, Polistes, *Vespa* 192
chinensis, Ponera nigrita 209
chinensis, Vespa crabro 194
Chionopompilus 171
Chionopterus 173
chionospilus, Sphecodes 289
chionura, Megachile analis 307
chipsani, Crabro 231
chittendenii, Anoxus 124
chlakovskii, Halictus 281
Chlorampulex 211
Chlorandrena subg. 265
Chloracheilus 161
Chlorodryinus 119
Chlorodynerus 180
chlorogaster, Nematus 60
chloronotus, Apathus 329
chloropus, Lasioglossum, *Halictus* 280
chloropygum, Hedydium
 chloropygum 129
chlorosoma, Rhogogaster, *Tenthredo* 88
chloroticus, Odynerus 180
chobauti, Colletes 258
chodjana, Anthophora khambana f. 324
cholodkovskii, Hedydium 130
chongar, Trypoxylon frigidum 244
chorasmius, Ammoplanellus,
 Ammoplanus 246
chorasmius, Ammoplanus 246
chosenensis, Chrysis 134
chosenensis, Chrysis nitidula 134
chosensis, Crossocerus palmipes 229
chosensis, Tiphia 150
chosrovicus, Harpactus 224
chotanensis, Halictus 284
christofi, Anthophora 324
christophi, Strongylognathus 207
christophi, Tapinoma 198
chromata, Pristiphora 71
Chrysallampus 127
Chrysidea 133
Chrysidae 126
Chrysidae 127
Chrysidiini 133
Chrysidoidea 118
Chrysis 133
chrysites, Ectemnius, *Crabro* 233
Chrysocleptes 126
Chrysogona 133
Chrysolarra 237
Chrysolyda 102
chrysomalla, Ceratina 310
chrysomelina, Osmia 299
chrysonota, Holopyga, *Ellampus*
 chrysonota 131
Chrysopheon 303

chrysoprasina, Chrysis 134
 chrysopus, Andrena 267
 chrysopyga, Andrena 275
 chrysopyga, Eucera 320
chrysopyga, Nomada 314
 chrysorrhoea, Hoplocampa,
 Tenthredo 67
 chrysocleles, Andrena, *Melitta* 271
 chrysostigma, Chrysis 134
chrysostomus, Crabro 231
 Chrysura 139
 chrysura, Macrophyia, *Tenthredo* 85
chrysura, Melitta 294
chrysura, Systropha 291
 chrysurus, Lithurgus 295
 Chthonolasius subg. 201
churchevillei, Tenthredopsis 100
Chyphonocheilus 161
Chyphonocheilus subg. 161
Chyphotinae 159
chyzeri, *Tenthredo* 94
 cicatrix, Holopyga 132
cicum, Pontania 66
Ciliaporus 174
 ciliaris, Arge, *Tenthredo* 23
ciliata, Andrena 268
ciliata, Apis 272
ciliata, Formica pratensis var. 200
ciliata, Mutilla 154
 ciliata, Poecilotiphia, *Meria* 150
ciliata, Prosopis 259
ciliatula, Andrena 268
 ciliatus, Agenioideus, *Pompilus* 167
 ciliatus, Dolerus 79
Cilissa 294
cilia, Allantus kietteri 90
 Cimbex 29
Cimbicidae 28
Cimbicinae 29
Cimbicisoma 30
cinclus, Halictus 286
 cincreus, Anthophora, *Podalirius* 325
cincta, Andrena 265
cincta, Dimorpha 219
cincta, Osmia 298
cincta, Pristiphora 71
cincta, Trachusa 303
cinctella, Bembix cinctella 221
cinctellus, Agenioideus, *Pompilus* 167
cinctipes, Blepharipus 228
cinctipes, Emphytus 38
cinctipleuris, *Tenthredo*, *Allantus* 92
cinctitarsis, Stenocrabro 228
cinctiventris, Nomada
 ferruginata var. 314
cinctus, Allantus, *Tenthredo* 38
cinctus, Crabro 221, 231
cinctus, Dioxygus, *Trachusa* 303
cinctus, Dolerus 38
cinctus, Nematus 69
 cineraria, Andrena, *Apis* 268
 cineraria, Eucera 320
cineraria, Eucera 321

cinerascens, Andrena 265, 275
cinerascens, Colletes 257
cinerascens, Dasypoda argentata 293
cinerea, Anthophora 324
cinerea, Apis 257
cinerea, Eucera 320
cinerea, Formica 200
cinereae, Euura 56
cinereae, Nematus 64
cinereipes, Birkha, *Tenthredo* 76
cinereipes, *Tenthredo* 76
cinereofusca, Formica 200
cinereofusca, Formica *cinerea* var. 200
cinereophila, Andrena 265
cinereus, Nematus 75
cinereus, Podalirius 325
cinereus, Pomphilus, *Sphex* 171
cineta, Nomada 313
cineticornis, Nomada 311
cingillipes, Allantus, *Emphytus* 38
cingillum, Allantus, *Tenthredo* 38
cingulata, Allantus 96
cingulata, Andrena auct. 272
cingulata, Holopyga 131
cingulata, Monostegia, *Poecilosoma* 43
cingulata, Nomada 268
cingulata, Phytetopoda, *Mutilla* 155
cingulata, Sphecius 170
cingulatum, Anthidium 300
cingulatus, Allantus, *Tenthredo* 38
cingulatus, Bombus 330
cingulatus, Dolerus 38
cingulatus, Pamphilus 104
cingulicornis, Chrysis cingulicornis 134
cingulifer, *Tenthredo*, *Allantus* 97
cingulum, *Tenthredo* 38, 97
cinnabarina, Arge 23
cinnabarina, Nomada 316
cinxia, Caliroa, *Tenthredo* 46
cinxius, Crossocerus, *Crabro* 228
circassica, Sciaapteryx 89
circe, Chrysis 134
circensis, Pterochilus
 bembeciformis var. 184
circularis, Athalia, *Tenthredo*
 circularis 44
circularis, Cerceris, *Philanthus* 253
circularis, Pristiphora 74
circulata, Crocisa 327
circumcincta, Megachile, *Apis* 307
circumcinctum, Anthidium
 quadriseriatum var. 300
circumscriptus, Nematus 59
cirrhostomus, Nematus 73
cirrhozonius, Halictus 281
cirsii, Sabulicola 287
ciscaspica, Andrena eversmanni ssp. 272
ciscaucasicus, Leptothorax
 tuberum 208
citreipes, *Tenthredo* 86
citreus, Nematus 63
citrine, Lasius bicornis var. 201
citrinus, Lasius 201
 Cladardis 46
Cladius 53
Cladius subg. 54
clara, *Tenthredella livida* 95
Claremontia 46
clarens, Kokujewia ectrapela 25
clarinicollis, Chrysis 134
clarinicollis, Chrysis ignita var. 134
clarior, Formica rufibarbis
 subpilosa var. 200
clarior, Formica sanguinea var. 200
clarior, Paravespa rex var. 188
claripennis, Ancistrocerus 179
claripennis, Andrena
 carbonaria var. 272
claripennis, Bethylus 125
claripennis, Coelioxys 305
claripennis, Dolerus germanicus 78
claripennis, Goniozus, *Bethylus* 125
claripennis, Hylaeus 261
claripennis, Hylotoma 25
clarissima, Formica 200
clarissima, Formica rufibarbis var. 200
claristernis, Diprion simile 33
claristernis, Pachynematus
 imperfectus 60
clariventris, Chrysis chosenensis f. 134
clarkella, Andrena, *Melitta* 263
clarkii, Cephalcia 102
clarum, Zarudnidium 129
clathratus, Hylaeus 261
Clavellaria 29
Clavellarius 29
Clavicera 310
clavicernum, Trypoxylon 244
clavicornis, Eucera 276
clavicornis, Melitturga, *Eucera* 276
clavicornis, Monosapta, *Apis* 145
clavimana, Didineis 219
clavipes, Andrena florea var. 272
clavipes, Gonatopus, *Gelis* 120
clavipes, Pomphilus 211
clavipes, Rhopalum, *Sphex* 236
claviventris, Hoplitis, *Osmia* 296
clematidis, Pseudodineura,
 Pelmatopus 75
clémenti, Priocnemis 164
cleopatra, Andrena ephippium var. 270
Cleptes 126
Cleptinae 126
clericalis, Arachnospila, *Pompilus* 170
clibrichella, Euura, *Nematus* 57
clipeata, Vespa 222
Clisodon subg. 323
clitellata, Euura, *Nematus* 57
clitellatus, *Nematus* 57
clivorum, Messor auct. 204
clivorum, Messor structor var. 204
cloëvorax, Crabro 229
clotho, Melecta 326
clusius, Halictus leucozonius 281
Clydostomus 89
clypealis, Allantus 96

- clypealis*, Cephus 109
clypealis, Odynerus 183
clypealis, Passaloecus 247
clypear, Eoanthidium, *Anthidium* 301
clypear, Lasioglossum, *Hylaeus* 282
clypearia, Sphex 234
clypearis, Andrena 264
clypearis, Crabro 229
clypearis, Hylaeus, *Prosopis* 261
clypeata, Andrena 264, 265
clypeata, Apis 234
clypeata, Ceratophorus,
 Pemphredon 247
clypeata, Eucera 320
clypeata, Euura, *Amauronematus* 58
clypeata, Formicapis 297
clypeata, Lestica, *Apis* 234
clypeata, Lyda 103
clypeata, Macrocerca 321
clypeatus, Crabro 226
clypeatus, Halictus 283
clypeatus, Panurgus 276
clypeatus, Philanthus 234
clypeatus, Temnothorax, *Myrmica* 208
Clypeocrabro 234
clypeopictus, Stenodynerus,
 Odynerus 190
Clytochrysus 231
Clytovorus 124
Cnemidandrena subg. 265
coactifrons, Machaerothrix 165
coactula, Pristiphora, *Nematus* 71
coangustata, Vespa 180
coarcata, Eucera 320
coarctata, Andrena 267
coarctata, Formica 209
coarctata, Ponera, *Formica* 209
coarctata, Vespa 181
coarctatum, Rhopalum, *Sphex* 235
coarctatus, Ammatomus, *Gorytes* 220
coarctatus, Eumenes, *Vespa* 181
coarctatus, Gorytes 220
cochlearietarsi, Halictus 277
cochlearipes, Apis 323
cochlearitarsis, Halictus, *Lucasius* 277
cockerelli, Tenthredo, *Tenthredella* 94
coecutiens, Apis 319
coecutiens, Epeoloides, *Apis* 319
coelebs, Sphecodes 289
coelostimontana, Eumenes 181
coelestimum, Hedydriidium 142
Coelioxys 304
Coelioxys subg. 305
Coelocrabro 227
Coeloeucus 247
coerulans, Mutilla 157
coerulea, Ceratina 310
coerulea, Hylotoma 23
coeruleipennis, Tenthredo 23
coeruleocephala, Dufourea 290
coeruleescens, Dolerus 80
coeruleescens, Tenthredo 23
coeruleus, Nematus 69
cogens, Cerceris 253
cognata, Andrena 273
cognata, Macrophyia 86
cognata, Myrmosa 153
cognata, Pteronidea 61
cognata, Stelis 303
cognata, Tenthredo 85
cognatus, Odynerus 184
cohaesus, Dolerus 78
coitana, Andrena, *Melitta* 271
colchica, Nylanderia, *Paratrechina* 202
colchica, Ponera coarctata
 coarctata 209
colchica, *Vespa norvegica saxonica*
 natio 194
colchicum, Lasioglossum littorale 284
colibri, Tenthredo 44
collectanea, Euura, *Nematus* 58
collare, Hedychrum 130
collaris, Allantus 83
collaris, Andrena 269
collaris, Colletes 257
collaris, Diodontus 247
collaris, Hartigia draconis 110
collaris, Lestica, *Crabro* 234
collaris, Monophadnooides 50
collaris, Nematus 70
collaris, Priophorus padi 54
collaris, Profenus 52
collaris, Sphecodes 289
collaris, Tachysphex 239
collaris, Tenthredo 80
collaris, Tenthredo maculata 92
collariventris, Odynerus 183
Colletes 257
Colletidae 257
colletiformis, Andrena 264
Colletinae 257
collina, Tapinoma 197
collinsonana, Melitta 273
collinus, Nematus 66
Colloliris 237
Colobopsis 199
colon, Tenthredo 94
colonialis, Andrena 265
colonoides, Tenthredella velox 96
Coloplitis 296
Coloptera 212
coloradensis, Pristiphora 71
colorata, Prosopis 262
coloratus, Dolerus pullulus 78
coloratus, Elampus 127
coloratus, Halictus 284
Colpa 147
Colpa subg. 147
Colpopyga 127
Colposelandria 75
colpostoma, Arachnospila, *Pompilus* 168
columbiae, Blepharipus 228
columna, Empria 39
coma, Sapyga 145
comari, Cladius 54
comata, Andrena nigriceps ssp. 265
comatum, Anthidium 300
combaella, Andrena 275
combinata, Andrena, *Apis* 273
combinata, Hoplomerus
 melanocephalus var. 187
combinatus, Sphecodes 290
comma, Tenthredo 86
commarginata, Myrmica 205
commixtus, Halictus 286
communis, Hylaeus 259
communis, Vespa 196
communista, Bothriomyrmex 197
communista, Bothriomyrmex
 meridionalis var. 197
commutabilis, Bombus lapponicus 330
compacta, Euura, *Amauronematus* 58
compactus, Mellinus 246
compar, Hylaeus 260
comparata, Andrena 272
comparata, Chrysis 134
comparata, Nomada 311
comparata, Nomada
 maculifrons var. 311
compedita, Solierella, *Silaon* 241
compeditus, *Silaon* 241
compressa, Pristiphora, *Nematus* 71
compressicornis, Cladius, *Tenthredo* 54
compressicornis, Pelopoeus 251
compressum, Chlorion, *Sphex* 211
compressus, Halictus, *Andrena*
 compressus 278
compressus, Janus, *Sirex* 110
compressus, Strongylogaster 82
compta, Nomada 311
comptum, Anthidium 301
comptus, Crabro 231
comptus, Pasites 319
comta, Andrena 268
comta, Chrysis 134
Conaspidia 84
Concavixyla 20
concessus, Monophadnooides 50
concinna, Amasis 32
concinna, Eucera 320
concinus, Anoplus, *Pompilus* 172
concinus, Entomericus 246
concinus, Entomosericus 246
concinus, Hylaeus 261
concinus, Oryttus, *Sphex*
 concinus 224
concinus, Sphex 224
concolor, Andrena atrorubricata var. 272
concolor, Andrena helvola var. 264
concolor, Chrysis fulgida var. 135
concolor, Miscophus 241
concolor, Pemphredon 249
concolor, Psenulus, *Psen* 252
concolor, Stizus 225
concolor, Tenthredo providus 98
concolor, Tenthredopsis thomsoni 99
concolora, Mutilla 154
concreta, Andrena tibialis ssp. 272
condei, Pristiphora 71

conductus, Nematus 62
confinis, Allantus 91
confinis, Andrena 273
confinis, Cryptocheilus 161
confinis, Ectemnius, *Crabro* 232
confinis, Hylaeus 259
confinis, Tetramorium 209
confluens, Halictus 278
conformis, Phyllotoma 46
confusa, Ampulicomorpha 122
confusa, Blennocampa 46
confusa, Eucera 320
confusa, Heriades 296
confusa, Hylotoma 23
confusa, Larra 237
confusa, Mutilla 157
confusa, Osmia 298
confusa, Pristiphora 71
confusor, Priocnemis 163
confusus, Bombus confusus 327
confusus, Crabro 230
confusus, Dolerus sanguinicollis 80
confusus, Hylaeus 261
confusus, Nematus 57
congener, Crossocerus, *Crabro* 227
congener, Nematus 71
congerens, Formica 199
Congodryinus 121
congruens, Andrena 273
congruus, Stigmus 247
congruus, Temnothorax,
Leptothorax 208
conica, Apis 304
conica, Apis 305
conica, Cerceris 253
coniceps, Pristiphora 74
conicus, Sphecius, *Stizus conicus* 225
coniensis, Tenthredo 90
conifer, Philoctetes, *Ellampus* 132
conigera, Cerceris 253
conjugata, Pristiphora, *Nematus* 71
conjugens, Dolichovespula
media var. 195
conjuncta, Andrena 272
conjungens, Andrena varians var. 264
conjungens, Halictus 278
conjungens, Nomada 311
conjungens, Tenthredopsis 98
connata, Pontania leucapsis 63
connatus, Cimbex, *Tenthredo*
connatus 29
connectens, Janus 110
connectens, Melitta 271
connexa, Andrena 273
connexa, Apis 316
connexus, Symmorphus, *Odynerus* 191
conoideus, Coelioxys, *Anthophora* 305
consanguinea, Chrysis
consanguinea 134
consanguineus, Gorytes 224
consimilis, Andrena 267, 269
consimilis, Cleptes 127
consimilis, Eucera 320

consimilis, Tachyagetes 174
consimilis, Tachysphex 239
consobrina, Andrena 272, 274
consobrina, Arachnospila, *Pompilus* 168
consobrina, Prosopis 260
consobrina, Sciaapteryx, *Tenthredo* 89
consobrinus, Bombus 328
consobrinus, Crabro 230
consobrinus, Monophadnoides 50
consobrinus, Nematus 61
consobrinus, Sphecodes 289
consocius, Tachysphex 238
consonus, Monophadnoides 50
conspersa, Coelioxys 304
conspersus, Coelioxys 304
conspersus, Monophadnoides 50
conspersus, Nematus 71
conspicua, Tenthredo 94
conspicuus, Monophadnoides 50
constans, Odynerus 179
constitutus, Monophadnoides 50
constricta, Lestica 234
constrictiventris, Pseudepipona 189
constrictus, Andrena 265
constrictus, Elampus, *Notozus* 127
constructor, Auplopus, *Agenia* 165
consueta, Dasylabris maura
nostras var. 153
consularis, Euchroeus purpuratus 144
Contemptevagetes 170
contemptus, Pompilus 171
contigua, Melitta 275
contigua, Thrinax 83
contiguus, Crabro 231
continentalis, Halictus 285
continentalis, Megachile maritima 308
continuus, Ectemnius, *Crabro* 232
continuus, Nematus 67
contractus, Nematus 75
contrario, Osmia adunca 297
controversus, Bombus 328
contusa, Andrena 267
contusa, Tenthredo, *Tenthredella* 94
convergatana, Tenthredo,
Tenthredella 91
convergens, Coelioxys 305
convergens, Rhogogaster 88
convergens, Sphecodes 288
convergens, Stigmus 250
conversa, Smicromyrme variabilis f. 155
convexa, Andrena 274
convexa, Apis 275
convexa, Xylocopa 310
convexitellaris, Macrophyta 85
convexuscula, Melitta 274
convexusculum, Lasioglossum,
Hylaeus 282
convexus, Hylaeus 284
convexus, Pamphilus 104
convexus, Tachysphex 238
convolvuli, Tenthredo 291
copetica, Heriades 296
copetica, Nomada 311

Copidoceros 25
Copoxyla subg. 309
copitis, Sphecodes 290
Coptoformica subg. 199
coquebertii, Tenthredopsis,
Tenthredo 98
cora, Macrophyta 85
coracina, Pemphredon 248
coracinus, Dolerus, *Tenthredo* 79
coracinus, Monophadnoides 50
coracinus, Smicromyrme 156
corallina, Arge 23
corallipes, Tenthredo 85
coranicus, Katamenes sichelii 185
corcyraea, Nomada 312
Cordandrena subg. 267
cordata, Athalia 44
cordata, Tenthredo 99
cordatoides, Athalia 44
cordialis, Andrena 267
cordigera, Tenthredo 38
cordivalvata, Priocnemis 163
coreana, Abia 28
coreana, Apareophora 45
coreana, Cimbex femorata 30
coreana, Tenthredo 96
coreanus, Dolerus 80
coreanus, Epeorus 317
coreanus, Orussus 117
coreanus, Pamphilus 104
coreanus, Phylloecus, *Hartigia* 110
coreensis, Cerceris 253
coreensis, Megalodontes 107
Corenius 247
Corenocrabro 227
coriacea, Arge 25
coriacea, Priocnemis 163
coriacea, Prosopis 262
coriacea, Tachytes 238
coriaceum, Hedychridium, *Hedychrum*
coriaceum 129
coriaceus, Nematus 60
coriaceus, Tachysphex 238
coriaceus, Tachysphex, *Tachytes* 238
coriarus, Hylaeus 284
corniculata, Mutilla 154
corniculus, Dolichurus, *Pompilus* 211
corniculus, *Pompilus* 211
cornifrons, Osmia, *Clalicodoma* 299
corniger, Belomicrus 242
corniger, Cladius 54
corniger, Passaloecus 248
cornigera, Apis 313
cornubiae, Athalia 44
Cornuchrysis 133
cornuta, Andrena 295
cornuta, Cerceris 253
cornuta, Eucera 321
cornuta, Euura, *Amauronematus* 58
cornuta, Osmia, *Megachile* 299
cornutum, Trypoxylon 244
cornutum, Trypoxylon frigidum 244
cornutus, Hylaeus 259

- cornutus*, Lithurgus auct. 295
cornutus, Lithurgus, *Andrena* 295
coronata, Blennocampa 47
coronata, Coelioxys 304
coronata, Ephutomma, *Eremomyrme* 155
coronata, Mutilla 155
coronata, Tarpa 107
coronata, Tenthredo 75
coronatum, Anthidium 301
coronatus, Aneugmenus, *Tenthredo* 76
coronatus, Eumenes, *Vespa* 181
coronatus, Oryssus 117
coronatus, Philanthus 255
coronatus, Philanthus, *Sphex* 255
coronatus, Sphex 255
Corporaalinus 50
corporica, Pseudoscolia 256
corpulentus, Lygaeonematus 71
corruetus, Eumenes coronatus 181
corsa, *Andrena nigroaenea* 269
corsica, *Andrena florea* ssp. 273
corsica, Bothriomyrmex
 meridionalis var. 197
corsicus, Bothriomyrmex 197
corsicus, Eumenes coarctatus 181
corticalis, Osmia 298
corticalis, Temnothorax, *Myrmica* 208
corticous, Pachynematus 66
corusca, Hybotoma 23
coruscans, Dolerus 80
corvina, *Andrena tibialis* ssp. 272
corvinum, Lasioglossum, *Halictus* 283
corvinus, Hylaeus 262
coryli, Platycampus 70
coryli, Rhogogaster 88
coryli, Tenthredo 95
Corymbas 84
Coryna 76
Corynia 22
Corynidinae 31
Corynis 31
Corynopus 235
corytus, Monophadnooides 50
Cosmiaporus subg. 166
costae, *Astata costae* 218
costae, Bothriomyrmex 197
costae, *Cryptoceroides*, *Salius* 161
costae, Mimesa 250
costae, Odynerus 185
costae, Phyllotoma 49
costae, Pontania 67
costae, Tachysphex, *Tachytes* 238
Costae, *Tachytes* 238
costalis, Allanthus 97
costalis, Monophadnooides 50
costalis, Sciaapteryx, *Tenthredo*
 costalis 89
costalis, Tenthredo 89
costaniana, Osmia 298
costata, Tenthredo 90
costatus, Monophadnooides 50
costiferellus, Halictus 284
costulatum, Lasioglossum, *Halictus* 280
cothurnatus, Dolerus 77
coxalis, Macrophyia, *Dolerus* 87
coxalis, Nomada 312
coxalis, Pterocheilus 187
coxalis, Tenthredo 39
Crabro 29
Crabro 225
Crabro subg. 226
crabro, *Vespa* 193
crabroneus, Mellinus, *Sphex* 246
Crabronidae 217
crabroniformis, Masaris 145
crabroniformis, Onychopterocheilus,
 Odynerus 187
crabroniformis, *Vespa* 193
Crabroninae 225
Crabronini 225
Craesus 69
craseundus, Temnothorax 208
Crasedopyga subg. 153
crassa, Hybotoma 23
crassa, Tenthredo 91
crassa, *Vespa* 195
crassepuncta, Holopyga 131
crassepunctatum, Lasioglossum,
 Halictus 283
crassicapitis, Priocnemis 163
crassicauda, Pachynematus 71
crassicornis, Cladius 54
crassicornis, Corynis, *Tenthredo* 32
crassicornis, Crocisa 327
crassicornis, Didineis 219
crassicornis, Ectemnius, *Crabro* 233
crassicornis, Eriocampa 46
crassicornis, Evagetes, *Pompilus*
 crassicornis 170
crassicornis, Hoplismenoides 224
crassicornis, Nematus 54, 71
crassicornis, Sphecodes 288
crassicornis, Stizoides, *Tiphia* 225
crassicornis, Symmorphus, *Vespa* 191
crassidens, Euura, *Amauronematus* 58
crassipes, Anthophora 325
crassipes, Dolerus 235, 243
crassipes, Euura, *Nematus* 58
crassipes, Mimesa 250
crassipunctatus, Eumenes
 subpomiformis 182
crassispina, Euura, *Nematus* 58
crassispina, Temnothorax
 nylanderii var. 208
crassispinus, Temnothorax,
 Leptocephalus 208
crassiventris, Nematus 74
crassula, Macrophyia, *Tenthredo* 85
crassuliformis, Macrophyia 87
crassulus, Nematus 61
crassum, Trichiosoma 30
crassus, Dolerus 80
crassus, Monophadnooides 50
crassus, Sphecodes 287
crataegi, Cimbex 31
crataegi, Hoplocampa, *Tenthredo* 67
crataegi, Nematus 71
crataegi, Priophorus 54
crataegi, Tenthredo 31
craterus, Halictus 281
Cratolarra 236
craverii, Hoplisoides, *Hoplitus*
 craverii 224
craverii, *Hoplitus* 224
craverii, Laurentia 83
crawfordi, Panurginus 276
Creightonella subg. 305
Crematogaster 204
crenata, Tenthredo, *Tenthredella* 94
crenulata, Heriades 296
creperus, Taxonus 43
cresoni, Pemphredon 249
cretanus, Sphecodes puncticeps 289
Cretembolemus 122
cretensis, Stenodynerus
 aequisculptus 189
cretensis, Strongylogaster 82
creticola, Halictus carinaeventris 279
creticola, Prosopis 261
cribraria, Sphecodes quatuorcineta 231
cribraria, Vespa 226
cribrarius, *Crabro*, *Vespa* 226
cribrata, Antepipona, *Lionotus* 180
cribrata, Cerceris 252
cribrata, Ceropales cribra 161
cribrosa, Sphecodes 291
crimeae, Mutilla 157
crimensis, Eumenes 181
crimensis, Eumenes dubius 181
crinipes, Anthophora 323
crinita, Eucera 320
Crioscolia subg. 147
crippae, Perineura 84
crispus, Amauronematus 67
cristata, Bembix 221
cristata, Megachile 307
cristatus, *Crabro* 232
cristatus, Sphecodes 287
Cristiger 33
cristula, Lasioglossum, *Halictus* 280
croatica, *Andrena* 267
croatica, Macrophyia erythropus 86
croaticus, Dolerus 80
croaticus, Sphecodes 287
crocea, Hemichroa, *Tenthredo* 67
crocea, Ponera japonica var. 209
crocea, Tenthredo 63
croceicornis, Euponera 209
croceipes, Anthophora 324
croceipes, Blennocampa 53
croceipes, Gorytes 221
croceiventris, Andrena 274
croceiventris, Tenthredo 51
croceus, Pamphilus 104
crocipennis, Tenthredo 25
Crocisa 326
Crossocerus 227
Crossocerus subg. 227
cruciata, Tenthredo 99

cruciatus, *Phylloecus* 111
cruciger, *Epeolus*, *Nomada* 317
crudelis, *Ammoplanus* 246
crudelis, *Andrena combinata* ssp. 273
cruenta, *Chrysis* 135
cruenta, *Dioxyss* 303
cruenta, *Nomada* 312
cruentatus, *Pachycephus*, *Cephus*
 cruentatus 110
cruentus, *Sphex* 223
Cryptandrena subg. 267
cryptarum, *Bombus*, *Apis* 327
Cryptocampus 55
Cryptocheilus 161
Cryptocheilus subg. 161
Cryptochilus 161
Cryptopone 209
Cryptus 22
csikiana, *Anthrena* 268
csikiana, *Chrysis fouqueti* 135
Ctenocerinae 160
Ctenonomia subg. 280
ctenophorus, *Tachysphex* 238
Ctenoplecta 319
Ctenoplectrini 319
Ctenopriocnemis 162
Ctenosmia 296
Ctenotilla 155
Ctenotillini 154
cubiceps, *Crabro* 232
Cubitalia 319
Cubitalia subg. 319
Cubitella 81
cucullata, *Andrena humilis* ssp. 265
cucullata, *Tenthredella* 94
cucurbitina, *Ceratina*, *Apis* 310
Cullumanobombus subg. 328
culmorum, *Chelostoma* 295
cultrarius, *Cephus* 109
cultratus, *Cephus* 109
Cuneala 89
cuneiformis, *Odynerus* 187
cunicularia, *Cerceris* 254
cunicularia, *Formica* 200
cunicularius, *Colletes*, *Apis* 257
cunyi, *Tenthredo* 94
cupes, *Cerceris* 253
Cuphopterus 227
cuparia, *Andrena* 298
cupratum, *Hedychridium* 142
cupratus, *Ellampus auratus* 132
cupratus, *Pseudomalus*, *Ellampus* 132
cuprea, *Apis* 298
cuprea, *Chrysura*, *Chrysis* 139
cupreum, *Hedychridium*,
 Hedychrum 129
cupreus, *Odynerus pubescens* var. 183
curinthicus, *Leptopus* 53
curiosus, *Monophadnoides* 50
curonica, *Andrena paucisquama* ssp. 268
curruca, *Spilomena*, *Celia* 249
cursor, *Sphex* 180
curticornis, *Nematus* 63

curtispina, *Nematus* 56
curtispinis, *Pteronus* 56
curtivalvis, *Andrena* 265
curtiventris, *Omalus* 132
curtulus, *Nysson* 220
curtus, *Fenusa* 48
curvatum, *Sceliphron*, *Pelopaeus* 214
curvicornis, *Eucera* 291
curvicornis, *Systropha*, *Eucera* 291
curvicrus, *Megachile* 308
curvipes, *Anthidium* 303
curvipes, *Tenthredo* 85
curvisscapa, *Prosopis* 259
curviscutum, *Crocisa* 327
curvitarsis, *Eucera* 320
curvithorax, *Myrmica* 206
curvungula, *Andrena* 268
cussariensis, *Andrena* 268
cyanea, *Ceratina*, *Apis* 310
cyanea, *Chrysis japonica* var. 136
cyanea, *Lyda* 101
cyanea, *Mutilla* 157
cyanea, *Sphex* 141
cyanea, *Trichrysis*, *Sphex* 141
cyaneata, *Chrysis* 134
cyanella, *Hylotoma* 23
cyanescens, *Osmia* 298
cyanescens, *Xylocopa* 309
cyaneum, *Hedychrum* 130
cyaneus, *Sphex* 214
cyanocrocea, *Arge*, *Tenthredo* 23
cyanura, *Chrysis* 140
cyanurum, *Stilbum*, *Chrysis* 141
Cyclostoma 192
cylindrica, *Chrysis* 134
cylindrica, *Meria* 149
cylindrica, *Perineura* 100
cylindrica, *Tenthredo* 85
cylindrica, *Tenthredo*, *Tenthredina* 96
cylindricornis, *Schizocera* 26
cylindricum, *Polochrum* 146
cylindricus, *Hylaeus* 282
cylindricus, *Nematus* 61
cylindricus, *Phylloecus* 111
Cymura 130
cynips, *Eura* 65
cynosbati, *Janus*, *Tenthredo* 110
cynthiae, *Philoctetes* 132
Cyperolerus 77
Cyperolerus subg. 77
cyphognatha, *Nomada* 316
Cyphona 26
cyphonota, *Priocnemis* 162
Cyphopyga 305
cypria, *Strongylaster lineata* 82
cypriaca, *Anthophora acervorum* 323
cypriaca, *Melecta* 326
cypriacum, *Anthidium florentinum* 300
cypriacus, *Miscophus* 241
cypriacus, *Odynerus*
 melanocephalus 187
cyprica, *Andrena* 270
cypicus, *Eumenes mediterraneus* 182
cypriina, *Andrena ocreata* ssp. 274
cyprius, *Tachysphex panzeri* 239
cyrenaicus, *Euodynerus notatus* 183
cyrenaicus, *Tachysphex* 239
cyrnea, *Pontania* 56
Cyrteuchridium 128
Cyrtogonatopoides 120
Cyrtogonatopus 120
Cystomutilla 153
cythica, *Pemphredon* 249
Cytisodaster 88
czekelii, *Halictus* 284
czerskiae, *Lapidariobombus sicheli* 329
czerskianus, *Bombus terrestris* 328
czerskiellus, *Agrobombus*
 anachoreta 331
czerskii, *Bombus* 328
czerskii, *Myrmosa* 153
dacica, *Cerceris* 253
dacica, *Megachile* 306
daedala, *Pontania* 63
daetekensis, *Tenthredo* 93
dagestanica, *Andrena* 270
dagestanica, *Chrysis indigotea* var. 136
dagestanicum, *Tapinoma*
 emeryanum 198
daghestanica, *Physetopoda*, *Mutilla* 155
daghestanicus, *Bombus* 332
daghestanicus, *Cephus*,
 Pseudocephus 109
dagmarae, *Myrmica moravica* var. 205
dahlbomi, *Diodontus* 247
dahlbomi, *Heptamelus*, *Caenoneura* 36
dahlbomi, *Mimumesa*, *Mimesa* 251
Dahlbomia 251
dahlbomii, *Dolichurus* 211
dahli, *Xyela* 20
dahlii, *Pleroneura*, *Xyela* 20
dahlii, *Tenthredo* 90
dailatorreana, *Nomada* 311
daisetsuzanus, *Crabro pubescens* 228
daisetsuzanus, *Evgates* 171
daisetusana, *Gilpinia* 33
daisi, *Sphecodes* 290
dalanicus, *Parodontodynerus*
 ephippium 188
dalecarlicus, *Amauronematus* 56
dalii, *Nomada* 316
dallatorreana, *Ceratina* 310
Dallatorreanus, *Crabro kollaris* 233
dallatorrei, *Andrena* 273
dalmatica, *Bembix* 221
dalmatica, *Eucera* 320
dalmatica, *Mutilla* 157
dalmaticum, *Lasioglossum*
 aegyptiellum 280
dalmatina, *Macrophya* 85
dalmatina, *Smicromyrme*
 curvitentris var. 156
damascena, *Hylotoma syriaca* 23
damascenum, *Lasioglossum*,
 Halictus 283

- dantici, Euodynerus, *Vespa* 183
 dantici, *Vespa* 183
danuvia, *Andrena* 268
danuvia, *Nomada goodeniana* 313
daphne, *Chrysis* 134
daphnis, *Chrysis daphnis* 135
daphnoideas, *Pontania acutifoliae* 55
dasiphorae, *Pristiphora*, *Pristicampus* 71
Dasyagetes 173
Dasylabrinae 153
Dasylabri 153
Dasylabris 153
Dasylabris subg. 153
Dasymegilla subg. 323
Dasypoda 293
Dasypodainae 293
Dasypodaini 293
dathei, *Dolerus* 79
dauriana, *Chrysis* 135
daurica, *Chrysis* 136
daurica, *Hoplitis*, *Osmia* 297
dauriensis, *Cleptes* 126
davei, *Tianmuthredo* 100
davidi, *Ctenoplectra* 319
davidsoni, *Crabro* 227
daviesanus, *Colletes* 257
davisi, *Plenoculus* 241
daw, *Odynerus* 187
dawsoni, *Pristiphora* 71
dealbata, *Tenthredo* 91
dealbata, *Tenthredo* 91
deandrenalis, *Andrena*
 hattorfiana ab. 265
deaurata, *Tenthredella* 91
debile, *Stenamma*, *Myrmica* 207
debilis, *Ammophila* 212
debilis, *Argo* 24
debilis, *Tachysphex* 240
debilitatus, *Syrmorphus*, *Odynerus* 191
Decanematus 55
Decatria 49
decemguttata, *Sapyga* 146
decemguttata, *Sapygina*, *Sapyga* 146
decemguttatus, *Pompilus* 161
decemmaculatus, *Nysson* 220
decemmaculatus, *Philanthus* 255
decempunctata, *Coelioxys* 305
decens, *Syrmorphus*, *Odynerus* 191
decens, *Tenthredo* 94
deceptoria, *Andrena* 267
deceptoria, *Megachile* 306
deceptorium, *Trypoxylon* 244
decimmaculata, *Vespa* 242, 243
decipiens, *Allantus* 96
decipiens, *Allantus carpini* 39
decipiens, *Andrena* 267
decipiens, *Halictus* 280
decipiens, *Hylaeus* 262
decipiens, *Macrophyia albicincta* 84
decipiens, *Nematus* 62
decipiens, *Pristiphora*,
Lygaeonematus 71
decipiens, *Sapyga* 145
decipiens, *Trichiosoma tibiale* 31
declinatus, *Nematus* 59
decolor, *Entodecta* 50
decolorans, *Ceratina* 310
decolorata, *Selandria* 76
decora, *Chrysis splendidula* 143
decora, *Chrysis* 135
decora, *Dasypoda* 293
decorata, *Andrena* 272
decorata, *Athalia* 44
decorata, *Nomada lineola* var. 313
decorata, *Scolia* auct. 148
decoratum, *Anthidium* 300
decorus, *Tachysphex* 239
decorus, *Thyreus*, *Crocisa* 326
dediticius, *Hemitaxonius* 39
dedzli, *Spilomena* 249
deficiens, *Nematus* 58
deflenda, *Antepipona*, *Ancistrocerus* 180
deforme, *Sceliphron*, *Pelopaeus* 214
degeeri, *Tenthredo* 55
degener, *Apis* 267
degener, *Tenthredo* 117
deiphobus, *Halictus* 281
deirambo, *Evagetes* 171
dejeani, *Scolia* 148
dejeanii, *Dufourea* 290
dejectus, *Allantus* 84
delagrangi, *Allantus flaveola* 90
delagrangi, *Clavellaria amerinae* 30
delicatula, *Pontania* 56
delicatula, *Tenthredo* 88
delicatulus, *Stromboceros*, *Tenthredo* 82
delicatulus, *Tenthredo* 82
delicatus, *Halictus* 285
dellei, *Asiemphytus*, *Macremphytus* 40
dellei, *Priophorus* 54
delmasi, *Halictus* 286
delphinalis, *Allodynerus*, *Odynerus* 178
delphinalis, *Halictus gavarnicus* 279
Delphinaloides 178
Delta 180
delumbis, *Taxonus* 43
deminuta, *Pontania* 56
Dendrolasius subg. 201
densepiosellus, *Ancistrocerus* 179
densopilosus, *Tetramorium* 209
dentata, *Andrena* 267
dentata, *Euura*, *Pachynematus* 58
dentata, *Teralonia* 321
dentata, *Tetraloniella*, *Eucera* 321
Dentathalia 43
dentatus, *Cemonus* 248
denticolle, *Lasioglossum*, *Halictus* 280
denticornis, *Andrena* 269
denticornis, *Crabro* 229
denticoxa, *Crossocerus*, *Crabro* 229
denticrus, *Anthophora* 325
denticrus, *Crossocerus* 229
denticulata, *Andrena*, *Melitta* 265
denticulata, *Coelioxys* 305
denticulata, *Apis* 276
denticulatus, *Crabro* 231
denticulatus, *Messor* 204
denticulatus, *Messor barbarus*
 meridionalis var. 204
dentifer, *Entomognathus*, *Crabro* 234
dentifer, *Odynerus* 187
Dentigera subg. 259
dentina, *Tenthredella* 94
dentipes, *Crabro* 226
dentipes, *Nomada* 312
dentipes, *Pterocheilus* 186
dentiventris, *Andrena* 272
dentiventris, *Dufourea*, *Halictoides* 290
dentiventris, *Osmia* 296
dentsukanus, *Crossocerus* 228
denudata, *Pristiphora pallidiventris* 73
denudatus, *Nematus* 71
deplanata, *Myrmica* 205
deplanata, *Myrmica lobicornis* var. 205
depressa, *Bethylus* 123
depressa, *Pristocera*, *Bethylus* 123
depressa, *Tenthredo* 105
depressata, *Strongylogastroidea* 39
depressus, *Nematus* 74
depressus, *Odynerus* 187
derasus, *Hylaeus* 281
derbentina, *Andrena* 275
Dermasothes 149
dernaensis, *Eumenes pomiformis* 181
derufata, *Pseudepipona*
 herrichii var. 188
derzavini, *Dolerus* 79
derzhavini, *Hortobombus*
 consobrinus 328
desantisi, *Nematus* 62
desbrochersi, *Strongylogaster* 82
deserta, *Tenthredo* 78
deserti, *Miscophus* 241
deserticola, *Ammophila* 212
deserticola, *Anthophora* 324
deserticola, *Diplorhoptrum* auct. 207
deserticola, *Solenopsis*,
Diplorhoptrum 207
Desertixyela 20
Desertodynerus 180
desertorum, *Chrysura*, *Chrysis* 139
desertorum, *Eremiasphecius*
Shestakovia 246
desertorum, *Megachile* 305
desertorum, *Philanthus* 255
desertorum, *Salius* 161
desertorum, *Sphex* 215
desertus, *Sphecodes* 289
despecta, *Dineura* 75
destillatorium, *Sceliphron*, *Pepsis* 214
destricta, *Pontania* 58
destructor, *Anoplonyx* 53
detonsus, *Eumenes coarctatus* 181
detorta, *Andrena braunsiana* ssp. 271
Deuteragenia 164
Deuterageniini 164
Deuterepipona subg. 188
Deuterocimbex 29
deuteronymus, *Bombus* 331

deutziae, *Macremphytus* 40
 devia, *Tenthredo, Allantus* 96
deviridatus, *Halictus tumulorum* 279
dexter, *Halictus zonulus* 281
diacantha, *Chrysis diacantha* 135
diacantha, *Melecta* 326
diacantha, *Nomada* 312
diacanthus, *Crabro* 229
diadema, *Anthidium* 300
diadema, *Vespa* 193
diakonovi, *Polistes* 192
dialeucus, *Allantus* 92
dianae, *Hortobombus consobrinus* 328
diaphanus, *Nematus* 57
dibowskii, *Macrophyta* 87
Dichodynerus 180
dichroa, *Apis* 294
dichroa, *Chrysura, Chrysis* 139
dichropsis, *Chrysura* 143
Dichrysis 133
Dicondylus 120
Dicrodolerus 77
Dicrodolerus subg. 77
Dicrogenium 123
Dicrostema 51
Dictenulus 132
Dictyolyda 102
Dicyrtomellus 170
Didineis 219
Didocha 22
Didonia subg. 267
didymus, *Allantus, Tenthredo*
 didymus 38
Dienopus 223
Diepyris 123
diervillae, *Nomada* 316
differens, *Odynerus* 183
differens, *Spilomena* 249
difficilis, *Eucera* 320
difficilis, *Nomada* 315
difficilis, *Odynerus* 190
difficilis, *Osmia* 298
diffinis, *Ferreola, Anoplius* 166
difformis, *Hylaeus, Prosopis* 261
difformis, *Lasius* 292
difformis, *Tenthredo* 54
digitalis, *Melitta* 273
diglypha, *Coelioxys* 304
Digonatopus 120
digoniensis, *Allantus amaenus* 97
Digoniozus 125
dilatatus, *Crabro* 228
dilatatus, *Hylaeus, Melitta* 260
dilecta, *Andrena* 270
dilobum, *Anthidium* 302
diluta, *Myrmica* 206
diluta, *Nomada* 315
diluta, *Nomada lineola* var. 313
diluta, *Tenthredopsis dorsalis* 99
dilutus, *Nematus* 59
dimidiata, *Andrena* 265
dimidiata, *Anthophora*
 acervorum var. 323

dimidiata, *Arge, Hylotoma* 23
dimidiata, *Athalia* 44
dimidiata, *Hoplocampa flava* 68
dimidiata, *Hylotoma* 24, 25
dimidiata, *Melitta* 294
dimidiata, *Osmia* 298
dimidiata, *Tenthredo* 99
dimidiaticlypeus, *Eumenes* 182
dimidiaventralis, *Megachile* 305
dimidiatus, *Crossocerus, Crabro*
 dimidiatus 230
dimidiatus, *Dolerus* 77
dimidiatus, *Katamenes, Eumenes* 185
dimidiatus, *Nematus* 64
dimidiatus, *Nysson* 220
dimidiatus, *Pamphilus* 103
dimidiatus, *Sphecodes* 288
Dimorpha 145, 218
Dinax 40
Dinetinae 245
Dinetomorpha 236
Dinetus 245
Dineura 55
Dineuridea 70
Dineurus 248
Dinevra 55
dinurum, *Anthidium* 301
diocrioides, *Tenthredo, Allantus*
 maculiger 91
Diodontus 247
Dioxoides 303
Dioxyini 303
Dioxys 303
Diphadnus 70
Diphaglossinae 257
Diphlebus 248
Diphysis 303
Diploplectron 218
Diplorhoptrum 207
Diplorrhos 132
Diprion 33
Diprionidae 33
Diprioninae 33
Dipteromorpha 89
discipennis, *Nematus* 71
disclosa, *Chrysidea, Chrysis*
 disclosa 133
disclosa, *Chrysis pumila* 133
Discoelius 181
discoidalis, *Mutilla* 157
discoidalis, *Nematus* 71
Discolia subg. 147
discolor, *Allantus* 90
discolor, *Cryptochilelus, Sphex* 162
discolor, *Pristiphora* 73
discolor, *Tachytes* 239
discoloratus, *Temnothorax*,
Leptothorax 208
disconotatus, *Euodynerus*,
Odynerus 183
discophora, *Tenthredo* 95
discors, *Priophorus* 54
discosignatus, *Odynerus*
 egregius var. 191
discrepans, *Cladius* 54
discrepans, *Halictus* 281
discrepans, *Prosopis* 260
discrepans, *Tenthredopsis* 98
discreta, *Macrophyta* 87
discretus, *Hylaeus* 259
discum, *Lasioglossum, Halictus* 280
discus, *Hylotoma* 24
discussa, *Meria* 149
disjuncta, *Osmia* 298
dispar, *Euura, Nematus* 58
dispar, *Monophadnus* 50
dispar, *Pompilus* 172
dispar, *Psen* 250
dispar, *Tenthredo* 90, 93
disparilis, *Arge, Hylotoma* 23
disparilis, *Camptopoeum friesei* var. 276
displacentia, *Myrmica* 205
dissecta, *Bembix* 222
dissectum, *Anthidium* 300
dissectus, *Mellinus* 220, 222
dissectus, *Oxybelus dissectus* 242
dissidens, *Andrena* 264
dissimilis, *Emphytus* 38
dissimilis, *Monophadnus* 45
dissimulans, *Tenthredo* 92
distans, *Hylaeus* 260
distigmus, *Goniozus* 125
distincta, *Andrena* 267, 274
distincta, *Chrysis distincta* 135
distincta, *Ferreola* 173
distincta, *Megachile* 307
distincta, *Mutilla* 154
distinctum, *Chelostoma, Eriades* 296
distinctus, *Cryptocampus* 70
distinctus, *Gonatopus* 120
distinctus, *Psen* 252
distinctus, *Psithyrus* 329
distineta, *Coelioxys* 305
distinguenda, *Chrysis* 143
distinguenda, *Andrena* 267
distinguenda, *Euura*,
Amauro nematus 58
distinguenda, *Hylotoma* 25
distinguenda, *Lasius umbrata* 201
distinguenda, *Nomada* 312
distinguendus, *Bombus* 331
distinguendus, *Crossocerus, Crabro* 229
distinguendus, *Gonatopus* 120
distinguendus, *Hoplisoides, Gorytes*
 gazagnairei 224
distinguendus, *Lasius, Formicina* 201
distinguendus, *Nysson* 220
distinguendus, *Pompilus* 172
distinguendus, *Priophorus* 55
distinguendus, *Rhophitoides* 291
distinguendus, *Sphecodes* 290
distinguendus, *Stizus* 225
divergens, *Andrena* 271
divergens, *Coelioxys* 304
divergens, *Halictus* 280

divergens, *Myrmica* 205
divergens, *Myrmica bergi* var. 205
diveri, *Strongylognathus* 207
diverseducta, *Macrophyia rufipes* 86
diversicolor, *Hylotoma rosae* 24
diversicornis, *Pseudoscolia*,
 Philanthus 256
diversipennis, *Tenthredo*, *Allantus*
 bifasciata 90
diversipes, *Bembix* 221
diversipes, *Cephus* 108
diversipes, *Crossocerus* 228
diversipes, *Janus luteipes* 110
diversipes, *Macrophyia*, *Tenthredo* 85
diversipes, *Nomia* 291
diversipes, *Nomiapis*, *Nomia* 292
diversus, *Bombus* 328
dives, *Ammophila* 213
dives, *Ectemnius*, *Solenius* 232
dives, *Eremochares* 213
dives, *Solenius* 232
divisa, *Andrena* 272
divisa, *Blennocampa* 46
divisa, *Crocisa* 327
divisa, *Melitta* 287
divisus, *Crabro* 232
divisus, *Sphecodes* 287
divitoides, *Crabro* 232
dlusskyi, *Formica* 199
dlusskyi, *Temnothorax* 208
dmitrijewi, *Halictus* 280
dmitrijewii, *Halictus* 280
dobrowlanensis, *Andrena* 267
dochmocera, *Pristiphora*, *Nematus* 71
Dociliris 237
docilus, *Dolerus* 79
Doderia 50
doderleini, *Phoenusa* 48
doderoi, *Andrena* 263
Doeringiella 317
dohrni, *Megachile* 295
dohrnii, *Fenusia*, *Kaliostrophinga* 48
dolens, *Tenthredo* 85
Dolerus 77
Dolerus subg. 77
dolichocephalus, *Hylaeus* 260
Dolichochelys 120
Dolichocrabro 227
Dolichoderinae 197
Dolichoderus 197
Dolichovespula 194
dolichura, *Euura*, *Nematus* 58
Dolichurus 211
dolorosa, *Andrena* 272
dolosa, *Tenthredo* 46
Dolus 123
domestica, *Sphecodes* 179
domesticus, *Sclerodermus* 124
dominiquei, *Corynis* 32
dominula, *Formica* 200
dominula, *Polistes*, *Vespa* 193
donatus, *Halictus cristula* 280
dongarica, *Halictus limbellus* 284

Donisthorpina 120
Dorhettenyx 90
Dorhettenyx subg. 90
doriae, *Eremochares* 213
dorni, *Ceropales bicoloripes* 160
dorni, *Hylaeus* 260
dorpatica, *Eriocampa* 42
dorsalis, *Abia* 29
dorsalis, *Andrena* 270
dorsalis, *Cerceris* 253
dorsalis, *Cerceris* 254
dorsalis, *Dineura* 68
dorsalis, *Dineura virididorsata* 55
dorsalis, *Megachile* 306
dorsalis, *Meria*, *Bethylus* 149
dorsalis, *Nematus* 61
dorsalis, *Schizocera scutellaris* 26
dorsalis, *Selandria* 81
dorsalis, *Tenthredo* 100
dorsata, *Andrena*, *Melitta* 273
dorsata, *Tenthredo* 33
dorsata, *Tenthredopsis raddatzi* 99
dorsatula, *Andrena* 270
dorsigera, *Tenthredo* 84
dorsivittatus, *Nematus* 62
dorsivittatus, *Tenthredopsis* 99
Dositheus 77
dossuarius, *Pteronus* 59
Dosytheus 77
dournovii, *Spinolia*, *Chrysis* 141
dovnari, *Sciaapteryx* 89
draconis, *Hartigia* 110
dragana, *Andrena* 267
drewseni, *Cladius* 55
dromedarius, *Sirex* 116
Druida 49
drupacearum, *Lyda* 103
dryas, *Tenthredo* 89
Dryinidae 118
Dryininae 119
Dryinopsis subg. 149
Dryinus 119
Dryphus 235
Dryudella 218
dschingis, *Arachnospila* 169
dialis, *Perineura* 43
dubia, *Anthophora* 324
dubia, *Bembix* 221
dubia, *Hylotoma* 24
dubia, *Larra* 241
dubia, *Nomada marshamella* var. 314
dubia, *Polistes* 192
dubia, *Sirex* 112
dubia, *Tenthredo* 47, 78, 95
dubia, *Tenthredopsis* 100
dubiosa, *Abia*, *Orientobia* 28
dubiosa, *Dufourea*, *Rophites graeca* 291
dubiosus, *Clytochrysus* 233
dubious, *Sphecodes gibbus*
 ephippium 289
dubitabilis, *Halictus* 283
dubitata, *Andrena* 273
dubius, *Aporus* 170

dubius, *Emphytus* 45
dubius, *Eumenes* 181
dubius, *Evagetes*, *Aporus* 171
dubius, *Gastrosericus maracandicus* 236
dubius, *Nysson* 220
dubius, *Odynerus* 188
dubius, *Oxybelus* 242
dubius, *Panurginus* 276
dubius, *Tachytes* 238
dubowskii, *Halictus* 283
ducalis, *Vespa* 194
duckei, *Lasioglossum*, *Halictus*
 duckei 283
dudai, *Hoplocampa minuta* 68
dudichi, *Tachyagetes* 174
Dufourea 290
Dufourea subg. 290
dufouri, *Macrocera* 322
dufouri, *Megachile* 305
dufouri, *Nysson* 219
dufouri, *Oxybelus* 242
dufouri, *Psen* 252
dufourianum, *Nysson* 219
dufourianus, *Odynerus* 190
dufourii, *Anthidium* 303
dufourii, *Anthophora* 323
dufourii, *Discoelius* 181
dufourii, *Nysson* 219
dufourii, *Odynerus* 187
dufourii, *Phileremus* 318
dumetorum, *Apis* 323
dumetorum, *Sphecodes* 180
dumetorum, *Tenthredo* 86
dumoucheli, *Bombus auct.* 331
duodecimguttatus, *Crabro* 231
duodecimmaculata, *Melecta*,
 Nomada 326
duodecimpunctata, *Macrophyia*, *Ten-*
 thredo *duodecimpunctata* 85
duplex, *Euura*, *Pristiphora* 58
duplex, *Pristiphora* 58
duplex, *Tenthredo* 87
duplicata, *Tenthredo* 94
duplicolor, *Dasylabris* 154
duplipunctatus, *Chrysanthampus* 127
duplipseudatus, *Sphecodes* 289
duskei, *Odynerus* 186
dusmeti, *Andrena limbata* ssp. 273
dusmetiolus, *Ancistrocerus*,
 Odynerus 179
dux, *Sirex*, *Paururus cyaneus* 112
dyari, *Hemicrooa* 67
dybovskij, *Nomada* 312
dybowskii, *Lasioglossum*, *Halictus* 283
dybowskii, *Megachile* 307
dybowskii, *Vespa* 194
dyozankeanus, *Sphecodes* 287
Dyscolocrabro 225
Dyscolocrabro 226
dzhingis, *Dienoplus* 223
dzieduszyckii, *Nomada* 316
dziuroo, *Diodontus* 247
dzynnanica, *Andrena* 272

Ebba 89
ebenina, *Podalonia*, *Ammophila* 213
ebeninus, *Hylaeus* 259
ebetsuense, *Rhopalum* 236
ebmeri, *Colletes* 257
ebmeri, *Hoplitis* 297
Ebolia 98
eborata, *Xiphydria* 115
eborina, *Tenthredo* 82
eburata, *Tenthredo* 94
eburnea, *Bembix* 221
eburneifrons, *Tenthredo* 94
eburneiguttatum, *Aglaostigma* 83
eburneofasciatus, *Oxybelus* 242
eburneofasciatus, *Oxybelus*
 _{dissectus} 242
eburneopictus, *Eumenes* 182
eburneus, *Phylloecus* 110
eburnus, *Euodynerus*
 _{quadrifasciatus} 183
ebusiana, *Bembix* 222
ecaphosus, *Halictus* 277
ecarinata, *Nomada* 312
echii, *Eucera* 320
Echinanthidium 300
echinatus, *Coelioxys* 304
echinulata, *Andrena* 263
Echthrodelpax 120
eckloni, *Onychopterocheilus*,
 _{Pterochilus} 187
ectemiformis, *Corenocrabro* 227
Ectemnius 231
Ectemnius subg. 231
ectrapela, *Kokujewia* 25
eczmiadzini, *Melecta* 326
Edapholigon 123
edentata, *Andrena* 265
edentata, *Euura*, *Pachynematus* 58
edenticulata, *Euura*, *Amauronematus* 58
eduardi, *Hypoponera*, *Ponera* 209
eduardi, *Tenthredo*, *Tenthredella* 92
effasciata, *Prosopis* 262
effossus, *Crabro* 231
egeria, *Pteronidea* 60
Eggysomma 167
eglantariae, *Tenthredo* 78
egregia, *Abia* 28
egregia, *Ceratina* 310
egregia, *Tenthredo reitteri* 96
egregius, *Cryptocheilus*, *Calicurgus* 161
egregius, *Euodynerus* *egregior* 191
egregius, *Nematus* 69
egregius, *Odynerus* 191
egregius, *Syneodynerus*, *Odynerus* 191
ehnbergi, *Andrena* 274
eichleri, *Aglaostigma langei* 83
eidmanni, *Myrmica* 205
Eidopompilini 173
einnersbergensis, *Nematus* 57
eisa, *Vespa analis* 193
eisukei, *Arachnospila*, *Pompilus* 170
eiteli, *Euura*, *Amauronematus* 58
Eitelius 55

elaeagnocola, *Euura*, *Pontania* 58
Elampini 127
Elampus 127
Elaphrosyron 174
elbrus, *Eura* 58
elbrusi, *Myrmica* 205
elbursa, *Prosopis annularis* 260
elderi, *Dolerus elderi* 78
elegans, *Andrena* 275
elegans, *Anthophilus* 255
elegans, *Cardiocondyla* 203
elegans, *Cardiocondyla auct.* 203
elegans, *Cardiocondyla schkaffi* 203
elegans, *Cerceris* 253
elegans, *Chrysis* 135
elegans, *Cleptes* 127
elegans, *Cryptocheilus*, *Pompilus* 161
elegans, *Emphytus* 38
elegans, *Epeolus cruciger* var. 317
elegans, *Gorytes* 222
elegans, *Harpactus*, *Arpactus*
 _{elegans} 223
elegans, *Lasioglossum*, *Halictus* 283
elegans, *Nematus* 62
elegans, *Odynerus* 190
elegans, *Odynerus* 191
elegans, *Oxybelus* *dissectus* 242
elegans, *Pachyprotasis* 87
elegans, *Thomsonia* 99
elegantissima, *Pseudapis*, *Nomia* 292
elegantula, *Tenthredo* 88
elegantulum, *Hedychridium* 129
elegantulus, *Oxybelus* 242
elelea, *Pteronidea* 62
eleonorae, *Andrena lepida* ssp. 273
Elinopsis 89
Elinora 89
Elinora subg. 90
elisabethae, *Psithyrus rupestris* 329
elisabethae, *Tenthredo* 92
Ellampus 127
elliptica, *Prosopis* 259
ellipticeps, *Lasioglossum*, *Halictus* 283
elongata, *Ammophila* 212
elongata, *Andrena* 265, 272
elongata, *Mutilla* 155
elongata, *Skorikovia*, *Mutilla* 155
elongatula, *Cladardis*, *Tenthredo* 46
elongatula, *Tenthredo* 46
elongatulus, *Crossocerus*, *Crabro*
 _{elongatulus} 229
elongatulus, *Lophyrus* 34
elongatum, *Anthidium* 301
elongatus, *Cephus* 108
elongatus, *Coelioxys* 305
elongatus, *Dolerus* 79
elongatus, *Evagetes*, *Anoplus* 171
elsei, *Coelioxys* 304
elysium, *Lasioglossum zonulum* 281
emanuelis, *Pseudoryssus* 117
emarginata, *Apis* 281
emarginata, *Blennocampa* 50
emarginata, *Cerceris* 254
emarginata, *Nomada* 312
emarginata, *Vespa* 179
emarginatus, *Biastes*, *Phileremus* 318
emarginatus, *Coelioxys* 304
emarginatus, *Crossocerus*, *Crabro* 229
emarginatus, *Lasius*, *Formica* 202
emarginatus, *Myrmus* 207
emarginatus, *Nematus* 59
emarginatus, *Sirex* 114
emarginatus, *Sphex* 214
Embolemidae 122
Embolemopsis 122
Embolemus 122
emelianoffi, *Halictus* 281
emendata, *Nomada* 311
emeryanum, *Tapinoma* 198
emesianus, *Halictus* 280
emichi, *Cephus* 110
Emphoropsis 325
emphytiformis, *Tenthredo* 97
Emphytina 39
Emphytus 37
Emphytus subg. 38
empoascae, *Aphelopus* 118
Empria 40
empriaeformis, *Strongylogaster*,
 _{Thrinax} 82
Encarsioneura 89
Endelomyia 46
Endemyolia 46
Endophytus 55
Endotethryx 90
Endotethryx subg. 91
enervis, *Nematus* 70
Engages 67
engelhardtii, *Anoplolyda* 105
Enisia 89
Enodia 215
enodis, *Argo*, *Tenthredo* 23
enodis, *Tenthredo* 22
ensicornis, *Nematus* 71
enslinella, *Andrena* 270
enslini, *Aprosthemra* 26
enslini, *Pemphredon*, *Dineurus* 248
enslini, *Pontania* 58
enslini, *Priocnemis* 163
enslini, *Pseudodineura*, *Pelmatopus* 75
enslini, *Spilomena* 249
enslini, *Tenthredella* 92
Entodecta 50
Entomognathus 234
Entomognathus subg. 234
Entomosericini 246
Entomosericus 246
eoaa, *Smicromyrme* 155
eoabnormis, *Arachnospila* 169
Eoanthidiellum 301
Eoanthidium 301
Eoanthidium subg. 301
Eoferreola 173
eomontanum, *Lasioglossum briseis* 282
Eopompilus 162
Eopsenulus 251

Eopsis 42
 eos, Lasioglossum 280
 eos, Myrmosa 153
 eos, Taimyrmosa, *Myrmosa* 153
Eosirex 113
Eotillinae 159
 eoum, Polochridium 145
 eous, Agenocimbex 29
 eous, Colletes 258
eous, Gorytes 223
Epagetes 173
Eparmatostethus 236
 epeoliformis, Nysson epeoliformis 220
Epeolini 317
 Epeoloides 319
 Epeolus 317
Ephemerocephus 108
 Ephemerocophus subg. 109
ephippiata, Apis 267
 ephippiata, Hoplocampa 67
 ephippiata, Osmia 298
 ephippiatus, Dolerus 80
 ephippiger, Anteon, *Gonatopus* 118
ephippiger, Nematus 73
Ephippionotus 110
ephippium, Andrena 270
ephippium, Anthophora 323
ephippium, Eumenes 188
ephippium, Eutomostethus, *Tenthredo*
ephippium 47
ephippium, Mutilla 155, 156
ephippium, Parodontodynerus,
Eumenes 188
ephippium, Sphecodes, *Sphex* 287
Ephutomma 155
Epibembex 221
Epicenematus 55
Epicnemius 192
Epicrossocerus 227
Epigonatopoides 120
epilosa, Euura, *Pteronidea* 58
epimelaena, Andrena 265
Epimyrra 207
epinotalis, Proformica 203
epiphron, Lasioglossum 283
Epipone 186
Episemus 125
Episuron 166
Episuronini 166
Epitactus 55
Epitaxonus 83
Epitritus 207
Epizuron 166
epovae, Megachile 308
Epyrinae 123
Epyrini 123
Epyris 123
equatia, *Pteronidea* 64
eques, *Tenthredo* 33
equestris, Trypoxyylon 250
equestris, Andrena 265
equestris, Bombus auct. 332
equestris, Chrysis 135

equestris, Mimesa, *Trypoxyylon* 250
equestris, *Tenthredo* 92
Equidolerus 77
Equidolerus subg. 78
equiseti, Ametastegia, *Tenthredo* 39
equiseti, *Tenthredo* 79
eradiatus, Cladius 55
erasa, *Tenthredo* 96
erasina, *Tenthredo* 96
Erasminus 69
erberi, Andrena 264
erberi, Cephus 108
erebia, *Tenthredo*, *Tenthredella* 97
erecta, Eurua, *Amauronematus* 58
Eremiaspheiinae 245
Eremiasphecium 245
eremita, Lophyrus 33
eremita, Passaloecus 248
eremocanus, Xenaporus 174
Eremochares 213
Eremodynerus subg. 184
Eremomyrme 155
Eremotilla subg. 156
ergenicus, Leptochilus, *Odynerus* 185
Eriades 296
ericetorum, Megachile 307
erichsoni, Empria 41
erichsonii, Pristiphora, *Nematus* 72
Erimyrme subg. 156
Erinys 180
Eriocampa 42
Eriocampoides 45
Eriocampopsis 42
eriophoroides, Psithyrus rupestris 329
eriophorus, Bombus 329
Eriops 276
eriphyle, Lasioglossum 283
ermak, Dinax, *Adamas* 40
ermak, Hedychrum rutilans 130
ermak, Sirex, *Paururus* 112
Ermilia 43
ermolajevi, Agrobombus adventor 331
ermolenkoi, Euura, *Nematus* 58
ermolenkoi, Kuriloagenia 164
errans, Nomada 312
errans, Stizus 221
errans, Stizus tridens 221
errata, Cerceris 253
erratica, Pachyprotasis erratica 87
erratica, *Tenthredo* 91
erraticum, Tapinoma 197
erraticum, Tapinoma, *Formica* 198
erschovi, Hedychriderium,
Hedychrum 129
erschowi, Anthophora 325
erubescens, Nomada lineola var. 313
erucarus, Odontopyris, *Parasierola* 125
erudita, *Tenthredo*, *Tenthredella* 94
eruditus, Halictus 278
eryihropus, Apis 303
eryngii, Cerceris 253
erythraea, Eoferreola, *Sphex* 173
erythraeus, Tachysphex 239

erythrocephala, Acantholyda,
Tenthredo 101
erythrocephala, Mutilla 153
erythrocephala, Nomada 312
erythrocephala, *Tenthredo* 101
erythrocerus, Epyris 123
erythrocema, Macrophyia 85
erythrocemis, Andrena 271
erythrocoele, *Tenthredopsis arrogans* 98
erythrogaster, Cephalcia, *Lyda* 102
erythrogaster, Nematus 74
erythrogaster, Tachytess 238
erythrogona, *Tenthredo* 79
erythromelas, Chrysis viridula var. 134
erythropareus, Nematus 59
erythropoda, Ceropales 160
erythropus, Dolerus aericeps 77
erythropus, Euura, *Amauronematus* 58
erythropus, Tachysphex, *Lyrops* 238
erythropus, *Tenthredo* 86, 87
erythropyga, Coelioxys 304
Erythrosmia subg. 298
erythrosoma, Nomada 314
erzurumensis, Megachile giraudi 306
esakii, Anoplius 172
esakii, Anteon 119
esakii, Macrophyia, *Pachyprotasis* 87
esakii, Nomada 311
esakii, Pasites 319
esakii, Pasites maculatus 319
esakii, Sphecodes 290
esakii, Vespa 194
esana, Nomada 312
esoense, Lasioglossum 280
esagnoli, Euodynerus 183
espanola, Andrena
truncatilabris ssp. 275
etizenensis, Crossocerus 228
etizenensis, Sphecodes 289
etnaensis, Megachile circumcincta 307
etorofensis, Phylloecus, *Hartigia* 110
Etrusca, Andrena 240
etrusca, Smicromyrme ausonia var. 156
etruscus, Tachytess, *Andrena*
etruscus 240
etzchmiadzini, Myrmilla, *Mutilla* 154
Eualastor 178
Euancistrocerus 178
Euandrena subg. 267
euboense, Lasioglossum, *Halictus* 283
Eucera 320
Eucera subg. 320
eucera, Cladius 54
Euceratina subg. 310
Eucerini 319
euchreiformis, Osmia 298
Euchridium 128
Euchroeus 140
Euchrum 128
Euclavellaria 30
eucnemidea, Eucera 321
eudokiae, Camponotus
herculeanus var. 198

- Eudryas* 54
Eueippona 186
eugeniae, Stizus 225
Euglages 318
Eugonatopus 120
Euleptochilus subg. 185
Eumegachile 305
Eumenes 181
Eumenes subg. 181
Eumenidion 181
Eumeninae 178
Eumetabolus 111
Eunomiooides 292
Euodynerus 183
Euodynerus subg. 183
Eoxybelus 242
Eupareophora 47
Euphilcremus subg. 318
epidus, Sphecodes longulus 288
Euplilis 235
eupodium, Nematus 59
Eupolistes 192
Eupomphiloides 168
Eupontania 55
eurasiaticus, Stigmus 250
Eurhadinoceraea 47
euris, Anthophora personata var. 323
europa, Megachile ericetorum 307
Eurogaster 89
Eurogaster subg. 91
euronotum, Lasioglossum zonulum 281
europaea, Ampulex 211
europaea, Cephalcia alashanica 102
europaea, Cimbex 29
europaea, Macropis 293
europaea, Mutilla 156
europaea, Syphingta 210
europaea, Tachytes 240
europaeum, Rygchium 189
europaeus, Agenioideus 173
europaeus, Stigmus 250
europea, Myrmica laevinodis var. 206
Eurrhynchium 189
Euryglossinae 257
Eurygnathilla subg. 154
eurygnathopsis, Halictus 278
eurygnathus, Halictus 278
euryscapus, Hylaeus 260
Euspongus 222
Eustenancistrocerus 184
Eustenancistrocerus subg. 184
Eutenthredopsis 98
Eutomostethus 47
Eutricharaea subg. 305
eutropis, Dolerus 80
Euura 55
euxinicum, Lasioglossum
euxinicum 280
Euxiphydria 115
Euzonia 222
Evagetes 170
Evgathes 170
evansi, Plastanoxus 124
evecta, Cerceris 255
evenensis, Euura, *Nematus* 58
eversmanni, Andrena 272
eversmanni, Cerceris 253
eversmanni, Dolerus eversmanni 79
eversmanni, Elampus, *Ellampus* 128
eversmanni, Megalodontes 107
eversmanni, Megalodontes, *Tarpa* 107
eversmanni, Melecta 326
eversmanni, Nematus 59
eversmanni, Nomada lineola var. 313
eversmanni, Pseudarge 25
eversmanni, Pseudoanthidium,
Anthidium 302
eversmanni, Stizus 225
eversmanni, Tenthredo 91
eversmanniana, Andrena 268
Evodia 257
Evura 55
Evylaeus subg. 282
ewaldi, Fenusia 48
exacutus, Lasius umbratus var. 201
Exagetus 173
exalbida, Tenthredo 86
exaltata, Priocnemis, *Sphex* 163
exaltata, *Sphex* 162
exannulata, Pachyprotasis antennata 87
exannulata, Tenthredopsis
fischbeini 100
exaratus, Pseneo, *Mimesa* 251
exarenaria, Dasylabris 154
exarmata, Blennocampa 47
exarmata, Eupareophora,
Blennocampa 47
excavata, Cerceris 253
excavata, Euura, *Pontania* 58
excavatus, Gonatopus 120
excellens, Amauronematus 63
excellens, Megachile 306
excellens, Sphecodes 289
excellens, Tenthredella 98
excellens, Tenthredo, *Allantus* 90
excelsa, Myrmica 205
excelsus, Tachysphex 238
excerptus, Tachysphex filicornis 238
excisa, Empria, *Poecilosoma* 41
excisa, Eucera 320
excisa, Euura, *Nematus* 58
excisa, Fenusia 49
excisa, Nomada 311
excisa, Perineura 99
excisa, Prosopis 261
excisa, Selandria 81
excisus, Agenioideus, *Pompilus* 167
excisus, Cephus, *Pseudocephalus* 109
excisus, Odynerus 179
excurrens, Hylaeus communis 260
exigua, Chrysis distincta 143
exiguum, Anteon, *Chelogynus* 119
exiguum, Trypoxylon clavicerum 244
exiguum, Trypoxylon exiguum 244
exiguus, Crossocerus, *Crabro* 229
exil, Bombus 331
exiliceps, Lasioglossum, *Halictus* 280
exilis, Cephus 110
exilis, Hylaeus 284
exilis, Macrophyia 87
exilis, Microdynerus, *Odynerus* 186
exilis, Odynerus 185
eximia, Macrophyia 85
eximia, Nannodynerus
chevrieranus var. 190
eximius, Philanthus 255
exoleta, Nematus fulvus 61
exoletus, Nematus 67
exorientis, Psen 251
exornata, Sapyga 145
exornata, *Tarpa* 107
expansa, Arge, *Hylotoma* 23
explanata, Tenthredo 91
exsecta, Formica 199
exsectus, Pseudohemiteanaxus 81
exspectata, Spilomena 249
extensis, Euura, *Nematus* 58
extranea, Chrysis rutilans 137
extraneus, Ammobates 318
extremus, Nematus 72
Eyagates 170
ezoana, Melitta 294
ezoensis, Alysson monticola 219
ezoensis, Andrena 274
ezonata, Habropoda 325
fabriciana, Nomada, *Apis* 312
fabriciella, Apis 312
fabricii, Cryptocheilus, *Pompilus* 161
fabricii, Hoplocampa 68
fabricii, Megalodontes, *Tarpa* 107
fabricii, Pempredon, *Cemonus* 248
fabricii, Selandria 81
facetus, Pamphilus 105
facialis, Cerobactrus 111
facialis, Euura, *Pteronidea* 58
facigera, Tenthredo 94
facilis, Nomada 312
faddei, Anthophora 324
fagi, Leptothorax muscorum var. 204
fagi, Tenthredo 94
fahraei, Euura, *Nematus* 58
fahringeri, Pseudochrysis 140
fairchildii, Echthrodelpach 120
fairmairei, Cataglyphis 199
falcata, Osmia 298
falcifera, Cerceris 253
fallaciosus, *Pompilus* 167
fallax, Agenia 165
fallax, Allodynerus delphinalis 178
fallax, Ammophila 212
fallax, Andrena 271
fallax, Camponotus, *Formica* 198
fallax, Chrysis subsinuata 138
fallax, Cleptes 127
fallax, Coelioxys 304
fallax, Cryptocheilus 161
fallax, Euura, *Nematus* 58
fallax, Gorytes 222

- fallax*, *Lasioglossum*, *Halictus*
fallax 280
fallax, *Lyda* 104
fallax, *Nematus* 58
fallax, *Nitela* 241
fallax, *Oxybelus* 243
fallax, *Priocnemis* 163
fallax, *Scolia* fallax 148
fallax, *Tenthredo* 67, 95
fallenii, *Lyda* 102
falsa, *Macrophyia militaris* 86
falsifica, *Andrena* 270
falsificissima, *Andrena* 270
famelica, *Aphaenogaster*,
Ischnomyrmex 203
famosa, *Funella* 48
fantoma, *Urocerus*, *Sirex* 113
fargeii, *Argogorytes*, *Gorytes* 221
Fargeii, *Crabro* 233
fasciale, *Anteon*, *Chelogynus* 119
fasciata, *Abia*, *Tenthredo* 28
fasciata, *Ampulex* 211
fasciata, *Andrena* 272, 275
fasciata, *Anthophora retusa* 325
fasciata, *Chrysis* 135
fasciata, *Chrysis* 141
fasciata, *Dioxys* 303
fasciata, *Discolia* 147
fasciata, *Euura*, *Amauronematus* 59
fasciata, *Hylotoma* 23
fasciata, *Larra* 225
fasciata, *Lyda* 103
fasciata, *Megachile* 307
fasciata, *Nomada ruficornis* var. 315
fasciata, *Tenthredo* 97
fasciata, *Xyphidria* 116
fasciatella, *Andrena* 267
fasciatellus, *Caliadurgus*, *Pompilus* 162
fasciatellus, *Halictus* 284
fasciatellus, *Pompilus* 162
fasciatipennis, *Lyda* 103
fasciatum, *Anthidium*
manicatum var. 300
fasciatum, *Pison Pseudonysson* 244
fasciatus, *Aporus* 168
fasciatus, *Asiemphytus*,
Macremphytus 40
fasciatus, *Ceratocolus* 233
fasciatus, *Crabro vagabundus* 227
fasciatus, *Dolerus* 38
fasciatus, *Epeorus* 317
fasciatus, *Halictus* 279
fasciatus, *Heterarthrus*, *Phyllotoma* 49
fasciatus, *Nomioides* 292
fasciatus, *Oxybelus* 242
fasciatus, *Panurgus* 276
fasciatus, *Pasites* 318
fasciatus, *Pompilus* 173
fasciatus, *Sphecodes* 288
fasciatus, *Stizus*, *Bembex* 225
fasciculata, *Andrena*
nigraenea var. 269
fasciculata, *Melecta* 326
fasciculata, *Nomada* 314
fasciipennis, *Tachysphex* 239
fascinata, *Scolia* 147
fastidiosissimus, *Eumenes* 182
fastidiosissimus, *Stenodynerus*
rufescens 190
fastidiosissimus, *Stenodynerus*,
Odynerus 190
fastidiosus, *Euodynerus*, *Odynerus* 183
fastigiata, *Priocnemis* 163
fastosus, *Pteronus* 59
faunus, *Phylloecus* 110
fausta, *Neurotoma*, *Lyda* 103
faustus, *Allantus* 90
faustus, *Sirex* 113
favosa, *Andrena* 267
favosa, *Eucera* 320
fedtschenkoi, *Camponotus* 199
fedtschenkoi, *Icteranthidium*,
Anthidium 301
feisthamelei, *Urocerus* 112
feminina, *Macrophyia diversipes* 85
feminina, *Pteronidea ribesii* 64
femoralis, *Aporus* 166
femoralis, *Belomicrus* 242
femoralis, *Bembix* 221
femoralis, *Cleptes* 126
femoralis, *Crabro* 226
femoralis, *Elampus* 127
femoralis, *Hylotoma* 24
femoralis, *Laelius*, *Bethylus* 124
femoralis, *Lygaeonematus pallipes* 72
femoralis, *Nematus* 58
femoralis, *Nomada* 312
femoralis, *Nomiapis*, *Apis* 292
femoralis, *Pompilus* 163
femoralis, *Strongylogaster* 82
femoralis, *Tenthredo* 85, 99
femorata, *Cimbex* 230
femorata, *Macrophyia* 87
femorata, *Tiphia* 150
femorata, *Xylocopa* 310
femoratum, *Hedychridium*,
Hedychrum 129
femoratus, *Cephus* 110
femoratus, *Cimbex*, *Tenthredo* 29
femoratus, *Dolerus* 79
femoratus, *Odynerus* 186
femoratus, *Pompilus* 165
femurrubrum, *Dolerus* 77
Fenella 47
fenestrata, *Priocnemis*, *Salius* 163
fenestrata, *Tenthredopsis* 99
fenestratum, *Diprion loricis* 34
fenestratus, *Emphytus* 41
fennica, *Formica* 199
fennica, *Gilpinia*, *Diprion* 33
fennica, *Megachile*
circumcincta var. 307
fennica, *Nomada* 316
fennica, *Pemphredon* 248
fennica, *Priocnemis* 163
fennicum, *Trichiosoma* 31
fennicus, *Dolerus* 78
fennicus, *Nematus* 70
fennicus, *Pontopristia* 61
fenniensis, *Chrysida medita* 138
Fentis 227
fentoni, *Tenthredo* 92
Fenusia 48
Fenusia subg. 48
Fenusella 48
fera, *Podalonia* 213
fera, *Tenthredo* 85
ferganensis, *Camponotus* 199
ferghanica, *Eucera* 320
feriata, *Selandria* 50
ferina, *Prosopis lineolata* 261
ferox, *Andrena* 267
ferox, *Oxybelus* 242
ferox, *Siobla*, *Macrophyia* 89
ferox, *Tetramorium* 209
feroxoides, *Tetramorium* 209
Ferreola 166
Ferreolina 166
Ferreoloides 168
ferrer, *Cerceris* 253
ferripennis, *Halictus tumulorum* 279
ferritincta, *Megachile* 308
ferrugatus, *Dolerus* 77
ferrugifer, *Bombus laesus* 331
ferruginata, *Andrena* 272
ferruginata, *Nomada auct.* 314
ferruginata, *Nomada*, *Apis* 312
ferruginatus, *Sphecodes* 288
ferruginea, *Andrena* 288
ferruginea, *Euura*, *Nematus* 59
ferruginea, *Hylotoma* 68
ferruginea, *Larra* 225
ferruginea, *Tenthredo* 94
ferrugineipes, *Halictus* 285
ferrugineitarsis, *Odynerus* 190
ferrugineociliatus, *Hortobombus*
consobrinus 328
ferrugineus, *Allantus* 68
ferrugineus, *Tachysphex* 238
ferruginosa, *Tenthredo* 38
fertoni, *Halictus* 280
fertoni, *Smicromyrme* 157
fertoni, *Tachysphex costai* 238
Fertoni 236
fervida, *Andrena* 267
fervida, *Holopyga*, *Chrysida fervida* 131
fervida, *Pheidole* 207
festiva, *Melecta* 326
festiva, *Tenthredopsis* 98
festivus, *Alysson* 219
festivus, *Amauronematus* 65
festivus, *Nematus* 63
festivus, *Pamphilus* 104
festivus, *Priocnemis* 163
Fethalia 89
fialai, *Sysphingta* 210
fidvicornis, *Nomada auct.* 316
fieuzeleti, *Trypoxylon* 244

figulus, *Sphex* 244
figulus, *Trypoxylon*, *Sphex* 244
figurata, *Andrena* 272
Fiharuxa 147
filamentosa, *Tenthredo* 94
filchnerae, *Anthophora* 324
filchnerae, *Bombus* 331
filiceti, *Blasticotoma* 27
filiceti, *Blasticotoma filiceti* 27
filicis, *Stromboceros* 75
filicis, *Strongylogaster*, *Tenthredo* 82
filicornis, *Ctenopriocnemis* 162
filicornis, *Evgates* 173
filicornis, *Nematus* 73
filicornis, *Tachyagetes*, *Evgates* 174
filicornis, *Tachysphex* 237, 238
filiformis, *Apethymus*, *Tenthredo* 40
filiformis, *Calameuta*, *Cephus* 108
filiformis, *Cephus* 108
filiformis, *Chrysura*, *Chrysia* 139
filiformis, *Euura*, *Pachynematus* 59
filippovi, *Carinostigmus*, *Stigmus* 247
filippovi, *Cephus* 109
Filixungulia 42
filum, *Calameuta*, *Cephus* 108
fimbriata, *Cerceris*, *Crabro* 253
finitima, *Blennocampa* 46
finitima, *Orientabia* 28
finitimus, *Crabro* 232
finschi, *Tenthredo finschi* 93
firssovi, *Leptothorax nassonovi* 208
fischeri, *Andrena* 265
fischeri, *Cryptochileius*, *Pompilus* 161
fischeri, *Nycteridium* 35
fissidens, *Coelioxys* 305
fissus, *Dolerus* 80
fissus, *Oxybelus* 243
flabellata, *Tarpa* 107
flabellicornis, *Megalodontes*, *Tarpa* 107
flabellifera, *Anthophora* 323
flabellifera, *Eucera* 320
flammula, *Athalia scutellariae* 45
flauicornis, *Tenthredo* 91
flauipes, *Crabro* 243
flava, *Cerceris* 254
flava, *Hoplocampa*, *Tenthredo* 67
flava, *Hylotoma* 24
flava, *Nomada* 312
flava, *Tenthredo* 91
flava, *Tenthredo bicincta* 97
flavata, *Vespa mongolica* var. 194
flavater, *Lygaeonematus ambiguus* 73
flavellus, *Eustenancistrocerus lobatus* 184
flavens, *Anthidium oblongatum* var. 301
flavens, *Brachythops*, *Tenthredo* 76
flaveola, *Dineura* 55
flaveola, *Tenthredo* 90
flavesca, *Andrena* 264
flavesca, *Cerceris* 253
flavesca, *Euura*, *Nematus* 59
flavesca, *Nomada fucata* var. 313
flavesca, *Selandria* 76

flavescens, *Stelis signata* 302
flavicans, *Nematus* 68
flaviceps, *Acantholyda*, *Tenthredo* 101
flaviceps, *Scolia* 148
flaviceps, *Vespula*, *Vespa* 196
flavicincta, *Vespa* 195
flavicincta, *Vespa* 221
flaviklypeata, *Tetralonia salicariae* var. 322
flaviklypeatus, *Odynerus* 183
flavicollaris, *Stethomostus*, *Atomostethus* 53
flavicolle, *Anthidium nigricolle* var. 302
flavicollis, *Blennocampa* 50
flavicollis, *Crabro* 232
flavicollis, *Heterarthrus*, *Phyllotoma* 49
flavicollis, *Hylotoma* 22
flavicollis, *Myzinella*, *Meria* 149
flavicomus, *Nematus* 73
flavicone, *Anteon*, *Gonatopus* 119
flavicornis, *Cerceris* 253
flavicornis, *Dufourea* 291
flavicornis, *Euodynerus notatus* 183
flavicornis, *Flavodufourea*, *Dufourea* 291
flavicornis, *Hylaeus* 285
flavicornis, *Megalodontes*, *Tarpa* 107
flavida, *Megachile pilicrus* 306
flavida, *Podalonia*, *Ammophila* 213
flavida, *Tenthredo* 91
flavidula, *Myrmica* 207
flavidus, *Bombus* 329
flavifasciata, *Tenthredo* 86
flavifrons, *Pamphilus* 105
flavifrons, *Scolia* 147
flavigularis, *Eumenes arbustorum* 185
flavigularis, *Katamenes*, *Eumenes* 185
flavilabre, *Anthidium* 303
flavilabris, *Andrena* 267
flavilabris, *Cerceris*, *Hylaeus* 253
flavilabris, *Tenthredo* 95
flavinervus, *Odontepyris* 125
flavior, *Selandria flavens* 76
flavior, *Vespa vulgaris* var. 196
flavipectus, *Tenthredo*, *Allantus* 93
flavipennis, *Andrena ephippium* var. 270
flavipennis, *Macrophya* 86
flavipennis, *Nematus* 56
flavipennis, *Sphex* 215
flavipennis, *Tenthredo* 97
flavipes, *Andrena* 275
flavipes, *Apis* 278, 279
flavipes, *Aprosthemma terebralis* 26
flavipes, *Ceratina* 310
flavipes, *Colpopyga*, *Hedychrum* 127
flavipes, *Crabro* 231
flavipes, *Eutomostethus*, *Tomostethus* 47
flavipes, *Euura* 63
flavipes, *Hedychrum* 127
flavipes, *Holcogneme* 66
flavipes, *Hylaeus* 260
flavipes, *Lyda* 105
flavipes, *Macrophya* 84, 85
flavipes, *Nematus* 64, 70
flavipes, *Nylanderia*, *Tapinoma* 202
flavipes, *Odynerus* 179
flavipes, *Schizocera* 26
flavipes, *Tenthredo* 25, 45, 71, 90, 99
flavipes, *Tiphia* 243
flavipicta, *Pristiphora* 73
flavipleuris, *Tenthredo* 96
flaviscapus, *Odynerus spinipes* ab. 187
flavissimus, *Crossocerus* 231
flavissimus, *Pachynematus clitellatus* 58
flavisternum, *Cephus* 109
flavistigma, *Cephalcia* 102
flavistigma, *Pemphredon* 248
flavistigma, *Selandria* 76
flavitarsis, *Hylaeus* 285
flavitarsis, *Crossocerus*, *Crabro* 231
flavitibia, *Dolerus gonager* 79
flaviventre, *Anthidium*, *nigricolle* var. 302
flaviventris, *Cephus* 110
flaviventris, *Hylotoma* 24
flaviventris, *Megachile* 308
flaviventris, *Nematus* 57
flaviventris, *Tenthredo* 103
flaviventris, *Tenthredo* 66
flavobarbatus, *Bombus agrorum* 331
flavobila, *Andrena* 271
Flavodufourea 291
flavofasciata, *Vespa* 194
flavogeniculatus, *Tachytes* 238
flavoguttata, *Nomada*, *Apis* 312
flavoguttata, *Perineura scutellaris* 99
flavoguttatum, *Anthidium*, *septemspinosum* var. 301
flavolimbata, *Nomada lineola* var. 313
flavomaculata, *Nomada* 311
flavomaculatum, *Anthidium*, *montanum* var. 300
flavomaculatus, *Tenthredopsis* 100
flavomandibulata, *Tenthredo*, *Tenthredopsis* 98
flavomarginatum, *Diprion laris* 34
flavomixta, *Arge*, *Hylotoma* 23
flavopicta, *Andrena* 292
flavopicta, *Nomada*, *Apis* 312
flavopleura, *Pristiphora* 73
flavopunctata, *Nomada*, *femoralis* var. 312
flavoscutellaris, *Tenthredella*, *mesomelas* 91
flavoscutellata, *Athalia lugens* 44
flavoscutellata, *Tenthredo moniliata* 95
flavoscutellatus, *Stenodynerus*, *tokyanus* 190
flavotectus, *Halictus carinaeventris* 279
flavovaria, *Ceratina* 310
flavoventralis, *Macrophya* 87
flavus, *Cryptochileius*, *Priocnemis* 161
flavus, *Lasius*, *Formica* 201
flavus, *Nematus* 68
fletcheri, *Dimorpha* 218
fletcheri, *Empria*, *Poecilosoma* 41

fletcheri, Nematus 71
fletcheri, Nomia 291
fletcheri, Taxonus 39
fletcheri, Tenthredo providus 98
floralis, Astatus 108
floralis, Colletes 258
floralis, Crabro 233
floralis, Hylaeus, *Prosopis* 261
floreæ, Amegilla, *Anthophora* 322
floreæ, Andrena 273
florentinum, Anthidium, *Apis* 300
floricola, Allodynerus, *Odynerus* 178
floricola, Andrena 270
floricola, Hylaeus 261
floricola, Odynerus 178
floricola, Tenthredo 87
florilegus, Bombus 327
floripetum, Pseudoanthidium 302
florisomne, Chelostoma auct. 295
florisomne, Chelostoma, *Apis* 295
florivaga, Andrena 268
Florkinus 234
florum, *Apis* 265
fluctuans, Eumenes 185
fluminensis, Macrophyia erythropus 86
fluvicornis, Taxonus 43
fodiens, Cerceris 253
fodiens, Colletes, *Apis* 258
foederata, Polistes 193
foersteri, Cephus 110
foersteri, Coelioxys 304
foersteri, Laelius 124
foersteri, Prosopis 259
foetidae, Euura, *Pontania* 59
folii, Tenthredo grossulariae 64
foliivolutor, Anthidium 303
Fongiba 149
fonscolombei, Scleroderma 124
fonscolombii, Andrena 267
forcipata, Myrmica 205
foreli, Formica 199
foreli, Formica exsecta
 pressilabris var. 199
forficula, Arge 25
Forficulotarpa 107
Formica 199
Formica subg. 199
formicaeformis, Myrtophorus 119
Formicapis subg. 297
formicaria, Mutilla 149
formicarius, Dryinus 120
formicarius, Gonatopus 120
formicarius, Gonatopus 121
Formicidae 197
formiciformis, Cephalonomia 124
Formicinae 198
formicinus, Gonatopus 121
Formicoidea 197
Formicoxenus 204
Formila 122
formosa, Pachyprotasis 87
formosana, Thrinax, *Hemitaxonius* 83
formosella, Blennocampa 48

formosicola, Coelioxys 304
formosus, Arpactus 223
formosus, Harpactus, *Arpactus*
 formosus 223
formosus, Janus, *Cephus* 110
formosus, Pterochilus 189
Forsia 47
forsii, Osmia 298
forsii, Tenthredo 95
forsiusi, Pontania 64
forsiusi, Pristiphora conjugata 71
forsslundi, Formica 199
forsythiae, Apareophora 45
forticarinatus, Oxybelus 242
fortior, Myrmica bergi var. 205
fortipes, Andrena 294
fortis, *Apis* 269
fortstriolatus, Palarus 243
fortunatus, Tachysphex panzeri 239
fosorius, *Apis* 258
fossilabris, Formica 199
fossoria, Megachile argentata var. 306
fossoria, Sphecodes 233
fossorius, Ectemnius 230
fossorius, Ectemnius, *Sphex* 233
fossulata, Tetralonia 322
Fossulocephus 109
foveatus, Elampus, *Ellampus* 128
foveifrons, Amauronematus tillbergi 65
foveivaginatus, Priophorus 54
foveolata, Osmia 296
foveolatum, Chelostoma, *Heriades* 296
foveolatus, Gorytes foveolatus 222
foveolatus, Halictus 286
foveolatus, Pachynematus 57
foveolatus, Psen 251
foveolatus, Symmorphus 191
Foveosmia subg. 295
foviefrons, Selandria 76
fragosa, Cerceris 253
fragrans, Bombus, *Apis* 331
franclemonti, Xylocelia 247
franconica, Andrena 271
frankenbergeri, Chrysis 135
franki, Tenthredopsis 100
fratellum, Lasioglossum, *Halictus*
 fratellum 283
fratellus, Anoplus, *Pompilus* 172
frater, Halictus 278
fratercula, Andrena 271
fraterculus, Eumenes 182
fraterculus, Nysson 220
fraterna, Coelioxys 305
fraterna, Eumenes 182
fraterna, Mutilla 154
fraternus, Nematus 70
frauenfeldi, Stenodynerus,
 Odynerus 190
fraxini, Nematus 74
fraxini, Tenthredo 53
frenalis, Euura, *Nematus* 59
freya, Andrena 265
freygessneri, Cryptocheilus,
 Priocnemis 161

freygessneri, Halictus 283
freygessneri, Stelis 302
freygessneri, Tachytes 240
Frey-Gessneri, Tachytes 240
freyi, Pachynematus 56
freyja, Euura 59
friederikae, Dioxyx cincta ab. 303
friesei, Bombus kirbyellus 327
friesei, Camptopoeum 276
friesei, Colletes 258
friesei, Diodontus 247
friesei, Nysson 220
friesei, Pristiphora, *Lygaeonematus* 72
friesei, Schizocera 26
friesei, Tenthredopsis, *Thomsonia* 98
frigida, Myrmica, *Nematus* 72
frigida, Siobla, *Encarsioneura* 89
frigidus, Colletes 257
frisoni, Psithyrus flavidus 329
frivaldszki, Aclastocera 145
frivaldszkyi, Eucera 320
frivaldszkyi, Allantus 97
frivaldszkyi, Arge, *Hylotoma* 23
frivaldszkyi, Chrysis frivaldszkyi 135
frivaldszkyi, Eumenes 184
Frivaldszkyiformis 22
frontale, Camptopoeum, *Prosopis* 276
frontalis, Andrena 268
frontalis, Prospis 275
frontalis, Tenthredella ferruginea 94
frontalis, Vespa 195
fronticalis, Pseudoscolia 256
fronticorne, Trypoxylon fronticorne 245
frontimacula, Anoplolyda 104
frontina, Corynis, *Amasis* 32
frontosa, Aphaenogaster famelica 203
fruhstorferi, Lipotriches, *Nomia* 291
frutetorum, Gilpinia, *Tenthredo* 33
fruticum, Nematus 75
fucata, Andrena 264
fucata, Nomada 313
fuciformis, Osmia 299
fuciformis, Scolia, *Sphex* 148
fucus, Halictus 277
fudakowski, Halictus 285
fuzdi, Sphecodes 290
fuerstenbergensis, Aneugmenus,
 Selandria 76
fugax, Diplorhoptrum auct. 207
fugax, Nomia, *Nomia* 292
fugax, Solenopsis, *Formica* 207
fugax, Tachysphex *Tachytes* 238
fuhescens, Megachile 308
fujii, Lasius 201
fujiiense, Euura, *Pikonema* 59
fukaii, Monophadnus 53
fukaii, Rhynchium 189
Fukpokta 149
fukuiana, Andrena 267
fukuiensis, Sphecodes 290
fukuokensis, Andrena 272
fulgens, Abia 28
fulgida, Chrysis 135

fulgidipennis, Monophadnus 50
fuliginata, Arge 24
fuliginipennis, Monophadnus 50
fuliginosa, Apis 267, 296
fuliginosa, Fenusa 48
fuliginosa, Tenthredo 53
fuliginosus, Lasius, *Formica* 201
fuliginosus, Pompilus 166
fuliginosus, Stethomostus, *Tenthredo* 53
fulminatrix, Chrysura 143
fulva, Andrena 268
fulva, Eumenes 185
fulva, Eumenes var. β 185
fulva, Hoplitis, *Osmia* 296
fulva, Saropoda 319
fulva, Tenthredo 91, 104
fulvago, Andrena, *Apis* 265
fulvescens, Andrena 265
fulvescens, Euura, *Pteronidea* 59
fulvescens, Tetraloniella, *Tetralonia* 322
fulviceps, Tenthredo 99
fulvicollis, Selandria 51
fulvicorne, Lasioglossum, *Melitta*
fulvicorne 283
fulvicornis, Andrena 271
fulvicornis, Colletes 258
fulvicornis, Hoplocampa, *Tenthredo* 68
fulvicornis, Hylaeus 285
fulvicornis, Mellinus 246
fulvicornis, Nomada 313
fulvicornis, Psenulus, *Psen* 252
fulvicornis, Spinarge, Arge 25
fulvicornis, Tenthredo 94
fulvicrus, Andrena 275
fulvicrus, Hylaeus 281
fulvicrus, Melitta 275
fulvida, Andrena 267
fulvimana, Megachile 307
fulvipennis, Pseudotiphia, *Tiphia* 150
fulvipes, Aglaostigma, *Tenthredo*
fulvipes 83
fulvipes, Ametastegia 39
fulvipes, Andrena 267
fulvipes, Anthidium punctatum var. 301
fulvipes, Anthophora 324
fulvipes, Apis 300, 303
fulvipes, Cerceris 253
fulvipes, Macropis, *Megilla* 294
fulvipes, Megilla 293
fulvipes, Nysson 220
fulvipes, Stizus 225
fulvipes, Tenthredo 38, 70, 105
fulvitarsis, Andrena 272
fulvitarsis, Andrena 275
fulvitarsis, Anthophora 323
fulvitarsis, Odynerus, *Hoplomerus* 187
fulvitarsis, Perineura 92
fulvitarsis, Psen 251
fulvitarsis, Tachysphex, *Tachytes* 238
fulvitarsis, Tachytes 238
fulvitellaris, Andrena gallica ssp. 269
fulviventre, Anteon, *Dryinus* 119
fulviventre, Anthidium 301

fulviventris, Apis 298
fulviventris, Epeoloides 319
fulviventris, Nysson 220
fulviventris, Planiceps 166
fulviventris, Tenthredo 94
fulvocincta, Melitta 282
fulvocincta, Phymatoceropsis 52
fulvocinerea, Anthophora
parietina var. 324
fulvo-crustatus, Campylogaster 264
fulvogastra, Mutilla rufipes var. 156
fulvohirta, Megachile lagopoda var. 307
fulvohirta, Osmia 297
fulvooides, Eumenes sichelii var. 185
fulvoleja, Pterocheilus sibiricus var. 189
fulvum, Aprosthemma 26
fulvus, Nematus 61
fumipennis, Arachnospila, *Pompilus* 170
fumipennis, Cephus 109
fumipennis, Crabro 233
fumipennis, Hylaeus 259
fumipennis, Nematus 68
fumipennis, Pamphilus, *Lyda* 104
fumipennis, Phyllotoma 49
fumosus, Dolerus 79, 81
funebris, Andrena 268
funebris, Athalia proxima 44
funebris, Stenodynerus, *Odynerus* 190
funeraria, Mutilla 153
funerarius, Palarus 243
funerarius, Sphecodes 215
funerea, Cerceris 254
funereipes, Pompilus 166
funereus, Allantus 90
funereus, Stethomostus, *Tenthredo* 53
funerula, Dasylabris kaszabi 153
funerulus, Nematus 72
fur, Nematus 60
furcata, Andrena 267
furcata, Anthophora, *Apis* 323
furcata, Sterictiphora, *Tenthredo* 26
furcata, Tenthredo 26
furcatus, Halictus 278
furcatus, Oxybelus 243
furnasensis, Halictus 283
furukawai, Megachile 307
furva, Nomada 313
furvensis, Nematus 73
furovoides, Nematus 313
furus, Monophadnus 50
fusca, Andrena 267
fusca, Apis 299
fusca, Formica 200
fusca, Nematus 74
fusca, Nomada 313
fusca, Nomioides minutissima 292
fusca, Osmia 299
fusca, Schizocera 26
fusca, Tenthredo 53
fusca, Vespa crabro 194
fuscarima, Pteronidea 65
fuscata, Andrena gelriae karelica f. 274
fuscata, Melitta 274

fuscata, Tenthredo 63, 90
fuscata, Xylocopa 310
fuscatus, Diphlebus 248
fuscicorne, Aprosthemma, *Schizocera* 26
fuscicornis, Apis 264
fuscicornis, Bethylus, *Omalus* 125
fuscicornis, Mesitius 124
fuscicornis, Nematus 70
fuscicornis, Nomada 313
fuscicornis, Omalus 125
fuscicornis, Sirex 113
fuscicornis, Tenthredo 95
fuscicornis, Tremex, *Sirex* 113
fuscifrons, Megachile lapponica 307
fuscinervis, Arge 23
fuscipennis, Arge, *Hylotoma* 23
fuscipennis, Bethylus 125
fuscipennis, Decatria 49
fuscipennis, Dichroa 287
fuscipennis, Dolerus germanicus 78
fuscipennis, Microdiprion, *Diprion* 34
fuscipennis, Nematitus, *Nematus* 68
fuscipennis, Nematus 68
fuscipennis, Pemphredon 249
fuscipennis, Phyllotoma 47
fuscipennis, Psen 251
fuscipennis, Psenulus *Psen* 252
fuscipennis, Solenius 233
fuscipennis, Tarpa 107
fuscipennis, Tenthredo 47
fuscipennis, Tenthredo, *Allantus*
marginella 96
fuscipes, Andrena, *Melitta* 265
fuscipes, Arge, *Hylotoma* *fuscipes* 23
fuscipes, Astata 218
fuscipes, Formica sanguinea var. 200
fuscipes, Psenulus 252
fuscipes, Symmorphus, *Odynerus* 191
fuscipes, Tenthredo 93
fuscitarsis, Crabro 232
fuscitarsis, Hylaeus immarginatus 285
fuscitarsis, Selandria serva 81
fuscitarsis, Tenthredo 39
Fuscitarsus, Crabro 232
fusciventris, Apis 272
fuscocostalis, Tenthredo 96
fuscohirta, Andrena 267
fuscoideas, Formica cunicularia 200
fuscoideas, Lasius flavus var. 201
fuscomaculata, Euura, *Nematus* 59
fuscomarginata, Arachnospila,
Pompilus 169
Fuscopolistes 192
fuscosa, Andrena 270
fuscoterminata, Tenthredo 92
fuscula, Pseudodineura, *Tenthredo* 75
fuscus, Cryptocampus 75
fuseipennis, Lithurgus 295
gabriellae, Hedyrhridium 129
gaditana, Vespa 225
gaetula, Andrena 272
gagates, Formica 200

gagathinus, Eutomostethus,
Tenthredo 47
 gagatoides, *Formica* 200
gagatoides, *Formica fusca* var. 200
Galactopterus 167
galactopterus, *Pompilus* 173
 galbula, *Colpa*, *Vespa* 147
galbula, *Scolia* 148
 galbula, *Vespa* 147
galeatus, *Leptothorax* 208
galericulatae, *Dentathalia* 45
galiciensis, *Allantus* 90
galileus, *Halictus* 278
 gallae, *Euura* 55
gallarum, *Nematus* 64
 gallarum, *Osmia* 299
gallarum, *Tenthredo* 56
gallica, *Amasis crassicornis* 32
gallica, *Andrena* 269
gallica, *Bembix* 222
gallica, *Polistes* 192
gallica, *Vespa* 192
gallicola, *Goniozus*, *Parasierola* 125
gallicola, *Nematus* 63
gallicola, *Pontania* 63
gallicum, *Episyron*, *Pompilus* 167
gallicus, *Bothriomyrmex* 197
gallicus, *Miscophus* 241
gallicus, *Odynerus* 185
gallicus, *Polistes*, *Vespa* 193
gallienii, *Myrmica* 205
galloisi, *Chrysis*, *Tetrachrysis* 135
gandzhensis, *Andrena*
 colletiformis ab. 264
ganglbaueri, *Bembix* 222
garbiglietti, *Trichiocampus* 54
 garrula, *Amegilla*, *Apis* 322
 garrula, *Scolia* 148
gascheti, *Andrena* 271
Gastrargyron 236
gastricus, *Monophadnus* 51
Gasteracanthus 236
Gastrosphaeria 215
 gaullei, *Anteon* 119
gaunitzi, *Pristiphora* 74
 gavarnicus, *Halictus* 279
gazella, *Ancistrocerus*, *Vespa* 179
geerii, *Vespa* 195
gegen, *Ammophila* 213
gegen, *Ammoplanus* 246
gegen, *Bembix oculata* 222
gegen, *Diodontus* 247
gegensimus, *Miscophus* 241
gehrsi, *Euura*, *Pachyneurus* 59
gei, *Fenusia* 50
gelida, *Empria*, *Tenthredo* 41
gelida, *Pristiphora* 72
gelidus, *Nematus* 66
gelriae, *Andrena* 274
gemellus, *Nematus* 73
gemina, *Pemphredon* 248
geminata, *Sterictiphora*, *Tenthredo* 26
geminatus, *Halictus* 279

geminatus, *Hoplisus* 223
geminus, *Hylaeus* 283
gemma, *Chrysis bidentata* var. 139
gemmarum, *Cryptocampus* 62
Gemmura 55
genalis, *Hylaeus* 261, 262
genalis, *Megachile* 306
genalis, *Stelis* 303
genata, *Endemyolia* 46
generosa, *Holopyga*, *Ellampus*
 generosa 131
genevensis, *Andrena* 273
genevensis, *Halictus puncticollis* 283
gengiskhan, *Pseudochrysis* 140
geniculata, *Lyrops* 238
geniculata, *Meria*, *Myzine* 149
geniculata, *Pristiphora*, *Nematus* 72
geniculata, *Schizocera* 26
geniculata, *Selandria* 53
geniculata, *Strongylogaster* 82
geniculata, *Tachytes panzeri* 238
geniculata, *Tenthredo* 50, 79
geniculatus, *Crabro* 226, 229, 233
geniculatus, *Nysson* 220
genimaculata, *Prosopis*
 clypearis var. 261
genistae, *Rhogogaster* 88
gentianae, *Andrena* 274
genualis, *Tenthredopsis stigma* 100
genucinctus, *Dolerus* 77
geoffrellus, *Sphecodes*, *Melitta* 288
geoffroyanus, *Odynerus* 179
geoffroyi, *Cladius* 54
geoffroyi, *Lyda* 110
geoffroyi, *Polistes* 192
georgica, *Cataglyphis aenescens* 199
georgiense, *Aprosthemma staudingeri* 26
georgii, *Stenamma* 207
geranii, *Ametastegia*, *Emphytina* 40
Gerbekas 124
germabensis, *Megachile* 305
germabensis, *Tachytes* 237
germabica, *Andrena* 272
germabicus, *Odynerus* 183
germania, *Vespa crabro* 193
germanica, *Andrena* 265
germanica, *Dasypoda* 293
germanica, *Eumenes dubius*
 sareptanus var. 182
germanica, *Nomada* 312, 314
germanica, *Vespula*, *Vespa* 196
germanicus, *Dolerus*, *Tenthredo*
 germanicus 78
germanicus, *Nematus* 73
germanicus, *Odynerus* 190
germari, *Chrysis germari* 135
gerstaekerii, *Bombus* 328
gerstaekerii, *Hedychrum*
 gerstaekerii 130
gerstaekerii, *Prosopis* 260
gerstaekeriformes, *Hedychrum* 142
gerszabiensis, *Megachile* 305
gertabi, *Chrysis* 135
gertrudis, *Passaloecus* 248
gerula, *Pristiphora*, *Lygaeonematus* 72
gessneri, *Dolerus* 78
Gestrodynerus subg. 188
Ghelynia 302
Ghilardocheilus subg. 187
ghilarovi, *Priocnemis* 163
ghilarovi, *Psenulus* 252
ghilarovi, *Tracheliodes* 236
ghilianii, *Dryudella* 218, 219
ghilianii, *Nematus* 71
gibba, *Andrena* 274
gibba, *Mutilla* 156
gibba, *Nomada* 287, 288
gibba, *Sphecodes* 287
gibba, *Tachysphex lativalvis* 238
gibberosa, *Tenthredopsis* 99
gibbomima, *Arachnospila*,
Psammochares 169
gibbosa, *Aphaenogaster*, *Formica* 203
gibbosa, *Apis* 288
gibbosum, *Aglaostigma*, *Tenthredo* 83
gibbosus, *Dolerus* 79
gibbulus, *Evagetes*, *Pompilus* 171
gibbus, *Bothriomyrmex* 197
gibbus, *Hylaeus* 261
gibbus, *Sphecodes*, *Sphecodes* 288
gibbus, *Tachysphex* 238
giffardi, *Solenius* 232
gifuensis, *Tenthredo* 94
gifui, *Tenthredo* 94
gigantea, *Coelioxys rufescens* var. 304
gigantea, *Megaxyela* 20
gigantea, *Melecta armata* var. 326
gigas, *Ichneumon* 113
gigas, *Tenthredo mesomelas* 91
gigas, *Urocerus*, *Ichneumon gigas* 113
gilanum, *Lasioglossum*, *Halictus* 283
Gilbertella 166
Gilbertellana 166
gillettei, *Blennocampa* 50
Gilpinia 33
gilvifrons, *Andrena* 270
gilvipes, *Dolerus*, *Tenthredo gilvipes* 79
giltus, *Agrobombus* 331
ginran, *Nomada* 313
giraudi, *Megachile* 306
giraudi, *Osmia* 299
giraudi, *Phyllaecus* 111
giraudii, *Halictus* 283
glabella, *Nomada* 315
glabellifer, *Cephus* 111
glaber, *Camponotus aethiops* var. 199
glaber, *Philanthus* 255
glaberrima, *Nomada* 313
glabra, *Apis* 288
glabrata, *Ametastegia*, *Tenthredo* 39
glabrata, *Myrmilla*, *Mutilla* 154
glabrata, *Pachyprotasis* 87
glabrigollis, *Athalia* 44
glabrifrons, *Euura*, *Pontania* 59
glabriuscum, *Lasioglossum*, *Halictus*
glabriuscum 283

glabriventris, *Andrena* 267
glacialis, *Bombus* 330
glacialis, *Dolerus konowi* 78
glacialis, *Epeolus* 317
glacialis, *Eumenes* 182
glacialis, *Euura, Amauronematus* 59
glasunovi, *Chrysis* 135
glasunovi, *Halictus morbillosus r.* 280
glasunowi, *Parnopes* 142
glauca, *Formica* 200
glauca, *Formica rufibarbis* var. 200
glauca, *Pristiphora* 72
glaucae, *Euura, Pontania* 59
glaukopis, *Fesunella, Phoenusa* 48
glaukopis, *Nomada* 313
glaukopis, *Tenthredo* 67
glenelgensis, *Nematus* 60
glesicornis, *Allantus* 93
glesipennis, *Pachynematus* 57
glinka, *Pontania* 64
globiceps, *Prosopigastra*,
Tachysphex 237
globosa, *Osmia* 298
globosa, *Smicromyrme* 156
globosiformis, *Osmia* 298
globulilabris, *Andrena* 273
globulus, *Hylaeus, Prosopis* 261
gloriosa, *Holopyga* 142
Glossamegilla subg. 322
Glossomyrme 157
glottianus, *Nematus* 59
glottianus, *Taxonius* 39
glutinans, *Apis* 259
glutinosae, *Nematus* 67
glutinosus, *Heriades* 299
glycybromifer, *Halictus* 280
gmelini, *Alpinobombus kirbyellus* 327
Gnathias 310
gobicola, *Dasylabris* 153
gobicola, *Pseudomyrmosa* 152
gobiensis, *Bembix* 222
gobiensis, *Diodontus major* 247
gobiensis, *Podalonia, Ammophila* 213
gobiensis, *Sphex* 215
gobiensis, *Stizus* 225
goesswaldi, *Epimyrma* 207
golosejevi, *Stenamma* 207
gonager, *Crossocerus* 227
gonager, *Dolerus, Tenthredo* 79
Gonagerformis 77
Gonatopodinae 120
Gonatopus 120
gongshanensis, *Vespula* 196
Gongylocorsia 103
Gonioxybelus 242
Goniozus 125
Gonius 243
Gonodontochrysis 133
Gonostigmus 249
gonymelas, *Nematus* 71
goodeniana, *Nomada, Apis* 313
gordiagini, *Myrmoxenus* 207
gorge, *Lasioglossum* 283

gorkaensis, *Athalia lineolata* 44
gorkiensis, *Halictus* 280
Gorystizus 221
Gorytes 222
gossypina, *Tenthredo* 42
gotarus, *Pachynematus* 57
gracile, *Rhopalum* 235
gracilenta, *Tenthredo* 95
gracilentus, *Leptocimbex*,
Clavellaria 30
gracilicornis, *Arge, Hylotoma* 23
gracilicornis, *Cladius* 54
gracilicornis, *Clameuta, Cephus* 108
gracilicornis, *Hylaeus, Prosopis* 260
gracilicornis, *Paracharactus*,
Selandria 51
gracilicornis, *Prosopis*
luteifrons var. 260
gracilicornis, *Pteronidea* 63
gracilidentata, *Euura, Pteronidea* 59
gracilior, *Sphecodes* 287, 288
gracilipes, *Anthophora* 324
gracilipes, *Stizus* 225
gracilis, *Chrysis angustula* 143
gracilis, *Andrena* 264
gracilis, *Astrotus* 218
gracilis, *Bembix* 221
gracilis, *Cephus* 109
gracilis, *Coelioxys* 305
gracilis, *Crabro* 229
gracilis, *Dolerus* 79
gracilis, *Halictus* 284
gracilis, *Mesitius* 124
gracilis, *Neostromboceros* 81
gracilis, *Palarus* 243
gracilis, *Passaloecus, Diodontus* 248
gracilis, *Priocnemis, Salius* 163
gracilis, *Smicromyrme* 155
gracilis, *Symmorphus, Odynerus* 191
gracilis, *Tenthredo* 39
gracilis, *Tenthredo, Tenthredella* 98
gracilis, *Tenthredopsis* 99
gracilissimus, *Crabro* 231
gracillima, *Ammophila* 212
gracillima, *Chrysis gracillima* 135
graeca, *Chrysis* 137
graeca, *Eucera* 320
graeca, *Mutilla* 157
graeca, *Myzine* 149
graeca, *Pseudomeria* 149
Graecandrena subg. 266, 267
graecus, *Allantus* 90
graecus, *Gonatopus* 121
graecus, *Halictus* 278
graeffei, *Hylotoma* 22
graeffei, *Osmia* 300
graelsii, *Chrysis graelsii* 135
grahami, *Polistes rothneyi* 192
grahami, *Vespula rufa* 196
graja, *Macrocerca* 321
graja, *Nomada* 313
graja, *Tetraloniella, Macrocerca* 322
gramineus, *Halictus* 279
graminicola, *Myrmecina, Formica* 205
graminis, *Cephus* 109
graminis, *Nematus* 62
grandiceps, *Tenthredo* 98
grandii, *Leptochilus alpestris* var. 185
grandii, *Mimesa* 250
grandii, *Tachysphex* 238
grandimargo, *Systropha* 291
grandior, *Parnopes* 142
grandior, *Argogorytes, Gorytes*
mystaceus 221
grandis, *Cladius, Nematus* 54
grandis, *Dineura* 68
grandis, *Hedychrum* 130
grandis, *Hylaeus* 277
grandis, *Lophopompilus, Pompilius* 173
grandis, *Megadineura, Dineura* 68
grandis, *Nematus* 54
grandis, *Osmia* 296
grandis, *Pompilus* 173
grandis, *Pseudomalus, Ellampus* 132
grandis, *Salius* 173
grandis, *Selandria* 81
grandis, *Siobla* 89
grandis, *Solenius* 233
grandis, *Sphecodes* 287
grandis, *Tachyagetes, Evagetes* 174
grandis, *Tachytes* 236
grangeoni, *Acantholyda* 101
granulatus, *Crabro* 232
granulosus, *Halictus* 283, 285
granulosus, *Tachysphex* 239
graphicus, *Odynerus* 183
grata, *Tenthredo* 90
gratirosus, *Tachyagetes, Aporus* 174
gratus, *Aneugmenus* 76
gravenhorsti, *Nematus* 68
gravida, *Andrena* 265
gravidula, *Ceratina* 310
gredan, *Andrena gelriae* 274
gredleri, *Hylaeus* 259
gredleri, *Leptocephalum* 204
Gredleri, *Lindenius* 232
Greeleyella 276
gregori, *Lestodryinus* 120
gregoriense,
Pseudoanthidium alpinum 302
gregoryi, *Smicromyrme pusilla*
septentrionalis f. 156
gressitti, *Polistes rothneyi* 192
greyi, *Anthidium* 301
gribodoi, *Chrysis* 143
gribodoi, *Chalepoxenus* 203
gribodoi, *Halictus* 284
gribodoi, *Tenthredo* 95
gribodoi, *Vespa crabro* var. 194
griffini, *Cimbex* 30
grisea, *Apis* 275
grisea, *Scolia* 147
griseipennis, *Halictus* 283
griseithorax, *Dasypoda argentata* 293

griseohirta, Megachile circumcincta var. 307
griseola, Andrena 273
griseola, Macrocera 322
griseolum, Lasioglossum, *Halictus griseolum* 283
grisescens, Halictus 285
grisescens, Tropidotilla, *Mutilla* 157
griseus, Nematus 57
groenblomi, Pristiphora, *Lygaeonematus* 72
groenblomi, Trichiosoma 31
groenlandica, Euura, *Amauronematus* 59
grohmanni, Chrysis grohmanni 143
grohmanni, Icteranthidium, *Anthidium* 301
Grosapis 305
grossa, Nomada 311
grossa, Siobla 89
grossa, Tiphia 147
grosseseratum, Aglaostigma, *Macrophyopsis* 83
grossulariae, Nematus 64
grossulariae, Pristiphora 70
grossulariae, Tenthredo 39, 64
gruenwaldti, Rophites 291
grumorum, Chrysis 135
guentheri, Pompilus 169
gueruensis, Leptochilus signatus 185
guichardi, Leptochilus 185
guichardi, Nysson 220
Gunnea 52
guntionensis, Konowia 115
gusenleitneri, Antepipona 180
gusenleitneri, Halictus 278
gussakovskii, Abia 28
gussakovskii, Andrena 269
gussakovskii, Empria 42
gussakovskii, Lindenius 235
gussakovskii, Priocnemis 163
gussakovskii, Sapyga 145
gussakovskii, Scolia 148
gussakovskii, Tachysphex 238
gussakovskij, Psenulus 252
gussakovskiji, Trypoxylon exiguum 244
gussakovskiji, Mimesa 250
gussakovskiji, Miscophus 241
gussakovskiji, Trypoxylon clavicerum 244
gussakovskijii, Priocnemis 163
guttata, Conaspida, *Eriocampa* 84
guttata, Poecilosoma 41
guttata, Tenthredo 41
guttator, Allantus brachycerus 93
guttatus, Crabro 231
guttatus, Dolerus aericeps 77
guttatus, Ectemnius 231
guttatus, Ectemnius, Crabro 232
guttatus, Nysson 220
guttatus, Sphecodes 245
guttulata, Nomada 313
guttulatus, Cryptocheilus, *Priocnemis* 161

gwynana, Melitta 267
gyllenhali, Pamphilus, *Lyda* 104
Gymnaphelopus 118
Gymnochares 167
Gymnomerus 184
Gymnonychus 70
Gymnus 302
gynandromorpha, Tenthredo 94
gyrifrons, Cryptocheilus, *Salius* 161
Gyrodroma 302
Gyrodromella subg. 296
Gyrostoma subg. 192
Haba 128
Habrophora 325
Habropoda 325
haematoches, Dolerus, *Tenthredo* 80
haematodes, Nomada jacobaeae var. 313
haematopoda, Apis 298
haematopus, Tenthredo 85
haematomygia, Hemichroa 67
haematurus, Bombus 330
haemorrhoa, Andrena, *Apis* 275
haemorrhoa, Apis 275
haemorrhao, Coelioxys 304
haemorrhao, Lyrops 239
haemorrhoidalis, Apis 275
haemorrhoidalis, Calameuta, *Tenthredo* 108
haemorrhoidalis, Lithurgus 295
haemorrhoidalis, Melitta, *Andrena* 294
haemorrhoidalis, Nematus 59
haemorrhoidalis, Oxybelus 242
haemorrhoidalis, Pristiphora pallidiventris 73
haemorrhoidalis, Psen 252
haemorrhoidalis, Scolia 147
haemorrhoidalis, Tenthredo 68
haemorrhoidula, Andrena 265
haemorrhous, Dolichurus 211
haemorroidalis, Trachelus 108
haesitans, Halictus 280
Hahiya 149
hainanensis, Polistes rothneyi 192
hainanensis, Vespula 196
hakiensis, Tenthredopsis 92
hakkari, Colletes 258
hakkarius, Halictus kessleri 279
hakodadi, Pompilus 167
hakonensis, Tenthredella 96
hakusana, Hinatara 49
hakusana, Nomada 314
hakusanus, Aproceros 25
hakusanus, Crossocerus 230
hakusanus, Psen 251
hakutozanus, Gorytes 223
hakutozanus, Symmorphus 191
haladai, Ammophila 212
halensis, Physetopoda, *Mutilla* 155
halensis, Tenthredo 85
halesus, Emphytus 39
haliartus, Emphytus 39
Halictidae 277
Halictinae 277
Halictini 277
Halictoides subg. 290
halictoides, Andrena 274
halictula, Dufourea, *Rhophites* 290
Halictus 277, 287
Halictus subg. 277
Halidamia 48
halidayellus, Mesitius 124
halinae, Arge 23
halinae, Cleptes 126
haloga, Andrena niveata ssp. 270
hamata, Osmia 297
Hamatodentiden 77
hammarstroemi, Nomada 314
hammeri, Neotrogaspidia, *Trogaspidia* 157
hammi, Cephalonomia 124
handlirschi, Ammoplanus 246
handlirschi, Diodontus 247
handlirschi, Gorytes 220
handlirschi, Nysson 220
handlirschanus, Bombus 329
hanedai, Sphecodes 288
hangacius, Ancistrocerus 179
hangani, Nomada 316
hankoi, Priocnemis 163
hannibalis, Trypoxylon 244
Hannonis, Crabro 232
hansenii, Anthophora 325
hanuman, Sphecodes 289
Haplocephalus 108
Haplogonatopus 121
harbecki, Pempredon 249
harbinensis, Alysson 219
harbinensis, Gorytes 223
harbinensis, Lindenius 235
harmandi, Lasioglossum, *Halictus* 280
Harpactes 223
Harpactopus 215
Harpactus 223
Harpagoxenus 204
harpicola, Euura, *Amauronematus* 59
Harpiphorus 42
Harpophorus 42
harrisoni, Pontania 64
hartigi, Cladardis 46
hartigi, Euura, *Amauronematus* 59
Hartigia 110
hartigii, Cephalcia, *Lyda* 102
hartigii, Dineura 55
hartigii, Dolerus 80
hartigii, Macrophya 90
hartigii, Nematus 74
hartmanni, Rophites 291
harukawai, Dolerus 80
harveyi, Nysson 220
harwoodi, Dolerus 80
hasanensis, Smicromyrme 156
hasdrubal, Amblyellus, *Pompilus* 168
hasdrubal, Pompilus 168
hasegawae, Arge 23
haslauica, Andrena florea f. 273

hasshanus, Sphecodes 289
hastatae, Euura 59
hastatae, Pontania 59
hastatavora, Euura 59
hasticornis, Megachile 308
hatogayuus, Sphecodes 290
hattorfiana, Andrena, *Nomada* 264
haueri, Cerceris 253
Hauffapis 332
haupti, Agenioideus 167
haustus, Emphytus 39
haworthana, Apis 325
hayashi, Lasius 202
healaei, Camponotus 70
healyi, Heterarthrus 49
hebes, Andrena 267
hebes, Euura, *Amauronematus* 59
hebescens, Coelioxys 304
hebraeus, Stizus 225
hederae, Colletes 258
hedgreni, Crabro 228
hedickei, Arachnospila,
Psammochares 169
hedikae, Andrena 263
hedini, Halictus hedini 278
hedstroemi, Euura, *Amauronematus* 59
Hedychrnidium 128
Hedychrum 130
heecheonparki, Pamphilus 104
hekouensis, Vespa 193
helena, Olgia 224
helpferanus, Bombus 331
heliaca, Prosopis 260
helicinus, Nematus 66
Helicosmia subg. 298
Heliophila subg. 323
helophilus, Tachysphex 239
helipolis, Andrena afzeliella var. 274
helipolites, Tachysphex 238
helleni, Euura, *Amauronematus* 59
helleni, Gonatopus, *Dicondylus* 121
hellenica, Myrmica 205
hellenica, Myrmica rugulosa var. 205
hellenica, Xylocopa 310
hellenensis, Chrysipus purpureifrons 139
Helleri, Ammosphecidium 241
hellmanni, Philanthus, *Anthophilus* 255
hellmanni, Salius 173
Hellus 145
helvetica, Amasis 32
helvetica, Ceropalae helvetica 160
helvetica, Mimesa 250
helveticae, Euura, *Pontania* 59
helveticus, Cryptocampus 56
helveticus, Miscophus 241
helveticus, Polistes 192
helveticus, Tachysphex 238
helvetius, Odynerus 186
helvola, Andrena, *Apis* 264
helvola, Apis 263
Hemibeleses 42
Hemibia 28
Hemiceropales subg. 160

Hemicalybion 214
Hemicroa 67
Hemiosmia subg. 298
Hemipogonius 162
Hemipterochilus 184
Hemisalius 166
Hemistenancistrocerus subg. 184
Hemitaxonus 83
Hemithyreopus 226
henschelii, Megachile 306
henschii, Pristiphora 74
henschii, Oryssus 117
henschii, Pseudoryssus, *Oryssus* 117
henschii, Xyela 21
Hensenia 214
hepaticae, Dineura 75
hepatimaculae, Pontania viminalis 56
Heptamelidae 36
Heptamelus 36
heptapotamiensis, Gorytes 222
Heptapotamius 42
hera, Nomada 314
heraclei, Allantus 96
heraclei, Crabro 232
herbaceae, Euura, *Nematus* 59
herculeanus, Camponotus, *Formica* 198
hercyniae, Amauronematus 61
hercyniae, Gilpinia, *Lophyrus* 33
heri, Eumenis mediterranea var. 182
Heriades 296
Heriades subg. 296
heringi, Pseudodineura, *Permatopus* 75
Heroecus 247
heroicus, Emphytus 39
heros, Ellampus 127
heros, Lestica, *Crabro* 234
herrichianus, Odynerus 191
herrichii, Odynerus 188
herrichii, Pseudepipona, *Odynerus* 188
herzi, Panurginus 276
herzi, Panurgus montanus 276
hesperia, Andrena 265
Hesperodryinus 120
Heterarthrinae 45
Heterarthrus 49
Heterelis subg. 147
Heterucera subg. 320
heterochromus, Tachysphex 238
Heterodasypoda 293
heterodoxa, Andrena 267
heteropenni, Xylocopa violacea ab. 310
Heterostelis subg. 302
hethiticus, Rophites algirus 291
hethiticus, Sphecodes rufiventris 290
Hexachrysis 133
Hexataenites subg. 277
heydeni, Ammophila 212
heydeni, Crossocerus 228
heydeni, Rhadinoceraea 52
hiaticula, Andrena humilis ssp. 265
hiatus, Emphytus 39
hiberniae, Megachile versicolor var. 307
hibernica, Megachile willughbiella var. 308
hibernica, Odynerus pictus var. 179
hibernicus, Nematus 73
iberus, Apis 275
hieroglyphica, Acantholyda,
Tenthredo 101
hierosolomita, Heriades 296
higashi, Halictus tumulorum 279
hikense, Anteon 119
hikosana, Andrena 270
hilaris, Pamphilus, *Lyda* 104
hillana, Apis 316
hilleckei, Thomsonia 100
himalaiensis, Busarbidea 76
himalayaensis, Anthophora
pulcherrima 324
Hinatara 49
hinganicus, Passaloecus 248
hipponensis, Nomada 314
hirashimai, Poecilagenia,
Poecilageniella 165
hircanus, Pompilus 164
hirowatarii, Euura, *Pachynematus* 59
hirsuta, Andrena 323
hirsuta, Chrysura, *Chrysis* 139
hirsuta, Myrmica 205
hirsuta, Nomada 312
hirsuta, Podalonia, *Sphece* 213
hirsutaffinis, Podalonia, *Ammophila* 214
hirsutifrons, Pompilus 169
hirsutilus, Phileremus 318
hirsutilus, Astatus 218
hirsutus, Sirex 112
hirta, Apis 293, 298
hirta, Scolia, *Apis* 148
Hirtanteon 118
hirtella, Andrena morawitzi var. 272
hirtellum, Trichiosoma 30
hirtellus, Halictus 287
hirticeps, Andrena 274
hirtipes, Andrena 293
hirtipes, Crabro 227
hirtipes, Dasypoda minor 293
hirtipes, Dasypoda, *Andrena* 293
hirtipes, Holopyga 132
hirtiventris, Halictus 282
hirtula, Myrmica 204
hirtula, Stenamma westwoodi var. 207
hirtus, Colletes 257
hirtus, Thyreus, *Crocisa* 327
hirundinaria, Apis 299
hispana, Eucera auct. 321
hispanica, Andrena nigriceps var. 265
hispanica, Andrena nitida var. 269
hispanica, Apis 240
hispanica, Bembix 222
hispanica, Prosopis difformis 261
hispanica, Stelis 302
hispanicum, Anthidium
florentinum var. 300
hispanicus, Alloodynerus 180
hispanicus, Colletes fodiens 258
hispanicus, Katamenes sichelii 185
hispanicus, Oxybelus 243

- hispanicus*, Tropidodynerus
 interruptus 192
- hispaniola*, Andrena nitida ssp. 269
- Hissarocephus* 110
- histrio*, Euura, *Nematus* 60
- histrio*, Nematus 60
- histrio*, Pamphilus 104
- histrio*, Tenthredo 100
- histrionica*, Andrena sillata ssp. 271
- histrionica*, Macrophyia 86
- histrionica*, Scolia, *Tiphia* 148
- histrionicus*, Thyreus, *Melecta* 327
- hoberlandti*, Eustenancistrocerus
 transitorius 184
- hoeppneri*, Nomada
 flavoguttata var. 312
- hoffmanni*, Lasioglossum, *Halictus* 283
- Hogardia* 224
- hohlbecki*, Cerceris 253
- hohlbecki*, Omalus, *Ellampus* 132
- hokkaidensis*, Pompilus 170
- hokkaidoensis*, Crossocerus
 annulipes 227
- hokkaidoensis*, Gilpinia 34
- hokkaidonis*, Tenthredo, *Allantus* 93
- Holandrena* subg. 267
- Holarctopompilus* 168
- Holcobleptes* 126
- Holcognema* 55
- Holcogneme* 55
- Holcognemis* 55
- Holepyris* 123
- hollandi*, Halictus 284
- holochlora*, Pedinomma
 rufescens var. 122
- Holochrysis* 139
- Hologambrus* 237
- holomelana*, Andrena 269
- holomelas*, Aphelopus 118
- Holonomada* 310
- Holopedina* 124
- Holopyga* 130
- holosericea*, Andrena 274
- Holotachytes* 240
- holosatica*, Megachile analis var. 307
- holosatica*, Vespa 195
- holtzi*, Halictus 277
- Homachthes* 319
- Homoeoneura* 83
- Homogambrus* 237
- Homoglenus* 123
- Homonotina** 166
- Homonotus* 161
- Homonotus* 166
- Homotus* 166
- hondoica*, Andrena 264
- hoodi*, Pristiphora 71
- hookeri*, Hylaeus 262
- Hoplammophila* 213
- Hoplandrena* subg. 267
- Hoplisoides* 224
- Hoplisus* 222
- Hoplitis* 296
- Hoplitis* subg. 297
- Hoplocampa* 67
- Hoplocampoides* 49
- Hoplocabro* 227
- Hoplocabron* 246
- Hoplopasites* 303
- Hoplosmia* subg. 298
- hoplostomum*, Dianthidium 301
- Horaia* 155
- horni*, Halictus 292
- hortensis*, Apis 277
- hortensis*, Nematus 65
- hortivaga*, Cerceris 253
- hortorum*, Bombus, *Apis* 328
- hortorum*, Pamphilus, *Lyda* 104
- hortorum*, Tenthredo 78
- hortulana*, Fesunella, *Tenthredo* 48
- horvathi*, Gonatopus 121
- horvathi*, Mesitius 124
- horvathi*, Philoctetes, *Ellampus* 132
- hospes*, Odynerus 178, 191
- hospitans*, Odynerus 191
- hospitus*, Emphytus 39
- hostia*, Tenthredo albiventris 96
- houskai*, Eumenes 182
- hoverlaensis*, Pristiphora 72
- huberti*, Eumenes 180
- hudsonicus*, Monophadnus 50
- hulteni*, Euura, *Amauronematus* 60
- humboldti*, Chrysura 140
- humboldti*, Pseudochrysis, *Chrysura* 140
- humboldti*, Cimbex 29
- humeralis*, Andrena 292
- humeralis*, Euura, *Nematus* 60
- humeralis*, *Hylotoma* 23
- humeralis*, Nematus 60
- humeralis*, Pseudotaxonus secundus 82
- humeralis*, Selandria 46
- humeralis*, Tenthredo 65
- humeralatum*, Aprosthemia, *Schizocera* 26
- humilis*, Andrena 265
- humilis*, Bombus 331
- humilis*, Pompilus 167
- hungaria*, Cilissa dimidiata var. 294
- hungarica*, Ammophila 212
- hungarica*, Astata 218
- hungarica*, Ceratina 310
- hungarica*, Empria, *Poecilosoma* 41
- hungarica*, Eucera 321
- hungarica*, Mutilla 157
- hungarica*, Nomada 314
- hungarica*, Nomia 292
- hungarica*, Pontania 67
- hungarica*, Stelis 302
- hungarica*, Tenthredo 100
- hungarica*, Tenthredopsis, *Tenthredo* 99
- hungarica*, Tetramorium caespitum 209
- hungaricum*, Tetramorium 209
- hungaricus*, Bembecinus, *Larra*
 hungaricus 221
- hungaricus*, Hylaeus, *Prosopis* 260
- hungaricus*, Mesitius 124
- hungaricus*, Nannodynerus 190
- hungaricus*, Sirex 113
- hungaricus*, Sphecodes 289
- hyacinthus*, Emphytus 39
- hyalina*, Conaspidia 84
- hyalina*, Messa 63
- hyalina*, Tenthredo 48
- hyalinialis*, Dosytheus 78
- hyalinata*, Priocnemis, *Sphex* 163
- hyalinatus*, Cephus 109
- hyalinatus*, Hylaeus 262
- hyalinatus*, Sphecodes 288
- hyalinella*, Andrena 268
- hyalinipennis*, *Odynerus* 188
- hyalinipennis*, Oryssus 117
- hyalinipennis*, Osmia 297
- hyalinoptera*, Aprosthemia 26
- hyalinopterus*, Cladius 54
- hyalinus*, Cabro 229
- hyalinus*, Eutomostethus 47
- hyalinus*, Pteronus 63
- hyalipennis*, Diodontus 247
- halopterus*, Cladius, *Priophorus* 54
- Hybonotus* 115
- hybrida*, Andrena 272
- hybrida*, Lasius 201
- hybrida*, Nomada ruficornis var. 315
- hybrida*, Osmia 297
- hybrida*, Tenthredo 41
- hybridus*, Cabro 233
- hybridus*, Sphecodes pellucidus 289
- hyemala*, Andrena 266
- hylaeiformis*, Colletes 258
- Hylaeinae** 257
- Hylaeinae** 259
- Hylaeus* 257, 259
- Hylaeus* subg. 259
- Hylocabro* 231
- Hylotoma* 22
- hylotomoides*, Tenthredo 68
- hymenaea*, Megachile 307
- Hypargyricus* 52
- hyperborea*, Ischyroceraea 96
- hyperborea*, Osmia 298
- hyperborea*, Pristiphora 74
- hyperboreus*, Bombus 327
- hypnorum*, Bombus, *Apis* 330
- hypobalius*, Nematus 70
- hypochlora*, Pedinomma
 rufescens var. 122
- Hypochrotaenia* 310
- Hypoclinea* 197
- Hypocrabro* 231
- hypocrita*, Bombus 327
- hypogastricus*, Nematus 69
- hypogastricus*, Nematus 70
- Hypolaepus* 69
- hypoleucus*, Nematus 66
- hypoleucus*, Pontania 67
- Hypomellinus* 224
- Hypomiscophus* 240
- hypopolia*, Andrena 274
- Hypoponera* 209
- Hypotaxonus* 43

hypotheticus, *Crabro cibrarius* 226
hypothrophica, *Lyda* 102
Hypothenemus 234
hypoxantha, *Euura, Nematus* 60
Hypsae, *Crabro* 232
hypsaes, *Ectemnius, Crabro* 232
Hypsiceraeus 160
hyrcanius, *Hylaeus* 261
hyrcanus, *Dolerus* 80
hystrix, *Andrena* 263

ianthe, *Fenusa* 42
iberica, *Andrena gallica* ssp. 269
iberica, *Dasylabris maura*
 nostras var. 153
iberica, *Mutilla* 157
iberica, *Nomada fucata* var. 313
ibericum, *Hedychridium* 142
ibericus, *Eumenes coarctatus* 181
ibericus, *Gorytes* 224
ibericus, *Tachytes* 239
iberobarbarus, *Leptochilus alpestris* 185
iberogallicus, *Ancistrocerus scoticus* 179
iberogallicus, *Eudrynerus dantici* 183
ibex, *Haliatus* 278
ibex, *Lindenius* 235
ibizensis, *Andrena flavipes* ssp. 275
Icalantica 159
Ichneumocheilus 161
Ichneumocheilus subg. 161
ichneumonea, *Apis* 277
ichneumonides, *Methocha* 149
ichneumonideus, *Ancistrocerus*,
 Vespa 179
ichneumoniformis, *Larra* 236
ichneumonoides, *Cryptocheilus*,
 Priocnemis 162
ichneumonoides, *Methocha* auct. 149
iconionus, *Evagetas* 171
Icronatha 150
Icteranthidium 301
icterica, *Apis* 267
icterica, *Tenthredo* 95
Icuma 224
idaei, *Tenthredo rubi* 26
identidem, *Pristiphora* 71
idiota, *Pristiphora* 71
idiotiformis, *Pristiphora* 71
Idolognathus 120
idolon, *Calameuta, Ichneumon* 109
idriensis, *Tenthredo* 85
ignava, *Macropha* 87
ignicollis, *Holopyga* 131
ignicollis, *Holopyga ovata* var. 131
ignifrons, *Chrysura, Chrysis* 139
ignita, *Chrysis, Sphex ignita* 135
ignita, *Sphex* 133
ignitus, *Bombus* 328
ignitus, *Cleptes, Ichneumon* 127
iiyamai, *Symmorphus* 191
ikonnikovi, *Bombus* 327
iliensis, *Andrena carbonaria* var. 272
iliensis, *Bembix* 221

ilinei, *Solenopsis* 207
illegalis, *Dasypoda* 293
illigeri, *Chrysis* 136
Illoswia 149
illyricum, *Lasioglossum* 286
illyricus, *Colletes* 258
imamiana, *Andrena carbonaria* var. 272
imayoshii, *Trypoxylon* 245
imbecilla, *Megachile* 306
imbecillus, *Omalus* 142
imitans, *Crabro* 229
imitans, *Formica* 200
imitator, *Macropha* 85
imitatrix, *Ceratina cyanea* var. 310
imitatrix, *Paraselandria* 81
immaculata, *Athalia rosae* 44
immaculata, *Chrysis* 136
immaculata, *Nomada* 314
immaculata, *Odynerus floricola* ab. 178
immaculata, *Stelis ornatula* var. 303
immaculata, *Tenthredo lachlaniana* 95
immaculatus, *Cephus* 109
immaculatus, *Hylaeus* 261
immaculatus, *Nematus* 64
immaculatus, *Oxybelus nigripes* 243
immaculatus, *Sirex antennatus* 113
immaculiventris, *Macropha*
 haemotopus 85
immaculosa, *Tenthredo moniliata* 95
immarginata, *Deuteragenia* 164
immarginatus, *Hylaeus* 285
immersa, *Empria, Tenthredo* 41
immundus, *Nematus* 65
immunis, *Cladus* 54
impatiens, *Oxybelus* 243
imperator, *Hylotoma* 25
imperatrix, *Chrysis* 143
impercepta, *Cerceris* 254
imperfecta, *Euura, Nematus* 60
imperfectus, *Tachysphex* 238
imperialis, *Nomada* 314
implicatus, *Ceropales* 160
impressa, *Chrysis* 136
impressa, *Tenthredo* 41
impressifrons, *Prospis* 262
impressus, *Sphecius* 224
impropria, *Anafenus, Entodecta* 45
impropria, *Entodecta* 45
impunctata, *Andrena* 267
impunctatus, *Colletes* 258
impunctatus, *Nematus* 56
impunctatus, *Sphecodes* 288
impura, *Tenthredo* 91
impurum, *Tetramorium, Myrmica* 209
inaequalis, *Allodynerus floricola* 178
inaequalis, *Chrysis inaequalis* 136
inaequalis, *Hylaeus* 261
inaequalis, *Schizocera* 26
inaequipunctata, *Chrysis* 136
inaffectata, *Tenthredo vespa* 97
inamoena, *Stelis scutellaris* 303
inanitus, *Pamphilus, Tenthredo* 104
Inbaltilla subg. 153
incana, *Andrena poupillieri* ssp. 274
incanus, *Nematus* 60
incerta, *Andrena* 264
incerta, *Chrysis analis* var. 135
incerta, *Colletes* 257
incerta, *Mutilla* 155
incerta, *Osmia* 298
incerta, *Tachytes* 238
incertus, *Dolerus* 79
incertus, *Nematus* 59
incertus, *Palarus* 243
incertus, *Sphecodes gibbus*
 ephippium 289
incertus, *Tachysphex, Tachytes* 238
incinctus, *Allantus* 91
incisa, *Andrena* 264
incisa, *Nomada* 314
incisoides, *Odynerus parietum* var. 179
incisum, *Anthidium variegatum* var. 300
incisus, *Odynerus parietum* var. 179
incisus, *Pachynematus* 72
Incoleipona subg. 183
incolorata, *Tenthredo* 23
incompleta, *Dasylabris sibirica* var. 154
incompleta, *Euura, Nematus* 60
incompleta, *Nomada ruficornis* var. 315
incompleta, *Prosopis* 260
incompletus, *Cephus pilosulus* 110
incompta, *Andrena austriaca* var. 267
incomptus, *Oxybelus* 242
incongruus, *Hylaeus* 262
inconspicua, *Andrena* 274
inconspicuus, *Nematus* 66
inconsueta, *Chrysis* 136
incrassata, *Pseudochrysis, Chrysis* 140
incurvata, *Myrmica* 205
inderskiana, *Nomada regalis* var. 311
indica, *Apis* 332
indica, *Scolia* 148
indicus, *Mesitius* 124
indicus, *Parvoculus, Mesitius* 124
indigena, *Andrena humilis* ssp. 265
indigena, *Mutilla* 157
indigotea, *Chrysis* 136
indobonensis, *Andrena* 272
indocilis, *Tenthredopsis raddatzi* 99
inerme, *Chelostoma* 296
inerme, *Tetramorium* 209
inermis, *Aporus* 174
inermis, *Coelioxys, Apis* 304
inermis, *Crabro* 228
inermis, *Dufourea, Halictoides* 291
inermis, *Osmia, Anthophora* 298
inermis, *Telostegus, Aporus* 174
inermis, *Tetramorium*
 caespitum var. 209
inxpectata, *Andrena hedikae* ssp. 263
inxpectatus, *Colletes* 258
infasciatus, *Emphytus* 38
infectum, *Anteon, Dryinus* 119
inferentia, *Poecilostoma* 43
infernalis, *Cephus* 108
infesta, *Ammophila* 212

infida, Lyda 105
infima, Tiphia 150
inflammata, Holopyga, *Ellampus*
 inflammata 131
inflammatus, Omalus
 sareptanus var. 132
inflatus, Coelioxys 304
inflatus, Nematus 64
inflexum, Sceliphron 214
infumata, Athalia, *Phyllotoma* 44
infumata, Macrophya 85
infumatus, Pachynematus 57
infundibuliformis, Sphecia 180
infuscata, Dasylabris maura
 nostras var. 154
infuscatus, Anoplius, *Pompilus* 172
infuscatus, Cephus 109
infuscatus, Colletes cunicularius 257
infuscatus, Emphytus 81
ingrica, Vespa 195
ingrica, Vespa norvegica saxonica
 morphe 195
ingricus, Crabro, *Thyreopus* 226
inhabilis, Selandria 47
inimicus, Vespa 221
ininterrupta, Sapyga 145
innominatus, Taxonus 39
innumerabilis, Odynerus 183
inocreata, Pristiphora 73
inopinatum, Trypoxylon 245
inopinatus, Liris 237
inornata, Pemphredon 248
inornata, Tenthredopsis raddatzi 99
inornatus, Crabro 228
inornatus, Crabro cribrarius 226
inornatus, Nematus, *Holcogneme* 69
inornatus, Tenthredopsis 99
inplana, Formica fusca picea var. 200
inquilina, Nomada 314
inquilinus, Monophadnus 51
insepultus, Polistes 193
insidiosa, Megachile
 circumcincta var. 307
insidiosus, Cleptes 127
insidiosus, Diodontus 247
insignata, Dasylabris siberica var. 154
insigne, Euura, *Nematus* 60
insigne, Pison 244
insignicornis, Ussurinus 100
insignis, Hylaeus 262
insignis, Passaloecus, *Pemphredon* 248
insignis, Pemphredon 247
insignis, Rhadinoceraea 47
insolata, Eumenes pomiformis f. 182
insolens, Vespa 178
insolita, Andrena 267
insperata, Chrysis 136
instabilis, Tenthredo 99
instratum, Aprosthemma, *Schizocera*
 instratum 26
insubrica, Apis 310
insubrica, Scolia 148
insubrica, Sphecia 147, 148

insubricus, Nematus 57
insueta, Mutilla marginata f. 157
insulana, Andrena colletiformis ssp. 264
insulare, Anthidium 301
insularis, Euura 62
insularis, Odynerus 190
insularis, Pachynematus 72
insularis, Physetopoda 155
insularis, Pristiphora 72
insularis, Vespa 193
insulicola, Odynerus 190
insulicolum, Lasioglossum, *Evyiaeus*
 baleicum 282
integer, Nematus ruficornis 74
integer, Philanthus 255
integra, Andrena 275
integra, Bembex 225
integra, Bembix 221
integra, Euura, *Pontania* 60
integra, Heriades 298
integra, Nomada 314
integrella, Chrysis 140
integrum, Anthidium 303
intercincta, Tenthredo, *Allantus* 98
interdichroa, Chrysura 143
interjecta, Chrysis interjecta 136
interjecta, Tenthredo 90
interjectus, Camponotus 198
intermedia, Pemphredon 247
intermedia, Agenia 164
intermedia, Andrena 272
intermedia, Andrena 274
intermedia, Anthophora 325
intermedia, Astata 218
intermedia, Blennocampa 46
intermedia, Cephalcia 102
intermedia, Dasylabris 153
intermedia, Dolichovespula,
 Vespula 194
intermedia, Fenusia 48
intermedia, Nomada 312
intermedia, Tenthredo 95
intermedia, Tenthredopsis nassata 100
intermedia, Thrinax 82
intermedia, Vespa sylvestris var. 194
intermedium, Aprosthemma,
 Schizocera 26
intermedium, Lasioglossum,
 Halictus 283
intermedius, Amauronematus 62
intermedius, Camptopoeum
 friesei var. 276
intermedius, Crabro 233
intermedius, Dolerus 80
intermedius, Monoctenus obscuratus 35
intermedius, Nomioides fasciatus 292
intermedius, Pachynematus 72
intermedius, Psen 252
intermedius, Sphecodes 288
intermixta, Anthophora
 acervorum var. 323
intermixta, Megachile 306
internalis, Eopompilus, *Sialus* 162
internalis, Sialus 162
interrogationis, Andrena 274
interrupta, Andrena 274, 275
interrupta, Cerceris, *Philanthus* 254
interrupta, Eucera 320
interrupta, Osmia 296
interrupta, Polistes 191
interrupta, Scolia 147
interrupta, Trachusa, *Apis* 303
interrupte-fasciata, Sphecia 231
interruptula, Andrena 275
interruptulus, Crabro 226
interruptum, Lasioglossum, *Hylaeus*
 interruptum 283
interruptus, Crabro 226, 233
interruptus, Halictus 279
interruptus, Megalodentes 107
interruptus, Nematus 62
interruptus, Nysson 219
interruptus, Nysson, *Mellinus* 220
interruptus, Oxybelus 243
interruptus, Polistes pamirensis var. 192
interruptus, Pterocheilus 189, 192
interruptus, Temnothorax, *Myrmica* 208
interruptus, Thyreopus 226
interruptus, Tropidodynerus,
 Polistes 192
interstinctus, Crabro 231
interstitiali, Selandria serva 81
interstitialis, Nematus 66
interstitialis, Selandria 81
intricans, Hoplisoides 224
intricatus, Crabro 230
inulae, Tetraloniella, *Tetralonia* 322
inventus, Tachysphex 238
inversus, Allantus 97
investiganda, Meria 149
investigator, Halictus 282
investigatoris, Halictus 282
invitabilis, Bombus ikonnikovi 327
invitus, Celonites abbreviatus 178
io, Chrysis 136
ione, Allantus 86
irakensis, Polistes nimpha 193
iranensis, Ancistrocerus 184
ireos, Anthophora, *Apis* 324
iridescens, Neurotoma, *Lyda* 103
iridifrons, Ectemnius, *Crabro* 233
iridipennis, Larra 237
iridipennis, Strongylogaster 82
iridis, Monophadnus 52
iris, Chrysis 136
iris, Elis 147
iris, Xylocopa, *Apis* 309
irkutensis, Allantus 93
irkutensis, Pteronus 75
irkutskiensis, Nomada 314
irradians, Euchroeus, *Pseudochrysis*
 purpuratus 140
irritans, Macrophya 84
irrogata, Selandria 45
irrorata, Lyda 102
ischiadica, Tenthredo 100

- Ischioceras* 152
Ischnolynthus 227
Ischyroceraea 89
ishiharai, Andrena 264
ishikawa, Megachile lapponica 307
ishikawai, Conaspidia 84
ishikawai, Evagetes 171
ishikawai, Symmorphus 191
islandica, Pontania 63
Isodontia 215
Isodyctium 51
isomera, Cladius 54
Isonotus 166
issikii, Nomada 314
italica, Dasypoda 293
italica, Mutilla 157
italica, Nomada 314
italica, Polistes 193
italica, Spinolia 149
italicus, Belomicrus 241
itelmena, Empria 41
itelmena, Euura, *Pontopristia* 60
itoi, Euura, *Pachynematus* 60
itoi, Pamphilus 105
Itycorsia 101
Itycorsia subg. 101
iucunda, Holopyga 129
iuncea, Vespa 179
Ivazuga 149
iwatai, Anoplus 172
iwatai, Dipogon 164
iwatai, Homonotus 166
iwatai, Passaloecus 248
iwatai, Pison 244
iwatai, Polistes rothneyi 192
iwatai, Symmorphus 191
Iwataia 231
izumindus, Sphecodes 288
- jacobi*, Andrena 267
jacobsoni, Cerceris 254
jacobsoni, Myrmica 205
jacoti, Tetramorium caespitum 209
jaculator, Astata 218
jacutensis, Tenthredo, *Allantus* 96
jacutica, Camponotus
 herculeanus var. 198
jacutica, Pristiphora 72
jagnobensis, Andrena 265
jailensis, Temnothorax, *Leptothorax* 208
jakobsoni, Hoplomerus 188
jakobsoni, Myrmecocystus
 altisquamis var. 199
jakovleffi, Trichiosoma 30
jakovlevi, Hartigia 111
jakovlevi, Hoplitis, *Pseudosmia* 297
jakovlevi, Trichiosoma 30
jakovlevi, Xiphydria 116
Jakovleviella 89
jakovlewii, Melecta 326
jakowleffi, Caenocephalus 108
jakowleffi, Caenocephalus 108
jakowleffi, Cerceris 254
- jakowleffi*, Dinax 40
jakowleffi, Nematus, *Holocneme* 69
jakowleffi, Tenthredopsis 99
jakowlevi, Messor rufitarsis 204
jakowlevi, Pristiphora,
 Lygaeonematus 72
jankowskii, Bombus 328
jankowskii, Parabatozonus,
 Pompilus 167
jankowskii, Pompilus 167
jankowskyi, Colletes 258
janssoni, Megachile
 centuncularis var. 308
Janus 110
janus, Pseudomalus 132
japanensis, Athalia spinarum 44
japonense, Trypoxylon fronticorne 245
japonensis, Dienoplus 224
japonensis, Nomada flavoguttata 312
japonibia, Andrena haemorrhoa ssp. 275
japonica, Abia 28
japonica, Aphaenogaster 203
japonica, Aphaenogaster schmitzi 203
japonica, Apis mellifera var. 332
japonica, Athalia, *Tenthredo* 44
japonica, Cephalcia lariciphila 102
japonica, Chrysis 136
japonica, Crocisa 326
japonica, Dasypoda 293
japonica, Formica 200
japonica, Formica fusca var. 200
japonica, Hylotoma 23
japonica, Jermakia 98
japonica, Lyroda 240
japonica, Lyroda nigra 240
japonica, Melitta 294
japonica, Mutilla 156
japonica, Neosyrista 111
japonica, Nomada 314
japonica, Pemphredon 248
japonica, Perineura 88
japonica, Ponera 209
japonica, Priocnemis 164
japonica, Selandria 82
japonica, Solenopsis 207
japonica, Solenopsis fugax var. 207
japonica, Tenthredo, *Allantus* 93
japonica, Tenthredopsis carinata 98
japonica, Vespa 193, 194, 196
japonicola, Halictus 281
japonicum, Chalybion, *Pelopaeus* 214
japonicus, Ancistrocerus, *Odynerus* 179
japonicus, Aneugmenus 76
japonicus, Camponotus 198
japonicus, Cimbex connatus 29
japonicus, Crabro alatus 234
japonicus, Discoelius 181
japonicus, Emphytus 87
japonicus, Goniozus 125
japonicus, Lasius 202
japonicus, Lasius emarginatus var. 202
japonicus, Lophyrus 34
japonicus, Macrocephalus 111
- japonicus*, Monoplopus 109
japonicus, Nematus, *Croesus* 69
japonicus, Nesodiprion, *Lophyrus* 35
japonicus, Odynerus 190
japonicus, Paracharactus,
 Neotomostethus 51
japonicus, Sphecodes 290
japonicus, Stigmus 250
japonicus, Stromboceros 82
japonicus, Tachysphex 239
jarkandensis, Halictus 278
jaroslavensis, Alysson fuscatus 219
jaroslawensis, Amauronematus 61
Jaroszewskyi, Crabro 226
jasnitskii, Nomada 313
jasnitzkii, Andrena 267
jasonia, Myrmilla 154
jaxartis, Chrysis 136
jaxartis, Chrysis sybarita var. 136
Jaynesia subg. 150
jeholensis, Coelioxys mandibularis 304
jemilleri, Nematus 61
jennyae, Myrmica 205
jensi, Lasius 201
Jermakia 89
jesoensis, Bombus 327
jesoensis, Dolerus 81
jessonicum, Rhopalum, *Crabro* 236
jezoensis, Xiphydria 116
joannis, Allantus scrophulariae 97
joergensenii, Pontania 56
johannae, Agonatopoides 121
johni, Apis 332
johni, Miscophus 241
johnsoni, Andrena jacobi var. 267
johnsonii, Calameuta 111
jonasi, Arge, *Hylotoma* 23
jonellus, Bombus, *Apis* 330
josephi, Thomsonia 99
jozana, Tenthredo, *Tenthredopsis* 90
jozana, Xiphydria 115
jozankeanum, Trichiosoma 31
Jucancistrocerus 184
Jucancistrocerus subg. 184
jucunda, Andrena 270
jucunda, Astata 218
jucunda, Dioxyx cincta 303
jucunda, Nomioides 292
jucunda, Osmia 298
jucunda, Siobla, *Encarsioneura* 89
jucundum, Hedychridium, *Holopyga* 129
jucundus, Odynerus 184
jucundus, Pamphilus 105
jucundus, Pamphilus, *Lyda* 105
judaica, Apterogyna 159
judaicus, Stenodynerus difficilis 190
jugalis, Allantus 92
jugicola, Euura, *Nematus* 60
juglans, Tomostethus 53
Jugurtha, Larra 237
juliae, Solenopsis, *Diplorhoptrum* 207
julliani, Xyela, *Pinicola* 21
julliani, Epeolus 317

julliani, *Tachysphex* 238
 julliani, *Tetraloniella*, *Tetralonia* 322
junci, *Dosytheus* 77
Juncilerus 77
juncivorus, *Tomostethus* 47
junipereti, *Leptothorax* 208
juniperi, *Monocetus*, *Tenthredo* 35
juniperi, *Tenthredo* 35
jurassica, *Nomada piccioliana* 315
jurinae, *Cimbex* 32
jurineanum, *Anteon* 118, 119
jurinei, *Holopyga* 131
jurinei, *Vespa* 194
Jurini, *Pison* 244
jurinii, *Larra* 240
Jurja 149
juvenculus, *Colletes* 258
juvencus, *Ichneumon* 112
juvencus, *Sirex*, *Ichneumon juvencus* 112
juvenilis, *Tenthredo* 83

kabyliana, *Macrophyia rustica* 86
kaiensis, *Nomada okamotonis* 315
kaiensis, *Sphecodes* 289
Kaliofenusa 48
Kaliofenusa subg. 48
Kaliostosphinga 48
kallipygus, *Tachysphex incertus* 238
Kalloombus subg. 328
kamafuse, *Sphecodes* 290
kamensis, *Camponotus marginatus* var. 198
kamikochensis, *Pamphilus* 105
kamikochiana, *Andrena* 271
kamtchatica, *Phyllotoma* 49
kamtchatica, *Pontopristia* 56
kamtchatica, *Pristiphora* 72
kamtchatica, *Pteronidea* 56
kamtchatica, *Thrinax*, *Hemitaxonus* 83
kamtchaticus, *Dolerus* 78
kamtchaticus, *Nematinus* 68
kamtchaticus, *Taxonus* 43
kamtschatica, *Ammophila sabulosa* 212
kamtschatica, *Andrena apicata* var. 263
kamtschatica, *Myrmica* 205
kamtschaticus, *Leptothorax acervorum* 204
kamtschatica, *Andrena* 264
kamtschatkensis, *Dolichovespula saxonica* 195
kamtshatica, *Xyela* 21
kamtshaticus, *Bombus lapponicus* 330
kamyschiensis, *Myrmica* 205
kankaucharis, *Lasioglossum*, *Halictus* 283
kanoi, *Tenthredella alboannulata* 93
kanshireana, *Crocisa* 326
kansuense, *Lasioglossum*, *Halictus* 280
kantarae, *Halictus pallens* 281
karafutonis, *Chrysis* 136
karafutonis, *Emphytus* 37
karafutonis, *Eumenes* 182
karafutonis, *Pompilus* 167

karaginus, *Bombus lapponicus* 330
karavajevi, *Myrmica*, *Symbiomyrma* 206
karavajevi, *Sifolinia* 206
karawajewi, *Plagiolepis* 202
karawajewi, *Strongylognathus* 207
kareli, *Allantus* 90
karelica, *Andrena gelriae* 274
karelicus, *Symmorphus* 191
karenkona, *Vespa* 196
karvoneni, *Lygaeonematus* 70
karvoneni, *Pteronidea* 60
kasakorum, *Messor* 204
kaschiriensis, *Mutilla* 157
kasczenkoi, *Myrmica* 206
kasczenkoi, *Myrmica scabrinodis* 206
kashgarensis, *Megachile* 308
kashmirensis, *Astata kashmirensis* 218
kasparyani, *Arachnospila* 169
kasparyani, *Holepyris* 123
kasparyani, *Lasioglossum* 280
kastsheevi, *Xiphydria* 116
kaszabi, *Ammoplanus* 246
kaszabi, *Ancistrocerus ichneumonideus* 179
kaszabi, *Arachnospila* 169
kaszabi, *Cerceris* 254
kaszabi, *Colletes* 258
kaszabi, *Dasylabris* 153
kaszabi, *Diodontus* 247
kaszabi, *Lindenius* 235
kaszabi, *Oryttus* 224
kaszabi, *Parapiagetia* 237
kaszabi, *Philanthus* 255
kaszabi, *Poecilotiphia*, *Meria brevicauda* 150
kaszabi, *Polistes missus* 193
kaszabi, *Pterocheilus phaleratus* 189
kaszabi, *Rhogogaster* 91
kaszabi, *Stizus* 225
kaszabi, *Temnothorax*, *Leptothorax* 208
kaszabi, *Tenthredo* 91
Katamenes 184
Kathepyris 123
katkovi, *Alysson* 219
katuniensis, *Formica rufibarbis* var. 200
Kaufmannia 224
kawabatai, *Rhopalum* 235
kawakami, *Xiphydria* 115
kazakhstanicus, *Odynerus albopictus* 187
kazbekianus, *Ancistrocerus Odynerus* 179
kazenasi, *Cerceris* 253
Kelidoptera 103
kengracensis, *Andrena* 275
kennedii, *Alyson* 219
keriensis, *Bombus* 329
kerketa, *Tenthredo* 96
kerriae, *Andrena* 273
kervilleanus, *Halictus* 284
kerzhneri, *Lasioglossum* 280
kessleri, *Halictus* 279
khabarovii, *Andrena* 267

khambana, *Anthophora* 324
khangmarensis, *Eumenes septentrionalis* 182
khankensis, *Andrena* 265
khasanensis, *Colletes cunicularius* 257
hasania, *Andrena* 272
hasianus, *Crabro* 233
kiangsensis, *Eumenes* 182
kiutschouense, *Lasioglossum Halictus* 283
kichizoi, *Masaakia* 49
kiefferi, *Allantus* 90
Kiesenwetteri, *Crabro* 235
kinburni, *Tapinoma* 198
kincaidi, *Monophadnoides* 50
Kincaidia 83
kindenmani, *Anthidium reticulatum* 302
kiotonis, *Aneugmenus* 76
kirbienus, *Epeolus* 318
kirbii, *Ammophila* 215
kirbii, *Prionyx*, *Ammophila* 215
Kirbya 294
kirbyanus, *Phileremus* 319
kirbyella, *Nomada* 311
kirbyi, *Andrena* 267
kirbyi, *Nematus* 57
kirchneri, *Pontania* 57
kirghisia, *Thyreomelecta* 326
kirgisica, *Ammophila* 213
kirgisica, *Bembix* 222
kirgisica, *Colletes* 258
kirgisica, *Myrmica rugosa* var. 205
kirgisicum, *Lasioglossum* 284
kirillovi, *Leptothorax tuberum* 208
kirishimensis, *Ametastegia* 39
kiritschenkoi, *Amblyellus Sophropompilus* 168
kiritschenkoi, *Myrmilla* 154
kiritschenkoae, *Smicromyrme* 156
kirschaumi, *Halictus* 283
kirschaumi, *Hylaeus* 261
kissi, *Anthidium florentinum* var. 300
kitamius, *Sphecodes* 288
kizilkumi, *Philanthus* 255
kizilkumii, *Dicyrtomellus*, *Pompilus* 170
kizilkumii, *Stizus* 225
Kjellia 40
klausnitzeri, *Monophadnoides Monophadnus* 50
klugi, *Anthidium* 301
klugi, *Anthophora* 322
klugi, *Diprion pini* 33
klugi, *Macrophyia* 86
klugi, *Nematus* 64
klugii, *Colpa*, *Scolia* 147
klugii, *Emphytus* 40
klugii, *Hylotoma* 24
klugii, *Lyda* 102
klugii, *Macrophyia* 85
klugii, *Nematus* 61
klugii, *Pterocheilus* 189
klugii, *Selandria* 41
klugii, *Tarpa* 107

klutchianus, Dolerus 78
klutschianis, Pratobombus
 hypnorum 330
knipovitschi, Temnothorax,
_{Leptothorax} 208
knipovitschi, Leptothorax tuberum 208
 kobayashii, Arge 23
kobensis, Colletes 258
kodorica, Camponotus vagus var. 198
koebelei, Cephalcia, *Cephaleia* 102
koebelei, Pteronus 62
koebelei, Stromboceros 82
koehleri, Tenthredo 90
koenigi, Tenthredo 95
kohatensis, Andrena
 cussariensis var. 268
kohli, Allantus 93
kohli, Crossocerus, Crabro 229
kohli, Diodontus 247
kohli, Mimesa 251
kohli, Nomada 314
kohli, Polistes 192
kohli, Psen 251
kohlii, Agenioideus, *Pompilus* 167
kohlii, Anthidium 301
kohlii, Gorytes 222
kohlii, Psen 251
koikense, Trypoxylon 245
koikensis, Nomada 314
koikensis, Sphecodes 288
Kokarevta 147
kokpeticus, Odynerus 183
kokuewi, Tenthredopsis 99
kokujevi, Myrmilla 154
kokujewi, Dolerus 80
Kokujewia 25
kolazyi, Chrysis 136
kolazyi, Trypoxylon 245
Kolazyi, Trypoxylon 245
Kollaris, Crabro 233
kolthoffi, Arge 22
kolymensis, Euura, *Nematus*
 olfaciens 62
koma, Chrysura, *Chrysis* 139
koma, Trypoxylon 245
koma, Trypoxylon figulus 245
komarovi, Stizus 225
komarowii, Paravespa 187
konakovi, Bombus 332
kongauana, Tenthredo 93
kongauensis, Arge 23
kongosana, Macrophyia 85
konoi, Selandria 81
konosuense, Trypoxylon 245
konowi, Alphastromboceros,
_{Strongylogaster} 75
konowi, Amauronematus 67
Konowi, Crabro 233
konowi, Empria 42
konowi, Lyda 105
konowi, Nematus, *Pteronus* 75
konowi, Pelmatopus mentiens 75
konowi, Perineura 100

konowi, Pteronidea ribesii 64
konowi, Strongylogaster 75
konowi, Tenthredo 93
konowi, Tenthredopsis 99
Konowia 115
konowiana, Nomada jacobaeae var. 313
kontkaneni, Pachynematus 73
Kontuniemiana 55
kontuniemii, Pamphilus 105
Koptodynerus 190
Koptogaster subg. 260
korabica, Tenthredo 96
koraiensis, Xyela 21
korbi, Lasioglossum, *Halictus* 280
koreana, Birka, *Nesoselandria* 76
koreana, Corymbas 84
koreana, Macrophyia 86
koreana, Mesoneura 68
koreana, Myrmica 206
koreana, Nitela 241
koreana, Senoclidea, *Monophadnus* 53
koreanum, Lasioglossum 281
koreanum, Polochridium 145
koreanum, Trypoxylon 245
koreanus, Diprion 33
koreanus, Gorytes laticinctus 222
koreanus, Lasius 202
koreanus, Passalocetus 248
koreanus, Polistes rothneyi 192
koreanus, Psen 251
koreense, Pison, *Paraceramius* 244
koreensis, Vespula, *Vespa* 196
korjak, Bombus lapponicus 330
korlevici, Tenthredopsis 98
korleviciana, Nomada errans var. 312
koropokkrus, Bombus 330
korotnevi, Megachile 308
kosensis, Halictus 284
koshevnikovi, Cardiocondyla 203
koshevnikovi, Pheidole pallidula 207
kostjuki, Gorytes 224
 kostylevi, Ancistrocerus trifasciatus 180
 kostylevi, Eumenes 182
 kostylevi, Trypoxylon 245
 kotenki, Lonchodryinus 119
 kotenki, Tenthredo 98
 kotokui, Myrmica 206
 kotokui, Myrmica rubra 206
 kotschy, Andrena thoracica ssp. 269
 kouchkiana, Eucera 321
 Koxinga 234
 kozakorum, Myrmica 206
 kozarowi, Andrena 264
 kozhantshikovi, Hedychrum 130
 kozhantshikovi, Psilglossa
 odyneroides 188
 kozhevnikovi, Didineis 219
 kozlovi, Andrena callopyrrha ssp. 274
 kozlovi, Colletes 258
 kozlovi, Dasylabris 153
 kozlovi, Gorytes 222
 kozlovi, Lasioglossum, *Halictus* 284
 krajniki, Hedychrudium krajniki 129
 kraloffi, Halictus 281
 krausi, Euura, *Amauronematus* 60
 kraussei, Andrena 275
 kriebaumeri, Hylaeus 261
 kriegeri, Halictus 284
 kriegeri, Pteronus 57
 krkiana, Chrysis grohmanni 135
 Krombeinella 152
 krombeini, Incertosulcus 124
 krombeini, Pemphredon 248
 Krombeiniellum 244
 kropotkini, Panurginus 276
 krylovi, Pratobombus modestus 330
 kuangsiana, Vespa analis var. 193
 kubesni, Pachynematus 61
 kuccharonis, Xiphydria 116
 Kudakrumiini 152
 kudiana, Andrena 267
 kudiana, Coelioxys 305
 kudiana, Tenthredo 93
 kudianus, Rhogogaster 88
 kudiensis, Colletes 258
 kudiensis, Megachile willughbiella 308
 kuehlhorni, Alastor 178
 kuehlhorni, Smicromyrme 156
 kuennemannii, Megachile 307
 kullenbergi, Eucera 320
 kumamotonis, Onycholyda, *Lyda* 103
 kunashirensis, Priocnemis 163
 kunasirensis, Euura, *Nematus* 60
 kuntzei, Dioxyx 303
 kunugi, Apethymus 40
 kupyanskaya, Formica 199
 kuramaensis, Crabro reiteri 234
 kurbati, Megachile lapponica var. 307
 kurdica, Aphaenogaster 203
 kurdica, Aphaenogaster
 subterranea var. 203
 kurdistanica, Andrena 271
 kurentzovi, Arachnospila 169
 kurentzovi, Symmorphus 191
 kuri, Apethymus 40
 kurilarum, Pachyprotasis longicornis 87
 kurilese, Episyron 167
 kurilese, Stenamma 207
 kurileensis, Bombus tersatus 329
 kurileensis, Nipponodipogon,
_{Dipogon} 164
 kurileensis, Nomada 312
 kurilensis, Temnothorax,
_{Leptothorax} 208
 kurilensis, Tenthredo, *Tenthredella* 94
 Kuriloagenia 164
 kurisuei, Okomotonius 30
 kuro, Nomada 314
 kuroiwae, Athalia colibri 44
 kurokii, Myrmica 206
 kurokii, Myrmica rubra 206
 kuroo, Diodontus 247
 kurzenkoi, Arachnospila 169
 kurzenkoi, Epyris 123
 kurzenkoi, Mimesa 250
 kusdasi, Myrmilla capitata 154

- kushkia*, Andrena hyemala ssp. 264
kussariense, Lasioglossum, *Halictus* 280
kussariensis, Allantus 96
kuwayamai, Arachnospila, *Pompilus* 169
kuwayamai, Bombus schrencki 332
kuwayamai, Rhopalum 236
kuzini, Calameuta 109
kuznetsovi, Philoctetes, *Ellampus* 132
kuznetzovi, Cerceris 254
kuznetzovi, Chrysis 141
kuznetzovi, Empria 41
kuznetzovi, Smicromyrme 155
kuznetzovorum, Pristiphora,
Lygaeonematus 72
kychtacensis, Megachile 303
kyutekparki, Pamphilus 105
- Labeo* 120
Labeola 120
Laberinus 120
Laberius 120
labialis, Andrena 265
labialis, Andrena, *Melitta* 267
labialis, Selandria 41
labiata, Andrena 271
labiata, Andrena 272
labiata, Apis 287
labiata, Cerceris 254
labiata, Macropis auct. 293
labiata, Megilla 293
labiata, Tenthredo 85, 98
labiatarum, Nomiooides 292
labiatula, Andrena 267
labiatus, Mellinus 262
labiatus, Panurginus, *Panurgus* 276
Labidia 89
labiosus, Dolerus 78
laborans, Odynerus 190
laboriosa, Hoplitis, *Osmia* 296
labratus, Nematus pallicornis 70
labrosa, Andrena 291
labrosus, Halictus 283
lacerticida, Parabatozonus, *Sphex* 167
lachesis, Melecta 326
lachlaniana, Tenthredo 95
lacinata, Caliroa 45
lacordairei, Hoplitis 223
lacrymosa, Tenthredo 85
lacrymosa, Tenthredo 85
lactarius, Thyreopus 226
lacteipennis, Pachynematus 56
lacteore, Aglaostigma aucupariae 83
lacteus, Nematus 63
lactiflua, Tenthredopsis, *Tenthredo* 99
lactipennis, Panurginus 276
lactipennis, Sphecodes 288
ladakhana, Anthophora parietina 324
Laelius 123
laesus, Bombus 331
laeta, Euura, *Cryptocampus* 60
laeta, Nomada 311
laeta, Sciapteryx 89
laeta, Scolia 148
- laeta*, Scolioneura 48
laeta, Tenthredo 32, 92
laetum, Pararhynchium paradoxum 187
laetus, Amauronematus viduatus 66
laetus, Crabro 232
laetus, Nematus 70
laetus, Pterocheilus 188
laeve, Lasioglossum, *Melitta* 284
laeviceps, Crabro 228
laeviceps, Pachyprotasis 87
laeviceps, Pemphredon 249
laeviceps, Tenthredo rossica 90
laevicollis, Pachyprotasis 88
laevifrons, Ceratina 310
laevifrons, Colletes 258
laevifrons, Epeolus 317
laevifrons, Halictus 280
laevifrons, Hoplitis, *Osmia* 297
laevifrons, Megachile 306
laevifrons, Osmia 296
laevifrons, Priophorus 55
laevigata, Chrysura, *Chrysis* 139
laevigata, Prosopis 260
laevigatum, Lasioglossum, *Melitta* 280
laevigatus, Ectemnius 232
laevigatus, Nematus 56
laevigatus, Oxybelus 242
laevigatus, Psenulus, *Psen* 252
Laevigorytes 222
laevinodis, Myrmica 206
laevipes, Gymnomerus, *Odynerus* 184
laevipes, Odynerus 184
laevis, Harpactus, *Mutilla laevis* 223
laevis, Lindenius 235
laevis, Nematus 66
laeviuscula, Andrena 267
laeviuscula, Formicoxenus 204
laeviusculus, Hylaeus 283
laeviventris, Odynerus 191
laevoidea, Lasioglossum 284
Lagidina 84
lagodechii, Chrysis 134
Lagonis 49
lagopoda, Megachile, *Apis* 307
lagostae, Odynerus dantici var. 183
lagrecai, Nomada 314
lama, Bembix 221
Lambdopsis subg. 260
lamellatus, Oxybelus 242
laminata, Cerceris 253
lampadum, Hedyrhridium 142
lamprechtii, Dolerus 77
Lamproapis 310
lamprosoma, Spinolia 141
lamprosoma, Spinolia, *Chrysis* 141
lanata, Chrysis 136
lanata, Xylocopa 310
lanatae, Euura 60
lanatus, Tachysphex 239
lanceata, Euura, *Amauronematus* 60
lanceolata, Chrysis 136
lanceolata, Nomada 311
lanceolatus, Coelioxys 305
- lanceolatus*, Metallus, *Blennocampa* 50
langei, Aglaostigma, *Rhogogastera* 83
languida, Tenthredopsis, *Tenthredo* 99
laniensis, Euodynerus disconotatus 183
lanifica, Pristiphora, *Nematus* 72
lanifrons, Melitta 265
laniger, Apis 293
lanigera, Mesoneura 68
lanionis, Chelogynus 119
lanuginosus, Gastrosericus 236
lapicida, Sphecodes 180
lapidarius, Bombus, *Apis* 329
lapidarius, Crabro 234
lapidarius, Ectemnius, *Crabro* 231
laportei, Crabro 232
lappo, Euura 60
lapponica, Andrena 264
lapponica, Eumenes
pedunculatus var. 182
lapponica, Euura 56
lapponica, Megachile 306, 307
lapponica, Pontania 58
lapponica, Rhogogaster viridis 88
lapponica, Tenthredo 39, 73
lapponicola, Pontania 58
lapponicus, Bombus, *Apis* 330
lapponicus, Crabro 226
lapponicus, Pachynematus 71
Lara 236
Larandrena subg. 267
largipes, Tenthredo 69
laricicola, Pristiphora 71
lariciphagus, Anoplonyx, *Nematus* 53
lariciphagus, Nematus 53
lariciphila, Cephalcia, *Cephaleia*
lariciphila 102
laricis, Acantholyda, *Lyda* 101
laricis, Gilpinia, *Pteronus* 34
laricis, Pristiphora, *Nematus* 72
laricivorus, Nematus 72
laricivorus, Pachynematus 72
Larinematus 55
Larra 236
Larra subg. 236
Larrada 236
Larrana 236
Larraxena 236
Larrini 236
Larropsis 236
Larropsis subg. 236
larvata, Podalirius 324
larvatus, Crabro 232
Lasanthidium 301
laserpitii, Tenthredo 86
Lasianteon 118
Lasioglossum 279, 287
Lasioglossum subg. 280
Lasius 201
Lasius 323
Lasius subg. 202
latebrosus, Crabro 235
lateluteus, Allantus amaenus 97
lateniger, Allantus amaenus 97

- laterale*, Icteranthidium, *Anthidium* 301
lateralis, Allantus 96
lateralis, Amauronematus 65
lateralis, Camponotus, *Formica* 198
lateralis, Corynis, *Amasis* 32
lateralis, Dolerus 78
lateralis, Nematus 71
lateralis, Nomada 312
lateralis, Tenthredo 94
Lateralisiformis 31
laterater, Dolerus pratensis 78
laterimaculatus, Cephus gracilis 109
lateritia, Andrena hystrix var. 263
lateritia, Apterogyna 159
lateritia, Tenthredo 77
lathamana, Melitta 265
lathburiana, Nomada, *Apis* 314
lathyri, Andrena 274
latialis, Tarpa plagiocephalia 107
laticarpus, Macrophyia 83
laticauda, Epeolus 317
laticaudatus, Sphecodes 288
laticeps, Brachyipona, *Odynerus* 180
laticeps, Lasioglossum, *Halictus*
 laticeps 284
laticeps, Osmia 298
laticeps, Pseudisobrachium 123
laticeps, Tachysphex 238
laticeps, Thomsonia 98
laticincta, Halictus rubicundus 278
laticincta, Osmia 298
laticinctus, Allantus 76, 94
laticinctus, Gorytes, *Euspongus*
 laticinctus 223
laticorne, Rhopalum, *Crabro* 235
laticrus, Nematus 69
laticrus, Nomada 314
latidens, Oxybelus 243
latifrons, Hoplisoides, *Gorytes* 224
latifrons, Pamphilus, *Lyda* 105
latifrons, Prosopigastra 237
latifrons, Tachysphex 239
latifrons, Tachytes 240
latilineata, Vespa magnifica var. 194
latipalpis, Pterochelius 187
latipes, Eumenes 185
latipes, Nematus 69
latiserra, Euura, *Pontopristia* 60
latistriga, Nomada lineola var. 313
latisulcata, Euura, *Amauronematus* 60
latitarsis, Lindenius 235
lativalvis, Tachytes 239
lativentre, Lasioglossum, *Hylaeus* 280
lativentris, Pristiphora, *Nematus* 72
lativentris, Tiphia 150
latizona, Tenthredo 92
latreillei, Sphecodes 287
latreillii, Anthidium 301
latreillii, Bembix 222
latreillii, Myrmecina 205
latreillii, Trichiosoma 30
latro, Oxybelus 242
latrooides, Solenopsis orbula var. 207
latronis, Melitta 294
latronum, Rhopalum, *Crabro* 236
Latrorhopalum 235
Latroxybelus 242
latus, Monophadnus 53
latus, Odynerus melanocephalus 187
latus, Taxoblenus, *Taxonus* 43
laudata, Caliroa 45
Laurentia 83
Laurentina 83
Lautara 241
lautus, Allantus 97
lavandulae, Andrena 267
Lavia 221
lavrushini, Hylaeus 261
lazoiana, Andrena 270
leachella, Megachile 306
Leachia 55
leachii, Chrysis 136
leachii, Hylotoma 24
leachii, Nematus 72
leaiana, Osmia, *Apis* 298
lebanensis, Ammobatoides 318
lebedewi, Colletes 258
lecana, Andrena niveata ssp. 270
lecerfi, Andrena 274
leclercqi, Crossocerus 231
lecontei, Ancistrocerus 178
lecontei, Lophyrus 34
lederi, Allantus 96
lederi, Andrena 268
lefebvre, Urocerus 113
lefebvrei, Crossocerus 227
lefebvrei, Megachile, *Chalicodoma* 305
lefebvrei, Polistes gallica var. 193
legirupus, Polvynematus 56
lehri, Bifidoceropales 160
lehri, Deuteragenia 164
Leiocleptes 126
Leiogorytes 222
leonota, Euura, *Pteronidea* 60
leleji, Ancistrocerus 179
leleji, Andrena 272
leleji, Epyris 123
leleji, Halictus, *Seladonia* 279
leleji, Hedychrnidium 129
leleji, Hylaeus 260
leleji, Pamphilus 105
leleji, Tenthredo 95
Lelejola 125
lemani, Formica 200
lenensis, Dolerus scoticus 78
lenis, Andrena 270
lenis, Aphanus 50
lentus, Pachynematus 57
Leocrabro 231
leoninus, Halictus 282
Leonomyra 203
lepeletieri, Andrena 269, 270
lepeletieri, Anthidium 301
Lepeletieri, Crabro 230
lepeletieri, Macrophyia 86
lepeletieri, Nomada 311
lepeletieri, Odontodynerus
 deflendus 180
levida, Andrena 273
levida, Anthophora 324
levida, Tenthredo 42
levida, Tenthredo 81
Lepidandrena subg. 268
lepidotus, Nematus 66
lepidula, Chrysis sardarica 137
lepidulus, Hylaeus 260
lepidus, Harpiphorus, *Tenthredo* 42
lepidus, Nematus 62
Lepisiota 202
leporina, Melitta, *Apis* 294
Leptepipona 188
lepticus, Strongylogaster
 desbrochersi 82
leptocephala, Euura, *Nematus* 60
leptocephalus, Hylaeus, *Prosopis* 260
Leptocerca 67
Leptocercus 67
leptocerus, Nematus 66
Leptochilus 185
Leptochilus subg. 185
Leptocimbex 30
Leptolarra 237
leptomandibularis, Chrysis 136
Leptopolistes 192
Leptopriocnemis subg. 162
Leptopus 69
Lepto thorax 204
lerouxi, Halictus 278
leseleuci, Sirex 112
leskii, Cephus 109
Lestica 234
Lestiphorus 224
Lestodryinus 120
Lestophorus 224
lethe, Euura 60
lethierryi, Pamphilus, *Lyda* 105
lethifer, Pemphredon, *Cemonus* 248
leucaheneus, Halictus leucaheneus 279
Leucandrena subg. 268
leucaspis, Euura, *Nematus* 60
Leucempria 40
Leuchimon 170
leucobasis, Dolerus 80
leucocarpus, Nematus 57
leucocephala, Chrysolyda,
 Pamphilus 103
leucocephalus, Pamphilus 102
leucocera, Tenthredo 94
leucocnemis, Nematus 72
leucodus, Nematus 61
leucogaster, Nematus 66
leucognathus, Halictus 278
leucogona, Tenthredo 42
leu colena, Euura, *Nematus* 60
leucomalla, Megachile 306
leucomelana, Hoplitis auct. 296
leucomelana, Hoplitis, *Osmia* 296

leucomelas, Tenthredo 87
leuconotus, Sphecius 215
leuconotus, Sphecius 216
leucophthalma, Nomada, *Apis* 314
leucopoda, Aproceros 25
leucopoda, Macrophyia 84
leucopoda, Phyllotoma 49
leucopodia, Pristiphora, *Nematus* 72
leucopodus, Dolerus 39
leucopogon, Halictus 278
leucopterus, Dolerus 80
leucopum, Lasioglossum, *Melitta* 284
leucopus, Pristiphora, *Nematus* 72
leucopus, Tenthredo 86
leucopyga, Euura, *Amauronematus* 60
leucopygus, Halictus 283
leucorhina, Andrena auct. 275
leucorhinum, Dianthidium 300
Leucostelis 302
leucosternus, Allantus 93
leucosticta, Euura, *Nematus* 61
leucostoma, Apis 281
leucostoma, Athalia 44
leucostoma, Crabro 228
leucostoma, Crossocerus, *Sphecius* 228
leucostoma, Lygaeonematus 73
leucostoma, Tenthredo 95
leucostomoides, Coelocrabro 228
leucostomus, Emphytus 38
leucostomus, Gonatopus 121
leucotrocha, Euura, *Nematus* 61
leucozonias, Tenthredo 43
leucozonica, Cerceris 253
leucozonium, Lasioglossum, *Apis* 281
leucurus, Pompilus 173
leueopogonata, Chalicodoma 305
leueoura, Dasypoda 293
levigatus, Halictus 280
levilabris, Andrena 275
levipes, Crabro 230
levis, Halictus 284
lewinella, Melitta 273
lewisi, Smicromyrme 156
lewisi, Abia 28
lewisi, Dolerus 81
lezginica, Myrmilla, *Mutilla* 154
libanensis, Trachelus, *Cephus* 111
libanicus, Eumenes
 unguiculatus var. 181
libanicus, Symmorphus elegans 191
liberta, Athalia, *Tenthredo* 44
liburnica, Nomada alboguttata var. 311
lichata, Andrena bimaculata ssp. 272
lichtwardti, Euura, *Pachynematus* 61
lichtwardti, Tenthredo 94
liciata, Macrophyia 84
lidmilae, Smicromyrme rufipes f. 156
lienterica, Euura, *Nematus* 61
ligata, Tenthredopsis 99
ligeris, Croesus varus 69
ligniperda, Camponotus, *Formica* 198
ligniperda, Formica 198
ligniseca, Megachile, *Apis* 307

lignisecula, Megachile 307
ligustrina, Tenthredo 85
llipit, Halictus 284
llipitianus, Tachysphex 237
limacina, Tenthredo 45
limanica, Myrmica rugulosa 205
limassolica, Dioxoides tridentata 303
limata, Andrena 269
limbata, Andrena 273
limbata, Tenthredo 43, 95
limbatus, Euchroeus limbatus 140
limbellum, Lasioglossum, *Halictus* 284
limbiferum, Anthidium 301
limbiferus, Allantus 92
limbiferus, Leptochilus, *Odynerus* 185
limbalbris, Tenthredopsis 98
limbonotatus, Odynerus 190
limissicus, Eumenes pomiformis 181
limissicus, Halictus pollinosus 279
limitaris, Pachyprotasis 87
limonii, Andrena 264
lindbergi, Sphecodes alternatus 287
lindemii, Odynerus 183
Lindenius 234
Lindenius, Crabro 233
lindqvisti, Euura, *Amauronematus* 61
Lindqvista 55
linearis, Phylloecus, *Tenthredo* 110
linearis, Tenthredo 82
lineare, Lasioglossum, *Halictus* 284
lineata, Dryudella 218
lineata, Meria auct. 149
lineata, Nomada 242
lineata, Smicromyrme variabilis f. 155
lineata, Tenthredo 82
lineatotarsis, Crabro 231
lineatus, Oxybelus, *Nomada* 242
lineicoxis, Pachyprotasis 87
Lineobirka 76
Lineobirka subg. 76
lineola, Nomada 313
lineola, Osmia 297
lineolata, Athalia 44
lineolata, Nomada succineta var. 313
lineolata, Periclista, *Tenthredo* 51
lineolata, Tenthredo 51
lineolata, Vespa media var. 195
lineolatus, Halictus 281
lineolatus, Hylaeus, *Prosopis* 261
lineolatus, Nysson 220
lineolellus, Hylaeus 261
linguaria, Apis 320
linnaeella, Apis 276
Linsleyixyla 20
liogaster, Dolerus liogaster 80
Lioalyda 102
Liometopum 197
Lionotulus subg. 185
lioparea, Blennocampa 46
Lipotriches 291
lippulum, Stenamma, *Myrmica* 207
Liris 236
Liris subg. 237
Liris 236
Liris subg. 237
lisbonensis, Anthophora
 acervorum var. 323
Lisconeura 68
Lisoca 147
lissonotum, Lasioglossum, *Halictus* 284
listerella, Melitta 265
litatus, Dicyrtomellus, *Pompilus* 170
Lithoryssus 48
Lithracia 87
Lithurge 295
Lithurgini 295
Lithurgus 295
Lithurgus subg. 295
litoralis, Mutilla 157
litoralis, Tropidotilla, *Mutilla* 157
litorana, Anthophora 322
litterata, Tenthredopsis, *Tenthredo* 99
littoralis, Diphlebus 248
littoralis, Evagetes, *Pompilus* 171
littoralis, Mimumesa *Mimesa* 251
littoralis, Myrmica lobicornis 205
liturata, Apis 302
liturata, Empria, *Tenthredo* 41
liturata, Tenthredo, Allantus 96
liturata, Tenthredopsis lactiflua 99
lituratum, Anthidium 302
lituratus, Crabro 233
lituratus, Ectemnius, *Crabro* 233
Liturgus 295
livida, Tenthredo 95
lividiventris, Tenthredo 88
lividiventris, Tenthredopsis 99
lividincinctus, Prionyx, *Enodia* 215
livonensis, Eriocampa 46
lixivia, Mimesa 250
lobata, Andrena 276
lobatus, Odynerus 184
lobatus, Okutanus 51
lobatus, Pamphilus 105
lobicornis, Myrmica 206
locularoides, Andrena 263
Loderus 77
Loderus subg. 79
loeischii, Scolia 148
loeckeniana, Megachile alpicola 306
loekenae, Dolichovespula 195
loewi, Crabro 226
loewii, Tarpa 107
loktni, Empria 41
loktonovi, Hedychridium 129
lombardica, Andrena 267
lonae, Myrmica 206
lonae, Myrmica scabrinodis 206
Lonchodryinus 119
longa, Sphecius cribaria 226
longicauda, Euura, *Nematus* 61
longiceps, Formica 199
longiceps, Halictus 285
longiceps, Lasius 201
longiceps, Osmia 299
longiceps, Passaloecus 248

- longicolle, *Hedychrum* 130
longicolis, *Ammophila* 212
longicollis, *Krombeinella*, *Myrmosa* 152
longicollis, *Microdynerus* 186
longicollis, *Xiphydria*, *Tenthredo* 116
longicornis, *Amauronematus* 56
longicornis, *Ametastegia*, *Emphytina* 40
longicornis, *Apis* 320
longicornis, *Arge* 23
longicornis, *Dolerus* 79
longicornis, *Empria*, *Poecilosoma* 41
longicornis, *Eucera*, *Apis* 320
longicornis, *Pachyprotasis* 87
longicornis, *Pristiphora*,
Pachynematus 72
longicornis, *Prosopis* 262
longicornis, *Pteronus* 64
longicornis, *Sphex* 221
longicornis, *Taxoblenus* 43
longicornis, *Tenthredo* 50
longidens, *Amauronematus* 66
longifacies, *Lasioglossum*
sulcatulum 286
longifoliae, *Athalia* 44
longiforceps, *Anteon* 119
longigena, *Hedychridium* 129
longigenae, *Halictus gavarnicus* 279
longipalpis, *Crabro* 231
longipennis, *Rocalia* 81
longipennis, *Tenthredo*, *Allantus* 98
longipes, *Tenthredo*, *Allantus* 90
longiscapus, *Myrmica* 206
longiserra, *Decanematus* 61
longiserra, *Euura*, *Nematus* 61
longispinus, *Nematus* 66
longispinus, *Sulcomesitus* 124
longiuscula, *Coelioxys* 304
longivaginata, *Masaakia* 49, 50
longula, *Andrena* 264
longula, *Chrysis ignita* var. 136
longula, *Chrysis longula* 136
longula, *Mimesa* 251
longula, *Xyela* 21
longuloides, *Halictus* 284
longulus, *Halictus* 284
longulus, *Psen* 252
longulus, *Sphecodes* 288
lonicerae, *Nematus*, *Pachynematus* 69
loochoensis, *Vespa tropica* 194
Lophagenia 165
Lophanthophora subg. 324
Lophocrabro 231
Lophopompilus 173
Lophyrus 33
Loreisomorpha 151
Lorrhoem 211
loti, *Anthidium* 300
lubricus, *Psenulus*, *Psen* 252
lucens, *Andrena* 271
lucens, *Dolerus* 80
lucens, *Nematus*, *Pteronidea* 69
lucens, *Nitela* 241
lucens, *Symmorphus*, *Odynerus* 191
lucens, *Tenthredopsis camtschatalis* 98
lucida, *Andrena* 267, 269
lucida, *Holopyga*, *Hedychrum* 131
lucida, *Mesoneura opaca* 68
lucida, *Onycholyda*, *Pamphilus* 103
lucida, *Ponera coarctata* var. 209
lucida, *Tenthredo* 69
lucidellus, *Halictus* 284
lucidicollis, *Halictus* 282
lucidiventre, *Hedychridium* 129
lucidula, *Chrysis* 130
lucidula, *Megachile* 308
lucidulum, *Lasioglossum*, *Hylaeus* 284
lucidus, *Dolerus* 79
lucidus, *Halictus* 286
lucidus, *Nematus*, *Tenthredo* 69
lucifera, *Nomada mutabilis* var. 314
lucorum, *Bombus*, *Apis* 328
lucorum, *Lyda* 103
lucorum, *Tenthredo* 30
lucorum, *Trichiosoma*, *Tenthredo* 31
Lucorumiformis 30
luctatus, *Dolerus* 77
luctifer, *Allantus*, *Macrophyta luctifer* 38
luctuosa, *Cerceris* 252
luctuosa, *Chalicodoma* 305
luctuosa, *Melecta*, *Apis* 326
luctuosa, *Pemphredon* 249
luctuosa, *Pteronidea curtispina* 56
luctuosa, *Tenthredo* 67, 84, 87
luctuosus, *Dolerus* 39
luctuosus, *Epeorus* 317
luctuosus, *Nematus* 66
luctuosus, *Pompilus* 170
ludens, *Nematus* 73
Ludita 150
Luea 55
luffi, *Podalonia*, *Ammophila* 214
lukanicus, *Halictus* 277
lugdunensis, *Nematus* 66
lugens, *Amauronematus viduatus* 66
lugens, *Athalia*, *Tenthredo* 44
lugens, *Diprion laricis* 34
lugens, *Pemphredon* 249
lugubrata, *Tenthredo kiefferi* 90
lugubrescens, *Andrena* 268
lugubris, *Andrena* 268, 269
lugubris, *Atelosphecia* 237
lugubris, *Crabro* 248
lugubris, *Dolerus* 77
lugubris, *Formica* 199
lugubris, *Melitta* 280
lugubris, *Pemphredon*, *Crabro* 249
lugubris, *Pontania viminalis* 58
lugubris, *Prosopis hyalinata* var. 262
lugubris, *Tenthredo* 84, 85
lugubris, *Vespa media media* var. 195
luminosus, *Allantus* 97
lunata, *Cerceris* 254
lunata, *Sphecius* 226
lunata, *Xylocopa* 310
lunatus, *Gonatopus* 121
lunatus, *Harpactus*, *Larra* 223
lundbladi, *Crossocerus*, *Crabro* 229
lundbohmi, *Amauronematus* 64
unicornis, *Pompilus* 219
unicornis, *Alyson* 219
unicornis, *Didineis*, *Pompilus* 219
luniger, *Stizus* 225
lunulata, *Eumenes* 181
lunulatus, *Caenocephalus*, *Cephus* 108
lunulatus, *Crabro* 29
luperus, *Diodontus* 247
lupinelli, *Halictus* 278
lurida, *Tenthredo* 95
luridicarpa, *Macrophyta* 85
luridiventris, *Platycampus*, *Tenthredo* 70
luridiventris, *Tenthredo* 51
luridus, *Campioniscus* 53
luristanensis, *Eumenes sichelii* var. 185
lusca, *Trichrysia*, *Chrysis* 141
lusitanica, *Perineura* 98
lussinicus, *Halictus* 278
lustrans, *Sphecodes* 288
lutaria, *Mimesa*, *Sphex* 250
lutea, *Tenthredo* 29
lutea, *Nomada* 314
lutea, *Tenthredo* 68
luteicornis, *Cladius* 54
luteicornis, *Tenthredo* 91
luteifasciatus, *Stenodynerus pappi* 190
luteiformis, *Nematinus bilineatus* 68
luteipalpis, *Crossocerus* 229
luteipennis, *Tenthredo* 95
luteipes, *Anthidium* 303
luteipes, *Janus*, *Cephus* 110
luteipes, *Pristiphora* 72
luteiscapus, *Pterocheilus phaleratus* 189
luteiventris, *Cladius* 54
luteiventris, *Ephippionotus* 110
luteiventris, *Eutomostethus*,
Tenthredo 47
luteiventris, *Pristiphora* 72
luteiventris, *Tenthredo* 47
luteocincta, *Tenthredo* 96
luteogaster, *Nematus* 74
luteogaster, *Urocerus gigas* 113
luteola, *Arge pyrenaica* 24
luteola, *Cimbex* 29
luteola, *Myrmica* 206
luteola, *Tenthredo* 42
luteola, *Tenthredo* 43, 67
luteolum, *Diprion frutetorum* 33
luteomaculata, *Macrophyta postica* 86
luteomarginatus, *Cephus* 109
luteonotata, *Macrophyta rustica* 86
luteostigmatellus, *Halictus* 285
lutescens, *Andrena* 267
lutescens, *Aulacomerus* 54
lutescens, *Bembix* 221
lutescens, *Cimbex betulae* 29
lutescens, *Diprion frutetorum* 33
lutescens, *Hoplomerus*
signaticollis var. 187
lutescens, *Smicromyrme rufipes* var. 156
lutescens, *Tenthredo* 103

luteus, Cimbex, *Tenthredo* 30
luteus, Eopompilus 162
luteus, Nematinus, *Nematus* 68
lutulenta, Apis 267
lutzenkoi, Halictus 280
luxorius, Odynerus magnificus 180
luxuriosa, Hoplomerus melanocephalus
 melanocephalus var. 187
luxuriosus, Crabro 233
luxuriosus, Leptochilus
 membranaceus 185
luzonicus, Anoplus 172
Lyda 104
lyda, Chrysis 136
lydiae, Chrysura lydiae, *Chrysis* 139
Lygaeonematus 70
Lygaeophora 70
Lygaeotus 70
lyncea, Tetraloniella 322
Lyroda 240
Lyrondon 240
Lyrola 25
Lyrops 236, 240
lysholmi, Bombus balteatus 327
Lyssepyris 123
lythri, Tetralonia 322
lyubae, Philoctetes 132

maadiensis, Megachile leachella 306
macedonica, Andrena 267
macedonica, Eumenes dubius
 dubius var. 182
Macgillivraya 67
Macgillivrayella 67
macgillivrayi, Andrena 268
Machaerothrix 165
macilenta, Andrena 268
macilentus, Polemistus 249
macilentus, Sirex 111
macrocephalus, Panurgus
 calcaratus var. 276
Macrocephus 110
Macrocerca 321
macrocerus, Nematus 64
Macrodiprion 34
macrodonta, Osmia 297
macroglossa, Eucera 321
macroglossa, Osmia 299
Macrophyta 84
Macrophyta subg. 84
Macrophyopsis 83
Macrophyopsis subg. 83
Macropis 293
Macropis subg. 293
macrops, Arge 24
macroserrata, Pteronidea 57
macrostigmatus, Dipogon 164
Macrovespa 193
Macroxyelineae 20
mactae, Andrena 275
macula, Strongylogaster, *Tenthredo* 82
macularis, Andrena 268
macularis, Vespa 196

maculata, Apis 300
maculata, Athalia 45
maculata, Ceropales, *Evania*
 maculata 160
maculata, Cimbex 29
maculata, Evania 160
maculata, Megascolia, *Sphex* 147
maculata, Osmia 296
maculata, Pasites 319
maculata, Paururus juvencus 112
maculata, Pseudoscolia 256
maculata, Smicromyrme
 orientalis var. 156
maculata, Smicromyrme variabilis f. 155
maculata, Tenthredo 92
maculata, Vespa 194
maculata, Vespa 196
maculativentris, Macrophyta
 diversipes 85
maculatum, Aprosthemia, *Cryptus*
 maculatum 26
maculatus, Allantus kareli 91
maculatus, Crabro 29
maculatus, Halictus maculatus 279
maculatus, Odynerus 183
maculatus, Pasites 319
maculatus, Polistes 193
maculatus, Sphecodes 289
maculatus, Sphex 220
Maculedo 89
Maculedo subg. 92
maculicornis, Gorytes, *Hoplisus* 223
maculidorsis, Agrobombus 331
maculifrons, Loderus pratorum 79
maculifrons, Lyda 103
maculifrons, Nomada 314
maculiger, Nematus 63
maculiger, Tenthredo, *Allantus*
 maculiger 91
Maculipennis 162
maculipennis, Hoplisus 224
maculipennis, Kelidoptera, *Lyda* 103
maculipennis, Lyda 103
maculipes, Oxybelus 243
maculipes, Psenulus 252
maculiscutum, Prosopis 262
maculitibia, Macrophyta 86
maculiventre, Hedychrnidium 142
maculiventris, Monopodadnus nigripes 51
maculiventris, Nematus 61
maculiventris, Oxybelus 243
maculosa, Apis 272
maculosa, Lyda 105
maculosa, Tenthredo 85
Maculosiclectes 126
maculosus, Nysson, *Sphex* 220
Madecagonatopus 120
maderae, Anthophora 322
maderi, Chrysis 136
madidus, Dolerus, *Tenthredo* 77
madraspatanum, Sceliphron, *Sphex* 214
maeklini, Crabro 226
maesta, Mutilla 157

maesta, Pristiphora, *Nematus* 72
maetai, Andrena 265
maetai, Sphecodes 289
maetoi, Aphelopus 118
magadanensis, Euura, *Pontania* 61
magna, Andrena 269
magnicornis, Dolerus 79
magnicornis, Tenthredo 84
magnifica, Ceropales 160
magnifica, Spinolia 141
magnifica, Vespa 194
magnificus, Brachyodynerus,
Odynerus 180
magnificus, Odynerus 180
magnilabris, Amegilla, *Anthophora* 323
magnini, Acantholida nemoralis 102
magniserrula, Rhogogaster 88
Magnixyla 20
magnocularis, Heptamelus 36
magnus, Bombus 328
magrettiana, Andrena 272
magrettiana, Nomada nobilis var. 315
magretti, Osmia 296
magretti, Pompilus 171
maguanensis, Vespa 193
magunta, Andrena 270
magus, Tremex, *Sirex magus* 113
mahrensis, Notogonia 237
maidli, Amauronematus 56
maidli, Ammoplanus 246
maidli, Colletes 258
maidli, Xiphydria 115
majalis, Emphytus 41
majalis, Sphecodes 288
major, Cladius 54
major, Aporus 174
major, Cephus 110
major, Cerobactrus 110
major, Crocisa 327
major, Diodontus 247
major, Dolerus pratensis 78
major, Dryudella, *Astata stigma* 218
major, Tiphia 150
major, Trypoxylon figulus 244
major, Vespa crabro 193
major, Xyela 20
Mäklini, Crabro 226
malachurellus, Halictus 282
malachuroides, Halictus 284
malachurum, Lasioglossum, *Melitta* 284
malaisei, Agenocimbex 29
malaisei, Aglaostigma, *Neurosiobia* 83
malaisei, Andrena sericea var. 268
malaisei, Bathanthidium, *Stelis* 301
malaisei, Crossocerus, *Crabro* 227
malaisei, Euura, *Amauronematus* 61
malaisei, Nysson 220
malaisei, Pristiphora, *Lygaeonematus* 72
malaisei, Psammocharies 170
malaisei, Siobla 89
malaisei, Tenthredo, *Tenthredella*
 bipunctula 94
malaisei, Terrestrisbombus terrestris 328

malaisei, *Trichiosoma* 31
malaisei, *Trypoxylon* 245
malaisei, *Xeris* 114
Malaygorytes 221
mali, *Arge*, *Hylotoma* 24
mali, *Hylotoma* 24
mallochi, *Mimesa* 250
malvae, *Apis* 321
malvae, *Tetralonia*, *Eucera* 321
malyshevi, *Crabro* 226
mama, *Auplopus* 165
manchu, *Formica* 199
manchu, *Formica exsecta* var. 199
manchuriana, *Cerceris* 254
manchurianus, *Cryptochelus* 161
manchurianus, *Discoelius* 181
manchurianus, *Eumenes*
 mediterraneus 182
manchurica, *Coelioxys* 304
manczhericus, *Camponotus herculeanus*
 japonicus var. 198
mandarinia, *Vespa* 194
mandarinus, *Halictus* 282
manderstiernii, *Dasylabris*, *Mutilla*
 manderstiernii 153
mandibulare, *Lasioglossum*,
 Halictus 284
mandibularis, *Allantus persa* 90
mandibularis, *Cephus* 111
mandibularis, *Coelioxys* 304
mandibularis, *Dolerus* 77
mandibularis, *Hoplomerus* 192
mandibularis, *Neurotoma*, *Lyda* 103
mandibularis, *Oxybelus* 243
mandibularis, *Pemphredon* 248
mandibularis, *Psen* 251
mandibularis, *Psenulus* 252
mandibularis, *Solenopsis* 207
mandibularis, *Strumigenys* 207
mandibularis, *Tenthredo* 95
mandschurica, *Nomiapis*, *Nomia* 292
mandschuricum, *Lasioglossum*
 satschauense 281
mandschuricus, *Allodynerus* 178
mandshurica, *Euura*, *Pontania* 61
mandshuricus, *Auplopus* 165
mandshuricus, *Nematus* 69
manflava, *Cerceris* 254
Mangesia 123
mangichlakensis, *Scolia*
 quadricincta 148
Manica 204
manicata, *Apis* 300
manicata, *Hoplitis* 297
manicata, *Phyllotoma* 308
manicatum, *Anthidium*, *Apis* 300
manipula, *Megachile* 307
manni, *Ammobates oraniensis* var. 318
Manoxyela 20
manticata, *Eoferreola*, *Sphex* 173
manticatus, *Dolerus* 78
mantivora, *Tachysphex* 238
Maracandica, *Tachytes* 238

maracanicum, *Podium* 215
Maracandicus, *Gastrosericus* 236
marathroicus, *Ammoplanus*,
 Hoplocrabron 246
marcescens, *Andrena* 265
marchica, *Andrena* 265
marchicus, *Ammobatoides*
 abdominalis ssp. 318
Margandrena subg. 268
margelanica, *Nomada* 314
marginalis, *Trichiosoma* 31
marginalis, *Andrena* 267
marginata, *Abia* 28
marginata, *Andrena* 268
marginata, *Andrena*
 hattorfiana var. 265
marginata, *Apis* 264
marginata, *Chrysis marginata* 136
marginata, *Clavellaria* 30
marginata, *Eriocampa* 89
marginata, *Hylotoma* 24
marginata, *Lyda* 105
marginata, *Macrophyta* 86
marginata, *Megachile* 306
marginata, *Mutilla* 157
marginata, *Tenthredo* 30, 91
marginata, *Vespa* 195
marginatum, *Anthidium* 300
marginatum, *Lasioglossum*,
 Halictus 284
marginatus, *Camponotus part*, auct. 198
marginatus, *Cephus* 108
marginatus, *Colletes* 258
marginatus, *Epeorus* 317
marginatus, *Hylaeus* 259, 261
marginatus, *Pamphilus*, *Lyda* 105
marginella, *Apis* 268
marginella, *Osmia* 298
marginella, *Tenthredo marginella* 96
marginellum, *Lasioglossum*,
 Halictus 284
margreiteri, *Bombus* 329
mariae, *Bombus ikonnikovi* 327
marianum, *Hedychrum* 130
marinovi, *Tenthredo* 92
maris, *Pompilus trivialis* var. 174
marisca, *Sirex* 113
maritima, *Osmia* 298
maritima, *Athalia* 45
maritima, *Hoplitis*, *Formicapis* 297
maritima, *Megachile*, *Apis* 308
maritimus, *Miscophus* 240
maritimus, *Oxybelus* 243
markewitschi, *Pseudoxiphidria* 115
Marlattia 53
maroccana, *Bembix* 222
maroccana, *Tachysphex mocsaryi* 239
maroccanus, *Eumenes coarctatus* 181
maroccanus, *Odynerus* 190
marqueti, *Halictus* 284
marshakovi, *Pseudoscolia* 256
marshalli, *Nematus* 73
marshamella, *Nomada*, *Apis* 314
martialis, *Macrophyta rustica* 86
martialis, *Tenthredopsis nassata* 99
martinella, *Chrysis martinella* 136
martini, *Monostegia* 43
Martjanowi, *Crabro* 233
martjanowi, *Ectemnius*, *Crabro* 233
maruyamai, *Cryptocheilus* 161
maruyamanus, *Sphecodes* 288
mariyandicus, *Nematus* 58
Masaakia 49
masao, *Pamphilus* 105
Masarinae 176
mascula, *Hylotoma serva* 81
mascula, *Pontania piliserra* 63
masculina, *Macrophyta diversipes* 85
masculina, *Tenthredo atra* 93
masculus, *Emphytus basalis* 38
Mashona 234
masii, *Acrepyris*, *Pristocera* 123
massiliensis, *Megachile* 305
Mastigocera 20
Mastigocerus 20
masudai, *Arge* 25
mathesoni, *Phlebatrophia* 49
matsumurae, *Osmia* 299
matsumurae, *Xoanon*, *Sirex* 113
matsumurai, *Megacampsomeris*, *Campso-*
 meris grossa 147
matsumurai, *Vespa* 194
matsumurai, *Zaraea lewisii* 28
maura, *Dasylabris*, *Mutilla maura* 153
maura, *Dioxyx* 303
maura, *Pontania* 56
maura, *Tenthredo* 46, 95
maura, *Tenthredopsis raddatzi* 99
maura, *Thrinax* 82
mauritanica, *Chrysis* 143
mauritanica, *Andrena rutila* ssp. 270
mauritanica, *Arge pyrenaica* 24
mauritanica, *Cardiocondyla* 203
mauritanica, *Cardiocondyla*
 nuda var. 203
mauritanica, *Nomada* 314
mauritanicus, *Eumenes*
 unguiculatus var. 181
mauritanicus, *Pterochilus* 185
mauritanensis, *Eustenancistrocerus*
 transitorius 184
mauropsis, *Dasylabris* 153
maurus, *Ceratocolus* 230
mavromoustakisi, *Gorytes* 220
maxillosa, *Apis* 295
maxillosa, *Megachile* 306
maxillosa, *Vespa* 180
maxillosus, *Bombus* 329
maxillosus, *Sphex* 215
maxim, *Arachnospila* 169
maxima, *Phyllotoma* 49
maximum,
 Anthidium pubescens var. 303
maximus, *Halictus quadricinctus* 277
maximus, *Lithurgus* 295
maximus, *Pamphilus* 105

mayacensis, Halictus 283
mayeti, Anthidium 301
mcluckiei, Euura, *Amauronematus* 61
meda, Myrmilla 154
medea, Chrysura 139
media, Andrena, *Ciliissa* 271
media, Cerceris 254
media, Dolichovespula, *Vespa* 194
mediadentata, Chrysis ignita var. 136
medialis, Pamphilus stramineipes 105
mediata, Chrysis ignita var. 137
mediata, Chrysis mediatata 137
mediata, Hylotoma 23
mediator, Dolerus pratensis 78
mediatrix, Tenthredopsis
 puncticollis 100
mediatus, Crabro 226
medinae, Emphytus viennensis 38
medinai, Halictus 287
mediobinotata, Tenthredo maculata 92
mediocrate, Hedychrifidium 129
mediocre, Hedychrifidium 129
mediocris, Macrocerca 322
mediocris, Tenthredo 55
mediocrum, Hedychrifidium 129
mediomaculatus, Allantus amaenus 97
medianotata, Tenthredopsis
 carbonaria 99
mediorufescens, Tenthredella
 mesomelas 91
mediterranea, Anthophora 322
mediterranea, Bembix 222
mediterranea, Melecta armata var. 326
mediterranea, Tachysphex 239
mediterraneus, Eumenes, *Eumenis* 182
mediterraneus, Tachysphex 239
medium, Trypoxylon 245
medius, Diodontus 247
medius, Megalodontes 107
medius, Vespa crabro 194
medon, Mutilla 157
medullarius, Nematus 56
medusa, Eucera 320
medvedevi, Holopyga 131
medvedevi, Myrmica stangeana 206
Megabombus subg. 328
Megacampsomeris 147
megacephala, Andrena 273
megacephala, Nomada 313
megacephala, Pontania 58
megacephala, Tarpa 107
megacephalus, Crossocerus, *Crabro* 228
megacephalus, Halictus 285
Megachile 305
Megachile subg. 306
Megachileoides 303
Megachilidae 295
Megachilinae 295
Megachilini 304
megachiloides, Stelis 306
Megadasypoda 293
Megadineura 68

megaera, Melecta 326
Megagonatopus 120
megala, Andrena labialis ssp. 267
Megalastor subg. 178
Megalochila 305
Megalodontes 107
Megalodontesidae 107
Megalodus 107
megalopterus, Dolerus 80
Megalostizus 225
Megaloxyla 20
megalpina, Pristiphora pallidiventris 73
Megapis 332
Megapoliste 192
megapolitana, Nomada lateralis var. 312
megapolitana, Xiphydria, *Konowia* 116
Megapterocheilus subg. 189
megapteroides, Dolerus asper 79
megapterus, Dolerus 80
Megascolia 147
Megascolia subg. 147
Megasiobia 89
Megastizus 225
Megaxyela 20
megerlei, Bembix 222
megerlei, Crabro 236
Megilla 323
Mehelyella 180
mehelyi, Andrena 273
melaclena, Smicromyrme
 compacta f. 155
melaena, Bembix 222
melaena, Tenthredopsis litterata 99
melaensis, Chrysis ignita 143
melaleuca, Megachile ericetorum 307
melaleucus, Aphelopus, *Gonatopus* 118
Melanapis subg. 270
Melanaporus 166
Melanaporus 174
melanaria, Mutilla 152
melanarius, Allantus, *Tenthredo* 38
melanarius, Cephus 110
melanarius, Crabro 228
melanarius, Hylaeus 261
melanarius, Palmodes, *Sphex* 214
melanaspis, Nematus 63
melanaspis, Tenthredella fagi 94
melanaspis, Tenthredopsis inornata 99
melancholicus, Oxybelus 242
Melandrena subg. 268
melandura, Andrena 269
Melanoanthidium 300
Melanobates 48
Melanobombus subg. 329
melanocarpa, Pristiphora, *Nematus* 72
melanocephala, Euura, *Nematus* 61
melanocephala, Melitta 269
melanocephala, Mutilla 152
melanocephala, Osmia 299
melanocephala, Tenthredo 51, 78
melanocephalum, Tapinoma,
 Formica 198
melanocephalus, Odynerus, *Vespa* 186
melanocera, Nomada lineola var. 313
melanoceraea, Tenthredo omissa 97
melanocerus, Nematus 58
melanocerus, Sirex 112
melanochra, Arge, *Tenthredo* 24
melanochra, Tenthredo 86
melanochra, Vespa 187
Melanochroiformis 22
Melanocleptes 126
melanocorne, Lasioglossum
 fulvicorne 283
Melanocrabro 231
melanodus, Nematus 58
melanogaster, Crossocerus 228
melanogaster, Megachile 306
melanogaster, Osmia 298
melanogaster, Phymatoceropsis 52
melanoleuca, Osmia 297
melanoleuca, Tenthredo 85
melanoleucus, Nematus 67
melanomeros, Tenthredo omissa 97
melanomerus, Allantus marginellus 96
melanomitratus, Halictus 287
melanonota, Rhogogaster viridis 88
melanopleura, Dasypoda 293
melanopoda, Phaenusa 48
melanoprotus, Halictus 283
melanoptera, Andrena
 ssp. 269
melanoptera, Aprosthemia peletieri 26
melanoptera, Athalia circularis 44
melanopterus, Stizoides, *Stizus* 225
Melanopus 107
melanopus, Emphytus serotinus 40
melanopyga, Megachile 307
melanopyga, Tachytes 240
melanopyga, Tenthredo 49
melanopygum, Trichiosoma 31
melanorrhoea, Tenthredo 99
Melanoselandria 52
Melanosmia subg. 298
melanosoma, Eucera 321
melanosoma, Macrophyia 84
Melanostelis 302
melanosterna, Selandria, *Tenthredo* 81
melanosterna, Tenthredo 81
melanosternus, Nematus 63
melanostigma, Nematus 54
melanostoma, Eucera 320
melanostoma, Pteronidea 62
melanostomum, Anthidium 303
melanostomus, Nematus 71
melanothorax, Nomada lineola var. 313
melanotica, Crabro vagabundus 227
melanoxantha, Dineura 55
Melanithidium 300
melanura, Andrena 272
melanura, Bembix 222
melanura, Melitta, *Kirbya* 294
melanura, Stelis 302
melanurum, Aprosthemia, *Hylotoma* 26
melanurum, Pseudoanthidium,
 Anthidium 302

melanurus, *Bombus* 331
melanurus, *Nematus* 74
melanus, *Dolerus elderi* 78
melas, *Nysson* 220
melas, *Tachysphex* 239
Melastola 70
Melea subg. 324
Melecta 325
Melecta subg. 325
melectiformis, *Epeorus* 317
Melectini 325
melectoides, *Phileremus* 318
Melicerta 36
meliloti, *Andrena* 274
Melinia 47
melinum, *Proceratum*, *Ponera* 210
Melisandra 81
Melissina 321
Melitta 294
Melittidae 293
Melittinae 293
Melittoxena 318
Melitturga 276
Melitturgini 276
Meliturgopsis 325
mellifera, *Apis* 332
Melliniae 246
Mellinogastra 224
Mellinus 246
mellita, *Phyllotoma* 48
melona, *Andrena albopunctata* ssp. 268
membranaceus, *Leptochilus*,
 Odynerus 185
Mendacibombus subg. 329
mendax, *Halictus* 284
mendicus, *Nematus* 62
menetriesi, *Scolia* 148
menetriesii, *Tenthredo* 95
menozzi, *Bothriomyrmex* 197
menozzi, *Formica truncorum* var. 200
mentiens, *Pseudodineura*,
 Blennocampa 75
menyanthidis, *Tenthredo moniliata* 95
mephisto, *Nomada* 314
mequignoni, *Allantus vespa* 97
meraca, *Andrena* 274
Meragenia 165
merceti, *Polistes* 193
Mercetia 120
Meria 149
meridiana, *Macrocerca* 322
meridianus, *Ancistrocerus scoticus* 179
meridianus, *Dolerus germanicus* 78
meridionale, *Anthidium* 300
meridionalis, *Anthophora*
 retusa var. 325
meridionalis, *Bembecinus* 221
meridionalis, *Crabro* 233
meridionalis, *Ectemnius irridifrons* 233
meridionalis, *Ectemnius*, *Crabro* 232
meridionalis, *Eutomostethus*
 gagathinus 47
meridionalis, *Formica rufa* var. 200

meridionalis, *Halictus* 279
meridionalis, *Hylaeus* 262
meridionalis, *Lasius*, *Formicina* 201
meridionalis, *Melitta* 294
meridionalis, *Nematus* 66
meridionalis, *Nomada lineola* var. 313
meridionalis, *Oxybelus* 243
meridionalis, *Psenulus* 252
meridionalis, *Vespa crabro* 194
merops, *Dolerus* 80
Merospis 231
merula, *Camponotus* 198
mervensis, *Ammophila* 213
mervensis, *Anthophora* 322
mervensis, *Mutilla* 157
Mesagenia 164
mesasiatica, *Chrysis rutilans* var. 135
Mesitiinae 124
mesoclerus, *Halictus* 284
Mesocrabro 231
Mesodryinus 120
mesoleus, *Halictus* 280
mesomela, *Tenthredo* 91
mesomelas, *Bombus* 331
Mesoneura 68
Mesonevra 68
mesonalis, *Dolerus variator* 78
mesopleuralis, *Dolerus variator* 78
mesopleuralis, *Lindenius*, *Crabro* 235
mesopleuralis, *Tenthredella*
 mesomelas 91
Mesopora 251
mesopyrrha, *Andrena* 273
mesosclerum, *Lasioglossum*,
 Halictus 284
mesostenus, *Ammatomus*, *Gorytes* 221
mesothoracica, *Pteronidea*
 melanocephala 61
mesoxantha, *Andrena* 264
Mesoxyela 20
messanensis, *Hylotoma cyanocrocea* 23
Messor 204
Metacrabro 231
Metactenius 231
Metagonatopus 120
metallescens, *Andrena* 269
metallica, *Abia* 28
metallica, *Andrena* 269
metallica, *Holopyga*, *Hedychrum* 131
metallica, *Spinarge*, *Hylotoma* 25
metallica, *Stromboceros* 81
metallicus, *Eutomostethus*,
 Tomostethopsis 47
metallicus, *Miscophus* 241
Metallinella subg. 299
Metallopeus 89
Metallus 50
metapleuris, *Tenthredopsis nassata* 99
Metastenancistrocerus 180
metatarsalis, *Ammoplanus* 246
Metavespula 194
Metepipona 188
Methocha 149
Methocha subg. 149
Methochinae 149
Metoca auct. 149
Metocha auct. 149
metopias, *Halictus* 282
Metrionotus 124
mexicana, *Isodontia*, *Sphex* 215
meyeri, *Nannodynerus* 189
micado, *Eumenes* 182
micado, *Vespa* 194
Micadophila 213
micans, *Dolerus* 80
micans, *Larra* 237
micans, *Leptochilus* 186
micans, *Philotetes* 132
micans, *Rhadinoceraea*, *Tenthredo* 52
micans, *Sphex* 215
micans, *Tachysphex*, *Tachytes* 239
micans, *Tenthredo* 52
micipsa, *Psammophila* 213
Micramegila subg. 322
Micrandrena subg. 270
micraulius, *Nematus* 66
microcellatus, *Myrmoxenus* 207
microcephala, *Formica* 197
microcephala, *Tenthredo* 99
microcephalum, *Liometopum*,
 Formica 197
microcephalus, *Heterarthrus*,
 Tenthredo 49
Microcephus 111
microcercus, *Nematus* 62
Microcrabro 227
Microdasypoda 293
Microdiprion 34
microdonta, *Coelioxys* 304
Microdrapetes 173
Microdynerus 185
Microdynerus subg. 186
microdynerus, *Microdynerus*,
 Odynerus 186
Microglossa 249
Microglossella 249
micromegas, *Tachysphex* 238
micronematica, *Pristiphora* 73
Micronematus 70
Microphadnus 173
microphyes, *Euura*, *Nematus* 61
microps, *Nematus* 57
microps, *Tenthredo* 91
Micropterocheilus subg. 189
micropunctata, *Cerceris* 254
microrubra, *Myrmica* 206
Microstelis 302
microstictus, *Crabro* 232
microstieta, *Nomada* 316
microstigma, *Andrena* 294
migrans, *Andrena erberi* var. 264
mikado, *Megaxyela* 20
mikado, *Mutilla* 157
mikagei, *Macrophya* 87
mikhno, *Panurginus* 276

miliaris, Euura, *Tenthredo* 61
 militaris, *Macrophya*, *Tenthredo* 86
 militaris, Nysson 220
millefolii, Meria 146
milliati, Allantus stecki 90
 miltonota, Euura, *Nematus* 61
miltotus, Camponotus japonica var. 198
 mimator, Euura 61
Mimesa 250
mimica, Cephaleia 102
 mimulus, Nysson 220
Mimumesa 250
mimus, Amauronematus 61
 mimus, Euura, *Pteronus* 61
 minamikawai, Crossocerus 228
miniatus, Nematus 62
 miniatus, Sphecodes 288
minima, *Apis florisomnis* 295
minima, Fenusa 48
minima, Melecta luctuosa var. 326
 minima, Stelis 303
minimus, *Apis* 287
minimus, Mesitius 124
 minimus, Metrionotus, *Mesitius* 124
minoensis, Ronisia barbara 157
 minomalis, Onycholyda, *Pamphilus* 103
 minor, Andrena, *Ciliissa* 271
 minor, Astata 218
minor, Crocisa affinis var. 327
minor, Dasypoda 293
 minor, Eopompilus 162
 minor, Halictus 278
minor, Lophyrus 34
minor, Mesitius 124
minor, Trypoxylon figulus 245
Minorixyela 20
minos, Prospis hyalinata 262
minuma, Holopyga 131
minus, Anthidium 301
 minus, Trypoxylon 245
minuscula, Nomada 316
minusculum, Anthidium 300
minusculus, Dolerus 79
minuta, Andrena 287
minuta, Coelioxys 304
 minuta, Dufourea 290
minuta, Dufourea auct. 290
minuta, Fenella, *Phyllotoma* 47
minuta, Hoplocampa, *Tenthredo* 68
minuta, Nomada 312
minuta, Osmia 298
minuta, Priocnemis, *Pompilus* 163
minuta, Pseudomyrmeca, *Myrmosa* 152
minuta, Stelis 303
minuta, *Tenthredo* 75, 94
minuta, Tiphia 150
minuta, Vespa 186, 259
minuta, Xylocopa 309
minutepunctatus, Jucancistrocerus 184
minutissima, Andrena 270
minutissima, Apis 292
minutissima, *Macrophya* 87
minutissima, Melinia 48

minutissimum, *Lasioglossum*,
Melitta 284
minutissimus, *Nomioides*, *Apis*
minutissimus 292
minutissimus, *Stigmus* 249
minutula, Andrena, *Melitta* 270
minutula, Arachnospila, *Pompilus* 169
minutula, *Nomada solidaginis* var. 316
minutuloides, Andrena 270
minutuloides, *Lasioglossum* 284
minutulus, *Tachysphex* 238
minutum, *Hedychrum* 128
minutum, *Monomorium* 205
minutus, *Armitarsus* 84
minutus, *Diodontus*, *Crabro* 247
minutus, *Diphlebus* 248
minutus, *Dolerus* 75
minutus, *Elampus* 133
minutus, *Halictus* 285
minutus, *Hoplisus* 224
minutus, *Parammobatodes*, *Phiarus* 319
minutus, *Phiarus* 319
minutus, *Pompilus* 169
minutus, *Psen* 252
minutus, *Tachysphex* 237
minutus, *Taxonus* 39
Miocephala 22
mioceras, *Tenthredo* 91
mioceras, *Tenthredo*, *Tenthredella* 91
miogramma, *Dasylobus italicica* var. 154
mirabilis, *Cerceris* 253
mirabilis, *Chrysis* 137
mirabilis, Euura, *Pontania* 61
miranda, *Nomada jacobaeae* var. 312
mirandus, *Microdynerus*,
Leptochilus 186
miricolor, *Dolerus thomsoni* 77
mironowianus, *Agrobombus*
schrencki 331
miscellus, *Hylaeus* 262
miscisis, *Smicromyrme oratoria* 155
Miscophini 240
Miscophinus 240
miscophoides, *Atelosphex* 237
miscophoides, *Soliarella* 241
Mischophus 240
Miscus 212
misellus, *Halictus* 283
Misopyris 123
miserabilis, *Bembix* 222
mishimana, *Nomada* 313
mistura, *Nomada* 315
mitaii, *Nomada* 314
mitakensis, Andrena 271
mitakensis, Priocnemis 163
mitis, Andrena 264
mitis, *Hoplitis*, *Osmia* 296
mitjaevi, *Gorytes* 220
mitratolus, *Halictus melanomitratus* 287
mitsuhashii, *Tenthredo*, *Macrophya* 98
mitsukurii, *Eriocampa* 42
mitsukurii, *Tetraloniella*, *Tetralonia* 322
mittrei, *Cephus* 109
mixta, *Andrena* 264
mixta, *Anthophora* 323
mixta, *Blennocampa* 46
mixta, *Cerceris* 253
mixta, *Dasypoda* 293
mixta, *Megachile* 305
mixta, *Nomada succineta* var. 316
mixta, *Prosopis* 261
mixta, *Strongylogaster*, *Tenthredo* 82
mixta, *Vespa* 196
mixtella, *Nomada* 315
mixtura, *Andrena limata* ssp. 269
mixtus, *Lasius*, *Formica* 201
miyabei, *Lasioglossum* 284
miyagino, *Psen* 251
miyamotoi, *Andrena* 267
mizuhō, *Crabro nigritarsus* 231
mizuhō, *Ectemnius* 231
mizuhonis, *Symmorphus* 191
mlokosewitzii, *Apterogyna* 159
mlokosewitzii, *Anthophora* 324
mlokosewitzii, *Dasylabris* 154
mlokosiewitzii, *Bombus* 331
mlokosiewitzii, *Holopyga*,
Hedychrum 131
mlokoszewiczi, *Colletes* 258
mlokoszewiczi, *Dasypoda* 293
mocsaryi, *Alastor*, *Odynerus* 178
mocsaryi, *Allantus* 90
mocsaryi, *Ammophila* 213
mocsaryi, *Athalia rufoscutellata* 45
mocsaryi, *Bombus* 331
mocsaryi, *Cephus* 111
mocsaryi, *Chelostoma* 295
mocsaryi, *Cleptes* 142
mocsaryi, *Crabro* 226
mocsaryi, *Dolerus* 77
mocsaryi, *Hoplitis*, *Osmia* 297
mocsaryi, *Metrionotus* 124
mocsaryi, *Nomada* 314
mocsaryi, *Prionyx*, *Sphex viduatus* 215
mocsaryi, *Spilomena* 249
mocsaryi, *Spintharina*, *Spintharis* 141
mocsaryi, *Tachysphex* 239
mocsaryi, *Tenthredo* 95
Moczarellus subg. 162
moczari, *Arachnospila* 169
moczari, *Sulcomesitus* 125
moczari, *Tachysphex* 239
moderna, *Anthophora* 324
modesta, *Apis* 300
modesta, *Argo* 24
modesta, *Nomada marshamella* var. 314
modesta, *Olbia* 224
modestius, *Priophorus* 54
modestum, *Rhopalum* 235
modestus, *Bombus* 330
modestus, *Bothriomyrmex* 197
modestus, *Characopygus* 110
modestus, *Dolerus liogaster* 80
moeschleri, *Ancistrocerus*
trifasciatus var. 180
moeschleri, *Nomada* 314

moeschleri, Rophites 291
moestus, Allantus 90
moestus, Aporinellus, *Pompilus* 168
mogadorensis, Megachile 306
Mogerus 51
moiwana, Strongylogaster 82
moiwasanus, Allantus 91
mokrousovi, Chrysis 137
molhusina, Andrena 268
mollesonae, Formica sanguinea var. 200
mollis, Nematus 73
mollis, Pristiphora, *Nematus* 73
moltonii, Polistes nimpha var. 193
moltrechti, Halictus 281
momidzizyama, Smicromyrme 156
monacha, Megilla 325
monachus, Oxybelus 242
Monardis 50
mondaensis, Andrena
 bimaculata var. 272
mondaensis, Halictus, *Seladonia* 279
Monepyris 123
monetierensis, Allantus *vespa* 97
mongol, Bombus fragrans 330
mongolensis, Ammophila 212
mongolensis, Ammoplanus 246
mongolensis, Argogorytes 221
mongolensis, Crabro yasumatsui 230
mongolensis, Ephutomma,
 Smicromyrme 155
mongolensis, Halictus rubicundus 278
mongolianae, Chrysis 135
mongolica, Arachnospila 169
mongolica, Bembix oculata 222
mongolica, Chrysis 135
mongolica, Dasylabris, *Mutilla* 154
mongolica, Empria, *Poecilosoma* 41
mongolica, Melitturga 276
mongolica, Proformica, *Formica* 203
mongolica, Pseudepipona variegata 188
mongolica, Scolia 147
mongolica, Taimyrmosa, *Myrmosa* 153
mongolica, Tenthredo, *Allantus* 96
mongolica, Vespa 194
mongolicum, Anteon 119
mongolicus, Ancistrocerus,
 Odynerus 179
mongolicus, Colletes 258
mongolicus, Diodontus 247
mongolicus, Dolerus, *Loderus* 77
mongolicus, Euchroeus 140
mongolicus, Eumenes 182
mongolicus, Eumenes
 coarctatus var. 182
mongolicus, Gorytes sulcifrons 223
mongolicus, Halictus 279
mongolicus, Leptothorax serviculus 208
mongolicus, Lindenius panzeri 235
mongolicus, Passaloecus 248
mongolicus, Philoctetes
 horvathi var. 132
mongolicus, Philoctetes, *Ellampus* 132
mongolicus, Polistes gallicus var. 193

mongolicus, Stizus 225
mongolicus, Temnothorax,
 Leptocephalus 208
mongolina, Discolia 148
mongolinfuscatus, Anoplus
 infuscatus 172
mongoloexigua, Pristiphora 73
mongolofausta, Pristiphora 73
mongololaricis, Pristiphora 72
mongolominutula, Arachnospila
 minutula 169
mongolopinata, Arachnospila 169
mongoloproximus, Evagetas
 proximus 171
mongolorufa, Arachnospila 170
mongolorum, Sirex, *Paururus* 112
mongolospissa, Arachnospila 169
Monia 257
Monidia 257
Monilapis subg. 277
moniliata, Tenthredo 95
moniliatus, Hylaeus 261
monilicornis, Fenella, *Phyllotoma* 48
monilicornis, Passaloecus 248
monilicornis, Sphecodes, *Melitta* 289
monoceros, Megachile 295
monochroum, Hedychridium 129
Monochrysis 139
Monoceteniae 35
Monocetus 35
Monogonatopus 121
monogyniae, Pristiphora, *Nematus* 73
Monomatium 236
Monomorium 205
monomorium, Monomorium 205
Monophadnoides 50
Monophadnus 50
Monoplomerus subg. 186
Monoplopis 108
monopunctata, Smicromyrme
 hasanensis f. 156
Monosapya 145
Monosmia 297
Monosoma 43
monostega, Arge 25
Monostegia 42
monostictus, Hylaeus 261
monozonus, Dolerus madidus 77
monozonus, Tenthredo, *Allantus* 92
Monsoma 43
monspliensis, Mutilla 157
monstrificum, Lasioglossum,
 Halictus 284
monstrosa, Nomia 291
monstrosus, Crabro 230
montana, Ametastegia 39
montana, Camponotus
 herculeanus var. 198
montana, Colletes 258
montana, Eumenes 185
montana, Euura, *Nematus* 61
montana, Larrada 238
montana, Macrophyta, *Tenthredo*
 montana 86
montana, Mutilla 155
montana, Nomadita 315
montana, Pemphredon 249
montana, Rhogogaster viridis 92
montana, Scaptomyza 89
montana, Sirex gigas 113
montana, Tenthredo 29
montana, Tenthredo 84
montana, Tenthredo mesomelas 91
montandoni, Coelioxys 304
montanum, Anthidium 300
montanus, Crabro 232, 233
montanus, Epeorus 317
montanus, Oxybelus quadrinotatus 243
montanus, Pison 244
montanus, Priophorus 54
montanus, Psen 252
montanus, Rophites algirus 291
montanus, Tachytes 238
montebovensis, Spatidariella
 hyalinata 262
monticola, Bombus 330
monticola, Diodontus 247
monticola, Eumenes papillarius 182
monticola, Euura, *Nematus* 61
monticola, Hylaeus 261
monticola, Monophadnus, *Tenthredo* 50
monticola, Tachytes 238
monticola, Trypoxylon 245
monticola, Vespa norvegica saxonica
 morpha 195
montislinguarum, Podalirius
 furcatus 323
montivaga, Osmia 296
montivaga, Vespa adulterina 194
montivagans, Crabro 232
montivagus, Hylaeus 260
mooreana, Siobla 89
mopsus, Nysson 220
moravica, Allantus sabariensis 91
moravica, Aprosthemma maculata 26
moravica, Myrmica 205
moravica, Pristiphora 74
moravica, Smicromyrme montata f. 155
moravica, Stelis 303
moravicium, Tetramorium 209
moravicus, Gorytes 223
moravicus, Miscophus 241
morawitsi, Amauronematus 60
morawitsi, Allantus 91
morawitsi, Ammophila 213
morawitsi, Andrena 272
morawitsi, Anthophora 324, 325
morawitsi, Dasypoda 293
morawitsi, Harpactus 223
morawitsi, Odynerus 180
morawitsi, Prosopis 260
morawitsi, Sapyga 145
morawitsi, Tenthredo 97
morawitzii, Mutilla 157
morawitzii, Spinolia, *Chrysis* 141
morbillousus, Halictus 280
mordax, Heriades 296

- moreana*, Scolia 148
Morgania 319
moricei, Amauronematus 59
moricei, Anthidium 302
moricei, Characopygus 110
moricei, Hedychrnidium 129
moricei, Hylaeus, *Prosopis* 260
moricella, Andrena 270
moricella, *Prosopis* 260
morio, Andrena 269
morio, Ceratophorus, *Pemphredon* 247
morio, Cladius 54
morio, Crossocerus 229
morio, Cubitalia, *Eucera* 319
morio, Lasioglossum, *Hylaeus* 284
morio, Nesolandria, *Tenthredo* 81
morio, Pemphredon 247
morio, Pemphredon 249
morio, Tenthredo 46
morio, Tiphia auct. 150
morionellus, Nematus 63
morisitai, Lasius 201
Morphanthidium 300
Morphota 240
mortifer, Pemphredon 249
morvandica, Macrophya ribis 86
morvandicus, Allantus scrophulariae 97
Morysmula 152
mosaicum, Anthidium 300
moscaryi, Anthidium 302
moscovita, Tenthredopsis, *Perineura* 99
moshkarareppus, Bombus beaticola 330
mosquense, Monosoma 42
mosquense, Nycteridium 35
Motes 237
mouffetella, Melitta 272
mowchowense, Trypoxylon 245
mucheii, Polistes gallicus 193
mucheii, Tenthredo sulphuripes 97
mucorea, Crocisa ramosa var. 327
mucoreus, Halictus, *Hylaeus* 279
mucronata, Euura, Nematus 62
mucronata, Tenthredo 91
mucronatus, Crabro 229
mucronatus, Oxybelus, *Crabro* 243
Muellerella 123
muinensis, Nomada 314
muliebris, Allantus cingulatus 38
muliebris, Dolerus haematodes 80
muliebris, Macrophya rufipes 86
muliebris, Nematinus bilineatus 68
mulleolus, Osmia veridata 299
multicincta, Tarpa 107
multicinctus, Allantus 97
multicolor, Tenthredo 86
Multidentiden 77
multifasciata, Strongylogaster, *Tenthredo* 82
multifasciata, Tenthredo 82
multifasciatus, Allantus 97
multiluteus, Allantus amaenus 97
multiplex, Nematus 61
multispinosa, Megachile 307
munakatai, Megachile willughbiella 308
munakatai, Nomada 312
munakatai, Stigmus 250
munditus, Priophorus 54
mundula, Melecta 326
mundus, Euura, *Amauronematus* 62
muraria, Apis 298, 305
murarius, Symmorphus, *Vespa* 191
muraviovi, Panurginus 276
murcaria, Apis 317
Murciana 90
murgabensis, Plenoculus,
Ptygosphe 241
murithianus, Sphecodes 288
murorum, Crabro 230
murotai, Conaspidea 84
murotai, Crossocerus 228
murotai, Rhynchium
quinquecinctum 189
murotai, Sphecodes 289
muscaria, Apis 317
muschtaidica, Aphaenogaster
gibbosa 203
muscorum, Bombus, *Apis* 331
muscorum, Leptothorax, *Myrmica* 204
muskaense, Rhopalum 236
mustelina, Osmia 299
mutabilis, Andrena 267
mutabilis, Chrysis mutabilis 137
mutabilis, Corynis, *Abia* 32
mutabilis, Nomada 314
mutata, Vespa dybowskii 194
mutica, Abia 28
mutica, *Mutilla erythrocephala* var. 154
mutica, *Myrmica* 204
mutica, Myrmilla, *Mutilla* 154
mutica, Nomada 315
muticus, Odynerus 187
utilata, Tenthredo 79
Mutilla 156
utilillaria, Methoca 149
utilillarius, Gonatopus 149
Mutillidae 152
Mutillinae 154
Mutillini 156
utilloides, Clytovorus,
Scleroderma 124
utilloides, Scleroderma 124
mutinensis, Odynerus sinuatus var. 191
mutini, Andrena 267
mutsu, Sphecodes 290
mutsuoides, Sphecodes 289
myosotidis, Euura, *Tenthredo* 62
myosotidis, Pteronus fallaciosus 62
Myrmecina 205
Myrmecomorphus 122
myrmecophila, *Myrmica* 206
myrmecophilus, Parvoculus 124
Myrmecosalius 162
Myrmentoma subg. 198
Myrmica 205
Myrmicinae 203
Myrmilla 154
Myrmilla subg. 154
Myrmillinae 154
Myrmosa 152
myrmosina, Mutilla 152
Myrmosinae 152
Myrmosini 152
Myrmoxenus 207
myrsiniticola, Euura, *Pontania* 62
myrtilli, Eriocampa 45
myrtilloidica, Euura, *Pontania* 62
mysta, Chrysis 137
mysta, Xoanon 113
Mystacanthophora subg. 324
mystacea, Andrena 272
mystaceus, Argogorytes, *Sphex*
mystaceus 221
mysticalis, Chrysis 137
mysticus, Nematus 64
mysticus, Tachysphex 238
Mystrophorus 119
Myzinella 149
Myzininae 149
nadezhdae, Messor kasakorum 204
nadia, Megachile 306
nadigi, Crocisa 327
nagasakiensis, Halictus 282
nagasei, Pompilus 169
nagatomii, Vespa analis 193
nagyi, Mesitius 124
nahariensis, Euodynerus posticus
punctatissimus var. 183
naiptchianus, Bombus 329
nakabusensis, Stenemphytus,
Allantus 43
nakayamai, Neurotoma 103
nambui, Crabro 230
nambui, Rhynchium
quinquecinctum 189
nana, Andrena, *Melitta* 270
nana, Anthophora
quadriasciata var. 322
nana, Anthophora
quadriasciata var. 323
nana, Fenusa, *Tenthredo* 48
nana, Larra 237
nana, Nomada 312
nana, Osmia 299
nana, Pontopristia 61
nana, *Prosopis minuta* var. 261
nana, Stelis 303
nana, Tetraloniella, *Tetralonia* 322
nanae, Trichiosoma 31
nanaeformis, Andrena 270
nanana, Andrena 264
Nannaporus 174
Nannodynerus 189
Nannopompilus 170
Nanopterochilus 189
Nanopompilus 170
nansetsurei, Symmorphus 191
nanula, Andrena 270
nanulus, *Hylaeus* 285

nanum, *Pseudoanthidium*,
Anthidium 302
nanus, *Nematus* 73
nanus, *Sirex* 114
napalkovi, *Agrobombus adventor* 331
napata, *Tenthredo* 99
naryschkini, *Discoelius* 181
nasale, *Anthidium* 302
nasalis, *Andrena* 265
nasalis, *Macrocerca* 276
nasicolle, *Anthidium*
 manicatum var. 300
Nasoxiphia 115
nassata, *Tenthredopsis*, *Tenthredo* 99
nassonovi, *Temnothorax*,
 Leptothorax 208
nasuta, *Anthophora* 323
nasuta, *Apis* 309
nasuta, *Cerceris* 254
nasuta, *Formica* 203
nasuta, *Stelis*, *Anthidium* 302
nasutum, *Trypoxylon varipes* 245
nasutus, *Colletes* 258
nasutus, *Homonotus* 166
nasutus, *Lindenius* 234
nasutus, *Lithurgus* 295
nasutus, *Phileremus* 318
navajo, *Ancistrocerus* 178
navasi, *Nomada* 311
navitatis, *Cerceris* 252
nawai, *Andrena* 264
nearctica, *Monostegia* 43
nearcticus, *Ancistrocerus* 178
Neateuchopus 111
nebrascensis, *Mimesa* 250
nebulosa, *Apis* 275
nebulosa, *Tenthredo* 91
nebulosum, *Aglaostigma*, *Macrophya* 83
nebulosus, *Halictus kessleri* 279
nebulosus, *Nematus* 68
neesi, *Eumenes mediterranea* var. 181
negativa, *Endemyolia*, *Caliroa* 47
neglecta, *Bembix* 222
neglecta, *Euura*, *Nematus* 62
neglecta, *Nomada* 315
neglecta, *Pseudochrysis*, *Chrysis* 140
neglecta, *Tenthredo* 84, 99
neglectus, *Diphlebus* 248
neglectus, *Emphytus* 38
neglectus, *Gorytes* 223
neglectus, *Lasius* 202
neglectus, *Sirex* 112
negoroi, *Hylaeus chasanensis* 259
negotiosus, *Dolerus* 78
Nematinae 53
Nematinus 68
Nematoberos 42
Nematus 69
nemesis, *Macrophya* 86
Nemka 155
nemorae, *Lasioglossum* 282
nemoralis, *Dasypoda* 293
nemoralis, *Formica* 199

nemoralis, *Lophyrus* 34
nemoralis, *Macrodiprion*, *Lophyrus* 34
nemoralis, *Mutilla* 156
nemoralis, *Neurotoma*, *Tenthredo* 103
nemoratus, *Heterarthrus*, *Hylotoma* 49
nemorus, *Dolerus* 78
nemorum, *Cephalcia* 105
nemorum, *Tenthredo* 34
neobesa, *Tenthredo* 90
Neoblepharipus 227
neobule, *Chrysis* 137
Neobusarbia 81
Neocharactus 51
Neocharactus subg. 51
Neochelogynus 118
Neocimbex 29
Neocleptes 126
Neoclista 51
Neodicrogenium 123
Neodiodontus 247
Neodiprion 34
Neodolerus 77
Neodolerus subg. 79
Neofoxia 251
Neogonatopoides 120
Neogonatopus 120
Neoleptochilus subg. 185
Neopareophora 70
Neopelmatopus 55
Neopristocera 123
Neopterocheilus subg. 187
Neopus 83
Neoscleroderma 124
Neosmia subg. 299
Neospinolia 140
Neostromboceros 81
Neosyrista 111
Neotomostethus 70
Neotrogaspidia 157
Neoxeris 114
Neoxyela 20
nepalensis, *Gonatopus*,
 Pseudogonatopus 121
Nephridia 244
Nepionema 70
nescia, *Pempredon* 248
nesiotica, *Megachile centuncularis* 306
nesiotis, *Halictus rubicundus* 278
Nesocabro 231
Nesodiprion 34
Nesohylaeus subg. 261
Nesoprosopis subg. 261
Nesolandria 81
nestor, *Pristiphora*, *Nematus* 73
nestorea, *Megachile* 305
Neurepyris 123
Neuropachys 112
Neurosiobia 83
Neurosiobia subg. 83
Neurotoma 103
neuter, *Halictus* 278
nevadensis, *Selandria* 49
nevelskiana, *Chrysis* 137
nidulans, *Apis* 322
nidulans, *Halictus* 278
nidulator, *Odynerus* 191
Nielseni, *Crabro* 232
niemelai, *Hedychrum* 130
nievesi, *Pristiphora* 74
niezabitowskii, *Athalia lineolata* 44
nigella, *Isodontia*, *Sphex* 215
nigella, *Pristiphora*, *Nematus* 73
nigella, *Selandria* 50
nigella, *Tenthredopsis* 99
nigellata, *Andrena* 271
niger, *Allantus arcuatus* 97
niger, *Cephus* 110
niger, *Crabro* 228
niger, *Crossocerus* 228
niger, *Dinetus* 236
niger, *Dolerus dubius* 78
niger, *Dolerus*, *Tenthredo* 80
niger, *Epyris* 123
niger, *Eumenes coarctatus* var. 181
niger, *Halictus* auct. 283
niger, *Hylaeus* 261
niger, *Lasius*, *Formica* 202
niger, *Liris*, *Sphex* 237
niger, *Miscophus* 241
niger, *Nysson niger* 220
niger, *Panurginus* 276
niger, *Phylloecus*, *Sirex* 110
niger, *Podalirius acervorum* var. 323
niger, *Psen* 250
niger, *Sphecodes* 289
niger, *Tachybulus* 244
Nigeriformis 77
nigerrima, *Mesoneura opaca* 68
nigerrima, *Sphecodes monilicornis* 289
nigerrima, *Sphex* 172
nigerrima, *Tenthredella fagi* 93
nigerrima, *Tenthredo* 53
nigerrima, *Tenthredopsis nassata* 99
nigerrimus, *Anoplus*, *Sphex* 172
nigerrimus, *Halictus* 286
nigerrimus, *Polyergus* 202
Nigliris 237
nigra, *Alysson pertheesi* 219
nigra, *Andrena scita* var. 273
nigra, *Apterogyna* 159
nigra, *Cimbex betulae* 29
nigra, *Dineura* 55
nigra, *Eucera* 320
nigra, *Euura* 56
nigra, *Formica* 201
nigra, *Macrophya* 87
nigra, *Macrophya neglecta* 84
nigra, *Macrophya quadrimaculata* 86
nigra, *Melecta* 326
nigra, *Mutilla* 156
nigra, *Myrmosa* 152
nigra, *Pachyprotasis* 87
nigra, *Sapyga* 146
nigra, *Sphex* 237
nigra, *Vespa crabro* 194
nigra, *Vespa parietum* β 179

nigra, Cilissa 294
nigrans, Dasypoda 293
nigrans, Vespa 193
nigrata, Formica 199
nigrata, Tenthredo 63
nigrata, Tenthredopsis coqueberti 98
nigratilobis, Tenthredopsis tessellata 100
nigratipennis, Tenthredopsis tessellata 100
nigratiscutis, Tenthredopsis tessellata 100
nigratus, Dolerus, *Tenthredo* 80
nigratus, Psen 252
nigrescens, Andrena marginata var. 268
nigrescens, Coelioxys rufescens 304
nigrescens, Dasypoda plumipes 293
nigrescens, Dolichovespula saxonica 195
nigrescens, Eucera 320
nigrescens, Leptothonax acervorum var. 204
nigrescens, Mutilla europaea f. 157
nigrescens, Odynerus dantici 183
nigrescens, Phyllotoma 49
nigrescens, Podalirius parietinus 324
nigrescens, Pontania bella 63
nigrescens, Sphecodes gibbus ephippium 289
nigrescens, Tachysphex 239
nigrescens, Tenthredopsis 99
nigricans, Dasypoda argentata 293
nigricans, Larrada 239
nigricans, Liris, *Larrada* 237
nigricans, Lyda 103
nigricans, Macrophyia superba 86
nigricans, Megachile lagopoda var. 307
nigricans, Melitta 294
nigricans, Priocnemis 163
nigricans, Pristiphora, *Nematus* 73
nigricans, Tenthredo 52, 55
nigricarpa, Periclista 51
nigricephala, Leptothonax tuberum var. 208
nigriceps, Andrena, *Melitta* 265
nigriceps, Anthidium 301
nigriceps, Anthophora 325
nigriceps, Dolerus 78
nigriceps, Hemibeleses 42
nigriceps, Hylaeus 261
nigriceps, Leptocercus 67
nigriceps, Myzine 149
nigriceps, Pristiphora, *Nematus* 73
nigriceps, Smicromyrme 156
nigriceps, Tenthredopsis 99
nigriclypeatus, Halictus vulpinus 285
nigriclypeatus, Odynerus 190
nigricollaris, Smicromyrme rufipes var. 156
nigricolle, Anthidium 302
nigricolle, Poecilosoma 46
nigricollis, Nematus 71
nigricollis, Tenthredo 94
nigricollis, Tenthredopsis 98

nigricolor, Camponotus maculatus quadrinotatus var. 198
nigricolor, Larra 239
nigricolor, Tachysphex, *Larra* 239
nigricoma, Trichiosoma 31
nigricornis, Amegilla, *Anthophora* 322
nigricornis, Ancistrocerus, *Odynerus* 179
nigricornis, Antepipona, *Lionotus* 180
nigricornis, Blepharipus 228
nigricornis, Euura, *Nematus* 62
nigricornis, Heriades 296
nigricornis, Hylaeus 259, 284
nigricornis, Macrocerca 321
nigricornis, Nematus 62, 73
nigricornis, Nomada 312
nigricornis, Oxybelus 242
nigricornis, Psen 252
nigricornis, Panurgus calcaratus var. 276
nigricorpus, Pachynematus 72
nigricoxae, Cephalcia, *Lyda* 102
nigricoxis, Emphytus viennensis 38
nigricula, Smicromyrme lewisi 156
nigridens, Crabro 234
nigridorsis, Amauronematus 75
nigrifacics, Prosopis lineolata var. 261
nigrifacies, Andrena cineraria var. 268
nigrifacies, Eucera 321
nigrifacies, Gorytes, *Hoplisus* 223
nigrifacies, Prosopis 260
nigrifacies, Prosopis annulata var. 259
nigrifacies, Prosopis pratensis var. 262
nigrifemur, Tenthredo atra 92, 93
nigrifemur, Tenthredo limbata 95
nigrifrons, Andrena 267, 270
nigrifrons, Crabro 231
nigrifrons, Eucera 320
nigrifrons, Macrophyia quadrimaculata 86
nigrifrons, Pontania ischnoceros 61
nigrigena, Meria 149
nigrihumerus, Polistes snelleni f. 192
nigrilabris, Allantus 97
nigrilabris, Eucera 320
nigribolis, Tenthredopsis parvula 99
nigrina, Macrophyia duodecimpunctata 85
nigrina, Nomada lineola var. 313
nigrina, Nysson 220
nigrinotum, Athalia 44
nigrinotum, Melitta leporina var. 294
nigrinotum, Polistes bimaculatus var. 192
nigrinum, Anthidium 301
nigrinum, Rhopalum 235
nigrinus, Cephus 109
nigrinus, Crabro 232, 235
nigrior, Crabro 228
nigrior, Eumenes formosensis 182
nigrior, Tenthredo marginella 96
nigripectus, Allantus 93
nigripectus, Lophyrus pini 33
nigripenne, Aprosthemma maculatum 26
nigripennis, Cephus, *Eumetabolus* 109
nigripennis, Sphecodes 288
nigripennis, Tenthredo 24
nigripes, Anthidium 301
nigripes, Anthophora 324
nigripes, Arge, *Tenthredo nigripes* 24
nigripes, Arge pyrenaica 24
nigripes, Dolerus pratensis 78
nigripes, Hinatara, *Fenusia* 49
nigripes, LasioGLOSSUM, *Halictus* 285
nigripes, Myrmica sulcinodis var. 206
nigripes, Nysson 220
nigripes, Odynerus 183
nigripes, Oxybelus 243
nigripes, Podalirius acervorum var. 323
nigripes, Sphecodes 287
nigripes, Sphecodes 150
nigripes, Tenthredo 51
nigripes, Tenthredo coreana 96
nigripes, Tenthredopsis thomsoni 99
nigripes, Trichiosoma 31
nigripleurus, Macrophyia postica 86
nigripleurus, Pristiphora amentorum 56
nigripleurus, Tenthredo velox 96
nigripleurus, Tenthredo, *Allantus* 96
nigricutis, Macrophyia militaris 86
nigricutum, Prosopis pictipes var. 261
nigrisoma, Tenthredo 39
nigristernis, Diprion pini 33
nigrita, Blepharipus 228
nigrita, Eucera 320
nigrita, Eucera caspica var. 320
nigrita, Fenella 47, 48
nigrita, Hoplocampa chrysorrhoea 67
nigrita, Macrophyia 87
nigrita, Mimesa 250
nigrita, Mutilla 156
nigrita, Nomada 272, 312
nigrita, Nomioides variegata 292
nigrita, Sirex 113
nigrita, Tachytes 237
nigrita, Tenthredo 53
nigritarsis, Allantus 90
nigritarsis, Cephus 111
nigritarsis, Hylotoma 23, 24
nigritarsis, Macrophyia diversipes 85
nigritarsis, Nematus 66
nigritarsis, Tenthredo 97
Nigritarsus, Crabro 231
nigritarsus, Ectemnius *Crabro* 231
nigritegulis, Athalia bicolor 44
nigritegulis, Megalodontes 107
nigritorax, Anthidium manicatum var. 300
nigritorax, Podalirhis albigena var. 323
nigritibiis, Eriocampa ovata 42
nigritulum, Anthidium kohlii var. 301
nigritulus, Telostegus 174
nigritus, Aphanisus 50
nigritus, Blepharipus 227
nigritus, Cephus 111
nigritus, Crabro 227

nigritus, Crossocerus, *Blepharipus* 228
nigritus, Dolerus 39
nigritus, Hylaeus, *Mellinus* 260
nigritus, Tomostethus, *Tenthredo* 53
nigriventris, Amasis crassicornis 32
nigriventris, Ammophila 213
nigriventris, Andrena 267
nigriventris, Apis 269
nigriventris, Dineura testaceipes 55
nigriventris, Euura, *Nematus* 62
nigriventris, Halictus 284
nigriventris, Larra 237
nigriventris, Megachile 308
nigriventris, Nomada rufipes 316
nigriventris, Nomioides variegata 292
nigriventris, Osmia, *Anthophora* 298
nigriventris, Oxybelus 243
nigriventris, Rhogogaster 88
nigriventris, Tenthredo colon 94
nigroaenea, Andrena, *Melitta* 269
nigroaenea, Ceratina 310
nigroaeneus, Oxybelus 242
nigrobarbata, Andrena 268
nigrociliata, Dasylabris maura f. 153
nigrociliatus, Hortobombus
 consobrinus 328
nigroclypeata, Halictus vulpinus 285
nigroclypeata, Onycholyda 103
nigroclypeata, Tenthredo 93
nigrocoerulea, Cephaleia 103
nigrocuneatus, Hylaeus 261
nigrodorsata, Tianmuthredo 100
nigrodorsatus, Nematus 69
nigroelypeata, Prosopis styriaca var. 261
nigrofasciata, Andrena scita var. 273
nigrofasciata, Tenthredo 87
nigromemorata, Tenthredo, *Tenthredella*
 velox 95
nigroflava, Dineura 55
nigrofulva, Anthophora 323
nigrolabiata, Ceratina 310
nigrolateralis, Tenthredo 92
nigrolineata, Macrophyta 87
nigrolineatus, Hylaeus 260
nigrolineatus, Nematus 64
nigrolineatus, Tenthredopsis
 sachalinensis 98
nigromaculata, Anthophora 324
nigromaculatus, Nematus 69
nigromongolica, Pristiphora 73
nigronodosa, Arge, *Hylotoma* 24
nigronota, Pteronidea 60
nigronotata, Pachyprotasis 87
nigronotata, Pteronidea miliaris 61
nigronotata, Tenthredopsis elegans 99
nigronotatus, Tenthredopsis 99
nigropicta, Tenthredo, *Macrophyta* 93
nigropilosa, Andrena
 truncatilabris ssp. 275
nigropuncticeps, Pristiphora 70
nigror, Crabro 228
nigroscutellata, Athalia
 rufoscutellata 45

nigroscutellata, Macrophyta
 brevilabris 87
nigroscutellata, Rhogogaster viridis 88
nigroscutellata, Tenthredopsis 98
nigroscutellatum, Diprion pini 33
nigroscutellatus, Allantus
 marginellus 96
nigrosericea, Andrena 269
nigrospina, Andrena 272
nigrosterna, Andrena 267
nigrosternatus, Nematinus 68
nigrosternum, Pachyprotasis rapae 87
nigrostigmata, Mesoneura 69
nigro-stincta, Andrena 267
nigrotibia, Eumenes coarctatus ab. 181
nigrotibialis, Halictus leucozonius 281
nigrovaginata, Arge 24
nigroventralis, Hortobombus
 consobrinus 328
nigroviolacea, Cimbex variabilis 29
nigrovittata, Tenthredella olivacea 92
nigrum, Anthidium oblongatum var. 301
nigrum, Lasioglossum auct. 283
nikiforuki, Andrena verae ssp. 267
nikitae, Leptothorax 208
nikkoensis, Halictus 278
nikkoensis, Psenulus 252
nikkoensis, Spilomena 249
nikolajevskii, Gorytes 221
nikolaji, Chrysura 139
nikolskajae, Heterocoelia 125
nikolskayae, Halictus, *Seladonia* 279
nilotica, Vespa 194
nimbatus, Hylaeus 260
nimpha, Polistes, *Vespa* 193
nipanicus, Euodynerus, *Lionotus* 183
nipanicus, *Lionotus* tomentosus var. 183
niponica, Bembix 222
nippon, Andrena 273
nippon, Hylaeus 261
nippon, Sphecodes 289
nippon, Tenthredo viridatrix 92
nipponense, Lasioglossum, *Halictus* 285
nipponensis, Arge 24
nipponensis, Formica fusca var. 200
nipponensis, Lasius 201
nipponensis, Lasius fuliginosus var. 201
nipponica, Ammophila sabulosa 213
nipponica, Deuteragenia 164
nipponica, Dolichovespula saxonica 195
nipponica, Megachile 307
nipponica, Myrmecina 205
nipponica, Myrmecina graminicola 205
nipponica, Nesoselandria 81
nipponica, Phymatocera 52
nipponica, Rhogogaster 93
nipponicola, Lasioglossum 280
nipponicola, Passaloecus 248
nipponicus, Alysson sibiricus 219
nipponicus, Hylaeus 260
nipponicus, Monoctenus 35
nipponicus, Neostromboceros 81
nipponicus, Pamphilus
 zhelochovtsevi 106

nipponicus, Pseudomalus, *Omalus* 133
nipponicus, Tachytes 240
nipponis, Argogorytes 221
nipponis, Crossocerus heydeni 228
Nipponodipogon 164
Nippononysson 219
Nippaponopson 251
Nitela 241
Nitela subg. 241
Niteliopsis 241
Nitellopterus 240
nitens, Andrena 269
nitens, Brachythops, *Selandria* 76
nitens, Dolerus 80
nitens, Nematus 64, 72
nitens, Prenolepis, *Tapinoma* 203
nitens, Tapinoma 202
nitida, Andrena 281
nitida, Andrena, *Apis* 269
nitida, Deuteragenia 164
nitida, Eriocampa 46
nitida, Pemphredon 249
nitida, Tenthredo 43
nitidiceps, Tenthredo, *Tenthredella* 91
nitidior, Allantus arcuatus 96
nitidior, Tachysphex 239
nitidipleuris, Euura, *Amauronematus* 62
nitidiuscula, Andrena 271
nitidiuscula, Prosopis 260
nitidiusculum, Lasioglossum, *Melitta*
 nitidiusculum 285
nitidiventris, Andrena 269, 270
nitidiventris, Eucera 320
nitidula, Ceratina 310
nitidula, Chrysis 135
nitidula, Hoplitis, *Osmia* 297
nitidula, Meria 149
nitidula, Myrmica 204
nitidulum, Anthidium 301
nitidulum, Lasioglossum, *Prosopis*
 nitidulum 285
nitidulus, Cleptes, *Ichneumon* 127
nitidulus, Formicoxenus, *Myrmica* 204
nitidulus, Halictus 285
nitidulus, Hylaeus 259
nitidulus, Sphecodes 288
nitidus, Megalodontes 107
nitidus, Tachysphex, *Astata* 239
nitobei, Clavellaria 30
nivalabnormis, Arachnospila,
Pompilus 169
nivalis, Euura, *Pontania* 62
nivalis, Nematus 58
nivalis, Tenthredo 96
niveata, Andrena 270
niveata, Osmia, *Anthophora* 298
niveatus, Bombus 330
niveatus, Prionyx, *Sphex* 215
niveatus, Sphecodes olivieri 289
niveipennis, Sphecodes pellucidus 289
niveocinctum, Lasioglossum,
Halictus 281
niveofasciatus, Colletes 257

niveopicta, *Pseudepipona* 188
nivosa, *Macrophyia* 84
Nobandrena subg. 271
nobile, *Hedychrum* 130
nobile, *Hedychrum*, *Sphex* 130
nobiliforme, *Hedychrum* 130
nobilis, *Andrena* 271
nobilis, *Cerceris* 253
nobilis, *Gorytes* 222
nobilis, *Nematus* 60
nobilis, *Nomada* 315
nobilis, *Ussurinus* 100
nobilis, *Vespa magnifica* var. 194
noctilio, *Sirex* 112
nodicorne, *Lasioglossum*, *Halictus* 285
nodicornis, *Rhadinoceraea* 52
nodicornis, *Sphecodes* 287
Nodulodentiden 77
nodus, *Cataglyphis*, *Formica* 199
Nogatopus 120
Nokbibula 147
Nomada 310
Nomadinae 310
Nomadini 310
Nomadita 310
Nomadosoma 311
Nomiapis 291
Nomiinae 277, 291
Nomineia 123
Nomioides 292
Nomioides subg. 292
Nomiodinae 277, 292
nomioidis, *Sphecodes* 289
nomurae, *Cimbex* 29
nonultimus, *Dolerus* 80
nonveilleri, *Apterogyna* 159
nonveilleri, *Smicromyrme* 156
noomen, *Hylaeus* 261
nordeneae, *Krombeinictus* 217
Norumbega 226
norvegica, *Andrena* 264
norvegica, *Megilla furcata* var. 323
norvegica, *Pontania* 64
norvegica, *Tentredella fuscicornis* 95
norvegicus, *Bombus*, *Psithyrus* 329
norvegicus, *Halictus* 283
norwegica, *Dolichovespula*, *Vespa* 195
norwegica, *Vespa* var. *adulterina* 194
noskiewiczi, *Nomada* 315
nostras, *Dasylobus* *maura* 153
notabilis, *Nematus* 72
notabilis, *Nysson* 220
notabilis, *Odynerus* 178
notanda, *Pteronidea pavida* 63
Notandrena subg. 271
notata, *Bembix* 222
notata, *Tenthredo* 64, 86
notatiformis, *Euodynerus posticus*
 posticus var. 183
notatus, *Crabro* 230
notatus, *Cryptochelius*, *Sphex* 161
notatus, *Euodynerus*, *Vespa* 183
notatus, *Nematus* 66

notha, *Tenthredo notha* 97
Nothocrabro 227
Nothosphecius 224
Notocyphinae 160
Notoglossa 242
Notogonia 236
Notogonidea 237
Notogonius 236
notomelas, *Mutilla* 157
notomelas, *Tenthredo* 93
Notorhopalum 235
Notozus 127
nova, *Andrena* 264
novaki, *Smicromyrme* 156
novellai, *Melecta* 326
novellus, *Odynerus* 178
novemguttata, *Macrophyia* 85
novioregensis, *Nomada* 315
novittola, *Athalia* 44
novoglabrenovi, *Dolerus* 81
novoglabrenovi, *Tenthredopsis* 83
nox, *Chrysis* 137
nubecula, *Agenioideus*, *Pompilus* 167
nubilus, *Cladius*, *Trichiocampus* 54
nucleolus, *Halictus smethmanellus* 285
nudatus, *Prionyx*, *Sphex* 215
nudigastra, *Andrena* 265
nudigastroides, *Andrena*
 nudigastra ssp. 265
nudipectus, *Euura*, *Pontania* 62
nudipes, *Priocnemis* 163
nudiuscula, *Melitta* 273
nugarius, *Eumenes coarctatus* 181
nugdunensis, *Microdynerus*,
 Odynerus 186
numida, *Anthidium* 301
numidica, *Megachile circumcincta* 307
nundius, *Dolerus* 78
nuortevai, *Euura*, *Pteronidea* 62
nupricola, *Lasioglossum* 283
nupta, *Cerceris* 254
Nurmiya 149
nursei, *Odynerus* 191
nursei, *Palarus* 243
nutricia, *Dolerus* 78
nyctea, *Tenthredo* 85
nyctelius, *Dolerus* 78
Nycteridium 35
nycthemera, *Andrena* 264
Nyctovespa 193
nyemitawa, *Arge* 24
Nygmapolistes 192
nylanderi, *Andrena* 271
nylanderi, *Formica rufa* var. 199
nylanderi, *Halictus* 285
nylanderi, *Psen* 252
Nylanderia 202
nymphae, *Bombus* 330, 331
Nyssa 220
Nysson 219
Nyssonini 219
Nyssonus 220
Oar 130
obducta, *Euura*, *Nematus* 62
oberbergeri,
 Smicromyrme variabilis f. 155
obensis, *Myrmica limanica* 205
oberthuri, *Vespa* 194
obesa, *Arge* 24
obesa, *Tenthredo* 43
obesus, *Allantus* 90
obesus, *Amauronematus* 64
obliquus, *Crabro* 229
oblita, *Euura*, *Nematus* 62
oblita, *Pristiphora* 55
oblitus, *Nematus* 62
oblongatum, *Anthidium*,
 Anthophora 301
oblongus, *Dolerus* 79
oblongus, *Nematus* 72
obnata, *Macrophyia* 87
obo, *Ammophila* 213
obo, *Diodontus* 247
obovata, *Melitta* 282
obscura, *Andrena florea* var. 273
obscura, *Apis* 268, 269, 287
obscura, *Athalia rosae* 44
obscura, *Coelioxys* 305
obscura, *Corynis*, *Tenthredo* 32
obscura, *Ctenotilla pectinifera* var. 155
obscura, *Dolerus violaceus* 79
obscura, *Emphytus perla* 39
obscura, *Megachile* 307
obscura, *Megachile*
 kuenemannii var. 307
obscura, *Mutilla* 157
obscura, *Nomada* 315
obscura, *Prosopis variegata* var. 262
obscura, *Tenthredo* 31
obscura, *Tenthredo* 93
obscura, *Thomsonia* 99
obscuraria, *Vespa* 196
obscuraria, *Tenthredella luteipennis* 95
obscurata, *Andrena* 267
obscurata, *Athalia rufoscutellata* 45
obscurata, *Lasius brunneus* var. 202
obscurata, *Nomioides minutissima* 292
obscurata, *Prosopis* 260
obscurata, *Tenthredo atra* 97
obscurata, *Tenthredopsis austriaca* 99
obscuratum, *Lasioglossum*, *Halictus*
 obscuratum 285
obscuratus, *Lasius* 202
obscuratus, *Monocetus*, *Lophyrus* 35
obscuratus, *Nematus* 56
obscuratus, *Paracharactus* 51
obscurilla, *Anthophora furcata* f. 323
obscuricanda, *Andrena* 273
obscuricornis, *Polistes*
 foederata var. 193
Obscuriformis 31
obscurior, *Nematus* 73
obscurior, *Paravespa rex* var. 188
obscurior, *Pteronidea salicis* 64

- obscuripennis*, *Tachysphex*,
Tachytes 239
obscuripennis, *Tachytes* 239
obscuripes, *Camponotus* 198
obscuripes, *Camponotus herculeanus*
 ligniperdus var. 198
obscuripes, *Euura, Nematus* 62
obscuriventris, *Mesoneura opaca* 68
obscurum, *Anthidium diadema* var. 300
obscurum, *Anthidium latreillei* var. 301
obscurus, *Allantus* 94
obscurus, *Allantus costatus* 90
obscurus, *Amauronematus* 65
obscurus, *Bombus lapponicus* 330
obscurus, *Crabro* 231
obscurus, *Lithurgus cornutus* var. 295
obscurus, *Mellinus* 246
obscurus, *Mesitius* 125
obscurus, *Misophus cypriacus* 241
obscurus, *Nematus* 73
obscurus, *Platycampus pectoralis* 53
obscurus, *Ukayakos, Mesitius* 125
obsidiana, *Aphaenogaster, Atta* 203
Obsoleta, *Apis* 240
obsoleta, *Neurotoma nemoralis* 103
obsoleta, *Tenthredo* 91
obsoletus, *Cleptes* 127
obsoletus, *Halictus* 285
obsoletus, *Tachytes, Apis* 240
obstrictus, *Stenancistrocerus*,
Odynerus 189
obtusa, *Tenthredo* 43
obtusalis, *Empria pulverata* 43
obtusata, *Coelioxys* 304
obtusatum, *Anthidium* 300
Obtusia 51
obtusidens, *Chrysis* 137
obtusidentatus, *Odynerus* 185
obtusifrons, *Nomada*, *Nomada* 315
obtusispina, *Coelioxys* 305
obtusiventris, *Pompilus* 163
obtusus, *Coelioxys* 304
obtusus, *Pompioides* 168
obversus, *Paracharactus* 50
occidens, *Lasioglossum, Halictus* 281
occidenta, *Holopyga amoena* 131
occipitalis, *Ammophila* 212
occipitalis, *Andrena* 269
occipitalis, *Nematus* 64
occipitalis, *Priocnemis* 163
occipitosum, *Aglaostigma*,
Macrophyopsis 83
occitanicum, *Chlorion* 214
occitanicus, *Palmodes, Sphecodes* 214
occitanicus, *Sphecodes* 214
occultodistinctus, *Bombus*
 lapponicus 330
occultus, *Nematus* 73
oceanicus, *Bombus* 330
oceanicus, *Leptocephalum*,
Mychothorax 204
ocellaris, *Pemphredon* 249
ocellata, *Bembix* 222
ochlerus, *Odynerus* 179
ochotica, *Abia, Orientobia* 29
ochotica, *Lestica, Crabro*
 subterranea 234
ochoticus, *Crabro* 234
ochracea, *Andrena* 273
ochracea, *Cryptopone, Ponera* 209
ochraceum, *Aglaostigma*,
Macrophyopsis 83
ochraceus, *Nematus* 63
ochroleuca, *Amegilla, Anthophora* 322
ochroleuca, *Melicerta* 36
ochroleucus, *Heptamelus, Melicerta* 36
ochroleucus, *Hortobombus*
 consobrinus 328
ochroneurus, *Dolerus* 80
ochropoda, *Heterarthrus, Tenthredo* 49
ochropoda, *Tenthredo* 49
ochroptera, *Mimesa* 250
ochroptera, *Tenthredo* 25
ochropus, *Arge, Tenthredo* 24
ochropus, *Nematus* 57
Ochropusiformis 22
ochropyga, *Andrena* 270
ochrostoma, *Apis* 316
ochrostoma, *Nomada* 311
Ocla 39
ocreata, *Apis* 274
ocreata, *Tenthredo* 85
octavonotatus, *Solenius* 232
octoguttata, *Scolia* 145
octomaculata, *Stelis* 302
octomaculatus, *Cryptocoelius*,
Sphex 162
octopunctata, *Bembix* 222
octosignata, *Megachile* 307
octostrigata, *Andrena* 274
oculata, *Bembix oculata* 222
oculata, *Dimorpha* 218
oculata, *Nomada* 315
oculata, *Scolia, Discolia* 148
oculata, *Solenopsis orbula* var. 207
oculatus, *Hylaeus* 260
Odontaporus 166
Odontopyris 125
Odontocrabro 235
Odontodynerus 180
odontogastra, *Dufourea* 290
Odontolarra 240
odontophora, *Cerceris* 254
odontophorus, *Belomicrus*,
Oxybelus 242
Odontophyes 20
Odontopterochilus 189
odontopyga, *Stelis* 302
odontostoma, *Piagetia* 237
odontura, *Megachile* 305
odoratus, *Aphanisus* 45
odoratus, *Lasius flavus* 201
odyneroides, *Psiliglossa*,
Raphiglossa 188
odyneroides, *Raphiglossa* 188
Odynerus 186
Odynerus subg. 186
oehlkei, *Hylaeus* 260
oertzeni, *Aneugmenus, Selandria* 76
ogasawarai, *Xiphydria* 116
ogivalis, *Coelioxys* 305
ogloblini, *Didineis* 219
ogloblini, *Selandria* 76
ogumae, *Pomilus* 172
ogumae, *Sirex* 113
ohdeyamanus, *Sphecodes* 287
ohnonis, *Psen* 251
ohtsukius, *Sphecodes* 290
oishii, *Caliroa, Eriocampoides* 46
okabei, *Crabro* 226
okamotonis, *Nomada* 315
Okamotonius 30
okeanskajensis, *Dolerus* 81
okeanskayanum, *Trypoxyylon* 245
Okutanius 51
okuyetsu, *Sphecodes* 289
olcesei, *Tiphia* 150
olfaciens, *Euura, Nematus olfaciens* 62
olgae, *Coelioxys mandibularis* 304
olgae, *Ellampus* 128
Olgia 224
olhae, *Nomada* 315
Oligonematus 70
oligospila, *Euura, Nematus* 62
oligotricha, *Andrena*
 bimaculata ssp. 272
olivacea, *Bembix olivacea* 222
olivacea, *Lasius flavus* var. 201
olivacea, *Tenthredo olivacea* 92
Olivacedo 89
Olivacedo subg. 92
olivaceus, *Diversobombus*
 ussuriensis 329
olivaceus, *Nematus* 62
olivieri, *Apterogyna* 159
olivieri, *Cimbex* 32
olivieri, *Eumenes* 180
olivieri, *Sphecodes* 289
olivieri, *Xylocopa* 310
Olochrysis 139
olympicum, *Lasioglossum, Halictus*
 bavaricum 282
Omachthes 319
Omalus 131
omega, *Macrophyia* 87
omega, *Tenthredo, Tenthredella* 98
omissa, *Andrena pubescens* ssp. 269
omissa, *Dolichovespula, Vespa* 195
omissa, *Polistula* 193
omissa, *Tenthredo, Allantus* 97
omissoides, *Allantus* 96
omissus, *Nysson* 220
omnicincta, *Tenthredo vespa* 97
Omphalius 225
Onchopterocheilus subg. 189
Oncodolerus 77
Oncodolerus subg. 79
oncogaster, *Stenocephalus* 111
Onerarius 192

ononides, Osmia 299
Onycholyda 103
Onychopterocheilus 187
Onychopterocheilus subg. 187
opaca, Andrena 273
opaca, Mesoneura, *Tenthredo* 68
opaca, Nomada 315
opaca, *Tenthredo* 68
opacella, Rhogogaster 92
opaciceps, *Tenthredo*, *Tenthredella* 91
opacicollis, Andrena 265
opacicollis, Lagonis, *Rhadinoceraea* 49
opacifovea, Andrena 274
opacifrons, Halictus 283
opacifrons, Sphecodes 288, 289
opacipleuris, *Tenthredopsis* 100
opaciventris, Andrena 263
opacomaculata, *Tenthredo* 41
opacum, Trichiosoma 31
opus, Ammobates 318
opus, Eumenes mediterraneus var. 182
opus, Hylaeus 261
opus, *Odynerus* 188
opus, Oxybelus 242
opaca, Formica 199
opalina, Scolia 147
opalipennis, Cerceris 253
opalipennis, Liris, *Notogonia* 237
Opandrena subg. 271
opima, Osmia 299
opinabilis, Polistes 193
opinata, Arachnospila, *Pompilus* 169
Oplomerus 186
Opoporus 186
optata, Andrena rufomaculata ssp. 275
optimellus, Halictus 282
opulenta, Eumenes coarctatus var. 181
opulenta, *Tenthredopsis* 98
oralis, Andrena 271
oraniense, Anthidium 300
oraniensis, Ammobates, *Phileremus* 318
oraniensis, Aencyla 319
oraniensis, Andrena 267
oraniensis, Megachile ericetorum 307
Oraniensis, Tachytes 239
oraria, *Tenthredo* 95
orarius, Pachynematus 73
oratoria, Phytospodida,
 Smicromyrme 155
oratorius, Haplogonatopus,
 Gonatopus 121
orbatus, Thyreus, *Crocisa* 327
orbiculatus, Ammoplanus 246
orbitalis, Antepipona, *Odynerus* 180
orbitalis, Lionotus 190
orbitalis, Nematus 60
orbitalis, *Odynerus* 190
orbitalis, Periclista albida 51
orbitalis, Pompilus 168
orbitalis, *Tenthredella* atra 93
orbitalis, *Tenthredo* 99
orchonensis, Dolerus 78
ordinaria, Tiphia 150

ordubadensis Cladius 54
ordubadensis, Eumenes pomiformis 181
ordubadensis, Polistes omissus var. 193
orenburgensis, Andrena 268
orenburgensis, Nomada bifida 316
orenburgensis, Stenodynerus,
 Odynerus 190
Oreocrabro 231
Oreomelissa subg. 271
oresterus, Psen 251
Orientabia 28
orientale, Liometopum 197
orientaliella, Andrena 267
orientalis, Allantus 90
orientalis, Anthophora 325
orientalis, Athalia spinarum 44
orientalis, Cephus 111
orientalis, Cladius 54
orientalis, Coelioxys 304
orientalis, Dolerus germanicus 78
orientalis, Eopompilus 162
orientalis, Eucera nigrilabris 320
orientalis, Evagetes 171
orientalis, Formica fusca picea 200
orientalis, Formicoxenus 204
orientalis, Halictus cylindricus 285
orientalis, Hemichroa 67
orientalis, Janus 110
orientalis, Lasius 201
orientalis, Lasius fuliginosus var. 201
orientalis, Liometopum
 microcephalum var. 197
orientalis, Macrophyia rufipes 86
orientalis, Mimesa 251
orientalis, Mychothorax acervorum 204
orientalis, Myrmica ruginodis 206
orientalis, Odontepyris, *Prosierola* 125
orientalis, *Odynerus* 180
orientalis, *Odynerus* trifasciatus 180
orientalis, Osmia 298
orientalis, Palmodes, *Sphex* 215
orientalis, Priocnemis fuscus 164
orientalis, Prosopigastra 237
orientalis, Prosopigastra
 punctatissima 237
orientalis, Psen 251
orientalis, Smicromyrme 156
orientalis, Solenopsis fugax 207
orientalis, Sphecodes 289
orientalis, Sterictiphora, *Schizocera* 26
orientalis, Thrinax contigua 82
orientalis, Urocerus gigas 113
orientalis, Vespa 194
orientausa, Arachnospila 169
orienticola, Andrena humilis var. 265
orientis, Anoplonyx 53
orientis, Euchroeus 144
orientis, Halictus morbillulosus r. 280
orientis, Tachytes panzeri 240
Orientoheriades 296
Orientosmia 297
Orientotrichusa 301
Orlovinga 147
ormillatus, Hylaeus 262
ornata, Andrena 273
ornata, Cerceris 254
ornata, Cimex 29
ornata, Cimex femorata 29
ornata, Polistes 193
ornata, Polistes gallica var. 193
ornata, *Tenthredo* 99
ornata, *Tenthredo*, *Allantus* 90
ornata, *Tenthredopsis*, *Tenthredo* 99
ornatipennis, Tachysphex 239
ornatrix, *Tenthredopsis* 99
ornatula, Stelis, *Gyrodroma* 302
ornatus, Loderus 79
ornatum, Anthidium 300
ornatum, Rhynchium 187
ornatus, Anoplonyx 53
ornatus, Crabro 233
ornatus, Crossocerus 229
ornatus, Philanthus 252
ornatus, Stizus 225
Ornicrabro 227
orontis, Halictus 280
Orthocrabro 227
Orthoxybelus 242
ortinga, Pristiphora 72
Orussidae 117
Orussinae 117
Orussoidea 117
Orussomorpha 117
Orussus 117
oryssoides, *Tenthredo* 95
Oryttus 224
Osirini 319
Osmia 295, 297
Osmia subg. 299
Osmiini 295
ossetica, *Proformica* 203
ostriaria, *Pristiphora* 73
osychniukae, Andrena 275
osychniukae, Andrena
 abbreviata ssp. 275
osychniuki, Andrena 275
osychniuki, Andrena abbreviata ssp. 275
Othyreus 226
ottomana, *Macrophyia* 86
ottomanus, *Hoplisus* 224
ouchii, Lasius 201
ouralensis, *Tenthredo*, *Allantus* 93
ovalis, Crossocerus 229
ovata, Andrena 274
ovata, *Eriocampa*, *Tenthredo* *ovata* 42
ovata, *Holopyga* var. b 131
ovata, *Sphex clypeata* 234
ovata, *Tenthredo* 42
ovatula, Andrena, *Melitta* 274
ovatus, *Anoplonyx*, *Leptopus* 53
ovatus, Crabro 229
ovicula, *Corynis* 32
ovidorsalis, *Tiphia* 150
ovina, Andrena 269
ovinella, Andrena 269
ovinigris, *Tiphia* 150

oviventris, *Ancistrocerus, Odynerus* 179
Oxaeinae 263
oxalata, *Pseudoselandria* 76
oxyacanthea, *Trichiosoma* 31
Oxybelini 241
Oxybeloides 241
Oxybelus 242
Oxybiastes 317
Oxycleptes 126
Oxycrabro 227
oxygona, *Anthophora* 324
zbeki, *Megachile circumcincta* 307

paccia, *Sirex* 145
Pachycephalus 110
pachycerus, *Dolerus* 77
pachycerus, *Nematus* 62
Pachygonatopus 120
Pachynematus 55
Pachyosmia 297
Pachyprotasis 87
pachysoma, *Sphex* 216
pachyvalvis, *Lygaeonematus* 71
pacifica, *Apis* 306
pacifica, *Argo ciliaris* 23
pacifica, *Blasticotoma filiceti* 27
pacifica, *Dolichovespula*, *Vespula* 195
pacifica, *Macrophyia* 89
pacifica, *Nomada* 315
pacifica, *Orientabia* 28
pacifica, *Pemphredon* 249
pacifica, *Polistes* 193
pacifica, *Polistes gallica* var. 193
pacifica, *Vespula norvegica* 195
pacificum, *Trypoxylon* 245
pacificus, *Agenioideus* 167
pacificus, *Amauronematus* 65
pacificus, *Auplopus* 165
pacificus, *Crabro* 229
pacificus, *Lestiphorus, Gorytes* 224
pacificus, *Simplemphytus* 39
pacis, *Leptothorax, Doronymyrmex* 204
padi, *Aneugmenus*, *Tenthredo* 76
paedemontana, *Mutilla* 155
Paedia 318
paedidus, *Lygaeonematus* 72
paehypoda, *Anthophora vulpina* 323
pagana, *Argo*, *Tenthredo pagana* 24
paganettii, *Andrena fimbriata* 267
Paganiformis 22
paganiformis, *Argo* 24
nahanganus, *Halictus* 287
Paidia 318
palaeantarctica, *Scolia* 148
palaeantarctica, *Xiphydria* 116
Palaeocimbex 29
palaestinensis, *Eumenes dubius*
 dubius var. 181
palaestinus, *Cephus pygmaeus* 109
Palarini 243
Palarus 243
palestinensis, *Anthophora*
 acervorum 323

palestinensis, *Apterogyna* 159
palestinensis, *Artiotilla biguttata* f. 157
Palisotius 192
paliuri, *Andrena* 271
Pallandrena subg. 271
pallasi, *Elampus, Notozus* 128
pallasi, *Onychopterocheilus*,
 Pterocheilus 187
pallens, *Cimbex* 29
pallens, *Euura, Pteronus* 62
pallens, *Lasiglossum, Halictus* 281
pallens, *Lygaeonematus* 56
palleola, *Dasypoda* 293
pallescens, *Colletes* 257
pallescens, *Crocisa* 326
pallescens, *Euura, Nematus* 62
pallescens, *Hylaeus* 260
pallescens, *Monophadnus, Tenthredo* 50
pallescens, *Plagiolepis* 202
pallescens, *Vespa* 222
palliatus, *Nematus* 60
pallicarpus, *Nematus* 56
palliceps, *Nematus* 57
pallicornis, *Nematus* 70
pallicoxa, *Pristiphora* 73
pallida, *Bembix* 222
pallida, *Cataglyphis* 199
pallida, *Cimbex* 29
pallida, *Dineura* 67
pallida, *Eucera caspica* var. 320
pallida, *Gilpinia, Lophyrus* 34
pallida, *Pristiphora, Lygaeonematus* 73
pallida, *Selandria* 67
pallida, *Sirex augur* 113
pallida, *Tenthredo* 67
pallida, *Tenthredopsis* 99
pallidenotata, *Nomada* 313
pallidicarpus, *Nematus* 56
pallidiceps, *Nematus* 57
pallidicincta, *Apis* 258
pallidicornis, *Aproceros, Aprosthemma* 25
pallidicornis, *Hylaeus* 259
pallididens, *Prosopis* 261
pallidilabris, *Macrophyia* 85
pallidinervis, *Argo* 24
pallidior, *Euura, Amauronematus* 62
pallidipalpis, *Crossocerus* 229
pallidipes, *Cephus* 109
pallidipes, *Emphytus* 39
pallidipes, *Halictus* 286
pallidipes, *Hylotoma* 22
pallidipes, *Lophyrus* 34
pallidipes, *Pamphilus* 105
palliditarsis, *Mimesa* 251
palliditarsis, *Nematus*
 brachyacanthus 66
pallidiventralis, *Nematus* 62
pallidiventris, *Pristiphora, Tenthredo* 73
pallidiventris, *Tenthredo fulva* 91
pallidocaudatus, *Bombus*
 lapponicus 330
pallidopicta, *Cerceris* 253
pallidula, *Cephalcia, Cephaleia* 102
pallidula, *Pheidole, Myrmica* 207
pallidula, *Pristiphora* 71
pallidula, *Tenthredo* 96
pallidulus, *Sphecodes* 282
pallidum, *Lasiglossum, Nomiooides* 285
pallidus, *Cimbex* 29
pallidus, *Gonatopus, Dicondylus* 121
pallidus, *Pamphilus* 105
pallilolum, *Lasiglossum, Halictus* 285
pallilomus, *Halictus* 286
pallimacula, *Dolerus* 40
pallimacula, *Dolerus* 41
pallimacula, *Empria, Dolerus* 41
pallimaculatus, *Emphytus* 41
pallior, *Tenthredella flavicornis* 91
pallipes, *Ametastegia, Tenthredo* 39
pallipes, *Andrena ephippium* var. 270
pallipes, *Ardis, Dolerus* 45
pallipes, *Birkha* 76
pallipes, *Calameuta, Astatus* 109
pallipes, *Cladius* 54
pallipes, *Cleptes* 127
pallipes, *Dineura* 68
pallipes, *Dolerus* 45
pallipes, *Emphytus* 39
pallipes, *Halictus* 285
pallipes, *Hylotoma* 22
pallipes, *Hylotoma* 34
pallipes, *Microdiprion, Hylotoma* 34
pallipes, *Monsoma, Poecilosoma* 43
pallipes, *Pamphilus, Lyda* 105
pallipes, *Pristiphora* 70
pallipes, *Pseneulus, Sphex* 252
pallipes, *Schizocera* 26
pallipes, *Tenthredo* 71
pallipes, *Trachelus* 111
pallispinosa, *Nomada* 315
pallitarsis, *Andrena* 271
palliventris, *Nematus* 62
palmaria, *Sphex* 229
palmata, *Tenthredo* 79
palmatus, *Crabro* 226
palmatus, *Crossocerus* 229
palmeni, *Nomada* 315
palmicornis, *Cladius* 54
palmipes, *Apis* 323
palmipes, *Crossocerus, Sphex*
 palmipes 229
palmipes, *Sphex* 150
palmipes, *Sphex* 227
Palmodes 214
paludosa, *Isodontia, Sphex* 215
palumboi, *Andrena* 264
palumbula, *Larra* 237
palustris, *Prosopis* 261
palustris, *Tenthredo* 77, 92
pamirensis, *Andrena nigriceps* ssp. 265
pamirensis, *Polistes* 192
pamirica, *Anthophora fuliginosa* 324
pamiricola, *Anthophora parietina* 324
pamiricus, *Euodynerus dantici* 183
pammelas, *Crossocerus* 227
Pamphiliidae 101

Pamphiliinae 102
Pamphilioidea 101
Pamphilus 104
pamyrensis, *Tenthredo* 97
pannonica, *Eucera* 321
pannonica, *Mimesa* 250
pannonica, *Osmia* 279
pannonicus, *Alastor biegelebeni* 178
pannonicus, *Halictus* 278
panurgimorpha, *Andrena* 265
Panurginae 263, 275
Panurgini 275
Panurginus 276
Panurgus 276
panzeri, *Bembix* 222
panzeri, *Chrysis* 127
panzeri, *Elampus*, *Chrysis* 128
panzeri, *Emphytus* 40
panzeri, *Hylaeus* 261
panzeri, *Lindenius*, *Crabro* 235
Panzeri, *Lyrops* 240
panzeri, *Megachile leucomalla* var. 306
panzeri, *Mutilla* 157
panzeri, *Myzine* 146
panzeri, *Nomada* 315
panzeri, *Nysson* 220
panzeri, *Osmia* 299
panzeri, *Tachysphex*, *Tachytes* 239
panzeri, *Tachytes* 240
panzeri, *Tachytes*, *Lyrops panzeri* 240
panzeriana, *Nomada* 315
pagorum, *Crabro* 231
papaveris, *Hoplitis*, *Apis* 297
papillarius, *Eumenes*, *Sphex* 182
papillosa, *Euura*, *Tenthredo* 63
pappi, *Metriondonotus* 124
pappi, *Metriondonus* 124
pappi, *Stenodynerus* 190
pappi, *Tenthredo* 92
Papuacrabro 231
par, *Euchroeus*, *Spinolia* 140
Paraanthidium subg. 303
Parabatozonus 167
Parabia 28
Paracalyza 123
Paraceramius 244
Paracharactus 51
paraclypeata, *Eucera* 321
Paracoelioxys 304
Paradicordylus 120
paradiformis, *Hylaeus* 261
Paradirus 110
paradoxa, *Bembix* 222
paradoxa, *Eucera* 321
paradoxa, *Solierella* 241
paradoxum, *Pararrhynchium*,
Odynerus 187
paradoxus, *Bombus confusus* 327
paradoxus, *Pararhabdepyris* 124
Paradryinus 119
Paraferrereola auct. 173
Paragonatopus 120
Paragymnomerus 187

Paralaelius 123
paralella, *Pristiphora*, *Nematus* 73
Paralelopis 236
paralienus, *Lasius* 202
parallela, *Vespa* 193
parallelus, *Apethymus*, *Emphytus* 40
parallelus, *Aphanisus* 45
parallelus, *Nematus* 58
Paralloma 83
Paramacrophyia 84
Paramacropis subg. 294
Paramasaakia 51
Paramegilla subg. 324
Parammabatodes 319
paramushirensis, *Hemicroa* 67
Paramyrme 155
Paramyrmosa 152
Paramyrmosa 152
paramythensis, *Andrena* 275
Parandrenella subg. 271
Paranematus 69
Paranoplus 172
Paranothyreus 225, 226
Paranothyreus 226
Paranteon 118
Paranthidellum 301
Paraphylotoma 47
Parapiagetia 237
Parapison 244
Paraprosopis subg. 261
Pararhabdepyris 124
Pararrhynchium 187
Paraselandria 76
Parasierola 125
Parasiobia 43
Parasphe 215
Parastalor subg. 178
Parastatis 89
Parastenancistrocerus subg. 184
Parastromboceros 75
Parasymmorphus subg. 190
Parataxonius 40
Parathredo 90
Parathredo subg. 92
parathoracica, *Andrena* 269
Parathyreopus 225, 226
Paratropancistrocerus subg. 189
Paravespa 187
Paravespa subg. 188
Paravespula 195
Paraxyela 20
parcivalvis, *Dineura*, *Pontania* 55
Pardesiya 147
pardo, *Bembix* 222
Pareixenius 173
Parellampus 127
Pareophora 51
Parepyris 123
Pareudynerus subg. 183
Parhypodynerus 189
paria, *Tenthredo moniliata* 95
parietina, *Apis* 324
parietina, *Megachile*, *Apis* 305
parietina, *Osmia* 299
parietinus, *Ancistrocerus*, *Vespa* 179
parietoides, *Odynerus parietum* var. 179
parietum, *Ancistrocerus*, *Vespa* 179
parietum, *Vespa* 178
parietum, *Vespa* 195
paripunctata, *Mutilla* 157
parisiensis, *Lyda* 103
parisinus, *Crabro* 243
Parisobrachium 123
parkanense, *Hedychridium* 129
parkanensis, *Crabro* 235
parkanensis, *Lindenius*, *Crabro* 235
parkeri, *Blepharipus* 227
parki, *Acantholyda* 102
Parna 51
parnasius, *Allantus* 90
Parnopes 142
Parnopini 142
Parodontodynerus 188
Paroxycrabro 227
parreyssi, *Cephus* 111
parreyssii, *Syrista*, *Cephus* 111
parumcrinitus, *Heriades* 295
parumpunctata, *Andrena* 268
parumpunctatus, *Halictus* 286
parva, *Nomada jacobaeae* var. 313
parva, *Pristiphora*, *Nematus* 73
parviceps, *Allantus* 91
parviceps, *Andrena* 268
parviceps, *Fenusia* 49
parvicornis, *Cubitalia*, *Eucera* 320
parvicornis, *Nematus* 72
parvicorpus, *Crossocerus wesmaeli* 230
Parvoculus 124
parvula, *Acantholyda* 101
parvula, *Apis* 292
parvula, *Coelioxys* 304
parvula, *Colletes* 264
parvula, *Empria*, *Poecilosoma* 41
parvula, *Eucera* 321
parvula, *Euura*, *Nematus* 63
parvula, *Larrada* 239
parvula, *Megachile* 306
parvula, *Melitta* 270
parvula, *Mutilla* 153
parvula, *Osmia* 296
parvula, *Priocnemis* 163
parvula, *Pseudodineura*, *Tenthredo* 75
parvula, *Tenthredo* 68
parvula, *Tenthredo* 75
parvula, *Tenthredopsis* 100
parvuloides, *Andrena* 270
parvulum, *Anthidium* 302
parvulum, *Lasioglossum*, *Hylaeus* 285
parvulus, *Belomicrus*, *Oxybelus* 242
parvulus, *Crabro* 232
parvulus, *Diodontus*, *Passaloecus* 247
parvulus, *Hylaeus* 259
parvulus, *Microdynerus*, *Odynerus* 186
parvulus, *Odynerus* 180
parvulus, *Temnothorax*, *Myrmica* 208
parvus, *Cephus*, *Pseudocephus* 109

parvus, Cladius 54
parvus, Pseudohemitanoxus 81
pascuorum, Bombus, *Apis* 331
Pasites 319
Pasitomachthes 319
Passaloecus 247
passerini, *Macrophyia diversipes* 85
pastoralis, *Nomada* 315
Patagiata sdubg. 261
patagiatus, *Bombus* 328
patagiatus, *Odynerus* 186
patellarius, *Sphex* 226
patellata, *Prosopis* 259
patellata, *Tenthredo* 39
patellatus, *Colletes* 258
patellatus, *Crabro* 226
patellatus, *Halictus patellatus* 278
patriarchalis, *Chrysis* 141
patricius, *Dolerus* 78
patrizii, *Myzinella* 149
patula, *Coelioxys* 305
patulum, *Lasioglossum*, *Halictus*
 pygmaeum 285
paucipunctatus, *Thrinax*,
 Hemitaxonus 83
paucisquama, *Andrena* 268
paula, *Andrena* 267
pauli, *Odynerus* 185
paulus, *Hylaeus* 260
pauperata, *Andrena* 268
pauperata, *Mutilla* 155
pauperatulellus, *Halictus* 287
pauperatus, *Blepharipus* 230
pauperculum, *Anthidiellum* 300
pauperculus, *Hylaeus* 259
pauperior, *Tenthredella temula* 93
Paururus 112
pauxillum, *Lasioglossum*, *Hylaeus* 285
pauxillus, *Crabro* 228
paveli, *Andrena* 272
pavesii, *Chrysis* 137
pavida, *Euura*, *Nematus* 63
pavida, *Tenthredo* 86
pavidus, *Nematus* 63
Pavlovskia 241
pavlovskii, *Crabro* 231
pavlovskii, *Holopyga* 131
Pavostelis 302
paxillorum, *Heriades* 302
pecki, *Embolemus* 122
Pectinia 52
pectinicornis, Cladius, *Tenthredo* 54
pectinicornis, *Tarpa* 107
pectinifera, *Mutilla* 155
pectinipes, *Evgates*, *Sphex* 171
pectoralis, *Andrena* 269
pectoralis, *Hylaeus* 261
pectoralis, *Monophadnus spinolae* 51
pectoralis, *Nematus* 57
pectoralis, *Nomada* 315
pectoralis, *Polistes* 193
pectoraloides, *Andrena* 269
peculiarus, *Pseudepipona*,
 Odynerus 188

pedata, *Anthophora* 325
pedatoria, *Pachyprotasis*, *Macrophyia* 87
pedestris, *Gonatopus* 121
pedestris, *Sapyga* 145
pedestris, *Tenthredo* 77
pedestris, *Tenthredo* 78
pedetes, *Cerceris* 254
pedicellaris, *Ectemnius*, *Crabro* 232
pedicellaris, *Pompilus* 171
pedicornis, *Osmia* 299
Pedinomma 122
pedunculatus, *Eumenes*, *Vespa* 182
pedunculi, *Euura*, *Nematus* 63
peineae, *Eriocampa* 42
pektusanense, *Prenanteon* 119
pelagius, *Halictus confusus* 278
peletieri, *Aprosthemia*, *Schizocerus*
 peletieri 26
peletieri, *Nematus* 70
pellucida, *Brugmoia* 141
pellucida, *Chrysis* 141
pellucida, *Tenthredo* 94, 95
pellucida, *Trichrysis*, *Chrysis* 141
pellucidus, *Euchroeus* 141
pellucidus, *Sphecodes* 289
Pelmatopus 75
Pelopoeus 214
pelops, *Astatus* 218
peltaria, *Sphex* 226
peltarius, *Crabro*, *Sphex* 226
Pemphilis 225, 226
Pemphredon 248
Pemphredoninae 246
Pemphredonini 246
pendelica, *Andrena potentillae* ssp. 273
pendulus, *Stigmus* 249, 250
penicillata, *Cerceris* 253
peninsularis, *Cerceris* 254
peninsularis, *Tenthredo* 95
pennata, *Anthophora* 323
penninum, *Tetramorium*
 caespitum var. 209
pennipes, *Apis* 325
Pentachrysis 140
Pentadentosmia subg. 297
pentandrae, *Nematus* 56
Pepsinae 160, 161
Pepsini 161
percitus, *Miscophus* 241
peregrinator, *Phymatoceropsis*,
 Neotomostethus 52
peregrinum, *Lasioglossum*, *Halictus* 285
peregrinus, *Hylaeus* 260
peregrinus, *Lamproclytus* 268
perézi, *Eucera* 320
perezi, *Megachile* 306
perfecta, *Meria* 149
perfodata, *Nomia* 291
perforator, *Colletes* 258
perforatus, *Odynerus* 185
pergibbus, *Sphecodes* 288
Periclista 51
Periclista subg. 51
Periclistoptera 45
Perineura 88
Perinevra 88
Perisemus 125
Perisierola 125
Perissostigmus 246
perkinsi, *Allantus* 97
perkinsi, *Amauronematus* 59
perkinsi, *Halictus confusus* 278
perkioemaekii, *Pachynematus* 61
perla, *Ametastegia*, *Tenthredo* 39
perlata, *Tenthredo* 99
perlautus, *Halictus villosulus* 287
permicus, *Halictus* 283
permixtus, *Entomognathus* 234
pernotatus, *Euodynerus notatus* var. 183
Perodryinus 120
Peronistilus 109
perplexa, *Anthophora* 325
perplexans, *Lasioglossum*, *Halictus* 285
perplexus, *Trichomyrmex*,
 Monomorium 209
perrini, *Chrysis* 137
perrisi, *Ammoplanus* 246
perrisi, *Chrysis analis* var. 137
perrisi, *Stizus* 225
perrisi, *Stizus perrisi* 225
perrisi, *Anthidium* 301
persa, *Allantus* 90
persiana, *Myrmica bergi* 205
persica, *Chrysidea* 133
persica, *Tenthredo*, *Allantus* 90
persicum, *Pamphilus* 105
persicus, *Bombus* 331
persicus, *Sphecodes* 289
personata, *Megilla* 323
perspicax, *Tetramorium* 209
perspicax, *Tetramorium caespitum* st.
 ferox var. 209
perspicillaris, *Nematus* 61
pertheesi, *Alysson* 219
perturbator, *Priocnemis*, *Sphex* 164
peruana, *Vespa* 195
perversus, *Sphecodes* 290
pervigil, *Apis* 300
pesenkoi, *Hylaeus* 260
Petalosternon subg. 325
petiolaris, *Anoplus* 172
petiolaris, *Mutilla* 154, 156
petiolata, *Dasylabris*, *Mutilla* 154
petiolatus, *Solenius* 233
petiolatus, *Stigmatodipogon*,
 Dipogon 165
petri, *Elampus*, *Notozus* 128
petrimagni, *Leptocimbex* 30
petroselini, *Andrena* 271
petrosus, *Halictus smethmanellus* 285
Peus 89
pfankuchi, *Hylaeus*, *Prosopis* 260
Phaenusa 48
phaeoptera, *Andrena* 269
phaeoptera, *Stelis*, *Apis* 303
phaeopterus, *Crabro* 226

phaeopterus, Priophorus 54
phalerata, Vespa 189
phaleratus, Pterocheilus, *Vespa* 189
phaneroleuca, Andrena 264
phaneromelas, Andrena 267
phantasma, Xanthosirex 113
phantoma, Hoplocampa 68
pharaonis, Halictus 285
pharaonis, Monomorium, *Formica* 205
pharaonum, Notogonia 237
Phauloclavelia 173
Peidole 207
Phi 180
Phiarus 318
philadelphica, Isodontia 215
philadelphicus, Sphecius 215
Philanthinae 252
Philanthini 255
Philanthinus 255
philanthoides, Crabro 234
Philanthus 255
Phileremus 318
philippi, Tachysphex 238
Philocetes 132
Philoponidea 256
Philoponoides 256
Philoponus 256
Phlebatrophia 49
phoenicia, Nomada ruficornis var. 315
Phoenusa 48
Phor 311
phryne, Chrysis phryne 137
pthiscus, Cephus 109
pthisica, Selandria 82
phylicoliae, Pontania 56
Phyllocolpa 55
phyllocolpa, Blennocampa 45
Phylloecus 110
Phyllopais 75
Phyllotoma 49
Phymatocera 52
Phymatoceriola 52
Phymatoceropsis 52
Phymatoceros 52
Physetopoda 155
Physoscelis 235
Physoscelus 235
piccioliana, Nomada 315
picciolii, Andrena 267
picea, Apis 305
picea, Formica 201
picea, Formica auct. 200
picea, Melitta 288
picea, Tenthredo 86
piceae, Lophyrus 34
piceae, Nematus 72
piceae, Pleroneura 20
piceanus, Symmorphus 191
picicornis, Andrena 267
piceus, Camponotus part, auct. 198
picicornis, Megachile 306
picicornis, Melitta 267
picicornis, Mimesa 251

picicornis, Sphecius 214
picicornis, Anthophora 322
picinus, Dolerus 79
picipes, Dolerus, *Tenthredo* 80
picipes, Melitta 274
picipes, Methocha, *Methoca* 149
picistigma, Colletes 258
picta, Apis 316
picta, Rhogogaster, *Tenthredo* 88
picta, Tenthredo 93
picta, Tiphia 149
picta, Xiphydria 116
picticeps, *Tenthredopsis* 100
picticollis, Bembix 222
picticollis, Nematus 64
picticornis, Dryudella, *Astatus* 218
picticornis, Stizus 225
picticornis, Tenthredo 98
picticrus, Stenodynerus, *Lionotus* 190
pictifrons, Pamphilus 105
pictior, Polistes 193
pictipes, Ancistrocerus 179
pictipes, Andrena 268
pictipes, Crabro 232
pictipes, Hylaeus 261
pictipes, Oxybelus 243
pictipes, Tenthredo 88
pictus, Crabro 232
pictus, Crabro 245
pictus, Cryptocampus 62
pictus, Dinetus, *Crabro* 245
pictus, Epeolus 317
pictus, Odynerus 178, 189
pictus, Passaloecus 248
pieli, Gorytes 223
pieli, Megachile lagopoda 307
pieli, Nematus 66
pielianus, Coelioxys 305
pigeoti, Tenthredo rossii 90
Pikonema 55
pileata, Andrena allosa ssp. 267
pilichi, Rophites 291
pilicornis, Cladius 54
pilicornis, Crossocerus 228
pilicornis, Hylotoma 23
pilicornis, Osmia 299
pilicornis, Sphecodes 288
pilicornis, Tenthredo 25
pilicruriformis, Megachile 307
pilicrus, Megachile 307
pilidens, Megachile 306
pilifrons, Sphecodes 289
pilipennis, Bethylus 150
pilipes, Andrena 272
pilipes, Apis 323
piloserra, Amauronematus 65
piloserra, Euura, *Nematus* 63
piloserra, Xyela 21
piliventris, Nomada 315
pilichi, Andrena 270
pilichi, Eucera 320
pilichi, Priocnemis 164
pilosa, Abia 28

pilosa, Anthophora 325
pilosa, Nomada 315
pilosella, Anthophora 324
pilosella, Vespa 195
pilosellus, Pomphilus pectinipes var. 171
pilosodorsata, Andrena coitana var. 271
pilosula, Melitta 267
pilosulus, Cephus 109
pilosus, Cenomus 249
pilosus, Colletes 257
pilosus, Epeolus 317
pilosus, Nematius willigkiae 69
Pilpomus 165
pineti, Lophyrus 34
pineti, Nematus 62
pineti, Tenthredo 33
pingshiangensis, Anthophora 323
pinguiculus, Sphecodes 289
pinguis, Dryudella, *Larra* 218
pinguis, Larra 218
pinguis, Phyllotoma 49
pini, Diprion, *Tenthredo* 33
pini, Tenthredo 33
pini, Tenthredo 70
Pinicola 20
Pinicolites 20
pinirufa, Tenthredo 34
pinivora, Acantholyda posticalis 102
pionganensis, Vespa japonica 196
piapiens, Anthophora 323
pirica, Acantholyda 102
Pironistilomorphus 109
pisarskii, Formica 199
pisarskii, Myrmica 206
pisinna, Tenthredopsis 100
Pison 244
Pisonitus 244
Pisonoides 244
pisonoides, Solierella, *Niteliopsis* 241
Pisum 244
Pitaliris 237
pithyusae, Leptothorax 208
placida, Andrena 268
placida, Chrysis 137
plaga, Tenthredopsis lactiflua 99
plagiata, Anthophora, *Megilla* 324
plagiata, Hoplocampa, *Tenthredo* 68
plagiata, Pteronidea ferruginea 59
plagiatus, Loderus ornatus 79
plagiocephalus, Megalodontes, *Tarpa* 107
Plagioceps 173
Plagiolepis 202
plana, Abia 28
plana, Empria, *Poecilosoma* 41
plana, Monardis, *Tenthredo* 50
plana, Myrmica lobicornis var. 205
plana, Tenthredo 50
planatus, Dolerus 79
planiceps, Gonatopus 121
planiceps, Selandria 76
planidens, Systropha 291

planifrons, *Crabro* 231
planifrons, *Pamphilius* 105
planiventris, *Discoelius* 181
plantaris, *Andrena* 274
plantaris, *Hylaeus* 259
Plastandrena subg. 272
Plastanoxus 124
Plastodryinus 120
Platycampus 69
platycera, *Osmia* 297
platyceros, *Nematus* 73
platyceros, *Tenthredo* 84
platycerus, *Lagidina*, *Tenthredo*
platycerus 84
platycerus, *Nematus* 75
platycerus, *Stauronematus*, *Nematus* 75
platycestus, *Halictus* auct. 280
Platyderes 166
Platygona 120
Platyhalictus subg. 278
platyodonta, *Osmia* 296
platyrhina, *Eucera* 320
platystigma, *Euura*, *Pteronidea* 63
platythorax, *Lasius* 202
Platyxiphidria 115
plebeja, *Tenthredo* 93
Plectrogonatopoides 120
Plectrogonatopus 120
Plenoculus 241
Pleroneura 20
plesius, *Priophorus* 54
plesius, *Stenocrabro* 229
pleuralis, *Athalia rufoscutellata* 45
pleuralis, *Halictus* 286
pleuralis, *Macrophypha rustica* 86
pleuripunctatus, *Gorytes*, *Hoplitus*
pleuripunctatus 223
pleuris, *Hoplocampa rutilicornis* 68
pleuritica, *Arge*, *Hylotoma* 24
pleuritica, *Blennocampa affinis* 48
pleuritica, *Platycampus luridiventris* 70
pleuritica, *Tenthredella cunyi* 94
pleuritica, *Tenthredopsis sordida* 100
Pleuriticiformis 22
pleuriticus, *Monophadnus* 51
pleuronotata, *Tenthredo semenovi* 95
pleurosternalis, *Tenthredopsis*
nassata 99
pleurostictus, *Nematus* 62
plicatus, *Hedychrum gerstaeckeri* 130
pliginskii, *Elampus*, *Notozus* 128
pliginskiji, *Skorikovia*,
Smicromyrme 155
Plistotrichia 319
plumbea, *Sphex* 171
plumicornis, *Hylaeus* 259
plumipes, *Andrena* 293
plumipes, *Anthophora*, *Apis* 323
plumipes, *Sphex* 216
plumosa, *Apis* 264
plurifasciata, *Prosopis* 260
plurinotata, *Hoplomerus*
armeniacus var. 187

plurinotata, *Melecta* 326
Pluschtschevskyi, *Crabro* 234
pluschtschevskyi, *Lestica*, *Crabro* 234
pluschtschewskii, *Stizus* 225
pluto, *Lasioglossum matianense* 284
Plutobethylus 123
podagra, *Anthophora* 324
podagrifica, *Pemphredon* 249
podagricus, *Crossocerus*, *Crabro* 227
Podalirius 323
Podalonia 213
Podasys 293
podkumokensis, *Rhogogaster*
chlorosoma 88
podalica, *Andrena* 274
podolicum, *Lasioglossum*, *Halictus*
podolicum 285
poecila, *Tenthredo* 95
Poecilagenia 165
Poecilageniella 165
Poecilandrena subg. 272
poecilechroa, *Tenthredo* 91
poecilonota, *Euura*, *Nematus* 63
poecilopus, *Tenthredo* 86
poecilopus, *Tenthredo* 95
Poecilosoma 40
Poecilostoma 40
Poecilostomidea 40
Poecilotiphia 149
poecilus, *Odynerus* 187
poggii, *Odynerus dantici* 183
Pogonius 164
polaris, *Bombus* auct. 327
polaris, *Euura*, *Nematus* 63
polaris, *Pontania* 59
polaris, *Rhogogaster* 88
Polemistus 249
polevoiae, *Hylaeus* 260
Policrabro 231
Polistarchus 192
Polistella subg. 192
Polistes 192
Polistes subg. 192
Polistinae 192
Polistoides 192
Polistula 192
polita, *Andrena* 273
polita, *Cerceris* 253
polita, *Euura*, *Nematus* 63
polita, *Megachile* 308
polita, *Myzine* 149
polita, *Nomia* 291
polita, *Psammophila* 213
polita, *Tiphia* 150
politum, *Lasioglossum*, *Hylaeus*
politum 285
politus, *Lophyrus* 34
pollinosa, *Eucera* 320
pollinosa, *Tetraloniella*, *Macrocera* 322
pollux, *Aporus*, *Pompilus* 166
Polochoridium 145
Polochromum 145
polonica, *Ancistrocerus nigricornis* 179
polonica, *Eucera* 320
Polybates 50
Polybiapis 311
polycentris, *Coelioxys* 304
polyctena, *Formica* 200
Polyergus 202
polygona, *Tenthredo* 109
polygoni, *Ametastegia* 40
Polyinematus 55
Polyodontus 141
Polyplanus 122
polypori, *Holopedia* 124
Polysealandria 75
polyspilus, *Pontania* 67
Polystichophagus 82
polytoma, *Gilpinia*, *Lophyrus* 34
polytomus, *Lophyrus* 33
pomiformis, *Eumenes*, *Vespa* 182
pomorum, *Bombus*, *Bremus* 331
Pompilidae 160
pompiliformis, *Larra* 237
pompiliformis, *Tachysphex*, *Larra* 239
Pompilinae 160, 166
Pompilini 167
Pompilinus 172
Pompiliodes 172
Pompilioides 172
Pompiloides 173
Pompilogastra 173
Pompiloidea 160
Pompilus 171
Ponera 209
Ponerinae 209
ponojense, *Nematus fastosus* 65
ponomarevae, *Anthophora* 325
Pontania 55
pontanioides, *Pteronidea* 68
Pontanthidium 300
pontica, *Astata* 218
pontica, *Lasius alienus* var. 202
pontica, *Solenopsis fugax* var. 207
ponticus, *Ancistrocerus claripennis* 179
ponticus, *Halictus* 278
ponticus, *Odynerus germanicus* 190
pontomoravicus, *Evagetes*,
Psammochares 171
Pontopristia 55
Poodolerus 77
Poodolerus subg. 79
popilliavora, *Tiphia* 150
popofiana, *Pontania* 64
popovi, *Biastes* 318
popovi, *Chrysis* 137
popovi, *Gonatopus* 121
popovi, *Xiphydria* 116
popovici, *Coelioxys* 305
popovii, *Parnopes* 142
Poppia 81
poppii, *Amauronematus* 57
poppii, *Pontania* 63
populi, *Cladius*, *Trichiocampus* 54
populi, *Nematus* 56
porcus, *Halictus* 283, 285

- porosicus*, Colletes 258
porphyrea, Chrysura, *Chrysis oraniensis* 139
portchinski, Crocisa 326
portchinskii, Bembix 222
portchinsky, Bombus 328
portschinskii, Physetopoda, *Mutilla* 155
porzana, Andrena tibialis ssp. 272
posthuma, Nomada 315
posthumus, Halictus 278
postica, Andrena 267
postica, Macrophyia, *Tenthredo* 86
posticus, Euodynerus, *Odynerus* 183
posticus, Nematus 55
postscutellatus, *Odynerus* 183
postumus, Nematus 71
postzonatus, Diversobombus ussuriensis 329
potanini, Arge 24
potanini, Euxiphidria, *Xiphydria* 115
potanini, Leptocimbex 30
potanini, Macrophyia 86
potanini, Scolia 148
potanini, Xiphydria 115
potentillae, Andrena 273
potulenta, Strongylogaststroidea 39
poupillieri, Andrena 274
poutziloi, Cerceris 254
pouzdranensis, Smicromyrme 156
praecociformis, Andrena 267
praecox, Andrena, *Apis* 264
praecox, Epitactus 57
praemarinus, Bombus 328
praestans, Hoplitis, *Osmia* 297
praestans, Melitturga 276
praeteritorum, Pseudocephaleia, Dictyolyda 106
praetexta, Andrena 272
praeustus, Epeolus 317
prahenda, Cerceris 253
Praia 30
prasina, Notozus spina ab. 128
prasinipes, Macrophyia 85
prasinus, Nematus 67
pratensis, Apis 269
pratensis, Dolerus, *Tenthredo* 78
pratensis, Formica 200
pratensis, Mellinus 246
pratensis, Tenthredo 102
pratensis, Vespa 193, 262
Pratensisformis 77
pratorum, Bombus, *Apis* 330
pratorum, Dolerus, *Tenthredo* 79
pratostepposus, Leptothorax 208
pravei, Arge 23
pravei, Cephus 109
pravei, Empria 41
pravus, Euura, *Amauronematus* 63
preaetermissus, Liris, *Notogonia festinans* 237
Prenanteon 119
Prenolepis 202
pressilabris, Formica 199
pretiosa, Andrena 268
pretiosa, Nomada fucata var. 313
pribaicalica, Euura, *Amauronematus* 63
priesneri, Discoelius 181
Priesnerius 160
primorensis, Cleptes 127
princeps, Hoplitis, *Osmia* 297
princeps, Nematus 69
Priocnemini 162
Priocnemiocheilus 161
Priocnemis 162
Priocnemis subg. 162
Priocnemissus 162
Priocnemus 162
Priocnemis 162
Priocnemocnemis 162
Priocnemocnemus 162
Priononyx 215
Prionophorus 54
Prionychini 214
Prionyx 215
Priophorus 53
Priophorus subg. 54
prisca, Cerceris 255
prisma, Scolia 145
prismatica, Megacampsomeris, *Scolia* 147
Pristicampus 70
Pristiphora 70
pristiphorinus, *Amauronematus* 59
pristiphoroides, Euura, *Amauronematus* 63
Pristis 68
Pristocera 123
Pristocerinae 123
Pristogonatopus 120
Pristola 70
Proanthidium subg. 301
proauriceps, Chrysis 137
problematica, Andrena 275
problematicum, Lasioglossum, *Hylaeus* 285
problematicus, Sphecodes 288
procera, Scolia 147
procera, Tenthredo 95
Proceratiinae 210
Proceratum 210
procerus, Allogonatopus 120
procerus, Psen 252
procerus, Synnevrus 220
Prochridium 132
Procimber 28
procrustes, Gorytes 223
prodichroa, Chrysura 143
productulus, Epeolus 317
productus, Epeolus 317
Profenus 52
Proformica 203
proformicarum, Rossomyrmex 203
Progoniozus 125
projectus, Tachysphex 239
proletarius, Crabro 232
prolongata, Xiphydria, *Tenthredo* 116
Pronaeus 211
propheta, Tremex 113
propinquia, Andrena 273
propinquia, Tenthredo 97
propinquia, Thyreomelecta, *Thyreus* 320
propinquus, Crabro 229
propinquus, Hylaeus 260
propinquus, Nematus 57
Propodea 89
Propodea subg. 92
propodeale, Hedyphridium 129
Prorhynchium 187
Prosalius 166
Prosanteon 118
Prosceliphron 214
Proscleroderma 151
Prosecris 40
proschalykini, Hedyphridium 129
Prosopigastra 237
Prosopis 259
Prosopis subg. 261
prospera, Tenthredo 94
Protelostegus 174
Protomphytus 39
Protomphytus subg. 39
protensus, Nematus 70
proteus, Bombus 328
proteus, Nomada 312
Protoctenius 231
Protodynerus 190
Protohalictus subg. 278
Protosmia 299
Protosmia subg. 300
Protostelis subg. 302
Prototaxonus 82
Protothyreopus 231
prototypus, Nematus 63
provancheri, Pemphredon 249
provecta, Megachile willughbiella 308
provida, Tenthredo, *Allantus* 98
providens, Atta 207
providens, Tenthredo 98
providentia, Tenthredo 98
provinciale, Anthidium 301
provincialis, Megachile 306
proviridis, Holopyga ovata 131
Proxiandrena subg. 273
proxima, Andrena 267
proxima, Andrena, *Melitta* 273
proxima, Athalia, *Tenthredo* 44
proxima, Eucera 320
proxima, Euura, *Nematus* 63
proxima, Hylotoma 24
proxima, Osmia 296
proximatum, Lasioglossum, *Halictus* 281
proximum, Chelostoma 296
proximus, Crabro 229
proximus, Evagetes part. 171
proximus, Evagetes, *Pomphilus* 171
proximus, Gorytes 223
proximus, Nematus 63
proximus, Odynerus 190

Proxylocopa subg. 310
prshewalskii, Anthophora 325
pruinosus, *Sphex* 216
pruni, Nematus 71
pruni, Pareophora, *Tenthredo* 51
pruni, Priophorus 54
pruni, *Trichiocampus* 55
prunicola, *Hoplocampa* 68
przewalskyi, *Argogorytes* 221
Psacas 130
Psammaecius 224
psammobiam, *Tachytes* 239
psammobius, *Oxybelus* 243
psammobius, *Tachysphex*, *Tachytes* 239
Psammochares 172
Psammocharoides 170
Psammoderini 173
Psammophila 213
psammophilus, *Lasius* 202
Psammophex 237
pschornwalcheri, Nematus 61
Psen 251
Pseneo 251
Psenia 251
Psenini 250
Psenulus 251
Psenus 251
pseudabnormis, *Pompilus* 169
Pseudagenia 165
Pseudapis 292
Pseudarge 25
Pseudbedychrum 130
Pseudepipona 188
Pseudepipona subg. 188
Pseudevagetes 167
Pseudisobrachium 123
Pseudoanthidium 301
Pseudoanthidium subg. 302
pseudoanticus, *Dolerus* 79
pseudoarmata, *Melecta* 326
pseudobaicalensis, *Bombus* 331
Pseudobrennoccampa 46, 47
pseudobrevitarsis, Chrysis 137
pseudobrevitarsis, Chrysis ignita var. 137
pseudocalidus, *Bombus cingulatus* 330
Pseudocalyoza 123
Pseudoccephaleia 106
Pseudocephus 109
pseudocerea, *Nomioides variegata* 292
Pseudochrysis 140
Pseudocilissa 294
pseudocinerascens, *Colletes* 258
Pseudoclavellaria 30
pseudocoenfluens, *Halictus* 278
Pseudocrabro 231
pseudodecipliens, *Pristiphora* 73
pseudodichroa, *Chrysura*, *Chrysis* 139
Pseudodineura 75
pseudodispar, *Euura*, *Pteronidea* 63
Pseudoeucera subg. 320
pseudofasciatus, *Sphecodes* 289
pseudofemoralis, *Dryudella* 218

Pseudoferreola 166
pseudofultus, *Katamenes sichelii* 185
pseudogeniculata, *Pristiphora* 73
pseudogermanica, *Vespa vulgaris* var. 196
Pseudogonatopoides 120
Pseudogonatopus 120
Pseudohemitaxonius 81
Pseudoheptamelus 36
pseudoignita, *Chrysis* 143
pseudojaponica, *Priocnemis* 164
pseudoklugi, *Empria* 41
pseudoleptocephala, *Euura*, *Amauronematus* 63
pseudoligusticus, *Hortobombus tichenkoi* 329
pseudolivacea, *Tenthredo* 98
Pseudomacrophya 84
Pseudomacrophya subg. 87
Pseudomalus 132
pseudomandibularis, *Tenthredella atra* 93
Pseudomegachile subg. 307
pseudomelanocarpa, *Pristiphora* 73
Pseudomicrodynerus subg. 186
Pseudomonophadnus 50
pseudomontana, *Smicromyrme punctata* 155
pseudomorbillosum, *Lasioglossum* 280
pseudomucoreus, *Halictus* 279
Pseudomutilla subg. 154
Pseudomyrmosa 152
pseudonotabilis, *Pteronidea* 57
pseudonotata, *Euodynerus quadrifasciatus* var. 183
Pseudo-Nysson 244
pseudoorientalis, *Smicromyrme* 156
pseudopacificus, *Polistes gallicus* 193
pseudopalidum, *Diprion polytomum* 34
pseudopalmarius, *Crossocerus*, *Crabro* 230
pseudopavida, *Pteronidea myosotidis* 62
pseudopilosulus, *Cephus* 109
Pseudoplisis 222
pseudopogonia, *Priocnemis* 163
Pseudopolistes 192
pseudopolita, *Andrena* 273
Pseudopterochilus 184
pseudopunctulatus, *Halictus* 285
pseudopyrrhina, *Chrysis cingulicornis* 134
pseudorejecta, *Arge* 24
pseudoreticulata, *Cephalcia* 106
pseudorossii, *Tenthredo* 90
Pseudoryssus 117
pseudosaxesenii, *Pristiphora* 74
Pseudoscolia 256
Pseudoscoliini 256
pseudoscotica, *Tenthredella atra* 93
Pseudoselandria 76
pseudosoror, *Vespa tropica* 194
Pseudospinolia 140
Pseudostelis subg. 303
pseudotabidus, *Cephus* 108
Pseudotaxonus 82
pseudothoracica, *Andrena* 275
Pseudotiphia 150
pseudotorneensis, *Euura*, *Amauronematus* 63
pseudounifasciata, *Scolia* 148
pseudovatula, *Andrena* 274
Pseudovespa 194, 195
Pseudovespula 194
pseudoviminalis, *Trichiocampus* 54
Pseudoxiphydria 115
Psilanteon 119
Psilepyris 123
Psiliglossa 188
psilopus, *Tachysphex* 239
Psithyrus 327
Psithyrus subg. 329
psylluis, *Sirex* 113
Pteneucera subg. 321
Pterocheilus 189
Pterogonatopus 120
Pteronidea 55
Pteronidea subg. 75
Pteronus 33
pterophorus, *Tenthredo* 35
pterotus, *Crabro* 226
Ptigospex 241
Ptyx 234
pubens, *Crabro* 227
puber, *Andrena* 275
pubescens, *Ammophila* 212
pubescens, *Andrena* 269
pubescens, *Anthophora*, *Apis* 323
pubescens, *Crabro* 228
pubescens, *Melitta* 265
pubescens, *Odynerus* 183
pubescens, *Pachynematus* 57
pubescens, *Periclista*, *Selandria* 51
pubescens, *Tenthredo*, *Allantus* 91
pubescens, *Trachusa*, *Anthidium* 303
pubescens, *Trichiosoma* 31
pubicorne, *Anteon*, *Gonatopus* 119
pucilii, *Cerceris* 254
puella, Nematus 62
puella, *Phylloboma* 76
pugnax, *Oxybelus* 243
pugnax, *Pamphilus* 105
pulawskii, *Cerceris* 253, 254
pulawskii, *Diploplectron* 218
pulawskii, *Holopyga* 142
pulawskii, *Spilomena* 249
pulchella, *Andrena* 292
pulchella, *Chrysis pulchella* 137
pulchella, *Coelioxys* 304
pulchella, *Ermilia* 43
pulchella, *Selandria* 43
pulchellus, *Hoplisus* 224
pulchellus, Nematus 60
pulchellus, *Oxybelus* 243
pulcher, *Cephus* 109
pulcher, *Pamphilus montanus* 105
pulcher, *Pompilus* 171

pulcher, Pompilus 171
pulcherrima, Andrena 267
pulcherrima, Anthophora 324
pulcherrima, Larra 225
pulchra, Megaxyela 20
pulchra, Nomada 315
pulchra, Vespa ducalis var. 194
pulchricornis, Lophyrus 34
pulla, Cimbex betulae 29
pulla, Paramyrmosa, *Myrmosa* 153
pullata, Arge, *Hylotoma* 24
pullata, Tenthredo discophora 95
pullatum, Hedydium simile f. 130
pullior, Dineura 55
pullulus, Crossocerus, *Crabro* 229
pullulus, Dolerus 78
pullus, Nematus 73
pullus, Stenodynerus 190
pulsata, Osmia 299
pulveracea, Eucera 321
pulveratum, Monsoma, *Tenthredo* 43
pulvereus, Halictus 279
pulverosus, Nomiooides 292
pulverosus, Tachytes 239
pulverulenta, Tenthredo 43
pumila, Chrysidea, *Chrysis* 133
pumila, Chrysis 133
pumila, Empria, *Poecilosoma* 41
pumila, Fenusa 48
pumila, Tenthredo 48
pumilae, Xyela 21
pumilio, Emphytus 50
pumilio, Euura, *Pachynematus* 63
pumiloides, Empria 41
pumilus, Cephus 110
pumilus, Crabro 232
pumilus, Metallus, *Tenthredo* 50
pumilus, Microphadnus, *Aporus* 173
pumilus, Nematus 71
punctata, Andrena aeneiventris var. 263
punctata, Apis 325
punctata, Calameuta, *Astatus* 109
punctata, Coelioxys 305
punctata, Crabro podagricus 227
punctata, Eumenes
 pedunculata var. 181
punctata, Nomada 243
punctata, Sapyga 145
punctata, Tenthredo 103
punctaticeps, Eumenes 182
punctatissima, Andrena 264
punctatissima, Holopyga
 punctatissima 131
punctatissima, Hypoponera, *Ponera* 209
punctatissima, Osmia 296
punctatissima, Prosopigastra 237
punctatissima, Pseudepipona
 postica 183
punctatissimum, Lasiosoma, *Hylaeus*
 punctatissimum 285
punctatissimus, Priocnemis, *Salius* 163
punctatissimus, Sphecodes 289
punctatum, Anthidium 301

punctatus, *Hoplisus* 224
punctatus, Ammobates, *Epeolus* 318
punctatus, Crabro 243
punctatus, Eumenes 182
punctatus, Eutomostethus,
 Tomostethus 47
punctatus, Hylaeus, *Prosopis* 262
punctatus, Pasites 318
punctatus, Pseudomalus, *Philoctetes* 133
punctatus, Selandria serva 81
puncticeps, Claremontia,
 Blennocampa 46
puncticeps, Nematus 70
puncticeps, Psen 252
puncticeps, Psenulus 252
puncticeps, Sphecodes 289
puncticolle, Lasiosoma, *Halictus* 285
puncticollis, Dolerus 80
puncticollis, Halictus 285
puncticollis, Polistes 192
puncticollis, Tenthredopsis 100
punctifemoratus, Armitarsus 84
punctifer, Pemphredon 249
punctifrons, Arge 25
punctifrons, Euura, *Pontania* 63
punctifrons, Macrophyia vacillans 87
punctifrons, Pristiphora, *Nematus* 73
punctifrons, Prosopis 259
punctifrons, Stenodynerus, *Lionotus* 190
punctigera, Tenthredo 68
punctilabris, Eucera 320
punctipleura, Notogonia 237
punctipleuris, Mimesa, *Psen* 250
punctipleuris, Nematus 66
Punctipsen 251
punctiscuta, Nematus fastosus 65
punctiventris, Sphecodes 287
Punctitiphia subg. 150
punctulata, Andrena 270
punctulata, Andrena
 hattorfiana var. 265
punctulata, Melitta 287
punctulata, Rhogogaster, *Tenthredo* 88
punctulata, Tenthredo 94
punctulatissima, Stelis, *Apis* 303
punctulatissimus, Hylaeus, *Prosopis* 260
punctulatus, Cephus 109
punctulatus, Ectemnius 233
punctulatus, Gorytes 224
punctulatus, Nematus 66
punctulatus, Psammaecius, *Gorytes* 224
punctulatus, Sphecodes 289
punctulifera, Nomada
 solidaginis var. 316
punctum, Crabro 229
punctum, Sapyga 145
punctum, Tenthredo 87
punctumalbum, Macrophyia,
 Tenthredo 87
punctuosus, Hoplisoides, *Hoplisus*
 punctuosus 224
punctuosus, Odynerus 190
punetiscuta, Nomada 316
punicea, Euura, *Tenthredo* 63
puniceus, Agenioideus,
 Psammochares 168
punjaubensis, Andrena 275
pura, Mutilla elongata var. 155
purpurascens, Hedydium,
 Hedydium 129
purpurata, Chrysis 140
purpuratus, Euchroeus, *Chrysis*
 purpuratus 140
purpurea, Osmia 298
purpurea, Tenthredo 95
purpureae, Euura, *Nematus* 64
purpureifrons, Chrysura, *Chrysis* 139
purus, Nematus 61
pushkiniana, Chrysis 137
pusilla, Chrysis chevrieri var. 135
pusilla, Nomada auct. 311
pusilla, Pontania 63
pusilla, Priocnemis, *Pompilus* 163
pusilla, Tenthredo 45
pusilla, Xyela 20
pusilla, Xyela 21
pusillum, Trichiosoma 31
pusillus, Cephus 109
pusillus, Crossocerus 230
pusillus, Dolerus 78
pusillus, Hylaeus 285
pusillus, Labeo 120
pusillus, Phiarus maculatus var. 319
pusillus, Pseudomalus, *Chrysis*
 pusillus 133
pusio, Cyrtachridium 129
pustulata, Mutilla 157
pustulator, Euura, *Pontania* 64
pustulatus, Amauronematus 61
putoni, Cleptes 127
putoniana, Dufourea 290
Pycnopompilus 168
Pycnotilla 157
pygialis, Auplopus, *Pseudagenia* 165
pygialis, *Omalus aeneus* var. 131
pygidiale, Rhopalum 235
pygidialis, Nomada 315
pygidialis, Stizus 225
pygidialis, Tachysphex 238
pygmaea, Ceropales pygmaea 174
pygmaea, Euura 64
pygmaea, Formica 202
pygmaea, Nomada 312
pygmaea, Pristiphora 70
pygmaea, Profenus, *Tenthredo* 52
pygmaea, Prosopis 259
pygmaea, Stelis 302
pygmaeum, Trypoxylon 244
pygmaeus, Diodontus 247
pygmaeus, Hylaeus 286
pygmaeus, Lindenius, *Crabro* 235
pygmaeus, Oxybelus 243
pygmaeus, Psen 252
pygmaeus, Sirex 109
pygmeus, Cephus, *Sirex* 109

pylnovi, Philoctetes, *Ellampus* 132
pyonganensis, Anteon 119
Pyramica 207
pyramidalis, Nematus 75
pyramidalis, Podalirhis 323
pyrenaica, Osmia 299
pyrenaica, Dioxyx 303
pyrenaica, Diphysis 303
pyrenaica, Hylotoma 24
pyrenaica, Plagiolepis barbara var. 202
pyrenaeicus, Mesitius 124
pyri, Psen 103
pyri, Tenthredo 67
pyrina, Megachile 307
Pyrobombus subg. 330
pyropygia, Andrena 269
Pyrosmia subg. 299
pyrosomus, Elampus, *Notozus* 128
pyrrha, Chrysis 134
pyrrha, Phylloecus, *Hartigia* 111
pyrrhina, Chrysis 138
Pyrrhomelecta 317
pyrrhonotus, Nematus 61
pyrrhopygus, Bombus 327

quadrannulata, Tenthredopsis 98
quadrata, Anthophora albigena 323
quadrata, Vespa 179
quadraticollis, Halictus 281
quadratus, Sphecodes monilicornis 289
quadriceps, Stigmus 250
quadricincta, Andrena 275, 294
quadricincta, Apis 277
quadricincta, Cerceris, *Philanthus* 254
quadricincta, Scolia, *Sphex*
 quadricincta 148
quadricincta, Tenthredo 78
quadricincta, Vespa 183
quadricinctula, Andrena 275
quadricinctus, Crabro 231
quadricinctus, Halictus, *Apis* 277
quadricolor, Bombus, *Psithyrus* 329
quadricolor, Cerceris 254
quadridentata, Tenthredo 95
quadridentata, Osmia 297
quadridentatus, Coelioxys, *Apis* 305
quadrifasciata, Amegilla, *Apis* 322
quadrifasciata, Apis 322
quadrifasciata, Cerceris, *Philanthus* 254
quadrifasciatus, Euodynerus, *Vespa* 183
quadrifasciatus, Gorytes, *Mellinus* 223
quadrifasciatus, Halictus 278
quadrifer, Crabro 234
quadriguttata, Scolia 145
quadriguttatus, Cephus 108
quadriguttatus, Hellus 145
quadriguttatus, Nysson 220
quadriguttulatus, Cephus 108
quadrilobum, Anthidium 301
quadrimaculata, Anthophora, *Apis* 323
quadrimaculata, Cerceris 253
quadrimaculata, Empria 41
quadrimaculata, Meria 149

quadrimaculata, Mutilla 153
quadrimaculata, Mutilla maura 154
quadrimaculata, Prosopis 260
quadrimaculata, Tenthredo 86
quadrinaculatus, Brachyodynerus,
Odynerus 180
quadrinaculatus, Crossocerus,
Crabro 230
quadrinaculatus, Hylaeus 283
quadrinotata, Apis 312
quadrinotata, Mutilla 157
quadrinotatum, Lasioglossum,
Halictus 285
quadrinotatum, Lasioglossum,
Melitta 281
quadrinotatus, Camponotus 198
quadrinotatus, Camponotus
 marginatus var. 198
quadripunctata, Andrena 265
quadripunctata, Cerceris 253
quadripunctata, Hylotoma 24
quadripunctata, Hypoclinea auct. 197
quadripunctata, Mutilla maura var. 154
quadripunctata, Scolia 147
quadripunctata, Scolia 148
quadripunctata, Tenthredopsis
 fenestrata 99
quadripunctata, Vespa 194
quadripunctatus, Dolichoderus,
Formica 197
quadriseriatum, Anthidium 300
quadrisignatum, Lasioglossum,
Hylaeus 286
quadrisignatus, Cephus 108
quadristrigata, Andrena 274
quadristrigata, Anthophora 325
quadristrigatus, Halictus 277
quadrivittata, Nomada lineola var. 313
quadrum, Cryptocampus 63
quartinae, Trypoxylon 245
quasiciliatus, Dolerus 79
quattuordecimpunctata, Melecta 326
quattuordecimpunctatus, Philanthinus,
Anthophilus 255
quatuordecimnotatus, Oxybelus 243
quatuormaculata, Sphex scutellata 226
Quebecensis, Larra 239
querqus, Fenus 48
querqus, Nematus 71
quetteae, Astata 218
quetteaensis, Eumenes affinissima
 race 182
quiescens, Palarus 243
quieta, Euura, *Nematus* 64
quinghaiensis, Melitta 294
quinquecincta, Colpa, *Scolia* 147
quinquecinctum, Rhynchium, *Vespa* 189
quinquecinctus, Allantus 97
quinquecinctus, Gorytes, *Mellinus*
 quinquecinctus 223
quinquecinctus, Megalodontes,
Tarpa 107
quinquecinctus, Mellinus 222

quinquefasciata, Cerceris, *Crabro*
 quinquefasciata 254
quinquefasciata, Vespa 187
quinquefasciatus, Cephus 110
quinquefasciatus, Gorytes, *Mellinus*
 quinquefasciatus 223
quinquemaculata, Dasylabris maura
 nostras var. 154
quinquemaculata, Mutilla 157
quinquemaculatus, Philanthus 246
quinquemontana, Euura,
Pachynematus 64
quinquepunctata, Mutilla 157
quinquepunctata, Sapyga, *Scolia* 145
quinquepunctata, Scolia 145
quinquespinosa, Nomada 311
quinquespinosus, Rophites 291
quittardi, Tenthredo maculata 92

raddatzi, Dolerus 80
raddatzi, Thomsonia 99
raddei, Ancistrocerus, *Odynerus* 179
raddei, Anthophora 325
radialis, Emphytus 46
radians, Chrysura, *Chrysis* 140
radiatus, Ectemnius, *Crabro* 231
radoszkovskii, Halictus 279
radoszkovskii, Katamenes
 sesquicinctus 185
radoszkovskii, Skorikovia,
Smicromyrme 155
radoszkowskyi, Andrena 272
radoszkowskii, Ammobatoides 318
radoszkowskii, Cleptes 127
radoszkowskii, Nomada 315
radoszkowskii, Tenthredo, *Allantus* 91
radoszkowskii, Tetralonia 321
radoszkowskyi, Anthidium 300
radoszkowskyi, Anthophora 325
radoszkowskyi, Gorytes 222
radoszkowskyi, Myrmosa 153
ragusea, Chrysis 137
rakosiensis, Nomada 313
ramicornis, Cladius 54
ramosus, Thyreus, *Crocisa* 327
ramulorum, Xylocopa 310
ranini, Euura, *Amauronematus* 64
ranunculi, Tenthredo 32
ranunculorum, Andrena 275
rapae, Pachyprotasis, *Tenthredo* 87
rapae, Tenthredo 87
Raptiformica subg. 200
raptor, Oxybelus 243
rapunculi, Chelostoma, *Heriades* 296
rarior, Bombus lapponicus 330
rasnitsyni, Arachnospila 169
rasnitsyni, Xyela 21
ratisbonensis, Andrena 274
ratzeburgi, Alysson 219
ratzeburgii, Macrophyia 86
ratzeburgii, Tenthredo 105
ravasinii, Myrmica 206
ravidus, Pachynematus 72

ravouxi, Hoplitis, *Osmia* 297
 ravouxi, Myrmoxenus,
 Formicoxenus 207
 ravulus, Colletes 258
ravus, Dolerus 79
reaumuri, Odynerus 187
 recedens, Myrmica 208
 recedens, Temnothorax, *Myrmica* 208
recepticus, Halictus 281
 recognata, Macrophyia 86
 recta, Hinatara, *Blennocampa* 49
rectangula, Andrena 264
rectangulata, Glossotilla 155
rectangulatus, Homonotus 166
 rectus, Auplopus, *Pseudagenia* 165
recurva, Coelioxys 305
redii, Nematus 63
redimita, Tarpa 107
redivivus, Tachysphex 240
reducta, Runaria 27
reductalba, Tenthredella mesomelas 91
reductenotata, Macrophyia rufipes 86
reductenotata, Tenthredo 85
reflexa, Coelioxys 305
reflexus, Pompilus 172
refractorius, Pachynematus 58
refulgens, Chrysura, *Chrysis* 140
regalis, Dasylabris, *Mutilla* 154
regalis, Nomada 311
regina, Andrena 272
Regiscolia sung. 147
regium, Trypoxylon regium 245
regularis, Anthidium 300
regulus, Leptochilus, *Odynerus* 185
 rehbinderi, Strongylognathus 207
reicherti, Dolerus sanguinicollis 80
reiecta, Prosopis communis var. 259
Reiseri, Tachysphex 238
Reiteri, Crabro 234
reitteri, Lestica, *Crabro* 234
reitteri, Allantus 96
reitteri, Rhadinoceraea 52
rejecta, Tenthredo 93
relicta, Pontania collactanea 58
relicta, Prosopis variegata 262
relicta, Tenthredo 93
relictana, Euura, *Pontania* 64
relictum, Trichiosoma 31
reliqua, Rhogogastera 92
remota, Anthocopa tergestensis 299
remota, Chrysura remota 137
remota, Megachile 305
reniformis, Odynerus, *Vespa* 187
repanda, Tenthredo 41
repandum, Polochrum 145
repandus, Mellinus 221
repertitus, Allantus scrophulariae 97
repentinus, Unitaxonus 39
repressa, Andrena hyemala ssp. 266
reptans, Pseudoanthidium,
 Anthidium 302
republicus, Crossocerus 229
reseri, Parna 51

resinana, Megachile 303
respersus, Rhogogastera 92
respondens, Euura, *Nematus* 64
resurgens, Halictus 277
Rethrax 89
reticulata, Caenolyda, *Tenthredo* 102
reticulata, Euura, *Nematus* 64
reticulata, Notogonia 237
reticulata, Tenthredo 102
reticulatum, Pseudoanthidium,
 Anthidium 302
reticulatus, Ceratoculus 233
reticulatus, Epyris 123
reticulatus, Sphecodes 290
retowskii, Parapsammophila 213
retusa, Anthophora, *Apis* 325
retusa, Pristiphora, *Nematus* 73
retusiformis, Anthophora 325
retusus, Elampus, *Notozus* 128
reuteri, Pristiphora, *Lygaeonematus* 74
rex, Amauronematus 59
rex, Paravespa, *Odynerus* 188
Rhadinoceraea 52
Rhadinoceraea subg. 52
rhaensis, Gonatopus 120
rhairopolis, Crabro 235
rhenana, Nomada 315
rhenana, Prosopis nigrafacies 260
rhenanum, Tetramorium 209
rhenanus, Halictus 281
rhenensis, Tenthredella solitaria 95
Rhineta 318
rhinoceros, Ammophila 213
rhinoceros, Osmia 297
rhinocerus, Tzustigmus 250
Rhinonitela 241
Rhinopsis 211
rhinotropis, Osmia 297
Rhipidioceros 107
rhodia, Andrena 269
rhodius, Euodynerus ypsilon 180
rhodius, Tachysphex 238
rhododaetyla, Dasypoda 293
rhodogaster, Dolerus 79
rhodosianus, Halictus 284
Rhodosmia 299
rhodostoma, Halictus cylindricus 282
Rhogogaster 88
Rhogogastera 88
rhombica, Eoferreola, *Sphex* 173
Rhombotilla subg. 156
rhopalocerus, Gorytes 221
Rhopalospiria 22
Rhopalum 235
Rhopalum subg. 235
Rhopitoides 291
Rhynchium 189
Rhynchogonatopus 120
ribauti, Passaloecus 248
ribesicola, Euura, *Pteronidea* 64
ribesii, Euura, *Tenthredo* 64
ribesii, Nematus 64
ribesii, Xiphydria, *Euxiphydria* 116
Ribinematus 55
ribis, Macrophyia, *Tenthredo* 86
ribis, Nematus 64
richardi, Athalia 44
richardsi, Cryptocheilus 161
Richardsdryinus 120
Ridestus 167
ridibundus, Andrena ventricosa ssp. 267
rifana, Pontania 67
rigiana, Chrysis rutilans 143
rimalis, Sphecodes 288
rimosicutum, Crocisa 327
rimosus, Hylaeus 260
rinki, Hylaeus, *Prosopis* 260
riparia, Apis 272
riparius, Halictus 284
riparius, Polistes 193
ritsemae, Stizus 225
rikuuiensis, Andrena 257
rijabovi, Ammoplanus 246
rijabovi, Cephus 109
robbinsi, Pontania 58
roberjeotiana, Nomada 315
robiniae, Nematus 65
robinsoni, Selandria 52
robortovskiyi, Colletes 258
robusta, Anthophora, *Megilla* 324
robusta, Cerceris 253
robusta, Cilissa 294
robusta, Coelioxys 304
robusta, Eucera 320
robusta, Hoplitis, *Heriades* 297
robusta, Mutilla 157
robusta, Nomada 313
robusta, Pristiphora, *Lygaeonematus* 74
robustior, Allantus amaena 97
robustus, Cryptocampus 56
robustus, Eumenes 182
robustus, Macrocephus 111
robustus, Odynerus melanocephalus 187
Rocalia 81
roettgeni, Passaloecus 248
rogenhoferi, Ammatomus, *Gorytes* 221
rogeri, Trichomyrmex 209
rolleri, Phyllocolpa 60
romana, Osmia 297
romani, Panurginus 276
romani, Pontopristia 61
romankovae, Andrena 267
romankovae, Deuteragenia,
 Dipogon 164
rondoui, Osmia 299
Ronisia 157
Rophites 291
Rophitinae 277, 290
roripae, Andrena 270
rosae, Andrena 267
rosae, Athalia, *Tenthredo rosae* 44
rosae, Selandria 46
rosarum, Blennocampa 50
rosarum, Hylotoma 24
roscidus, Pachynematus 66
roscipes, Andrena 267

rosea, Chrysis carnea var. 129
rosea, Tenthredo discophora 95
roseni, Eurhadinoceraea,
Rhadinoceraea 47
roseni, Rhadinoceraea 47
roseum, Hedychridium, *Chrysis* 129
rosincola, Arge 24
rosmarinifolia, Eupontania
collectanea 58
rossica, Cerceris 254
rossica, Ferreola 173
rossica, Nomada lineola var. 313
rossica, Sysphingta 210
rossicum, Hedychridium 129
rossicus, Epeolus variegatus var. 317
rossicus, Gonatopus 121
rossicus, Halictus 278
rossicus, Nipponodipogon, *Dipogon* 164
rossicus, Psen 250
rossii, Allodynerus, *Odynerus* 178
rossii, Bembix 222
rossii, Tenthredo, *Allantus* bifasciata 90
Rossomyrmex 203
rostellata, Andrena 271
rostralus, Pasites 318
rostrata, Apis 221
rostrata, Apis 222
rostrata, Bembex 221
rostrata, Bembix, *Apis rostrata* 222
rostrata, Nomada 311
rostratum, Lasioglossum, *Hylaeus* 281
rothi, Ammophila 213
rothneyi, Plagiolepis 202
rothneyi, Polistes 192
rotundarius, Crabro 230
rotundata, Apis 323
rotundata, Megachile 295
rotundata, Megachile, *Apis* 306
rotundatus, *Hylaeus* 259
rotundiventris, Megachile 303
roubali, Nysson 220
roubali, Pseudisobrachium 123
Royanthidium subg. 302
royoi, Anthidium 302
rozenburgensis, Epeolus 317
rubecula, Tenthredo 94
rubelloides, Halictus albipes 282
rubellus, Cryptocheilus, *Pompilus* 162
rubellus, Halictus 283
rubellus, *Hylaeus* 282
rubens, Formica exsecta var. 199
rubens, Halictus 282
rubeola, Tenthredella livida 95
ruberrima, Monophadnus spinolae 51
rubescens, Chrysis analis var. 133
rubescens, Nomada chrysopyga var. 312
rubi, Allantus 88
rubi, Empria 41
rubi, Fenusa 50
rubi, Metallus 50
rubi, Monophadnoides, *Selandria* 50
rubi, Perineura, *Allantus* 88
rubi, Priophorus 54

rubi, Selandria 50
rubi, Trypoxylon figulus minus 245
rubicanus, Dolerus elderi 78
rubicola, Ectemnius, *Solenius* 232
rubicola, Empria 41
rubicola, Heriades 296
rubicola, Odynerus 184
rubicola, Prosopis 259
rubicola, Psenulus 252
rubicolor, Camponotus
marginatus var. 198
rubicunda, Pseudarge 25
rubicundus, Halictus, *Apis* 278
rubicundus, Sphecodes 290
rubida, Andrena 264
rubida, Cerceris, *Philanthus* 254
rubida, Coelioxys erythropygia var. 304
rubida, Formica 204
rubida, Manica, *Formica* 204
rubidus, Crabro alatus 234
rubigidorsum, Psammochares 167
rubiginosa, Euura, *Amauronematus* 64
rubiginosa, Pseudepipona,
Odynerus 188
rubiginosa, Tenthredo 90, 99
rubiginosum, Anthidium 301
rubiginosum, Trypoxylon 245
rubivorus, Priophorus 54
ruborum, Halictus lerouxi 278
ruborum, Osmia 299
rubra, Andrena florea var. 273
rubra, Formica 205
rubra, Myrmica, *Formica* 206
rubra, Nomada 311, 313
rubra, Strongylogaster 82
rubricans, Calicurgus 165
rubricans, Mutilla 153
rubricans, Poecilagenia, *Calicurgus* 165
rubricata, Andrena 273
rubricollis, Nomada 316
rubricornis, Eumenes papillarius 182
rubricosa, Nomada 314
rubricosa, Nomada 316
rubricoxis, Tenthredo, *Tenthredella* 95
rubrifasciata, Chrysis 135
rubrimana, Megachile 306
rubripes, Macrophyia 85
rubripes, Odynerus 180
rubripes, Osmia 297
rubripes, Sphecodes 287
rubripes, Tenthredella livida 95
rubripes, Tenthredo 85
rubripes, Tenthredo lachlaniana 95
rubripyga, Chrysis 137
rubriventris, Andrena
ephippium var. 270
rubriventris, Mischophus 241
rubriventris, Philanthus 255
rubriventris, Tenthredopsis parvula 99
rubrocaudata, Tenthredella 94
rubrocaudatus, Nematinus 69
rubrofasciatus, Strongylogaster 43
rubromaculatus, Hoplomerus
tinniens var. 184
rubronotatus, Eumenes 182
rubropictus, Crabro 233
rubrosignata, Andrena 271
rubrosignata, Mutilla 154
rubrosignatus, Euodynerus
quadriseriatus 183
rubtzovi, Megachile 306
ruddii, Chrysis ruddii 137
ruddii, Embolemus 122
ruddii, Embolemus 122
rudepunctata, Tiphia 150
ruderarius, Bombus, *Apis* 331
ruderatus, Bombus, *Apis* 328
Rudia 154
rudis, Dasypoda 293
rudis, Prosopis lineolata 261
rudolfae, Andrena 267
rudolfae, Smicromyrme 155
rudolphae, Anthophora 325
rudowi, Tenthredo 94
rufa, Andrena hattorfiana var. 265
rufa, Anthophora 324
rufa, Apis 299
rufa, Arachnospila, *Psammochares* 170
rufa, Crocisa 327
rufa, Formica 199, 200
rufa, Holcocneme lucidus 69
rufa, Hylotoma 34
rufa, Psammochares cinctellus f. 167
rufa, Tenthredo 30, 67,
rufa, Tenthredo pectinata 34
rufa, Tenthredopsis 100
rufa, Vespa, *Vespa* 196
rufata, Tenthredopsis nassata 99
rufatus, Dolerus haematoches 80
rufescens, Apis 287
rufescens, Coelioxys 304
rufescens, Eumenes samurayi 182
rufescens, Formica 202
rufescens, Hylotoma 22, 23
rufescens, Myrmecomorphus 122
rufescens, Nematus 60
rufescens, Notozus superbus var. 127
rufescens, Polistes gallicus var. 193
rufescens, Polyergus, *Formica* 202
rufescens, Tetralonia 321
rufibarbis, Formica 201
ruficapilla, Tenthredo 67
ruficeps, Agenioideus, *Pompilus* 168
ruficeps, Allantus, *Emphytus* 38
ruficeps, Cystomutilla, *Mutilla* 153
ruficeps, Mutilla 153
ruficeps, Mutilla europaea f. 157
ruficeps, Nematus 62
ruficeps, Xiphydria 115
ruficinctus, Coelioxys 305
ruficollis, Ammophila 212
ruficollis, Eucera, *Macrocera* 321
ruficollis, Evania 223
ruficollis, Nematus 72
ruficollis, Smicromyrme, *Mutilla*
ruficollis 156
ruficornis, Andrena 270

ruficornis, *Apis* 311
ruficornis, *Athalia rosae* 44
ruficornis, *Bembix* 225
ruficornis, *Bethylus* 150
ruficornis, *Cerceris*, *Philanthus* 254
ruficornis, *Ceropales* 174
ruficornis, *Ectemnius*, *Crabro* 231
ruficornis, *Emphytus succinctus* 38
ruficornis, *Eriocampa* 89
ruficornis, *Gorytes* 223
ruficornis, *Lonchodryinus*,
 Gonatopus 119
ruficornis, *Macrocerca* 322
ruficornis, *Mellinus* 246
ruficornis, *Nomada*, *Apis* 316
ruficornis, *Nomia* 291
ruficornis, *Odynerus* 187
ruficornis, *Pompilus* 167
ruficornis, *Pristiphora*, *Nematus* 74
ruficornis, *Siobla*, *Allantus* 89
ruficornis, *Sphecodes* 289
ruficornis, *Stizus*, *Vespa ruficornis* 225
ruficornis, *Tenthredo* 95
ruficornis, *Vespa* 225
ruficurus, *Monophadnoides*,
 Selandria 50
ruficrus, *Andrena* 267
ruficrus, *Nomada fabriciana* var. 312
ruficrus, *Sphecodes* 289
rufifrons, *Melanapis* 270
rufilabris, *Nomada* 313
rufilata, *Andrena hystric* var. 263
rufimaculatum, *Trypoxylon* 245
rufimana, *Tenthredo* 98
rufina, *Leptochilus alpestris*
 iberobarbarus var. 185
rufina, *Paravespa rex* var. 188
rufinotis, *Empria abdominalis* 43
rufipe, *Sphecodes* 150
rufipectus, *Monophadnus spinolae* 51
rufipennis, *Hylaeus*
 nigrocuneatus var. 261
rufipennis, *Sphecodes* 287
rufipennis, *Tenthredo* 94
rufipes, *Andrena* 277
rufipes, *Apis* 303, 323
rufipes, *Astata rufipes* 218
rufipes, *Bembex* 225
rufipes, *Bethylus* 124
rufipes, *Cerceris* 255
rufipes, *Cladius* 54
rufipes, *Crossocerus*
rufipes, *Dasypoda argentata* 293
rufipes, *Dolerus aericeps* 77
rufipes, *Epeolus* 317
rufipes, *Episyron*, *Sphecodes* 167
rufipes, *Eucera* 320
rufipes, *Euodynerus quadrifasciatus* 183
rufipes, *Hylaeus* 286
rufipes, *Leptopus* 70
rufipes, *Macrophyta*, *Tenthredo* 86
rufipes, *Mutilla* 156
rufipes, *Nematus* 70

rufipes, *Nomada* 315
rufipes, *Nomada* 316
rufipes, *Nysson* 219
rufipes, *Odynerus oviventris* var. 179
rufipes, *Orussus* 117
rufipes, *Oxybelus* 242
rufipes, *Phauloclavelia* 173
rufipes, *Pristiphora* 74
rufipes, *Sapyga* 145
rufipes, *Smicromyrme*, *Mutilla* 156
rufipes, *Sphecodes gibbus*
 ephippium 289
rufipes, *Sphecodes* 166
rufipes, *Tachytes* 238
rufipes, *Tenthredo* 39, 86, 93, 95
rufipus, *Dolerus* 78
rufispinosus, *Sphecodes gibbus* 288
rufispinum, *Anthidium* 300
rufisternis, *Tenthredo ferruginea* 94
rufitarse, *Lasioglossum*, *Halictus* 286
rufitarsis, *Ammobates* 318
rufitarsis, *Chrysis* 137
rufitarsis, *Elampus*, *Notozus* 128
rufitarsis, *Eucera nigrilabris* 320
rufitarsis, *Gastrosericus* 236
rufitarsis, *Megachile* 307
rufitarsis, *Melitta* 267
rufitarsis, *Messor auct.* 205
rufitarsis, *Osmia* 299
rufitarsis, *Oxybelus* 242
rufitarsis, *Selandria* 76
rufithorax, *Arachnotheutes*,
 Pompilus 173
rufithorax, *Pompilus* 173
rufithorax, *Prospis* 262
rufithorax, *Sphecodes* 289
rufiventralis, *Tachysphex* 238
rufiventris, *Ammobates* 318, 319
rufiventris, *Andrena* 267
rufiventris, *Apis* 314
rufiventris, *Arge segmentaria* 24
rufiventris, *Cephus parreyssi* 111
rufiventris, *Crabro* 235
rufiventris, *Crabro* 236
rufiventris, *Epeolus cruciger* var. 317
rufiventris, *Gastrosericus* 236
rufiventris, *Halictus* 282
rufiventris, *Lyrops* 239
rufiventris, *Megachile* 307
rufiventris, *Miscophus* 241
rufiventris, *Nomia* 291
rufiventris, *Phileremus* 319
rufiventris, *Pristiphora* 74
rufiventris, *Scolia* 147
rufiventris, *Sphecodes*, *Tiphia* 290
rufiventris, *Stizus rufiventris* 225
rufiventris, *Tenthredo* 94
rufizona, *Andrena* 268
rufocincta, *Andrena ephippium* var. 270
rufocincta, *Apis* 312
rufocincta, *Arge* 24
rufocincta, *Lyda* 103
rufocincta, *Tenthredo* 38

rufocinctella, *Strongylogastroidea* 39
rufocinctus, *Allantus*, *Tenthredo* 38
rufocinctus, *Sphecodes* 215
rufocingulatus, *Allantus* 90
rufoclypeata, *Andrena* 265
rufofasciata, *Onycholyda*, *Lyda* 103
rufofemorata, *Tenthredopsis*
 austriaca 99
rufohirta, *Osmia* 297
rufohispida, *Andrena* 274
rufolineatus, *Polistes* 192
rufomacula, *Euura*, *Amauronematus* 64
rufomaculata, *Andrena* 275
rufomaculata, *Dolerus gessneri* 78
rufomaculata, *Formica exsecta*
 pressilabris var. 199
rufomaculata, *Myrmosa*
 melanocephala f. 152
rufomaculata, *Nomada lineola* var. 313
rufoniger, *Monophadnus* 47
rufo-niger, *Tachysphex* 239
rufonodis, *Cerceris* 255
rufonotalis, *Tenthredo* 93
rufonotatus, *Dolerus liogaster* 80
rufopicta, *Apis* 316
rufopictus, *Nippononysson* 219
Rufopompilus 167
rufopratensis, *Formica rufa* var. 199
rufoscutellata, *Arge captiva* 22
rufoscutellata, *Athalia* 45
rufoscutellata, *Vespa* 195
rufosignata, *Vespula* 196
rufospinosus, *Hoplomerus* 187
rufotegularis, *Halictus* 287
rufoterminalis, *Tenthredella solitaria* 95
rufotruncicola, *Formica rufa* var. 200
rufula, *Mutilla europaea* f. 157
rufula, *Nomada succineta* var. 316
rufulocollis, *Anteon* 119
rufus, *Cladius* 54
rufus, *Euura*, *Amauronematus* 64
rufus, *Halictus* 279
rufus, *Miscophus gallicus* 241
rufus, *Phiarus abdominalis* var. 318
rufus, *Psammochares* 167
rufus, *Pseudogonatopus* 121
rugicollis, *Hylaeus* 262
rugifer, *Ectemnius*, *Crabro* 232
rugifer, *Pempredon*, *Cemonus* 249
ruginodis, *Myrmica* 206
rugithorax, *Nomada* 312
rugosa, *Calameuta* 108
rugosa, *Tiphia* 150
rugosopunctata, *Poecilotiphia*,
 Myzine 150
rugosopunctatus, *Crabro* 232
rugosula, *Hylaeus* 285
rugosulus, *Halictus* 279
rugosus, *Crabro* 228
Rugosus, *Crossocerus* 228
rugosus, *Pamphilus* 105
rugosus, *Sphecodes* 287
rugosus, *Tachysphex* 238

Rugovespula 195
rugulosa, Andrena 270
rugulosa, Myrmica 206
rugulosa, Andrena 271
rugulosoides, Myrmica
 scabrinodis var. 206
rugulososcabrinodis, Myrmica
 rugulosa var. 205
rugulosus, Euchroeus, *Chrysis* 140
rugulosus, Oxybelus latro 242
ruida, Aphaenogaster famelica 203
ruiescente, Anthidium
 florentinum var. 300
rumelica, Nomia 291
rumicis, Unitaxonus 39
runari, Pseudoheptamelus 36
Runaria 27
rupestris, Bombus, *Apis* 329
rupestris, Prosopis 259
rupestris, Vespa 192
rupico, Allantus 91
ruralis, Emphytus rufocinctus 38
ruralis, Tenthredo 38
russa, Tenthredo 29
russica, Chrysis 137
russky, Odynerus 186
rustica, *Apis* 299
rustica, Arge, *Tenthredo* 24
rustica, Osmia 298
rusticanus, Nematus 72
Rusticiformis 22
ruthena, Tarpa 107
ruthena, Tenthredo 95
ruthenica, Anthophora 325
ruthenica, Cilissa 294
ruthenica, Didineis 219
ruthenicus, Crabro 233
ruthenicus, Epeolus 317
ruthenicus, Odynerus parvulus var. 180
ruthenus, Ellampus 133
rutila, Andrena 270
rutilans, Anthophora 324
rutilans, Chrysis rutilans 137
rutilans, Hedychrum rutilans 130
rutilicornis, Tenthredo 68
rutiliventris, Chrysis rutiliventris 137
ruzskyi, Camponotus
 marginatus var. 198
ruzskyi, Strongylognathus huberi 207
rybyensis, Cerceris, *Sphecodes* 254
rybyensis, Sphecodes 252
Rysepyris 123
ryukyuensis, Anoplus 172
ryukyuensis, Euodynerus notatus 183

sabandus, Allantus amaenus 96
sabariensis, Tenthredo, *Allantus* 91
sabatia, Smicromyrme 156
sabularum, Lasius, *Formicina* 201
sabuleti, Myrmica 206
sabulosa, Ammophila, *Sphecodes* 212
sabulosa, *Apis* 267
sabulosa, Cerceris, *Philanthus* 254

sabulosa, Ceropales bogdanovii 160
sabulosus, Crabro 246
sabulosus, Lasioglossum, *Halictus* 286
sabulosus, Sphecodes 288
sabulosus, Sphecodes 212
sachalinense, Aglaostigma,
 Neurosiobla 84
sachalinense, Trichiosoma 31
sachalinensis, Abia 29
sachalinensis, Andrena florea ssp. 267
sachalinensis, Anoplus 172
sachalinensis, Bombus sporadicus 328
sachalinensis, Calameuta, *Cephus* 109
sachalinensis, Camponotus 198
sachalinensis, Camponotus
 herculeanus var. 198
sachalinensis, Cephalcia, *Lyda* 102
sachalinensis, Cimbex
 quadrimaculata 29
sachalinensis, Dolerus 81
sachalinensis, Gilpinia 34
sachalinensis, Leptothorax tuberum 208
sachalinensis, Pseudepipona 183
sachalinensis, Rhadinoecraea 52
sachalinensis, Tenthredo 98
sachalinensis, Urocerus 113
sagemeqli, Nomada lineola var. 313
sagittata, *Notoglossa* 242
sagmaria, Tenthredopsis raddatzi 99
sagmarius, Euura, *Amauronematus* 64
sagulatus, Pachynematus 72
sah, Urocerus, *Sirex* 113
sahlbergi, Andrena 273
sahlbergi, Cardiocondyla 203
sahlbergi, Cardiocondyla
 elegans var. 203
Sahlbergi, Crabro 234
sahlbergi, Entomognathus, *Crabro* 234
sahlbergi, Evagetes, *Pompilus* 171
sahlbergi, Halictus 285
Sahlbergia 83
saigusai, Anoplus 172
Sainiella 90
saishiuensis, Palarus 243
sajanicus, Pamphilus 103
sajoi, Halictus 278
sakagamii, Andrena 264
sakagamii, Bombus ardens 330
sakagamii, Lasioglossum 286
Sala 70
salebrosa, Vespa koreensis 196
salentina, Mutilla 157
salicacea, Andrena 268
salicariae, Tetralonialla, *Macrocerca* 322
saliceti, Cimbex 30
saliceti, Hylotoma 24
saliceti, Nematus 64
saliceti, Tenthredo 56
salicicola, Euura 56
salicicola, Pachynematus albipennis 57
salicis, Andrena 275
salicis, Cimbex variabilis 30
salicis, Euura, *Tenthredo* 64

salicis, Tenthredo 61, 64
salicis, Trichiosoma 31
saliciscinereae, Euura, *Tenthredo* 64
salicislapponicae, Euura 56
salicispentandrae, Tenthredo 56
salicivora, Pristiphora,
 Lygaeonematus 74
salicivorus, Nematus 62
salina, Leptothorax tuberum var. 208
salina, Myrmica 206
salina, Myrmica scabrinodis var. 206
Salius 161
saltensis, Mutilla 157
saltuarius, Bombus, *Hortobombus* 328
saltuum, Neurotoma, *Tenthredo* 103
salviae, Amegilla, *Anthophora* 323
salviae, Anthophora, *Lasius* 323
samarensis, Halictus 278
samaricum, Lasioglossum, *Halictus* 286
samariensis, Lophopompilus, *Sphecodes* 173
sambuci, Allantus 86
sambucicola, Crabro 228
sameshima, Psen 251
samolad, Euura, *Pontania* 64
samurai, Polyergus 202
samuray, Eumenes 182
samurayi, Rhynchium
 haemorrhoidale 189
sancta, Myrmica scabrinodis var. 206
sanctus, Spatulocephus 110
sanguinatus, Sphecodes
 chionospilus 289
sanguinea, Camponotus
 japonicus var. 198
sanguinea, Formica 200
sanguineus, Phiarus
 abdominalis var. 318
sanguinicollis, Dolerus, *Tenthredo* 80
sanguinicollis, Eurhadinoecraea,
 Blennocampa 47
sanguinicollis, Hylotoma 24
sanguinipes, Macrophyta 85
sanguiniventris, Andrena erberi var. 264
sanguinolenta, Arge 22
sanguinolenta, Macrophyta,
 Tenthredo 86
sanguinolenta, Sphecodes 166
sanguinolentus, Homonotus, *Sphecodes* 166
sannio, Polymematus 56
sanvitali, Tengyra 149
sanzii, Elampus 128
sapellonis, Andrena 268
sapidus, Stenodynerus 190
sapphirina, Chrysis inaequalis 136
sapphirus, Chrysis 136
sapporensis, Cimbex 29
sapporensis, Crabro 234
sapporensis, Pamphilus, *Lyda* 105
sapporensis, Pompilus 166
sapporensis, Tenthredo, *Allantus* 93
sapporoensis, Bombus 327
sapporoensis, Crabro 230, 233
sapporoensis, Crabro dimidiatus 230

sapporoensis, Sphecodes 289
sapporonis, Aglaostigma, *Macrophya* 84
Sapyga 145
Sapyga subg. 145
Sapygidae 145
Sapygina 146
Sapyginae 145
sarachensis, Eucera 321
sarafschani, Bembix 222
sardarica, Chrysis 137
sardinica, Andrena
 _{colletiformis} var. 264
sardinensis, Nematus 69
sardoa, Aphaenogaster 203
sardoa, Prosopis brevicornis 259
sardous, Allodynerus delphinalis 178
sareptana, Ammophila 212
sareptana, Cerceris 253
sareptana, Pristiphora 74
sareptana, Scolia 147
sareptana, Tenthredo 90
sareptana, Tenthredopsis 99
sareptana, Tetrachrysis
 _{grohmanni} var. 136
sareptana, Tiphia 150
sareptana, Tropidotilla, *Mutilla* 157
sareptanus, Astatus 218
sareptanus, Cephus 109
sareptanus, Eumenes 182
sareptanus, Gorytes 224
sareptanus, Halictus 284
sareptanus, Odynerus 187
sareptanus, Onychopterocheilus,
 Pterocheilus 187
sareptanus, Philoctetes, *Ellampus* 132
sareptanus, Podalirius retusus 325
sareptanus, Tachysphex panzeri 239
sareptensis, Sphecodes 289
sarkissiani, Tetramorium
 _{caespitum} var. 209
sarmatica, Myrmilla 154
sarmatica, Nomada zonata 317
sarmaticus, Stizus 225
sarnia, Andrena nigroaenea 269
Sarochilus subg. 185
saropoda, Anthophora 324
Sasmarila 150
sassai, Symmorphus 191
satanas, Tachysphex
 _{schmiedeknechti} 239
satanas, Tremex 114
satoi, Ceratina 310
satonis, Beleses, *Abelteses* 40
satschauensis, Halictus 281
satschouanus, Crabro 235
satschouanus, Lindenius, *Crabro* 235
satsumanus, Rhynchium 183
satunini, Hedychriddium 129
satunini, Plagiolepis 202
satunini, Temnothorax, *Leptocephalus* 208
satyrus, Astatus 110
saundersella, Andrena 271
saundersi, Crabro 231

saundersi, Macrophya 85
saundersi, Tenthredopsis 99
saundersii, Halictus 285
saussurei, Anthophora 324
saussurei, Cleptes 127
saussurei, Megachile 307
saussurei, Vespa 196
sauterianus, Nematus 61
savignyi, Megachile 305
Savignyi, Oxybelus 242
sawai, Perineura 94
Saxatilipomphilus 168
saxatilis, Camponotus 198
saxatilis, Camponotus herculeanus 198
saxatilis, Camponotus
 _{pennsylvanicus} 198
saxatilis, Dolerus 77
saxesenii, Pristiphora, *Nematus* 74
saxialis, Hoplitis, *Anthocopa* 297
saxicola, Lyda 102
saxonica, Dolichovespula, *Vespa* 195
saxonica, Tenthredo 95
sayi, Crabro 232
sayi, Rhogogaster 87
scabiosae, Tetaloniella, *Eucera* 322
scabricollis, Sphecodes 290
scabrinodis, Myrmica 206
scabrilavalis, Nematus 67
scabrosa, Andrena 265
scabrosus, Eumenes sareptanus 181
scalaris, Brachystegus, *Nysson* 219
scalaris, Nysson 219
scalaris, Rhogogaster, *Tenthredo* 88
scalesii, Trichiosoma 31
scamni, Leptocephalus 204
scamni, Leptocephalus muscorum var. 204
scandinavus, Odynerus 184
scaposus, Crossocerus 229
Scapteroides 276
scapulare, Anteon, *Dryinus* 119
scapularis, Astata, *Astatus miegii* 218
scapularis, Selandria 55
scapularis, Tenthredo 76
scardicus, Halictus 277
scariosus, Sphecodes gibbus similis 287
Sceliphrinae 214
Sceliphron 214
schaefferella, Apis 316
schaefferi, Cimbex 29
schaefferi, Mutilla 157
schaefferi, Tenthredo 86
schaefferi, Tenthredo 97
schelkovnikovi, Halictus 282
schencki, Andrena 271
schencki, Chrysis 138
schencki, Crabro dives 232
schencki, Mutilla 155
schencki, Myrmica 206
schencki, Myrmica rubra var. 206
schencki, Nomioides minutissima 292
schencki, Psenulus, *Psen* 252
schenckiana, Chrysis ignita 138
schenckii, Hylaeus 260
schenckii, Sphecodes 290
schenckioides, Myrmica 206
schenkella, Andrena 270
schenkii, Anthophora parietina var. 324
scheviakovi, Megachile 308
scheviakovi, Nomada 313
shevrevi, Sapyga caucasica 145
schewyrewi, Nematus, *Pteronus* 75
schiodtei, Priocnemis 163
schirmeri, Tenthredella atra 93
schirmeri, Tenthredo omissa 97
Schistonyx 171
Schistosphex 237
Schizagenia 165
Schizanoplus subg. 167
Schizoceros 26
Schizocerus 26
Schlettereri, Crabro 232
schlettereri, Ectemnius, *Crabro* 232
schlueteri, Euura, *Amauronematus* 64
schmidti, Andrena transitoria ssp. 274
schmidtii, Crematogaster, *Acrocoelia* 204
schmidtii, Dolerus 80
schmidtii, Parammobatodes 319
schmidtii, Pseudepipona 180
schmidtii, Nematus 59
Schmiedeknechti, Crabro 234
schmiedeknechti, Entomognathus,
 Crabro 234
schmiedeknechti, Gorytes 223
schmiedeknechti, Nomada 313
schmiedeknechti, Tachysphex 239
schmiedeknechti, Tenthredopsis 83, 239
schmiedeknechti, Eremiasphecium 245
schmiedeknechti, Hylotoma 23
schneideri, Dolerus liogaster 80
schneideri, Teleutomyrmex 208
schoenherri, Tenthredo 38
schottii, Nomada 318
schottii, Pasites 319
schränkii, Tenthredo 94
schränkella, Melitta 268
schrencki, Bombus 331
schrenckii, Scolia 148
schrenckii, Vespa 196
schulthess, Cerceris 253
schulthessi, Ellampus 132
schulthessi, Halictus 285
schulthessi, Leptolarra 237
schulthessi, Odynerus 188
schulthessii, Belomicrus 242
schummeli, Epeolus 317
schwarzii, Andrena 267
schwarzii, Smicromyrme 156
schwarzii, Sphecodes 290
Schyrocera 26
sciaphilus, Psen 250
Sciapterina 89
Sciapteryx 89
scintillans, Chrysis 138
Sciopteryx 89
scirpaceum, Lasioglossum, *Halictus* 286
scissa, Tenthredo 90

scita, Andrena 273
 scita, Arge, *Hylotoma* 24
 scita, Hoplitis, *Osmia* 297
scita, Nomada 314
scita, Tenthredo 95
Scitandrena subg. 273
scitulum, *Lasioglossum*, *Halictus* 281
Sclerochroa 124
Scleroderma 124
Sclerodermi 124
Sclerodermus 124
Scolia 147
Scolia subg. 147
Scoliidae 147
Scoliinae 147
Scoliini 147
Scolioidea 145
Scoliodes 147
Scolioneura 52
scopipes, *Andrena ephippium* var. 269
scopipes, *Andrena morio* var. 269
scopolii, *Tenthredo* 93
scotapsis, *Euura*, *Nematus* 64
scotica, *Andrena trimmerana* var. 267
scotica, *Pemphredon* 249
scotica, *Tenthredo* 93
scoticus, *Ancistrocerus*, *Odynerus* 179
scoticus, *Dolerus* 78
scoticus, *Nematus* 72
Scotomphales 225
scripta, *Tenthredo* 87
scrrophulariae, *Tenthredo* 90, 97
sculpticollis, *Chrysallampus*,
Omalus 127
sculptilis, *Tachysphex* 239
sculpturata, *Chrysis* 138
sculpturata, *Chrysis ignita* var. 138
sculpturata, *Notogonia* 237
sculpturata, *Poecilagena*,
Pseudagenia 165
sculpturatum, *Hedychridium*,
Hedychrum 129
sculpturatus, *Panurginus* 276
scutatum, *Trypoxylon scutatum* 245
scutatus, *Crabro* 227
scutatus, *Crabro* 229
scutellare, *Hedychridium*,
Hedychrum 129
scutellariae, *Athalia scutellariae* 45
scutellaris, *Allantus* 97
scutellaris, *Allantus kochleri* 90
scutellaris, *Blennocampa* 51
scutellaris, *Chrysis* 128
scutellaris, *Chrysis* 138
scutellaris, *Crabro* 229
scutellaris, *Crocisa* auct. 327
scutellaris, *Formica* 204
scutellaris, *Hoplomerus spinipes* var. 187
scutellaris, *Hylaeus* 260
scutellaris, *Hylotoma* 26
scutellaris, *Lyda* 102
scutellaris, *Macrophyia rustica* 86
scutellaris, *Nomada* 326

scutellaris, *Osmia* 298
scutellaris, *Oxybelus* 243
scutellaris, *Physetopoda*, *Mutilla* 155
scutellaris, *Stelis* 303
scutellaris, *Stizus* 225
scutellaris, *Tenthredopsis dubia* 99
scutellaris, *Tenthredopsis*,
Tenthredo 100
scutellaris, *Thyreus*, *Nomada* 327
scutellata, *Euura*, *Nematus* 65
scutellata, *Macrophya* 85
scutellata, *Sphecius* 226, 230
scutellata, *Tenthredo*, *Allantus* 93
scutellata, *Crabro* 230
scutellatus, *Crabro*, *Sphecius* 226
scutifera, *Cerceris* 255
scutigerum, *Trypoxylon* 245
scutularia, *Sphecius* 226
scythes, *Fervidobombus* 331
scythia, *Arachnospila* 169
scythicus, *Characopygus* 110
scythicus, *Megalodontes* 107
seabrai, *Solerella* 241
sebetia, *Caliroa* 45
secunda, *Andrena thoracica* var. 269
secunda, *Strongylogaster*,
Pseudotaxonus 82
sedankiana, *Tenthredo* 92
sedilloti, *Megachile* 306
sedula, *Eucera* 321
seiboldi, *Tenthredo solitaria* 95
seesana, *Tenthredo* 82
segmentaria, *Tenthredo* 90
segmentarius, *Cryptus* 24
segmentarius, *Nematus* 62
segmentata, *Tenthredo* 43
segnis, *Anthophora* 324
seguro, *Tenthredo finschi* 93
segusina, *Anthophora* 323
seitzii, *Trachusa byssina* var. 303
sejensis, *Pachyprotasis sejensis* 88
sekerai, *Tenthredo olivacea* 92
Seladonia subg. 278
seladonius, *Halictus*, *Apis* 279
Selandria 81
Selandria subg. 75
Selandriidae 76
selandriiformis, *Dineura* 68
Selandriinae 75
selandrioides, *Nematus* 74
Selandropha 75
sellata, *Mutilla* 156
sellata, *Pseudepipona*, *Lionotus* 188
semenovi, *Acantholepis*
frauendorfii var. 202
semenovi, *Anthophora* 324
semenovi, *Hartigia* 111
semenovi, *Lepisiota*, *Acantholepis* 202
semenovi, *Prenanteon* 119
semenovi, *Pseudoclavellaria*,
Clavellaria 30
semenovi, *Tachysphex* 238
semenovi, *Temnothorax*,
Leptocephalus 208
semenoviana, *Abia* 29
semenoviellus, *Bombus* 328
semenowi, *Allantus* 95
semenowi, *Hoplomerus* 187
semenowi, *Sciapteryx* 89
semiaeneus, *Halictus* 282
semiate, *Podalirius simplicipes* 324
semiaurata, *Sphecius* 126
semiauratus, *Cleptes* auct. 127
semiauratus, *Sphecius* 127
semicincta, *Tenthredo* 46, 93
semicinctus, *Hylaeus* 280
semicircularis, *Megachile* auct. 306
semicircularis, *Omalus* 132
semicoerulea, *Arge*, *Tenthredo* 25
semicornis, *Tenthredo* 94
semicyaneus, *Cleptes* 127
semiflava, *Andrena* 264
semiflava, *Lyda* 102
semilactea, *Euura*, *Nematus* 65
semilaeve, *Lasioglossum*, *Halictus* 286
semilaevis, *Andrena* 271
semilucens, *Lasioglossum*, *Halictus* 286
semilunata, *Cerceris* 255
seminiger, *Allantus amaenus* 97
seminigra, *Anthophora retusa* 325
semigrisea, *Bembix* 222
semigrisea, *Brachythops* 76
seminitidus, *Psen* 251
seminuda, *Andrena* 273
seminuda, *Eucera* 321
semiopaca, *Euura*, *Pteronidea* 65
semiorbitalis, *Nematus* 63
semipleta, *Megachile* 307
semirubra, *Andrena* 273
semirufa, *Cerceris* 255
semirufus, *Armitarsus* 84
semirugosa, *Andrena* 271
semistriatus, *Gorytes* 224
semistrigosa, *Eucera* 320
semitectus, *Halictus* 279
semitomentosus, *Halictus* 283
semperi, *Anthophora* 324
sempersolis, *Euura*, *Amauronematus* 65
semseyi, *Tenthredo* 92
seneciorum, *Andrena* 265
senescens, *Anthophora* 323
senex, *Andrena* 269
senex, *Hylaeus* 278
senex, *Megachile* 307
senile, *Anthidium* 301
senilis, *Bembix* 222
senilis, *Colletes*, *Andrena* 258
senilis, *Halictus*, *Hylaeus* 277
Senoclidea 53
sensitivus, *Embolemus* 122
seorsa, *Pristiphora* 71
seoulensis, *Cerceris quinquefasciata* 254
seoulensis, *Symmorphus* 191
separanda, *Ammophila* 212
separanda, *Andrena* 273
separata, *Andrena* 267
separatus, *Paranoplius* 172

- separatus, *Pompilus* 172
sepsoides, *Gonatopus* 120
septemdentatus, *Gonatopus* 120
septemspinous, *Anthidium* 301
septentrionalis, *Arachnospila* 168
septentrionalis, *Crabro* 231
septentrionalis, *Eumenes* 182
septentrionalis, *Euura*,
 Amauronematus 65
septentrionalis, *Nematus*, *Tenthredo* 69
septentrionalis, *Smicromyrme*
 pusilla 156
seraxense, *Anthidium* 300
seraxensis, *Epeorus transitorius* var. 317
seraxensis, *Palarus* 243
serena, *Chrysis serena* 138
serica, *Arachnospila anceps* 168
sericans, *Megachile* 305
sericans, *Tenthredo* 50
sericata, *Abia* 29
sericata, *Andrena* 267
sericata, *Prosopis* 261
sericatus, *Oxybelus* 243
sericea, *Abia*, *Tenthredo* 29
sericea, *Apis* 268
sericea, *Tenthredo* 28
Sericiformis 30
sericeiventris, *Mutilla* 157
sericeomaculata, *Pompilus* 168
sericeum, *Pison* 244
sericeum, *Trichiosoma* 31
sericeus, *Agenioideus*, *Pompilus* 167
sericeus, *Anteon* 119
sericultorum, *Nomada* 314
seriepunctata, *Euura*, *Pteronidea* 65
sermola, *Pristiphora* 74
serotina, *Andrena* 269
serotina, *Nomada flavoguttata* var. 312
serotina, *Tenthredo* 40
serotinella, *Andrena*
 bimaculata var. 272
serotinus, *Apethymus*, *Tenthredo* 40
serotinus, *Crabro* 232
serotinus, *Nematus* 73
Serpapinta 150
serradifera, *Tenthredo* 95
serraticeps, *Anthidium* 302
serraticornis, *Psen* 251
serratulae, *Anthidium* 303
serratulae, *Trachusa* 303
serratus, *Ammoplanus* 246
serratus, *Aphelopus* 118
serricus, *Odynerus*, *Oplomerus* 187
serrilabris, *Osmia* 297
serripes, *Crabro* 230
serripes, *Odynerus* 187
serripes, *Pterocheilus* 186, 187
serrisquama, *Bombus cullumanus* 328
sertata, *Onycholyda*, *Pamphilus* 103
sertifer, *Neodiprion*, *Tenthredo* 34
sertus, *Smicromyrme*, *Mutilla* 156
serva, *Selandria*, *Tenthredo* 81
serva, *Tenthredo* 81
serviculus, *Temnothorax*,
 Leptocephalus 208
Serviformica subg. 200
servulellus, *Halictus* 283
servus, *Crabro* 228
setigera, *Andrena* 273
setosa, *Andrena* 275
setosus, *Ammobates* 318
setteri, *Nomada* 316
setulellum, *Lasioglossum*, *Halictus* 286
seuli, *Rhynchium* 183
sexannulata, *Tenthredo* 97
sexcincta, *Apis* 313
sexcincta, *Vespa* 196
sexcinctus, *Ectemnius*, *Crabro* 231
sexcinctus, *Halictus*, *Apis*
 sexcinctus 277
sexdentata, *Chrysis* 138
sexfasciata, *Nomada* 316
sexfasciata, *Vespa* 187
sexguttata, *Scolia* 145
sexmaculata, *Megachile* 306
sexmaculata, *Mutilla* 157
sexmaculata, *Scolia* 147
sexmaculata, *Scolia*, *Vespa* 148
sexmaculata, *Vespa* 147
sexmaculatum, *Anthidium* 301
sexmaculatus, *Aporinellus*,
 Pompilus 168
sexmaculatus, *Crabro* 230
sexnotatum, *Lasioglossum*,
 Halictus 281
sexnotatum, *Lasioglossum*, *Melitta* 281
sexpunctata, *Empria*, *Tenthredo* 41
sexpunctata, *Meria* 150
sexpunctata, *Tenthredo* 41
sexpunctata, *Trachusa* 302
sexpunctata, *Vespa* 179
sexpunctata, *Vespa mongolica* var. 194
sexpunctatus, *Hellus* 145
sexsignata, *Stelis* 302
sexsignatus, *Halictus* 285
sexspinosa, *Nomada* 312
sexstrigatum, *Lasioglossum*,
 Halictus 286
shaman, *Cerceris* 253
shaposhnikovi, *Bombus* 329
shaposhnikovi, *Tenthredo*, *Allantus* 92
sharificus, *Halictus malachurus* 284
Shariphora 70
sharpi, *Nematus* 61
shawella, *Melitta* 271
shawi, *Monophadnoides* 50
shawii, *Pemphredon* 249
sheppardana, *Nomada*, *Apis* 316
shestakovi, *Ancistromma* 236
Shestakovia 245
shestakoviiana, *Cerceris* 255
Shestakoviella 255
shestoperovi, *Allantus* 90
shibuyai, *Crossocerus*, *Crabro* 227
shibuyai, *Odynerus* 180
shidai, *Priocnemis* 163
shimizui, *Poecilagenia* 165
shimoyamai, *Trypoxylon* 245
shinanoensis, *Vespa* 195
shinoharai, *Nematus*, *Craesus* 69
shirakii, *Calicurgus* 162
shiroyamai, *Syrmophilus* 191
shirozui, *Andrena* 264
shirozui, *Sphecodes* 288
shishkini, *Halictus* 283
shitkovi, *Camponotus*
 herculeanus var. 198
shuckardi, *Cemonus* 248
Shuckardi, *Crabro* 229, 233
shuckardi, *Mimesa* 250
shuckardi, *Passaloecus* 248
shukardi, *Nysson* 220
shumagensis, *Pteronus* 63
shur, *Cerceris* 253
Sitalus 162
siberica, *Dasylabris*, *Mutilla* 154
siberiensis, *Megalodontes* 107
sibiriacum, *Lasioglossum*, *Halictus* 286
sibirica, *Abia* 29
sibirica, *Amasis lateralis* 32
sibirica, *Andrena* 275
sibirica, *Arachnospila* 169
sibirica, *Blennocampa* 45
sibirica, *Cerceris* 255
sibirica, *Ceropales* 160
sibirica, *Chrysis* 138
sibirica, *Didineis* 219
sibirica, *Dufourea paradoxa* 290
sibirica, *Euura saliceti* 64
sibirica, *Euura*, *Pontania* 65
sibirica, *Holopyga* 131
sibirica, *Macrophya* 86
sibirica, *Melecta* 326
sibirica, *Melitta*, *Ciliissa* 294
sibirica, *Mimesa*, *Psen* 250
sibirica, *Neurotoma* 103
sibirica, *Nomada errans* var. 312
sibirica, *Nomada lineola* var. 313
sibirica, *Rhogogaster* 88
sibirica, *Siobla* 89
sibirica, *Tachytes* 240
sibirica, *Tenthredo velox* 96
sibirica, *Tenthredo*, *Allantus* 98
sibirica, *Thyreomelecta*, *Crocisa* 326
sibirica, *Vespa* 196
sibirica, *Xyela* 21
sibiricana, *Mimumesa* 251
Sibiricobombus subg. 330
sibiricola, *Macrophya* 86
sibiricola, *Phylloecus* 111
sibiricola, *Rhadinoceraea* 52
sibiricola, *Tenthredo*, *Tenthredella* 92
sibiricum, *Aprosthemma* 26
sibiricum, *Banthanidium*,
 Anthidium 301
sibiricum, *Hedychridium* 129
sibiricum, *Trichiosoma* 31
sibiricus, *Ammoplanus* 246
sibiricus, *Bembecinus*, *Stizus*
 hungaricus 221

sibiricus, Bombus, Apis 330
sibiricus, Celonites 178
sibiricus, Cimbex 30
sibiricus, Crabro 226
sibiricus, Dicondylus 121
sibiricus, Dolerus germanicus 78
sibiricus, Dolichoderus 197
sibiricus, Dolichoderus quadripunctatus 197
sibiricus, Formicoxenus, *Leptothorax* 204
sibiricus, Hylaeus 260
sibiricus, Nematus 62
sibiricus, Odynerus 184
sibiricus, Philanthus 255
sibiricus, Psen 251
sibiricus, Pterocheilus, *Odynerus* 189
sibiricus, Sphecodes 290
sibiricus, Tachytes etruscus 240
sicanius, Microdynerus patagiatus 186
sicanus, Smicromyrme, *Mutilla* 156
sicelis, Microdynerus nugdunensis 186
sichelii, Anthophora 325
sichelii, Bombus 329
sichelii, Katamenes, *Eumenes* 185
sichotana, Megachile 308
sichuanensis, Spinarge 25
siciliaca, Chrysis pyrrhina 138
sickershusanus, Polyplanus 122
sickmanni, Ammophila 212
sickmanni, Chrysis 138
sicula, Anthophora 323
siculus, Ancistrocerus oviventris 179
siculus, Colletes tuberculatus 259
Siculus, Ectemnius 233
siculus, Evagetes, *Pompilus* 171
siculus, Hylaeus 261
siculus, Odynerus 185
sidemii, Bombus mocsaryi 331
sidemii, Colletes 258
sidorenkoi, Apethymus 40
sieboldii, Nematus 63
Sierocolpa subg. 150
Sierola 125
Sierolomorpha 151
Sierolomorphidae 151
sieversi, Osmia 298
sigma, Tenthredo 96
signata, Osmia 298
signata, Pteronidea segmentaria 60
signata, Stelis, *Anthidium* 302
signata, Tenthredo 85, 102
signaticeps, Pompilus 169
signaticollis, Paragymnomerus, *Hoplomerus* 187
signaticrus, Crabro 226
signatifrons, Discolia 147
signatum, Chelostoma 297
signatus, Colletes daviesanus 257
signatus, Crabro 230
signatus, Hylaeus, *Sphex* 262
signatus, Leptochilus 185
signatus, Triepeolus 317

signifer, Cephus haemorrhoidalis 108
sikhotealinensis, Evagetes, *Anopilus* 171
sikhotealinensis, Platyxiphidria 115
sikkimensis, Conaspidia 84
sikkimensis, Polistes rothneyi 192
sikorae, Tachysphex 238
silaceum, Proceratum 210
silantjewi, Bombus 328
Silaon 241
silaos, Odynerus 180
silensis, Tenthredo 95
sillata, Andrena 271
silvaticus, Chelogynus 119
silvester, Nematus 65
silvicola, Nomada 316
silvicola, Sphecodes 288
Simandrena subg. 273
similaris, Nematus 65
similator, Pseudopolistes sulcifer 193
similatus, Hylaeus 261
simile, Hedychrum 130
similis, Allantus 97
similis, Andrena 274
similis, Arge, *Hylotoma* 25
similis, Cephus 110
similis, Colletes 258
similis, Diprion, *Lophyrus* 33
similis, Dryudella, *Astatus* 218
similis, Epeolus 317
similis, Eucera 320
similis, Halictus 281
similis, Hylaeus 268, 285
similis, Hylotoma 24
similis, Meria 149
similis, Mutilla 157
similis, Odynerus 179
similis, Phystopoda, *Paramyrme* 155
similis, Pristiphora 74
similis, Prosopis 262
similis, Pteronus 66
similis, Rhogogaster 89
similis, Sapyga, *Sirex* 145
similis, Siobla, *Encarsioneura* 89
similis, Sphecodes 287
similis, Syrista 111
similis, Tenthredo 84
similis, Vespa 195
simillima, Andrena 265
simillima, Arge, *Hylotoma* 25
simillima, Nomada 316
simillima, Scolia 147
simillima, Stelis 303
simillima, Vespa 194
simillimus, Halictus 285
simillimus, Odynerus 187
simillimus, Sphecodes 290
simini, Heptapotamus 42
simius, Bombus lapponicus 330
Simplephytus 39
simplex, Ammoplanellus, *Ammoplanus* 246
simplex, Chrysura, *Chrysis simplex* 140
simplex, Coelioxys 305
simplex, Halictus 278
simplex, Hylotoma 68
simplex, Nomioides variegata 292
simplex, Oxybelus 242
simplex, Psen 252
simplex, Pteronidea pseudonotabilis 57
simplex, Vespa 183
simplicia, Mutilla 155
simpliciceps, Scrapter 319
simplicicornis, Pseudoscolia, *Philanthus* 256
simplicifrons, Pristiphora 74
simplicior, LasioGLOSSUM, *Halictus* 286
simplices, Anthophora 324
simplices, Corynopus 235
simplices, Pterocheilus 184
simulans, Dineura 75
simulans, Pachyprotasis, *Tenthredo* 88
simulatrix, Arge 25
sinaiticus, Tachysphex 238
sinense, Tapinoma 198
sinensis, Ammophila 213
sinensis, Andrena 269
sinensis, Aphaenogaster smythiesii 203
sinensis, Apis 332
sinensis, Hemipepsis 174
sinensis, Melitta 294
sinensis, Scolia 148
singula, Chrysis 138
singularis, Conaspidia, *Eriocampa* 84
singularis, Passaloecus 248
singularis, Rhadinoceraea 47
singularis, Sphecodes 290
singularis, Tiphia 150
sinica, Myrmica 205
sinicola, Hylaeus 261
sinicum, Hedychrum 142
sinister, Halictus zonulus 281
sinkiangensis, Melitta 294
sinobirmanica, *Leptocimbex potanini* 30
Sinostigma 248
sinuata, Heriades 296
sinuata, Vespa 191
sinuatissimus, Symmorphus 191
sinuofasciatus, Odynerus 184
sinuatum, Anthidium 302
sinuatus, Crabro 231
sinuatus, Hylaeus, *Prosopis* 261
sinuatus, Stizus 221
Siobla 89
Siobloides 84
Sirex 112
Siricidae 112
Siricinae 112
Siricoidea 112
Siricomorpha 101
sischkai, Andrena simillima ssp. 267
sisymbrii, Andrena 271
sixii, Selandria 81
sjoeblomi, Euura, *Amauronematus* 65
sjoestedti, Prosopis 260
skorikovi, Abia 29

- skorikovi, *Myrmilla* 154
Skorikovia 155
slavonicus, *Temnothorax nylanderi* 208
slossonae, *Xylocabro* 232
slovaca, *Dasylobris italicica* f. 154
slovaca, *Myrmica* 206
slavicka, *Nomada cinnabarina* var. 316
smaragdinus, *Euchroeus*

 purpuratus var. 140

smaragdinus, *Nematus* 60
smeathmanellus, *Halictus* auct. 285
Smicromyrme 156
Smicromyrme subg. 156
Smicromyrmini 155
smithae, *Euura*, *Pachynematus* 65
smithella, *Melitta* 264
smithii, *Cerceris* 254
smithii, *Hylaeus* 260
smyrnae, *Halictus* 284
smyrnensis, *Pachycephus* 110
smyrnensis, *Pachycephus*
 smyrnensis 110
smyrnensis, *Sphecodes* 289
Snappania 124
snelleni, *Polistes* 192
Snoflaki, *Crabro* 227
sobrina, *Tenthredo* 95
socia, *Gilpinia*, *Lophyrus* 34
socia, *Tenthredo* 81
sociabilis, *Eucera* 320
sociabilis, *Odynerus* 179
socius, *Crossocerus* 227
sodalitis, *Chrysis* 140
sodalitia, *Macrophyia*
 duodecimpunctata 85
soederbomi, *Melecta* 326
soenderupi, *Fenusella* 48
sogdiana, *Andrena* 271
sogdiana, *Eucera* 321
sogdiana, *Smicromyrme* 155
sogdianoides, *Arachnospila*,
 Pompilus 170
sogdianus, *Gorytes* 222
sogdianus, *Halictus* 279
soikai, *Eumenes arbustorum* 185
soikai, *Meria* 149
soikai, *Polistes pamirensis* var. 192
solea, *Nematus* 75
soleanus, *Anoplus* 173
Solenius 234
Solenopsis 207
solida, *Chrysis* 138
solidaginis, *Nomada* 316
solidus, *Eumenes* 181
solidus, *Gonatopus*, *Allogonatopus* 121
Solierella 241
solitaria, *Tenthredo solitaria* 95
solitaris, *Cladius* 54
solitarius, *Nematus* 61
solitarius, *Pteronus* 65
solivagus, *Cemonus* 249
solovieffi, *Halictus* 286
solowiyofka, *Arge*, *Hylotoma* 25
solowiyofkae, *Ammophila* 213
solowiyofkae, *Bombus* 328
solskii, *Cerceris* 253
Solskii, *Oxybelus* 243
solskyi, *Osmia* 298
solskyi, *Stigmus* 250
solstitialis, *Bombus* 331
somalica, *Vespa orientalis* var. 194
somotorensis, *Cerceris* 255
sonani, *Vespa magnifica* 194
soonii, *Chrysis* 138
Sophropompilus 170
sorbi, *Neurotoma* 103
sorbi, *Trichiosoma* 31
Sorbiiformis 30
sordida, *Apis* 275
sordida, *Tenthredopsis*, *Tenthredo* 100
sordidula, *Crematogaster*, *Myrmica* 204
sordidus, *Sphecodes* 216
soreli, *Halictus* 281
soreensis, *Bombus*, *Apis* 328
soror, *Anthophora* 323
soror, *Arge* 24
soror, *Chrysis* 138
soror, *Selandria* 46
soror, *Tenthredo* 94
sororia, *Tenthredopsis* 100
sortitor, *Tenthredo* 97
sotschica, *Halictus lativentris* 280
soyoensis, *Tenthredo* 93
Spadicocabro 231
spadix, *Dasypoda argentata* 293
Spanolarra 236
sparsiciliata, *Andrena* 270
sparsus, *Symmorphus* 191
spathepus, *Lasius* 201
Spatulariella subg. 262
specioides, *Myrmica* 206
speciosa, *Chrysis* 138
speciosa, *Euura*, *Amauronematus* 65
speciosa, *Tarpa* 107
speciosissima, *Nomada* 314
speciosissimus, *Tachysphex* 240
speciosus, *Epeorus* 317
speciosus, *Platycampus* 70
speciosus, *Sphecodes* 224
speciosus, *Sphecodes* 224
spectabilis, *Andrena* 272
spectabilis, *Cephus* 111
spectabilis, *Colletes* 257
spectabilis, *Eucera* 321
spectabilis, *Tenthredo* 98
spectrum, *Gonatopus*, *Dryinus* 121
spectrum, *Ichneumon* 114
spectrum, *Xeris*, *Ichneumon* 114
specularis, *Cerceris* 255
speculicaudatus, *Halictus* 283
speculiferus, *Halictus minutulus* 286
speculigerus, *Halictus minutulus* 286
speculinum, *Lasioglossum*, *Halictus* 286
speculinus, *Halictus perplexans* 286
speetabilis, *Dasypoda* 293
spenceri, *Ancistrocerus antilope* 178
Sphaeropthalminae 153
Sphaeropthalmini 153
Sphecidae 212
Sphecienus 224
Spheciiformes 211
Sphecinae 214
Sphecini 215
Sphecius 224
Sphecodes 287
sphecodimorpha, *Andrena* 273
Sphecodina 277
sphecodoides, *Tachysphex* 239
sphecoidea, *Melitta* 288
sphegoidea, *Apis* 268
Sphex 215
sphinx, *Vespa crabro* 193
Spilomena 249
Spilopompilus 166
spina, *Elampus*, *Hedychrum* 128
Spinarge 25
spinarum, *Tenthredo* 43
spinarum, *Tenthredo* 44
spineventris, *Osmia* 298
spiniceps, *Crabro* 228
spinicollis, *Crabro* 232
spinicollis, *Diodontus* 247
spinicollis, *Lestica* 234
Spinicoxa subg. 186
spinifera, *Jermakia* 92
spinigera, *Dasypoda* 293
spinigera, *Dioxys* 303
spinigera, *Melitta* 267
spinipectus, *Cerceris* 255
spinipectus, *Crabro* 230
spinipes, *Cephus*, *Banchus* 109
spinipes, *Crabro* 233
spinipes, *Ectemnius*, *Crabro* 233
spinipes, *Odynerus*, *Vespa* 187
spinipes, *Vespa* 186
spiniserra, *Amauronematus* 62
spinolae, *Dryudella* 219
spinolae, *Monophadnus*, *Tenthredo* 51
spinolae, *Nitela* 241
spinolae, *Osmia* 297
Spinolia 141
spinolosa, *Alysson*, *Pompilus* 219
spinous, *Celonites* 178
spinous, *Ceropales* 220
spinous, *Crabro* 220
spinous, *Nysson*, *Sphex* 220
spinous, *Pompilus* 219
spinous, *Sphex* 220
Spintharina 141
spinulosa, *Osmia*, *Apis* 298
spinulosus, *Sphecodes* 290
spiraeeae, *Blennocampa* 46
spiraeeae, *Euura*, *Nematus* 65
spiraeeae, *Megalodontes*, *Tarpa* 107
spiralis, *Andrena* 291
spiricornis, *Odynerus* 187
spirifex, *Sceliphron*, *Sphex* 214
spirifex, *Sphex* 214
spissa, *Arachnospila*, *Pompilus* 170

- spissicauda*, Coelioxys 305
spissicornis, Tarpa 107
splendens, Andrena 269
splendida, Aphaenogaster, *Atta* 203
splendida, Apis 264
splendidula, Chrysis 138
splendidus, Cleptes, *Ichneumon* 127
spoliata, Tachytes 238
sponsa, Coelioxys 305
spooneri, Mimumesa, *Mimesa* 251
sporadensis, Eumenes
 unguiculus var. 181
sporadicus, Bombus 328
spreta, Tenthredo 100
spretus, Halictus 285
springatei, Claremontia 46
spurcus, Amauronematus 57
spuria, Larra 241
spuria, Tenthredo 41
spurius, Miscophus, *Larra* 241
squalens, Anthophora
 personata var. 323
squalida, Anthophora 324
squalidus, Nematus 60
squamea, Andrena 273
squamigera, Andrena 264, 268
squamosus, Oxybelus 242
squamotus, Amauronematus 64
stachi, Tachysphex 240
stackelbergi, Philoctetes, *Ellampus* 132
stambuloffii, Cardiocondyla 203
stancici, Apterogyna 159
stangeana, Myrmica 206
stangeana, Myrmica bergi 206
staudingeri, Pristiphora, *Nematus* 74
Stauronema 75
Stauronematus 75
stchurovskyi, Polyodontus 141
stchurovskyi, Spinolia, *Polyodontus* 141
stecki, Astata 218
stecki, Cerceris 253
stecki, Lygaeonematus 73
steckianus, Odynerus
 dantisquama var. 190
steckianus, Stenodynerus,
 Odynerus 190
steckii, Allantus 90
stefaniana, Andrena
 chrysopyga var. 275
stefanii, Andrena 274
steini, Dolerus 77
steini, Emphytus succinctus 38
steini, Nematinus 69
Stelidina 302
Stelidium 302
Stelidomorpha subg. 302
Stelis 302
Stelis subg. 302
stella, Omalus, *Ellampus* 132
stellata, Tenthredo 87, 102
Stenagenia 165
Stenamma 207
Stenancistrocerus 189
Stenancistrocerus subg. 189
Stenanthidiellum subg. 301
Stenanthidium 300
Stenemphytus 43
Stenocephus 111
Stenoclavelia 161
Stenocabro 227
Stenodynerus 189
stenogaster, Euura, *Nematus* 65
Stenoglossa 188
Stenomelissa subg. 274
Stenomelinus 251
Stenomesoneura 68
Stenopolistes 192
stentoriscapus, Hylaeus 260
stenura, Andrena labialis ssp. 267
stephensi, Croesus 69
stepposa, Andrena 267
stepposus, Rophites clypealis 291
stepposus, Smicromyrme 156
Stercobata 252
Sterictiphora 26
Sterictiphorinae 25
Sterictophorinae 22
sternalis, Amauronematus vittatus 67
sternalis, Athalia rosae 44
Stethomostus 53
Stevenia 53
stevensi, Prosopis 260
stichi, Euura, *Pteronidea* 65
sticticus, Taxonus, *Tenthredo* 43
Stictoptila 227
stigma, Apis 303
Stigma, Crabro 229
stigma, Dryudella, *Dimorpha*
 stigma 218
stigma, Hemichroa 67
stigma, Lyda 104
stigma, Nomada 316
stigma, Nomada auct. 314
stigma, Tenthredo, *Tenthredella* 95
stigma, Tenthredopsis, *Tenthredo* 100
Stigmaporus 173
stigmatica, Andrena 269
stigmatica, Arge ustulata 25
stigmatica, Coelioxys 305
stigmatica, Pristiphora pallidiventris 73
stigmatica, Tenthredo vespa 97
stigmatical, Anisoneura 40
Stigmatodipogon 164
Stigmatosona 83
Stigmus 249
stilata, Dineura, *Tenthredo* 55
Stilbum 141
stipeus, Leptocephalothorax tuberum 208
stitia, Ametastegia equiseti 39
Stizoides 225
stizoides, Vespa germanica var. 196
Stizolarra 225
Stizomorphus 221
Stizus 225
stoeckhertella, Andrena 271
stoeckherti, Nomada 316
straminea, Pteronidea 62
stramineipes, Pamphilus, *Lyda* 105
stramineipes, Tenthredo 76
strandii, Lygaeonematus 58
strandii, Pison 244
stratiotes, Cerceris 255
Streptosella 170
stria, Tenthredo vestita 92
striata, Ammophila 213
striata, Nomada 316
striata, Tenthredo vestita 92
striatipleuris, Cleptes 127
striatula, Myrmica 205
striatum, Stenamma 207
striatum, Stenamma
 westwoodi var. 207
striatus, Blepharipus 233
striatus, Crabro 230
striatus, Crossocerus 230
striatus, Ceratocolus 233
striatus, Crabro 230, 233
striatus, Crossocerus 230
striatus, Gonatopus 121
striatus, Halictus 285
stribligata, Mutilla 157
stridula, Tiphiidae 155
strigata, Stelis 302
strigata, Tenthredo vestita 92
strigata, Tetraloniella, *Macrocerus* 322
strigata, Trachusa 300
strigatum, Anthidiellum, *Trachusa* 300
strigatus, Cemonus 248
strigosa, Andrena 267
strigosa, Tenthredo 86
strigosus, Cryptocheilus, *Pompilus* 161
strigosus, Tachytes 238
strigulosus, Hylaeus 260
strigulosus, Palmodes, *Sphex* 215
striola, Aphaenogaster 203
striolata, Cerceris 253
stroganova, Aprosthemma 26
stroganova, Arge 25
Strombocerina 82
Stromboceros 82
Strongylogaster 82
strongylogaster, Nematus 59
Strongylogastroidea 43
Strongylognathus 207
structor, Messor, *Formica* 204
Strumigenys 207
struthiocamelus, Sirex 113
struthiopteridis, Sahlbergia 83
struthiopteridis, Strongylogaster,
 Thrinax 82
strymonia, Andrena 263
stubbei, Hylaeus 260
stubbei, Tenthredo 94
stulta, Tenthredo 92
sturmii, Siobla, *Tenthredo sturmii* 89
stygius, Dolerus 80
stygius, Thyreopus 228
Stylopoza 81
styriacus, Hylaeus 261

styrius, Crossocerus, *Crabro* 228
 styx, Euura, *Pachynematus* 65
suadus, Nematus 58
suarezi, Pterocheilus phaleratus 189
suavis, Nematus 56
suavis, Sphecodes 215
subaenescens, Lasioglossum, *Halictus*
 subaenescens 286
subaeneus, Lindenius 235
subalatus, Dolerus 80
subalkenellus, Halictus 278
subalpina, Smicromyrme 155
subapicalis, Larra 225
subarctica, Pristiphora,
 Lygaeonematus 74
subarctica, Rhadinoceraea 52
subarcticus, Dolerus 78
subauratus, Ellampus
 schulthessi var. 132
subauratus, Halictus, *Apis*
 subauratus 279
subbaicalensis, Bombus auct. 331
subbaicalensis, Bombus helleranus 331
subbalteatus, Alpinobombus
 kirbyellus 327
subbifida, Pristiphora, *Nematus* 74
subboreale, Tapinoma 198
subcana, Selandria 46
subcarpathica, Tenthredopsis
 litterata 99
subcollaris, Alpinobombus
 kirbyellus 327
subcollaris, Bombus ignitus 328
subcomata, Mutilla 155
subcoriacea, Chrysis 138
subcornuta, Apis 313
subcornuta, Mutilla 154
subcostatus, Allantus 90
subcyaneum, Pseudisobrachium,
 Epyris 123
subcylindricus, Cephus 109
subdentata, Crematogaster 204
subdentata, Melitta 264
subdentatus, Tachysphex 240
subeburnea, Bembix 221
suberiophoroides, Bombus czerskii 328
subexcisus, Hylaeus 259
subfasciata, Andrena 272
subfasciata, Prosopis 261
subfasciatus, Dolerus 79
subfasciatus, Halictus 283
subflavicornis, Anteon 119
subfulvicorne, Lasioglossum, *Halictus*
 subfulvicorne 286
subfuscata, Tenthredo 78
subfuscatus, Prionyx, *Sphex* 215
subgalerita, Nomioides 292
subgemma, Euura 65
subgibbosa, Cerceris 254
subglaber, Evagetes, *Psammochara* 171
subglobosa, Apis 323
subglobosus, Campylogaster 275
subincana, Melitta 273

subintermedia, Deuteragenia,
 Pogonius 164
subinterruptus, Camptopoeum
 friesei var. 276
subinterruptus, Crabro 233
subita, Lyroda 240
subjecta, Tenthredo 43
sublaevis, Harpagoxenus, *Myrmica* 204
sublaevis, Lithurgus 295
sublaevis, Myrmica 204
sublaevis, Prosopis 261
sublineare, Lasioglossum 284
sublongula, Chrysis longula 136
submarginatus, Hylaeus 259
submediterraneus, Halictus,
 Seladonia 279
submicans, Osmia 299
submimetica, Paravespa rex var. 188
submonochromos, Hortobombus
 consobrinus 328
submutica, Poecilosoma 41
subnitens, Colletes 259
subnitens, Pteronidea 63
subnudus, Leptothonax
 nassonovi var. 208
subnudus, Psammochara 171
subnudivaca, Tenthredo, *Tenthredella* 92
subopaca, Abia candens 28
subopaca, Andrena 271
subopaca, Pristiphora 74
subopaca, Prosopis 261
subovalis, Sphecodes 290
subpilosa, Formica 201
subpilosa, Formica rufibarbis 201
subpilosa, Formica ruzskyi 200
subpomiformis, Eumenes 182
subpunctatus, Crabro 230
subpunctatus, Hylaeus 259
subpunctulatus, Sphecodes 289
subquadratus, Hylaeus 262
subquadratus, Sphecodes 289
subrenimacula, Ancistrocerus
 claripennis 179
subrufa, Cephus 111
subserrata, Blennocampa 46
subsolanus, Dolerus germanicus 78
subspinous, Anthidium 300
subspinous, Oxybelus 243
subsquamarialis, Andrena 264
subterranea, Aphaenogaster,
 Formica 203
subterranea, Lestica, *Crabro*
 subterranea 234
Subterraneobombus subg. 330
subterraneoides, Aphaenogaster 203
subterraneoides, Aphaenogaster
 pallida var. 203
subterraneus, Bombus, *Apis* 331
subterraneus, Crabro 234
subtilis, Crabro 232
subtilis, Hylaeus 259
subtrifida, Euxiphidria 115
subtruncata, Eriocampsapis 42
subulata, Pleroneura 20
subulata, Tenthredo 99
subulatus, Crossocerus, *Crabro* 230
subvinogradovi, Hortobombus
 consobrinus 328
subvittata, Arachnospila, *Pompilus* 169
succinata, Andrena 267
succincta, Apis 257
succincta, Chrysis 138
succincta, Nomada 316
succincta, Tenthredo 38, 92, 96
succinctula, Andrena 267
succinctus, Colletes, *Apis* 259
suchovi, Haplolygonotopus 121
sucineiceps, Paururus 112
sudaghensis, Halictus 286
sudai, Crossocerus verhoeffi 228
sudai, Nomada 311
sudai, Sphecodes 288
suecica, Andrena 268
suecica, Colletes 258
suecica, Formica 199
suecicum, punctatum var. 301
suecicus, Odynerus 191
suerinensis, Nomada
 alboguttata var. 311
suessionensis, Athalia 44
suessionensis, Nematus 74
suessionensis, Athalia 44
suffusa, Lyda 104
sugare, Dolichovespula media 195
sugiharae, Crabro 228
sugonjaevi, Priocnemis 163
suigenensis, Phymatoceriola 52
sulcata, Xiphydria 116
sulcatulum, Lasioglossum, *Halictus*
 sulcatulum 286
sulcatus, Dolerus leucopterus 80
sulcatus, Pachynematus 59
sulcifer, Polistes 193
sulcifrons, Dineura 55
sulcifrons, Gorytes sulcifrons 223
sulcifrons, Hoplisus 223
sulinodis, Myrmica 206
sulinodoscabrinodis, Myrmica
 sulinodis var. 206
sulcipes, Nematus 57
Sulcomesitius 124
Sulcopolistes 192
sulcus, Crabro 229
sulfuripes, Lionotus 183
sulphurata, Tenthredo 86
sulphureipes, Pamphilus
 sulphureipes 105
sulphureus, Nematus 63
sulphuripes, Tenthredo, *Allantus*
 sulphuripes 97
sumizome, Andrena lapponica ssp. 264
sumptuosa, Vespa silvestris var. 195
sundukovi, Anoplus 172
sungora, Dasylabris, *Mutilla maura* 153
sunna, Andrena alfkennella ssp. 270
superba, Chrysis 135

superba, *Macrophyia* 86
 superba, *Mimesa* 251
 superba, *Nomada* 314
 superba, *Pseudepipona, Odynerus* 188
 superbus, *Apis* 298
 superbus, *Labus* 182
 superbus, *Nematus* 60
 superbus, *Omalus* 127
 superbus, *Stenemphytus* 43
 superequester, *Agrobombus* 331
 superus, *Leptothorax*

 suprafulgens, *Halictus* 283
 suprunenkoi, *Vespa* 193
 surensis, *Allantus* 94
 suripes, *Dasypoda, Apis* 293
Susanowo 248
susiana, *Myrmilla* 154
spectus, *Hylaeus* 259
suspicax, *Arge* 25
suspicax, *Pison* 244
susterai, *Andrena* 274
susterai, *Nysson* 220
susterai, *Priocnemis* 164
susterai, *Smicromyrme montana* f. 155
sutor, *Sphecodes* 288
sutshanicum, *Lasioglossum* 281
sutshanicus, *Crabro* 228
suwonae, *Xyela* 21
suzukii, *Cimbex* 29
suzukii, *Crabro monstrosus* 230
suzukii, *Crossocerus, Crabro* 230
sveae, *Pteronidea* 63
swammerdamella, *Melitta* 293
swanetiae, *Pseudomeria* 149
syam, *Tzustigmus* 250
sybarita, *Nomada* 316
Sylaon 241
sylvarum, *Bombus, Apis* 332
sylvarum, *Tenthredo* 29
sylvatica, *Cimbex* 32
sylvatica, *Tenthredo* 104
sylvaticum, *Trichiosoma* 31
sylvaticus, *Pamphilus, Tenthredo* 105
sylvestris, *Bombus, Psithyrus* 329
sylvestris, *Dolichovespula, Vespa* 195
sylvestris, *Euura, Nematus* 65
sylvia, *Tenthredo* 95
symbalophthalma, *Abia* 28
Symmorphpha 325
Symmorphus 190
Symmorphus subg. 190
Symphta 20
sympyti, *Andrena* 267
Synaerema 88
Synairema 88
Synchalonia subg. 321
Synemphytus 37
Syneuodynerus 191
syngenesiae, *Bethylus* 125
Synhalonia subg. 321
Synneurus 220
Synorhopalum 227

Synothyreopus 226
syriaca, *Hylotoma* 23
syriaca, *Nomada* 314
syriacola, *Scolia fallax* 148
syriacus, *Halictus* 285
syriacus, *Trachelus* 108
syriensis, *Chrysis daphnis* 135
syringae, *Cephalosoma* 111
Syrista 111
syrius, *Halictus subauratus* 279
syrmiente, *Aprosthemma, Schizocera* 26
Systropha 291
szaboi, *Hedychrum* 130
szelenyi, *Metrionotus* 124
szepligetii, *Dryinus* 120
tabida, *Eumenes* 184
tabidus, *Sirex* 111
tabidus, *Trachelus, Sirex* 111
Tachyagetes 173
Tachyagetes subg. 174
Tachybulus 244
Tachynana 240
Tachyoides 240
Tachyplena 240
Tachyptera 240
Tachysphex 237
Tachystizus 225
Tachytes 240
tacitus, *Halictus* 281
taczanowskii, *Chrysis* 138
taczanowskii, *Praia* 30
tadzhicus, *Halictus* 281
tadzhicus, *Hoplomerus*
 melanocephalus 187
tadzhicorum, *Eumenes* 181
tadzhika, *Cerceris* 255
tadzhika, *Chrysis* 140
tadzhika, *Pavlovskia* 241
tadzhikus, *Tachysphex* 238
taediosa, *Myrmica* 206
Taeniandrena subg. 274
Taeniaporus 174
taeniata, *Nomada fucata* var. 313
taeniatus, *Dolerus* 77
taeniatus, *Harpiphorus* 43
taeniatus, *Nematus* 65
tahitense, *Rhynchium* 189
Taialia 249
taicho, *Sphecodes* 290
taiganus, *Nematus* 65
taiganus, *Urocerus gigas* 113
Taimyrmosa 153
taishanensis, *Melitta* 294
Taiwagenia 165
taiwanensis, *Coelioxys* 304
takachihoi, *Auplopus, Pseudagenia* 165
takagii, *Pristiphora* 72
takanensis, *Psen* 251
takedae, *Tenthredo olivacea* 92
takenoi, *Anteon* 119
takeuchii, *Cladius* 55
takeuchii, *Crossocerus* 228
takeuchii, *Gorytes* 223
takeuchii, *Pamphilus* 105
takeuchii, *Siobla* 89
takeuchii, *Thrinax, Hemitaxonius* 83
talyshensis, *Tenthredo* 92
tamanukii, *Prosopis* 259
tamanukii, *Tenthredo, Tenthredella* 92
tamara, *Chrysis* 138
tamarae, *Formica* 199
Tanaemyrmex subg. 199
tanaiticus, *Cephus* 109
tanakai, *Crossocerus* 228
tanakai, *Psenulus* 252
tancrei, *Myrmecocystus cursor* r. 199
tanoi, *Sphecodes* 290
tantillus, *Aphadnurus* 48
taorminae, *Andrena nana* var. 270
taorminicus, *Halictus patellatus* 278
Tapinoma 197
Taranga 244
taraxaci, *Andrena* 265
tardigradus, *Sirex* 113
tardum, *Aprosthemma, Hylotoma* 26
tardus, *Epyris* 123
Tarpa 107
tarraconensis, *Dryinus* 120
tarsalis, *Crabro* 228
tarsalis, *Epeolus* 317
Tarsandrena subg. 274
tarsata, *Andrena* 274
tarsata, *Tenthredo* 40
tarsata, *Tenthredopsis, Tenthredo* 100
tarsata, *Tetralonia* 320
tarsata, *Tetralonia* 325
tarsatum, *Lasioglossum, Halictus* 286
tarsatus, *Crossocerus, Crabro* 229
tarsina, *Tachytes* 240
tarsinus, *Tachysphex, Tachytes* 240
tartara, *Colpa, Elis* 147
tartarea, *Vespa crabro* var. 194
tataricus, *Halictus* 279
tataricus, *Messor* 204
tataricus, *Nematus* 69
tatjanae, *Andrena* 273
tauushi, *Cimbex connatus* 29
tauksiensis, *Cimbex* 29
tauriae, *Eumenes sichelii* 185
taurica, *Andrena gallica* ssp. 269
taurica, *Anthophora* 324
taurica, *Apterogyna* 159
taurica, *Cardiocondyla stambuloffii* 203
taurica, *Chrysis* 137
taurica, *Eucera* 321
taurica, *Hoplitis, Pseudosmia* 297
taurica, *Mutilla* 157
taurica, *Mutilla* 158
taurica, *Plagiolepis* 202
taurica, *Plagiolepis maura* var. 202
taurica, *Ponera* 209
tauricola, *Protosmia* 300
tauricum, *Aprosthemma* 26
tauricum, *Lasioglossum* 286
tauricus, *Agonatopoides* 121

tauricus, Celonites 178
tauricus, Celonites abbreviatus 178
tauricus, Chalepoxenus 204
tauricus, Colletes cunicularius 257
tauricus, Dolerus melanopterus 80
tauricus, Echthrodelpachus 120
tauricus, Elampus, *Notozus* 128
tauricus, Embolemus 122
tauricus, Harpactus 224
tauricus, Heterarthrus 49
tauricus, Katamenes, *Eumenes* 185
tauricus, Leptothonax
 unifasciatus var. 208
tauricus, Odynerus signaticollis 187
tauricus, Temnothorax, *Leptothonax* 208
tauridis, Tapinoma 198
tauriensis, Andrena hybrida ssp. 272
taurocaucasicum, Tetramorium 209
taurus, Nannodynerus
 aequisculptus 189
taurus, Osmia 299
tavastiensis, Nematus wahlbergi 69
Taxoblenus 43
Taxonus 43
Tea 173
teberdicus, Eutomostethus 47
tectus, Dolerus 79
tectus, Halictus 279
tegulare, Anthidium 302
regularis, Eustenancistrocerus,
Lionotus 184
regularis, Loderus genucinctus 77
tekkenensis, Ammophila 213
Teleutomyrmex 208
telmenicus, Hylaeus 260
Telostegus 174
tembotovi, Temnothorax 208
temesiensis, Emplytus 40
Temnothorax 208
temporalis, Aneugmenus, *Selandria* 76
temporalis, Coelioxys 305
temporalis, Euura, *Amauronematus* 65
temporalis, Tenthredella flavicornis 91
temporalis, Tenthredella stulta 92
temula, Tenthredo 93
Temuleo 90
Temuleo subg. 93
temuloides, Allantus 93
tenax, Pemphredon 248
tenebris, Eumenes
 lunulatus var. 181
tenebrosa, Andrena vitrea var. 272
tenebrosa, Pachyprotasis variegata 88
tenebrosa, Vespa analis var. 193
tenebrosus, Dolerus 79
tenella, Apis 267
tenella, Chrysis 138
tenella, Nomada 316
tenella, Parna, *Tenthredo* 51
tenella, Phyllotoma 49
tenella, Tenthredo 51
tenella, Tetralonia 322

tenellum, Pseudoanthidium,
Anthidium 302
tenellus, Hylaeus 284
tener, Cladius 54
tener, Nematus 57
tenera, Ametastegia, *Tenthredo* 39
tenera, Pemphredon 248
teneriffensis, Anthophora
 quadrifasciata var. 322
Tenila 241
tenius, Tenthredo 38
Tenthredella 89
Tenthredella subg. 93
Tenthredina 89
Tenthredina subg. 96
Tenthredinidae 37
Tenthredininae 83
Tenthredinoidea 22
Tenthredinomorpha 22
Tenthredo 89
Tenthredopsis 98
tenuicingulatus, Monophadnus 51
tenuicornis, Ammobates oraniensis 318
tenuicornis, Anoplus, *Pompilus* 172
tenuicornis, *Astatus* 111
tenuicornis, Claremontia, *Tenthredo* 46
tenuicornis, Entodecta 50
tenuicornis, Larrada 236
tenuicornis, Lygaeonematus 73
tenuicula, Andrena 263
tenuipennis, Tenthredo 98
tenuipunctus, Tachysphex 239
tenuis, Andrena 263
tenuis, Pachyprotasis 88
tenuis, Passaloecus 248
tenuis, Sphecodes 289
tenuiserra, Euura, *Amauronematus* 65
tenuiserra, Pelmatopus 55
tenuiserra, Pristiphora,
Micronematus 74
tenuivaginata, Tenthredo,
Tenthredella 92
tenuivittata, Cerceris 255
teranishii, Lasius 201
terebrator, Halictus 282
tergale, Icteranthidium 301
tergestensis, Andrena
 nycthemera var. 264
tergestensis, Osmia 299
tergestina, Blennocampa 46
Tergosmia subg. 299
terminale, Anthidium 300
terminalis, Amauronematus 60
terminalis, Anthophora furcata 323
terminalis, Eucera 320
terminalis, Stizus 225
terminata, Ammophila 213
terminata, Chrysis 138
terminater, Dolerus pratensis 78
terrestris, Apis 327
terrestris, Bombus, *Apis* 328
terrificola, Odynerus 184
terrificus, Leptocimbex 30

tersa, Megilla 325
tertium, Aglaostigma 84
tertius, Pseudotaxonus 82
tessellata, Tenthredo 98
tessellata, Tenthredopsis, *Tenthredo* 100
tessinensis, Pteronidea segmentaria 60
testacea, Andrena gwynana var. 267
testacea, Cephaleia 102
testacea, Cimbex lutea 30
testacea, Cimbex saliceti 30
testacea, Cryptopone 209
testacea, Ponera 210
testacea, Ponera coarctata var. 210
testacea, Pristiphora, *Pteronus* 74
testacea, Tenthredo 51
testaceicornis, Ellampus 132
testaceicornis, Nematus 74
testaceicornis, Pristiphora 74
testaceipes, Anthophora, *Podalirius* 325
testaceipes, Dineura, *Tenthredo* 55
testaceipes, Empria, *Poecilosoma* 41
testaceipes, Eriocampa 46
testaceipes, Euura, *Cryptocampus* 65
testaceipes, Sierola 125
testaceipes, Sphecodes gibbus
 ephippium 289
testaceum, Eciton 207
testaceus, Nematus 61, 65
testaceus, Pteronus 70
testaceus, Strongylognathus, *Eciton* 207
testaceus, Tomostethus 68
testudinea, Hoplocampa, *Tenthredo* 68
Tetrachrysis 133
Tetradontochelys 120
tetraedrus, Crabro 231
tetraedrus, Solenius 233
Tetralonia 321
Tetralonia, part. 320
Tetraloniella 321
Tetramorium 209
Tetranthidium 300
Tetratneura 40
tetazonianellus, Halictus 278
tetazonius, Halictus, *Hylaeus* 278
tetrica, Pristiphora, *Nematus* 74
tetrodonta, Osmia 299
Tetradontochelys 120
teutona, Macrophyia, *Tenthredo* 86
teutonicus, Odynerus 190
teutonus, Pompilus 236
thalenhorsti, Pristiphora 74
thalhammeri, Chrysis distincta 135
thalictri, Diphadnus 74
thalictri, Pristiphora, *Nematus* 74
thalictricola, Pristiphora 74
thalictrivora, Pristiphora 74
Thao 227
thargitai, Dolerus 78
thauma, Ampulicomorpha 122
Theleter 187
theresae, Dolerus aericeps 77
theresae, Macrophyia annulata 84
theresae, Sapyga 145

thersites, *Nomada* 316
theryi, *Megachile centuncularis* 306
thomsoni, *Ancistrocerus* 179
thomsoni, *Andrena* 273
thomsoni, *Colletes* 258
thomsoni, *Dasypoda* 293
thomsoni, *Dolerus* 77
thomsoni, *Gonatopus* 121
thomsoni, *Hylotoma* 24
thomsoni, *Lophyrus* 34
thomsoni, *Nematus* 57
thomsoni, *Pamphilius* 103
thomsoni, *Pristiphora* 71
thomsoni, *Profenusia, Fenusia* 52
thomsoni, *Thomsonia* 99
Thomsonia 98
thor, *Megalodontes* 107
thoracica, *Andrena, Apis* 269
thoracica, *Cilissa* 275
thoracica, *Dasypoda* 293
thoracica, *Tenthredo* 79, 99
thoracica, *Tenthredo opaca* 79
Thoracobombus subg. 331
thornleyi, *Tenthredopsis* 100
thracia, *Megachile sexmaculata* 306
Thrinax 83
thuringiaca, *Melitturga*
 clavicornis var. 276
thuringiacus, *Odynerus* 179
Thyreocerus 231
Thyreocnemus 225
Thyreomelecta 326
Thyreopus 225
Thyreosphe 211
Thyreus 234
Thyreus 326
Thysandrena subg. 274
Tianmuthredo 100
tianschanicus, *Odynerus tristis* 187
tianshanica, *Chrysis* 138
tibetanus, *Polistes rothneyi* 192
tibialatus, *Cladius* 54
tibiale, *Trichiosoma* 31
tibiale, *Trypoxylon* 244
tibialis, *Andrena, Melitta* 272
tibialis, *Astatas* 218
tibialis, *Athalia* 44
tibialis, *Cephus brachycercus* 109
tibialis, *Crabro* 227, 231, 233, 235
tibialis, *Euura, Nematus* 65
tibialis, *Pterocheilus phaleratus* 189
tibialis, *Selandria* 46
tibialis, *Tenthredo* 40
tibialis, *Vespa* 227
tichenkoi, *Bombus, Hortobombus* 328
tichomirovi, *Camponotus* 199
ticinensis, *Phoenusa* 48
tiflensis, *Protosmia, Osmia* 300
tigris, *Trachelus, Neateuchopus* 111
tiliae, *Allantus* 99
tiliae, *Blennocampa* 51
tiliae, *Selandria* 50
tilingi, *Bombus hypnorum* 330

Tilkuya 149
tillbergi, *Euura, Amauronematus* 65
timanensis, *Katamenes sichelii* 185
timida, *Macrophya* 87
timida, *Tenthredo* 78
timidus, *Microdynerus, Odynerus* 186
timidus, *Oxybelus* 243
tinctipennis, *Dolerus* 77
tinctipennis, *Pempredon* 249
tinctipennis, *Trichiosoma jakovlevi* 30
tinctus, *Odynerus* 183
tingitana, *Arachnospila* 170
tingitanus, *Dicyrtomellus*,
 Arachnospila 170
tinitinensis, *Halictus* 285
tinnunculus, *Halictus* 281
Tiphia 150
Tiphia subg. 150
Tiphidae 149
tiphiiformis, *Platyxiphidria* 115
Tiphinae 150
Tirolensis, *Crossocerus* 227
tirolensis, *Empria* 41
tirolensis, *Epeolus tarsalis* 317
tirolensis, *Hoplisus* 223
tirolensis, *Tenthredopsis dorsalis* 99
tischbeini, *Crabro* 228
tischbeini, *Nematus* 70
tischbeinii, *Tenthredopsis*,
 Tenthredo 100
tisiphone, *Melecta* 326
titania, *Tenthredo notha* 97
t-magnum, *Pteronidea*
 pseudonotabilis 57
tobiasi, *Arachnospila* 169
tobiasi, *Hedychrum* 130
toenita, *Euura, Nematus* 65
togashii, *Aglaostigma* 84
togata, *Mutilla* 155
togata, *Tenthredo* 37
togatus, *Allantus, Tenthredo* 38
tohi, *Gilpinia* 34
tokyanus, *Stenodynerus, Odynerus* 190
tolli, *Amauronematus* 55
tomentosa, *Anthophora* 324
tomentosa, *Eucera* 320
tomentosus, *Hylaeus* 278
tomentosus, *Odynerus* 183
Tomognathus 204
tomostethi, *Eutomostethus, Forsia* 47
Tomostethopsis 47
Tomostethus 53
Toncahua 234
tonkinensis, *Euodynerus notatus* 183
tonnaitchana, *Cimbex* 29
tonsus, *Gorytes* 221
Topcobius 124
torkai, *Andrena* 264
tormentillae, *Nomada roberjeotiana* 315
tormentillae, *Phyllotoma* 48
tornense, *Proanthidium* 301
toroki, *Dasypoda* 293
torosa, *Ronisia* 157
tortumensis, *Megachile genalis* 306
totus, *Dolerus pratensis* 78
tournieri, *Elampus* 127
tournieri, *Tiphia* 150
Towada 227
trabeata, *Tenthredo* 92
Trachandrena subg. 275
Trachelastatus 111
tracheliformis, *Tsunekiola* 236
Tracheliodes 236
Trachelosimus 234
Trachelus 111
Trachogorytes 246
Trachusa 303
trachyderma, *Andrena* 267
Trachyocheilus 161
tragica, *Chrysis* 138
tranquilla, *Apis* 275
tranquilla, *Euura* 65
transbaicalica, *Andrena* 272
transbaicalicus, *Eumenes* 182
transbaicalicus, *Evagetes* 171
transbaicalicus, *Harpactus* 224
transbaikalensis, *Halictus* 279
transcaspica, *Andrena* 273
transcaspica, *Crocisa* 326
transcaspica, *Osmia* 297
transcaspicum, *Anthidium* 301
transcaspicus, *Miscophus* 241
transcaspicus, *Nysson* 219
transcaspiensis, *Andrena*
 ovatula ssp. 274
transcaucasica, *Aphaenogaster*
 splendida 203
transcaucasica, *Formica* 201
transcaucasicus, *Cryptocheilus* 161
transfuga, *Apis* 273
transiens, *Crabro* 233
transiens, *Harpactus* 224
transigens, *Pachynematus clitellatus* 57
transigens, *Tenthredo atra* 93
transiliensis, *Harpactus* 224
transitoria, *Andrena* 274
transitoria, *Megachile* 305
transitoria, *Nomada* 312
transitorius, *Epeolus* 317
transitorius, *Odynerus* 184
transpositum, *Lasioglossum*,
 Halictus 286
transsibirica, *Myrmica* 206
transsylvatica, *Nomada* 312
transversalis, *Crabro* 229
transversalis, *Euura, Pteronidea* 65
transversalis, *Hylaeus* 261
transversalis, *Prosopis* 261
transvolgensis, *Halictus compressus* 278
trapezicolle, *Hedychrum* 130
trapeziformis, *Nomada* 316
trautmanni, *Euura, Amauronematus* 65
Trearea 83
Tremex 113
Tremicinae 113
tremula, *Tenthredo* 77

- tremulae, Caliroa 46
trialbata, Tenthredo 94
triandrae, Euura, *Pontania* 65
triangularis, Smicromyrme, *Mutilla* 156
triangulifer, Priophorus 54
triangulifer, Pseudomalus, *Omalus* 133
triangulomaculata, Dasylabris
italica var. 154
triangulum, Philanthus 217
triangulum, Philanthus, *Vespa* 255
triangulus, Hylaeus 259
triareolatus, Perisemus 125
tricarinata, Pseudepipona,
Odynerus 188
tricarinatus, Odynerus herrichii var. 188
Trichelobrachium 123
Trichepipona 188
Trichiocampus 53
Trichiocampus subg. 54
Trichiosoma 30
trichiosoma, Cimbex 31
trichocera, Tenthredo 52
trichogastor, Pemphredon 248
Trichogonatopus 120
Trichomyrmex 209
trichopsis, Halictus 287
trichorhinum, Lasioglossum,
Halictus 286
Trichosyron 170
trichroma, Tenthredopsis nassata 99
Trichrysis 141
triciliatus, Bombus tricornis 332
tricincta, Andrena 275
tricincta, Dasylabris maura
nostras var. 154
tricincta, Eucera 321
tricincta, Macrocerca 321
tricincta, Melitta 294
tricincta, Tenthredo 86
tricinctum, Lasioglossum, *Halictus*
tricinctum 286
tricinctus, Mellinus 220
tricinctus, Odynerus 180
tricingulum, Lasioglossum 279
tricingulum, Lasioglossum 281
tricolor, Aglaostigma, *Macrophyopsis* 83
tricolor, Alysson 219
tricolor, Arpactus 223
tricolor, Astata 218
tricolor, Belomicrus 242
tricolor, Cuneala 90
tricolor, Dryudella, *Astata* tricolor 219
tricolor, Eutomostethus, *Tomostethus* 47
tricolor, Lonchodryinus 119
tricolor, Miscophus 241
tricolor, Pamphilus 105
tricolor, Pompilus 240
tricolor, Pseudepipona 188
tricolor, Sphecodes 145
tricolor, Sphecodes 240
tricolor, Tenthredo 24
tricolor, Tropidodynerus interruptus 192
tricolor, Vespa 148
tricolorata, Andrena tibialis var. 272
tricoloratus, Sphecodes 240
tricoloripes, Emphytus 38
tricornis, Bombus 332
tricornis, Empria 41
tricuspidata, Andrena 265
tricuspidata, Coelioxys 305
tricuspis, Hylaeus 262
tridens, Bembecinus, *Vespa* tridens 221
tridens, Crabro 243
tridens, Empria, Poecilosoma 42
tridens, Nysson 220
tridens, Vespa 195
tridentata, Aglaopis, *Coelioxys* 303
tridentata, Andrena, *Melitta* 267
tridentata, Hoplitis, *Osmia* 297
tridentata, Nomada 312
tridentata, Parapiageta 237
tridentata, Vespa 193
tridentatus, Amauronematus 65
tridentatus, Microdynerus 186
tridentatus, Stizoides, *Crabro*
tridentatus 225
tridenticulata, Coelioxys 305
tridentirostris, Nomada 316
tridentis, Empria 42
Tridryinus 120
Trielis 147
Triempria 40
Triepeolus 317
trifasciata, Conaspidea 84
trifasciata, Mutilla 157
trifasciata, Tenthredo 86
trifasciata, Vespa 179
trifasciatus, Ancistrocerus, *Vespa* 179
trifasciatus, Hylaeus 281
triforis, Tenthredopsis 98
trifurcatus, Nematus 65
trigaria, Onycholyda, *Pamphilus* 103
trigeminus, Dolerus 79
Trigonotopus 120
trigonicus, Amauronematus 56
triuinterrupta, Sapyga 145
trilineata, Vespa 220
trilineatus, Nematus 65
trimaculata, Bembix 222
trimaculata, Chrysura, *Chrysis* 140
trimaculata, Prospis 261
trimaculatus, Dolerus 77
trimaculatus, Nematus 61, 64
trimaculatus, Nematus luteus 68
trimaculatus, Nysson, *Crabro*
trimaculatus 220
trimarginata, Allantus stecki 90
trimarginatus, Odynerus 179
trimmerana, Andrena, *Melitta* 267
trinacria, Coelioxys 304
trinacia, Heriades 297
trinacriae, Odynerus 185
tringa, Andrena 271
trinotata, Hylotoma 24
trinotatus, Crabro 231
tripartitum, Anteon 119
triplicatus, Dolerus, *Tenthredo* 77
tripolitanus, Halictus 280
Tripoxilon 244
tripunctata, Tiphia 149
tripunctata, Vespa 195
tripunctatus, Eumenes, *Sphex* 182
tripunctatus, Pompilus 161
tripunetata, Nomada 311
trisignatus, Nematus 57
trisinuosa, Mutilla 153
trispine, Lasioglossum, *Halictus* 286
trispinis, Halictus 287
trispinosa, Apis 243
trispinosa, Nomada 316
trispinosus, Evagetes, *Pompilus* 171
trispinosus, Lasioglossum, *Halictus*
interruptum 283
trispinosus, Oxybelus, *Apis* 243
trispinosus, Rophites algirus 291
Trissomalus 125
Tristactoides 107
Tristactus 107
tristior, Tenthredopsis 99
tristis, Amauronematus 65
tristis, Andrena apicata var. 263
tristis, Athalia lugens 44
tristis, Bombus 331
tristis, Cladius 54
tristis, Cubitalia, *Eucera* 319
tristis, Diodontus, *Pemphredon* 247
tristis, Elampus bidens 127
tristis, Macrophya 87
tristis, Mellinus 246
tristis, Nomioides minutissima 292
tristis, Odynerus 178
tristis, Odynerus, *Hoplomerus* 187
tristis, Pemphredon 247
tristis, Pontania piliserra 63
tristis, Prosopis 261
tristis, Prosopis annulata var. 259
tristis, Smicromyrme 156
tristis, Sphecodes 216
tristis, Tenthredo 29, 77, 81, 99
tristis, Triepeolus, *Epeolus* 317
trisulcatus, Aphelopus 118
tritici, Pachynematus 58
Tritikreion 20
Tritodiscoelius 181
trivialis, Arachnospila, *Pompilus* 169
trivittata, Eucera 321
trivittatus, Allantus 96
trivittatus, Nematus 62
trochanterata, Euura, *Pteronidea* 66
trochanterica, Macrophya 86
trochanterica, Pristiphora,
Lygaeonematus 74
trochantericum, Anthidium 301
Trogaspidia 157
Trogaspidiini 157
troglodyta, Trachelus, *Sirex* 111
troglodytes, Spilomena, *Stigmus* 249
troglodytes, Stigmus 249
Trogloembolemus 122

trogodermatis, Laelius 124
tromsouensis, Lygaeonematus 71
Trophocleptria 317
Tropidepipona subg. 192
Tropidodynerus 191
Tropidodynerus subg. 192
Tropidotilla 157
trossolus, Hedychridium,
 Euchridium 129
truchmenus, Batozonellus, *Pompilus* 166
Truncandrena subg. 275
truncata, Andrena 265
truncata, Coelioxys 305
truncata, Formica 199
truncaticolle, Lasioglossum,
 Halictus 286
truncatiformis, Andrena 265
truncatilabris, Andrena 275
truncatiserra, Pristiphora 74
truncatooides, Allantus 38
truncatula, Cerceris 254
truncatula, Osmia 298
truncatus, Allantus, *Tenthredo* 38
truncatus, Biastes, *Nomada* 318
truncatus, Camponotus auct. 199
truncatus, Colobopsis, *Formica* 199
truncatus, Halictus 283
truncatus, Nematus 70
truncatus, Pachynematus 58
truncatus, Philoctetes, *Chrysis* 132
truncatus, Thyreus, *Crocisa* 327
truncicola, Formica 200
truncorum, Apis 296
truncorum, Formica 200
truncorum, Heriades, *Apis* 296
trunculi, Tenthredo 39
Trypargilum 244
Trypetes 296
Trypoxilon 244
Trypoxylini 244
Trypoxylon 244
Trypoxylon subg. 244
Trypoxylyum 244
tschekanovskii, Myrmica 206
tscheki, Camponiscus 53
tschelcarica, Anthophora monacha 325
tscherkessicum, Stenamma
 westwoodi var. 207
tschertkoviana, Dasypoda 293
tschinggischananensis, Tenthredo 94
tsherepanovi, Rhogogaster 88
tsherskii, Abia 29
tsherskii, Pamphilus 103
tshettui, Cephus 110
tshitscherini, Bombus 331
tshiliensis, Myrmica sancta 206
tsingtauensis, Halictus 278
tsingtauensis, Hylaeus, *Prosopis* 260
tsingtauensis, Megachile 307
tsingtouensis, Halictus 278
tsukubana, Andrena 264
tsunekii, Ceropales, *Bifidoceropales*
 pygmaea 160

tsunekii, Colletes 257
tsunekii, Hedychridium 129
tsunekii, Megalodontes 107
tsunekii, Miscophus 241
tsunekii, Myrmosa 153
tsunekii, Psen 251
tsunekii, Sphecodes 288
Tsunekiola 236
tsushima, Halictus 278
tsushima, Tetramorium 209
tsushima, Anthophora 322
tuberculata, Apis 320
tuberculata, Cerceris, *Crabro* 255
tuberculata, Eucera auct. 320
tuberculata, Euura, *Nematus* 66
tuberculata, Hoplitis, *Osmia* 297
tuberculata, Melecta 326
tuberculata, Prosopis 260
tuberculatus, Colletes 259
tuberculatus, Halictus 279
tuberculatus, Monophadnoides 50
tuberculatus, Pterocheilus
 phaleratus 189
tuberosa, Formica 208
tuberum, Temnothorax, *Formica* 208
tubifex, Pelopaeus 214
Tubponentia 55
tulunensis, Nematus 69
Tumagenia 165
tumida, Tenthredo, *Allantus tumida* 98
tumidosus, Evagetes, *Pompilus* 171
tumidus, Harpactus, *Pompilus*
 tumidus 224
tumulorum, Halictus, *Apis*
 tumulorum 279
tunensis, Coelioxys atra var. 304
tunensis, Crocisa 326
tungusicum, Lasioglossum 281
tunicata, Euura, *Nematus* 66
turana, Dasylabris 154
turana, Anthophora 324
turana, Coelioxys rufescens var. 304
turana, Didineis 219
turanicum, Chalybion, *Sceliphron* 214
turanicus, Messor structor 204
turaniformis, Eumenes
 pedunculatus var. 181
turanum, Euchrum 129
turanus, Eumenes pedunculatus 182
turbida, Andrena 274
turca, Bembix 222
turca, Odynerus 184
turcarum, Tenthredo 68
turcestanica, Andrena 270
turcestanica, Eucera 321
turcestanica, Mutilla 155
turcestanica, Osmia 296
turcestanica, Prosopis 260
turcestanicum, Anthidium
 diadema var. 300
turcestanicus, Hoplomerus tinniens 184
turcica, Vespa 194
turcicum, Hedychridium 129
turicum, Hedychridium valesiense 129
turcicus, Eumenes pomiformis 182
turcicus, Lasius 202
turmenicus, Elampus, *Omalus* 128
turmenicus, Harpactus 224
turcomanica, Nomia aureocincta 291
turcomanicum, Tetramorium caespitum
 st. 209
turcomenicus, Bothriomyrmex 197
turcomenicus, Bothriomyrmex
 syrius 197
turgaiensis, Euura, *Nematus* 66
turgajensis, Physetopoda,
 Paramyrme 155
turgensis, Megachile analis 307
turdidus, Nematus 57
turonum, Passaloecus 248
turkestanica, Cerceris 255
turkestanica, Holopyga 131
turkestanica, Holopyga
 punctatissima var. 131
turkestanicus, Camponotus 199
turkestanicus, Camponotus sylvaticus
 st. 199
turkestanicus, Sphecodes gibbus 288
turkestanus, Camponotus 199
turkestanus, Camponotus
 sylvaticus var. 199
turkomannus, Halictus 277
tutihensis, Halictus 286
tydei, Podalonia, *Ammophila* 214
Typhoctinae 159
typica, Alysson pertheesi 219
typica, Claremontia 46
typica, Zaphymatocera 53
tyrannica, Vespa 193
tyrolensis, Panurginus 276
tyrolensis, Panurgus montanus 276
tyrrhena, Andrena 273
Tytthaliptus subg. 279
tyuzendzianus, Crossocerus 228
Tzustigmus 250
uberior, Allantus viennensis 38
uberior, Tenthredella temula 93
uchidae, Tenthredo 93
uchidai, Crabro 29, 230
uchidai, Megacampsomeris 147
ucrainicum, Stenamma 207
ucrainicus, Leptothorax
 unifasciatus 209
udegeicus, Agenioideus 168
udmurtica, Melitta 294
udos, Euura, *Nematus* 66
Ukayakos 125
ulanbatorensis, Tenthredopsis
 coqueberti 99
Ulandrena subg. 275
ulangomensis, Megachile analis 307
ulbrichti, Holcocneme 66
ulbrichti, Pristiphora conjugata 71
ulianini, Cardiocondyla 203
ulianini, Cardiocondyla elegans var. 203

- uliginosae*, Amauronematus 64
ulmariae, Macrocephus 111
ulmi, Cladius, *Tenthredo* 55
ulmi, Fenusa 48
ulmi, Janus foersteri 110
ulmi, *Tenthredo* 81
ulrikae, Colletes 259
ulterior, Halictus calceatus 282
umbellatarum, *Tenthredo* 91
umbilicalis, Odynerus 178
umbra, Sirex 113
umbraculatus, Lithurgus 295
umbrata, Euura, *Nematus* 66
umbratica, Eriocampa, *Tenthredo* 42
umbratus, Lasiurus, *Formica* 201
umbricola, Aproceros 25
umbrina, Coelioxys 304
umbrinus, Nematus 61
Umbripennis subg. 162
umbripennis, Nematus 57
umbrosa, Conospidia 96
umbrosa, *Tenthredo* 75
umbrosus, Schistonyx, *Pompilus* 172
uncata, Apis 300
uncinata, Osmia 299
uncinata, Cladius 54
uncta, Claremontia, *Tenthredo* 46
undata, Bembix 222
undulata, Poecilosoma 41
undulatafasciatum, Anthidium manicatum var. 300
unga, Pontania 60
unguicularis, Pompilus 168
unguiculatum, Delta, *Vespa* 180
unguinosus, Halictus 284
ungularis, Odynerus 183
ungyo, Crossocerus 230
unicincta, *Tenthredo* 90, 92
unicolor, Aporus 166
unicolor, Cimbex femorata 29
unicolor, Dineura 67
unicolor, Formica pratensis 199
unicolor, Mimumesa, *Psen* 251
unicolor, Pasites 318
unicolor, Pelopoeus 251
unicolor, Priocnemis, *Salius* 163
unicolor, Sphecius 249
unicolor, Spinolia, *Chrysis* 141
unicolor, Stelis minima f. 303
unicolor, Tachysphex, *Larra* 240
unicolor, Tiphiidae 150
unicolor, Panurgus 276
unicoloricornis, Abia fulgens 28
unicus, Bombus 328
unidentata, Nomia auct. 292
unidentatum, *Tenthredo* 31
unifasciata, Andrena 273
unifasciata, Nomioidea variegata 292
unifasciata, *Tenthredo* 92, 94
unifasciatus, Leptothorax, *Formica* 208
uniformis, Pseudochrysis, *Chrysis* 141
unigallae, Euura, *Pontania* 66
uniglumis, Crabro 242
uniglumis, Oxybelus, *Vespa* 243
uniglumis, Vespa 242
unimaculatus, Odynerus 191
uniocellatus, Anoplus 237
unipetiolaris, Mutila 154
Unitaxonus 39
universitatis, Andrena 274
Upaterka 149
upinense, Lasioglossum, *Halictus* 281
uralensis, Allantus 93
uralensis, Formica 201
urbaini, Aphelopus 118
urbis, Selandria 76
urgaensis, *Tenthredo* 91
Urocerites 112
Urocerus 113
Urocerus 115
ursina, Apis 276
ushinskii, Empria 42
ushinskii, Trichiosoma 31
ussurensis, Bombus 329
ussuriana, Macropis, *Ctenoplecta* 294
ussuricus, Goniozus, *Parasierola* 125
ussuriense, Pison 244
ussuriense, Stenamma 207
ussuriense, Trypoxylon 245
ussuriensis, Apis indica sinensis 332
ussuriensis, Arachnospila,
Psammochares 170
ussuriensis, Belomicrus 242
ussuriensis, Busarbidea, *Canoniades* 76
ussuriensis, Caliadurgus, *Calicurgus* 162
ussuriensis, Crabro 226
ussuriensis, Dipteronomorpha 93
ussuriensis, Elampus, *Notozus* 128
ussuriensis, Heptamelus 36
ussuriensis, Machaerothrix 165
ussuriensis, Obtusia 51
ussuriensis, Pamphilus 105
ussuriensis, Parancistrocerus 190
ussuriensis, Phymatocerola 52
ussuriensis, Praia 30
ussuriensis, Priocnemis 164
ussuriensis, Psen 251
ussuriensis, Tenthredella 93
ussuriensis, Tenthredo, *Allantus*
ussuriensis 93
ussuriensis, Xyela 21
Ussurinus 100
Ustilatiformis 22
ustulata, Andrena 267
ustulata, Arge, *Tenthredo* 25
usurata, Arachnospila 170
Utapitoca 159
utensis, Thyreopus 228
utinamius, Sphecodes 290
uvarovi, Chrysis 134
uvarovi, Hedydriidium 130
vacciniellus, Nematus crassipes 58
vacillans, Macrophyia 87
vafer, Pamphilus, *Tenthredo* 105
vaga, Andrena 269
vaga, Euura, *Tenthredo* 66
vaga, Nomada 316
vaga, Sphecius 168
vagabunda, Ammophila 213
vagabundica, *Tenthredo* atra 93
vagabundus, Cephus 108
vagabundus, Crossocerus, *Crabro* 227
vagans, Cerceris 255
vagans, Chrysis 141
vagans, Crabro 232
vagans, Heterarthrus, *Hylotoma* 49
vagans, Spilomena 249
vagatus, Crabro 232
vaginosa, Euura, *Pachynematus* 66
vagus, Camponotus, *Formica* 198
vagus, Nematus 66
vagus, Sphecius 168, 246
Valco 45
valeriana, Andrena 267
valesiacus, Nematus 57
valesiana, Chrysis 138
valesiana, Chrysis sybarita var. 138
valesianus, Sphecodes 287
valesiense, Hedydriidium 129
valesina, Chalicodoma 305
valga, Xylocopa 310
validicornis, Nematus 61
validus, Crabro 232
valkanovi, Amasis 32
vallator, Nematus 75
vallei, Prosopis 259
vallisnieri, Nematus 63
van der Lindenii, Crabro 228
vandeli, Myrmica 206
vandeli, Myrmica scabrinodis var. 206
vanderlindeni, Astata 218
vanduzeei, Emphytina 39
vanduzeei, Pteronidea 62
vanduzeei, Selandridaea 76
vankovitzii, Osmia 299
vanus, Bombus ikonnikovi 327
vapida, Aphaenogaster 203
vara, Anthophora 323
Varentzowi, Crabro 233
varentzowi, Ectemnius, *Crabro* 233
varia, Apis 300, 303
varia, Lyda 105
varia, Nomada 313
varia, Pontania 64
varia, Sapyga 145
varia, *Tenthredo* 99
variabilis, Andrena 267
variabilis, Bombus 331
variabilis, Cephus 108
variabilis, Cerceris 254
variabilis, Cimbex 29
variabilis, Cryptocheilus, *Sphex* 161
variabilis, Eoferreola, *Pompilus* 173
variabilis, Episemus 125
variabilis, Euura, *Amauronematus* 66
variabilis, Megachile muraria var. 305
variabilis, Nysson 220
variabilis, Smicromyrme 155

variana, *Tenthredopsis litterata* 99
varians, *Andrena*, *Melitta* 264
varians, *Anthophora acervorum* var. 323
varians, *Apis* 305
varians, *Cimbex* 29
varians, *Euura*, *Amauronematus* 66
varians, *Prosopis* 260
varianus, *Amauronematus* 62
variator; *Dolerus* 78
variator, *Euura*, *Nematus* 66
varicornis, *Chrysura*, *Chrysis* 140
varicornis, *Tenthredo* 40
variegata, *Apis* 300
variegata, *Cephaleia*, *Cephaleia* 102
variegata, *Ceropales*, *Evania* 160
variegata, *Deuteragenia*, *Sphex* 164
variegata, *Gilpinia*, *Lophyrus* 34
variegata, *Lyda* 105
variegata, *Pachyprotasis*, *Tenthredo* 88
variegata, *Sapyga* 145
variegata, *Sphex* 164
variegata, *Tiphia* 243
variegatus, *Ammobates* 319
variegatus, *Cephus* 109
variegatus, *Ceylalictus*, *Apis* 292
variegatus, *Dolerus* 78
variegatus, *Epeolus*, *Apis* 317
variegatus, *Hylaeus*, *Mellinus* 262
variegatus, *Odynerus* 188
variegatus, *Oxybelus* 243
variegatus, *Palarus*, *Tiphia* 243
variegatus 243
variegatus, *Philanthus* 255
variegatus, *Philanthus* 256
variegatus, *Sphecodes* 287
varifrons, *Prosopis* 259
variicolor, *Tenthredo* 98
variipennis, *Cryptochileius* 161
variipes, *Eriocampoides* 46
variipes, *Lygaeonematus* 74
variolatus, *Nysson* 220
varipes, *Caliroa*, *Tenthredo* 46
varipes, *Crossocerus* 229
varipes, *Dienoplus* 223
varipes, *Dolerus* 38
varipes, *Prosopis* 261
varipes, *Tenthredo* 88
varipes, *Trypoxylon* 245
varipictus, *Nematus* 56
varius, *Crossocerus* 230
varius, *Nematus* 55
varius, *Palarus flavipes* 243
varius, *Palarus* 243
variegatus 243
varius, *Pamphilus*, *Lyda* 105
Varna 55, 67
varus, *Crabro* 230
varus, *Crossocerus* 230
varus, *Nematus* 69
vastatrix, *Euura*, *Pteronidea* 66
vastidens, *Euura*, *Amauronematus* 66
vaticinia, *Smicromyrme oratoria* 155
vechti, *Deuteragenia*, *Dipogon* 164

vectis, *Coelioxys* 305
vedicus, *Harpactus* 223
velatus, *Nematus* 74
veles, *Tomostethus* 53
velocissima, *Amegilla*, *Anthophora* 322
velox, *Bombus*, *Agrobombus* 332
velox, *Odynerus* 187
velox, *Tenthredo velox* 95
velutina, *Andrena* 269
velutina, *Eucera*, *Tetralonia* 321
velutinus, *Cyphononyx* 171
velutinus, *Euodynerus* 183
veneficus, *Amauronematus* 65
veneticus, *Halictus* 278
ventraliformis, *Colletes* 258
ventralis, *Andrena* 267
ventralis, *Colletes* 258
ventralis, *Dineura* 55
ventralis, *Eurhadinoceraea*,
Tenthredo 47
ventralis, *Hemibeleses* 42
ventralis, *Hylotoma* 51
ventralis, *Nomada* 313
ventralis, *Triepeolus*, *Epeolus* 317
ventricosa, *Andrena* 267
ventricosa, *Tenthredo* 64
ventricosus, *Nematus* 64
ventriosa, *Hylotoma* 24
venusta, *Cimbex* 29
venusta, *Eumenes* 182
venusta, *Euura*, *Cryptocampus* 66
venusta, *Siobla*, *Ercarsioneura*
venusta 89
venustus, *Lindenius* 235
venustus, *Oxybelus* 243
venustus, *Pachynematus* 57
venustus, *Pamphilus*, *Lyda* 106
venustus, *Philanthus*, *Crabro* 256
verae, *Andrena* 267
verae, *Chrysis* 137
Veratra 52
Veratra subg. 52
vergesi, *Nannodynerus* 190
verhaereni, *Ancistrocerus* 179
verhoeffii, *Alysson* 219
verhoeffii, *Archianthidium*
pubescens 303
verhoeffii, *Cerceris* 255
verhoeffii, *Crabro* 232
verhoeffii, *Crossocerus* 228
verhoeffii, *Gorytes* 222
verhoeffii, *Stigmus* 250
Verna 55
verna, *Chrysis* 139
verna, *Tenthredo* 68
vernalis, *Anthophora* 325
vernalis, *Dolerus* 80
vernalis, *Emphytus* 41
vernalis, *Nematus* 56
vernalis, *Pteronidea* 56
vernalis, *Tenthredo* 42
vernalis, *Tiphia* 150
vernus, *Pachynematus* 57
verrucosae, *Pteronidea* 66
versicolor, *Anoplonyx* 53
versicolor, *Ceropales* 161
versicolor, *Cryptochileius*, *Sphex* 162
versicolor, *Megachile* 307
versicolor, *Nomioides minutissima* 292
versicolor, *Osmia* 299
versicolor, *Selandria* 51
versicolor, *Spintharina*, *Chrysis* 141
versuta, *Tenthredo* 96
verticalis, *Gilpinia* 33
verticalis, *Sciaapteryx* 89
verticalis, *Sphecodes* 289
verticata, *Tenthredo* 67
verticosa, *Arge* 25
verzhutskii, *Strongylogaster* 82
vescus, *Amauronematus* 65
vesicator, *Euura*, *Nematus* 66
Vespa 193
vespa, *Tenthredo* 97
vespertilio, *Sirex* 117
Vespiae 175
vespiformis, *Crabro* 231
vespiformis, *Tenthredo* 97
Vespinae 193
Vespoidea 175
Vespomorpha 118
Vespula 195
vestalis, *Bombus*, *Apis* 330
vestalis, *Psithyrus* auct. 329
vestigialis, *Dolerus*, *Tenthredo*
vestigialis 77
vestita, *Megachile* 307
vestita, *Tenthredo* 92
Vestitohalictus subg. 279
vestitus, *Crabro* 233
vestitus, *Halictus* auct. 279
veteranus, *Bombus*, *Apis* 332
vetuberosa, *Ammophila* 213
vetula, *Andrena* 273
vetula, *Scolia* 147
vetus, *Vespula rufosignata* 196
vexator, *Vespa* 193
vexillatus, *Crabro* 234
v-flaxum, *Nematus* 55
viaticus, *Anoplius*, *Sphex* 172
viator, *Mesitius* 124
viator, *Phylloecus*, *Cephus* 111
vicana, *Chrysis* 139
vicaria, *Pachyprotasis* 88
vicina, *Euura*, *Nematus* 66
vicina, *Megachile* 306
vicina, *Tetraloniella*, *Tetralonia* 322
vicinus, *Crossocerus* 227
vicinus, *Dolerus* 38
vicinus, *Euspongus* 223
vicinus, *Hylaeus* 261
vicinus, *Nematus* 66
victor, *Astata* 218
victor, *Oxybelus* 242
victoria, *Platycampus* 54
victorius, *Euchroeus* 140
victorovi, *Dryinus* 120

- vidua*, Bembix 222
vidua, Nomada 316
vidua, Osmia 298
viduata, Euura, *Tenthredo* 66
viduata, Mutilla 155
viduata, Nemka, *Mutilla viduata* 155
viduatinus, Euura, *Amauronematus* 66
viduatooides, Euura, *Amauronematus* 66
viduatus, Prionyx viduatus 215
viduus, Odynerus 179
viennensis, Allantus, *Tenthredo* 38
Viereckella 319
vigora, Holopyga 131
viktorovi, Smicromyrme 156
villiersi, Halictus 287
villipes, Dasypoda 293
villipes, Nomada stigma 316
villosa, Anthophora 324
villosa, Crocisa affinis var. 326
villosa, Dasypoda 293
villosa, Euura, *Nematus* 66
villosa, Hoplitis, *Megachile* 297
villosa, Megachile 306
villosa, Pseudotiphia, *Tiphia* 150
villosa, Siobla 89
villosa, Tenthredo 90
vilosula, Anthophora 323
vilosulopsis, Halictus 287
vilosulum, Lasioglossum, *Melitta* 287
villosum, Lasioglossum albipes 282
villosum, Trichiosoma, *Cimbex* 31
viminalis, Euura 60
viminalis, Euura, *Cynips* 66
viminalis, Nematus 60
viminalis, Tenthredo 54, 97
vincit, Ammobates 318
vindobonensis, Mimesa 250
vinokurovi, Chrysis 139
violacea, Apis 309
violacea, Hylotoma 23
violacea, Tenthredo 23, 39
violacea, Tenthredo, *Allantus*
 bifasciata 90
violacea, Xylocopa, *Apis* 310
violaceipennis, Allantus 90
violaceipennis, Ammophila 213
violaceipennis, Andrena 269
violaceipennis, Melanapis 270
violaceipennis, Odynerus dantici 183
violaceus, Dolerus 79
violaceus, Pseudomalus, *Sphex* 133
violascens, Andrena 267
violascens, Cimbex 29
violascens, Nomioides minutissima 292
violascens, Tenthredo, *Allantus* 90
virago, Paururus juvencus 112
virago, Tenthredo luteocincta 96
viratus, Eumenes latipes var. 185
virens, Gilpinia, Lophyrus 34
virens, Hedydium virens 130
virescens, Ellampus auratus var. 133
virescens, Halictus 279
virescens, Nematus 56
virescens, Pamphilus 106
virescens, Pseudomalus auratus 142
virescens, Selandria 82
virgatus, Bombus lapponicus 330
virginalis, Pachynematus 59
virgineus, Tenthredopsis 99
virginianus, Priophorus 54
viridana, Osmia 299
viridana, Pristiphora 73
viridans, Tenthredella 92
viridator, Banchus 109
viridatrix, Tenthredo viridatrix 92
virideglaucum, Lasioglossum 287
viridellum, Lasioglossum, *Halictus* 287
virideocincta, Chrysis succinata ab. 134
viridescens, Nematus 63
viridiaeneus, Halictus 282
virididorsata, Dineura, *Tenthredo* 55
viridifrons, Rhogogaster genistae 88
viridilineolatum, Hedydium 130
viridipes, Pamphilus 104
viridis, Euura, *Nematus* 67
viridis, Notozus 127
viridis, Pachyprotasis 87
viridis, Rhogogaster auct. 88
viridis, Rhogogaster, *Tenthredo* 89
viridis, Strongylogaster 82
viridis, Tenthredo 88
viridis, Tenthredopsis 100
viridissima, Euura, *Nematus* 67
viriditibialis, Onycholyda,
 Pamphilus 103
viriditibialis, Pamphilus 103
viridiventris, Omalus 127
viridiventris, Pseudomalus, *Ellampus*
 auratus 132
viridula, Chrysis 139
viridulus, Pamphilus 106
virilis, Dineura stilata 55
virilis, Euura, *Pontania* 67
virilis, Nematinus bilineatus 68
vitellina, Trichiosoma, *Tenthredo* 31
Vitelliniformis 30
vitis, Tenthredo 38
vitosaensis, Allantus arcuatus 96
vitoshensis, Tenthredo solitaria 95
vitrea, Andrena 272
vitreipennis, Andrena 272
vitreipennis, Nematus 72
vittata, Acantholyda
 erythrocephala 101
vittata, Euura, *Nematus* 67
vittata, Tenthredopsis raddatzi 99
vittatum, Aprosthemia, *Schizocera* 26
vittatus, Cephus 111
vittatus, Nematus 67
viticollis, Cerceris 255
vittifrons, Carinoscolia, *Scolia* 147
vittizi, Tenthredo vespa 97
vittulata, Eucera 321
vivida, Tenthredo 92
vividula, Formica 202
vividula, Nylanderia, *Formica* 202
vocifera, Andrena geliae 274
vogti, Colletes 258
voigti, Fenella 47
volatilis, Pamphilus, *Lyda* 106
volatilis, Sphecodes 289
volgadon, Stigmatorus 173
volgensis, Andrena 263
volgensis, Aprosthemia 26
volgensis, Apterogyna 159
volgensis, Leptothorax nassonovi 209
volgensis, Temnothorax,
 Leptothorax 209
vollenhoveni, Nematus 66
vollenhoveni, Selandria 76
vopiscus, Eutomostethus, *Tomostethus*
 ephippium 47
vulgaris, Dufourea 290
vulgaris, Halictus 284
vulgaris, Hylotoma 23
vulgaris, Priocnemis, *Pompilus* 164
vulgaris, Vespa, *Vespa* 196
vulgata, Vespa crabro 194
vulgatus, Notozus productus var. 128
vulneratus, Dolerus 80
vulnericeps, Mutilla 157
vulpecolor, Megachile melanopyga 307
vulpecula, Andrena 267
vulpecula, Osmia 298
vulpina, Andrena 282
vulpina, Apis 323
vulpinus, Halictus 285
vulsum, Lasioglossum, *Halictus* 287
vutshetitshi, Myrmilla 154
Waagenia 211
wacki, Colletes 259
wagneri, Tenthredo maculata 92
wahlbergi, Nematus wahlbergi 69
wahrmani, Apterogyna 159
waldheimii, Claremontia, *Tenthredo* 46
waldstaetterense, Empria 41
walishanovi, Andrena schwarzi ssp. 267
Walkeri, Crabro 228
walkeri, Crossocerus, *Crabro* 228
walkeri, Vespa 194
walteri, Chalybion, *Pelopoeus* 214
Walteri, Crabro 233
walteri, Ectemnius, *Crabro* 233
waltlii, Gastrosericus 236
wanda, Dolerus 80
wankowiczi, Cilissa 294
wankowiczi, Dasypoda 293
warnckeii, Colletes 259
washingtonia, Hemichroa 67
watanabei, Arge captiva 22
watasei, Andrena 270
watsoni, Katamenes 184
watsoni, Katamenes 185
weberi, Bembix 221
weiffenbachi, Pteronidea 63
weiffenbachiella, Euura 67
weiffenbachii, Euura 67
werestschagini, Crabro 226

wernerii, Leptothorax 208
werzhutskii, Pristiphora 75
wesmaeli, Ammoplanus 246
wesmaeli, Arachnospila, *Pompilus* 169
wesmaeli, Crossocerus, *Crabro*
 wesmaeli 230
wesmaeli, Ellampus 132
wesmaeli, Nysson 220
wesmaeli, Pemphredon, *Cemonus* 249
wesmaeli, Pristiphora, *Nematus* 75
Wesmaelinus 166
Wesmaelinus 166
westerlundi, Arachnospila, *Pompilus* 169
westerlundi, Chrysis 139
westermannii, Nematus 64
westwoodii, Fenella 48
westwoodi, Stenamma 207
westwoodii, Stenamma auct. 207
whitei, Nematus 73
Wickhamii, Dolichocrabro 227
wilkella, Andrena, *Melitta* 274
willigiae, Nematinus 68
willoughbyi, Dolerus 78
willughbiella, Megachile, *Apis* 308
wilmattae, Hylaeus 261
winkleyi, Andrena 274
winnipeg, Nematus 71
winnipegensis, Nematus 71
winteri, Pteronidea 57
wittenbourgi, Halictus 281
wittenburgi, Hortobombus
 consobrinus 328
wolfi, Arachnospila 169
wolfi, Dicyrtomellus 170
wolfi, Gonaporus 173
wolfi, Krombeinella, *Myrmosa* 152
wolhynica, Osmia 299
Wollmania 130
wttewaalli, Nematus 63
wuczeticzi, Microdynerus 185
wuestneii, Brachythops, *Selandria* 76
wuestneii, Didineis 219
wuestneii, Fenusia 48
wuestneii, Fenusella, *Fenusia* 48
wuestneii, Heterarthrus, *Phyllotoma* 49
wuestneii, Nematus 72
wuestneii, Tenthredopsis 100
wui, Camponotus japonica 198
wulungshanensis, Andrena 268
wuorentausi, Ametastegia 39
wurflenii, Bombus 327
wymperi, Camponotus
 herculeanus var. 198

Xambeui, Sylaon 241
Xantepyrus 123
xanthispis, Pontania viminalis 66
xanthispis, Tenthredo temula 93
Xanthidium 310
xanthobaptus, Nematus 65
xanthocarpus, Nematus 59
xanthocera, Arge 24

xanthocera, Strongylogaster,
 Tenthredo 82
xanthocerus, Megalodontes 107
xanthocerus, Nematus 55
xanthocnemis, Hylaeus 261
xanthogaster, Nematus 57
xanthogastria, Apis 296
xanthomelaena, Tenthredo 93
xanthomelana, Osmia 299
xanthomelas, Stenodynerus,
 Odynerus 190
xanthometopa, Apis 259
xanthomus, Nematus 73
xanthophorus, Nematus 64
xanthoptera, Vespa 194
xanthopterus, Nematus 56
xanthopus, Caenocephus 108
xanthopus, Dosytheus 78
xanthopus, Lasioglossum, *Melitta* 281
xanthopus, Melitta 280
xanthopus, Nematus 55
xanthopyga, Andrena 271
xanthopyga, Tenthredo 22
xanthorius, Allantus 90
Xanthosarus subg. 307
xanthoscelis, Andrena 270
xanthosensis, Halictus 284
Xanthosirex 113
Xanthosphecius 245
xanthosticta, Apis 312
xanthostoma, Phylloecus, *Cephus* 111
xanthostomus, Pteronus 61
xanthostylos, Nematus 58
xanthotarsus, Tenthredo 98
xanthothorax, Andrena 259
xanthotrichela, Pristiphora 73
xanthura, Melitta 274
xanthurum, Anthidium oblongatum 301
xanthurum, Aprosthemia 26
xema, Andrena coitana ssp. 271
Xenanteon 118
Xenaporus 174
Xenaporus subg. 174
Xenocrabro 231
Xenoglossodes 321
Xeris 114
Xeromalissinae 257
Xestobethylus 123
Xestocrabro 231
Xilocopa 309
xinjiangensis, Dolichovespula 195
Xiphidion 115
Xiphium 115
Xiphydria 115
Xiphydriidae 115
Xiphydriinae 115
Xiphydriola 116
Xoanon 113
Xyela 20
Xyelatana 20
Xyelidae 20
Xylinae 20

Xyeloidea 20
Xyelomorpha 20
Xylocelia 247
Xylocopa 309
Xylocopa subg. 310
Xylocopinae 309
Xylocopini 309
Xylocrabro 231
Xylocematum 113
Xylocucus 247
xylopedis, Lasioglossum zonulum 281
xylostei, Hoplocampoides, *Tenthredo* 49
xylostei, Selandria 49
xylota, Tenthredo, *Allantus solitaria* 95
Xyloterus 113
xylurgus, Crabro 231
Cyphydria 115

yakuticum, Lasioglossum, *Evlaeus* 287
yamamotoi, Anoplus 168
yamanei, Ancistrocerus 179
yamato, Crossocerus capitosus 228
yamatonis, Trypoxylon frigidum 244
Yambal 227
yanoi, Athalia 45
yanoi, Smicromyrme lewisi 156
Yanoniush 231
yarrowi, Metrionotus 124
yarrowi, Nomada 317
yasha, Nomada 313
yasumatsui, Arachnospila 169
yasumatsui, Auplopus 165
yasumatsui, Colletes 258
yasumatsui, Crossocerus, *Crabro* 230
yasumatsui, Priocnemis 163
yasushii, Urocerus, *Sirex* 113
yayeyamae, Polistes rothneyi 192
yequasicus, Pterochilus phaleratus 189
yessensis, Formica 200
yessensis, Formica truncicola var. 200
yezo, Crossocerus capitosus 228
yezo, Trypoxylon figulus 244
yezoanus, Symmorphus mutinensis 191
yezoensis, Acantholyda aglaia 101
yezoensis, Bombus 328
yezoensis, Evagetes 171
yezoensis, Onycholyda 104
yezoensis, Priocnemis 163
yezoensis, Tenthredo 98
yichunensis, Vespula 196
yokohamensis, Nematus, *Holcoecneme* 69
yosemite, Ectemnius 231
ypsilone, Chlorodynerus, *Odynerus* 180
ypsilone, Dasylabris rubrosignata 153
Yuchiha 227
yukonensis, Dolerus 78
yulongensis, Vespula 196
yunlongensis, Vespula 196

zablockii, Sphecodes 287
zacharovi, Empria 41
zaddachi, Dolerus 81
zaddachi, Dolerus aeneus 79

zaddachi, Pachynematus 57
zaddachi, Schizocera 26
zahaikewitshi, Cephus, *Pseudocephus* 110
zaidamensis, Crabro 228
zaitzevi, Chrysis 136
zaitzevi, Euura, *Amauronematus* 67
zakakica, Megachile melanopyga 307
Zalagium 84
zalinda, Prosopigastra 237
Zamacrophya 89
Zaphymatocera 53
Zarabayca 159
Zaraea 28
Zarudnium 128
Zarudnium 128
zarudnyi, Arge 24
zarudnyi, Ellampus 131
zarudnyi, Tachysphex 238
zavadili, Mesitius 124
zavadili, Smicromyrme rufipes f. 156
zavadili, Spilomena 249
zavattarii, Vespa orientalis var. 194
zaysanicus, Gorytes 222
Zebramegilla subg. 322
zella, Pristiphora 73
zelleri, Hedychridium, *Hedychrum* 130
zelleri, Pompilus 173

Zelorhopalum 235
zenaida, Siobla 89
Zendalia subg. 185
zeraldae, Megachile willughbiella 308
Zeteumenooides subg. 181
zetterstedti, Amauronematus 65
zetterstedti, Chrysis 139
zetterstedti, Nematus 62
zetterstedti, Vespa norwegica var. 195
zeyanense, LasioGLOSSUM 281
zhelochovtsevi, Dolerus 80
zhelochovtsevi, Pamphilus
zhelochovtsevi 106
zhelochovtsevi, Platycampus 70
zhelochovtsevi, Taxonus 43
zhelochovtzevi, Abia, *Zaraea* 29
zhelochovtzevi, Arge 25
ziegleri, Ceratocolus 230
zimini, Didineis 219
zimini, Solierella 241
Zimmermannia 126
zinojevi, Fenus, *Kaliofenus* 48
zinovjevi, Onycholyda 104
zinovjevi, Pamphilus 106
zojae, Myrmica 207
zonalis, Discoelius, *Vespa* 181
zonalis, Hylotoma 24

zonalis, Melitta 267
zonalis, Vespa 181
Zonandrena subg. 275
zonaria, Andrena rutila ab. 270
zonata, Apis 296
zonata, Nomada 311
zonata, Nomada 317
zonata, Tenthredo 38, 90, 92
zonatula, Habropoda 325
zonatus, Bombus 332
zonatus, Crabro 231
zonsteini, Arachnospila 169
zonula, Tenthredo 97
Zonuledo 90
Zonuledo subg. 97
zonulum, LasioGLOSSUM, *Halictus* 281
zukaensis, Pachyprotasis 88
zwakhalsi, Ceratina 310
zwickoviensis, Allantus consobrinus 89

Submitted September 20, 2017;
accepted October 27, 2017

**ANNOTATED CATALOGUE
OF THE HYMENOPTERA OF RUSSIA**

VOLUME I

SYMPHYTA AND APOCRITA: ACULEATA

Edited by S.A. Belokobylskij and A.S. Lelej

ТРУДЫ ЗООЛОГИЧЕСКОГО ИНСТИТУТА
РОССИЙСКОЙ АКАДЕМИИ НАУК

ПРИЛОЖЕНИЕ № 6

Компьютерная верстка *К.Г. Самарцева*

Подписано в печать 15.11.2017. Формат 84x 108/16
Бумага офсетная и меловая. Печать цифровая.
Усл. печ. л. 49.98. Тираж 300 экз.

Издательство ООО «Русская коллекция СПб»
199178, Санкт-Петербург, В.О., 13 линия, д. 30, оф. 4